



**HAL**  
open science

## Immediate targets of ETTIN suggest a key role for pectin methylesterase inhibitors in the control of *Arabidopsis gynecium* development

Amélie Andres Robin, Mathieu C Reymond, Géraldine Brunoud, Martin-Magniette Marie-Laure, Françoise Moneger, C.P. Scutt

### ► To cite this version:

Amélie Andres Robin, Mathieu C Reymond, Géraldine Brunoud, Martin-Magniette Marie-Laure, Françoise Moneger, et al.. Immediate targets of ETTIN suggest a key role for pectin methylesterase inhibitors in the control of *Arabidopsis gynecium* development. *Plant Signaling and Behavior*, 2020, 15 (7), pp.1771937. 10.1080/15592324.2020.1771937 . hal-03068443

**HAL Id: hal-03068443**

**<https://hal.science/hal-03068443v1>**

Submitted on 16 Dec 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1  
2  
3  
4 **Immediate targets of ETTIN suggest a key role for Pectin Methylesterase**  
5 **Inhibitors in the control of *Arabidopsis* gynoecium development.**  
6  
7  
8  
9

10  
11 Amélie Andres-Robin<sup>1\*</sup>, Mathieu C. Reymond<sup>1\*</sup>, Géraldine Brunoud<sup>1</sup>, Marie-Laure  
12 Martin-Magniette<sup>2,3,4</sup>, Françoise Monéger<sup>1</sup> and Charles P. Scutt<sup>1</sup>  
13  
14  
15

16  
17 Author Affiliations

18 *<sup>1</sup> Laboratoire de Reproduction et Développement des Plantes, Université de Lyon,*  
19 *ENS de Lyon, UCBL, INRAE, CNRS, 46 Allée d'Italie, 69364 Lyon Cedex 07, France.*  
20  
21

22 *<sup>2</sup> Université Paris-Saclay, CNRS, INRAE, Univ Evry, Institute of Plant Sciences Paris-*  
23 *Saclay (IPS2), 91405, Orsay, France.*  
24

25 *<sup>3</sup> Université de Paris, CNRS, INRAE, Institute of Plant Sciences Paris-Saclay (IPS2),*  
26 *91405, Orsay, France*  
27

28 *<sup>4</sup> UMR MIA-Paris, AgroParisTech, INRAE, Université Paris-Saclay, 75005, Paris,*  
29 *France*  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 \*Equal contributions.

40 Authors for correspondence: [francoise.moneger@ens-lyon.fr](mailto:francoise.moneger@ens-lyon.fr) and [charlie.scutt@ens-](mailto:charlie.scutt@ens-lyon.fr)  
41 [lyon.fr](mailto:charlie.scutt@ens-lyon.fr)  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Abstract

The control of gynoecium development in *Arabidopsis thaliana* by the auxin response factor ETTIN (ETT) correlates with a reduction in the methylesterification of cell-wall pectins and a decrease in cell-wall stiffness in the valve tissues of the ovary. Here, we determine the list of genes rapidly regulated following the *in-vivo* activation of an ETT fusion protein, and show these to be significantly enriched in genes encoding cell-wall proteins, including several pectin methylesterases (PMEs) and pectin methylesterase inhibitors (PMEIs). We also perform a genome-wide scan for potential ETT-binding sites, and incorporate the results of this procedure into a comparison of datasets, derived using four distinct methods, to identify genes regulated directly or indirectly by ETT. We conclude from our combined analyses that PMEIs are likely to be key actors that mediate the regulation of gynoecium development by ETT, while ETT may simultaneously regulate PMEs to prevent exaggerated developmental effects from the regulation of PMEIs. We also postulate the existence of one or more rapidly-acting intermediate factors in the transcriptional regulation of PMEs and PMEIs by ETT.

## Introduction

The *Arabidopsis thaliana* gynoecium, or female floral whorl, consists of a pistil of two fused carpels. This structure is divided longitudinally into stigma, style and ovary tissues and includes a short stalk termed the gynophore. The ovary consists of two valves, attached to a central replum, along which placental tissues form to generate the ovules.

Loss-of-function mutants of the auxin response transcription factor *ETTIN* (*ETT*), also known as *AUXIN RESPONSE FACTOR3* (*ARF3*), show defects in tissue patterning along the longitudinal axis of the gynoecium, including an increase in stigma and gynophore tissues and a reduction in the length of the ovary, including the valve tissues (1). These changes are accompanied by defects in abaxial/adaxial patterning, in agreement with the known role of *ETT* and its paralog *ARF4* in the definition of abaxial tissue identity both in floral organs outside the gynoecium and in leaves (2). *ETT* and *ARF4* expression is restricted to the abaxial domain of these organs by the presence in the adaxial domain of tasi-RNAs generated from *TAS3* by the activity of *miRNA390* (3).

1  
2  
3  
4  
5  
6 ARF proteins are known to bind DNA motifs termed auxin response elements  
7 (AuxREs), which are found in the promoter regions of numerous auxin-responsive  
8 genes, as reviewed by Guilfoyle and Hagen (4). To quantitatively characterise ETT's  
9 interactions with these motifs, position-specific scoring matrices (PSSMs) have been  
10 generated from Protein Binding Microarray (PBM) data (5), based on variants of the  
11 two top-scoring sequences among all possible 10-mers. The recovery of more than  
12 one top-scoring matrix is common in analyses of transcription factor binding  
13 preferences, and can be caused by the presence of interdependency among  
14 nucleotide positions in the binding site (6).  
15  
16  
17  
18  
19  
20  
21  
22  
23

24 The cellulose microfibrils of the plant cell wall are embedded in a mixture of  
25 polysaccharides termed pectin. The major component of pectin, homogalacturonan, is  
26 synthesized in a highly methyl-esterified form that contributes to cell-wall stiffness. A  
27 recent study by Andres-Robin et al. (7) showed that the regulation of gynoecium  
28 patterning by ETT correlated with changes in pectin methyltransferase (PME) activity in  
29 the cell wall. This study supports a model in which ETT contributes to normal  
30 gynoecium patterning by causing an increase in PME activity that results in a decrease  
31 in the level of methylesterification of cell-wall pectins. This decrease was found to  
32 correlate with a reduction in cell-wall stiffness and an increase in valve-length in the  
33 ovary. Strikingly, the artificial overexpression of PMEs or PME inhibitors (PMEIs)  
34 respectively rescued and phenocopied *ett* loss-of-function mutants, supporting a  
35 possible role for PMEs and/or PMEIs in the mechanism of action of ETT. The findings  
36 of Andres Robin et al. (7) were in accord with those of previous studies which found  
37 PMEs to promote cell wall loosening and developmental processes at the stem apex  
38 (8).  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 Here, we use a transcriptomics approach to identify genes that are rapidly regulated  
54 following the induction of ETT activity in transgenic plants, and also perform *in-silico*  
55 scans for potential ETT-binding sites in a genome-wide dataset of *A. thaliana* promoter  
56 sequences. Our analyses, combined with those of a study by Simonini et al. (9),  
57 suggest distinct roles for PMEIs and PMEs in mediating the role of ETT in the  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 gynoecium, and indicate the possible presence of rapidly-acting intermediate factors  
5 in the transcriptional regulation of these genes.  
6  
7  
8

## 9 **Results**

### 10 Putative immediate targets of ETT are enriched in genes involved in cell-wall 11 remodeling. 12

13  
14 To identify immediate targets of ETT, we generated transgenic plants in which the  
15 translocation to the nucleus of constitutively produced ETT, translationally fused to the  
16 hormone-binding domain of the rat glucocorticoid receptor protein (GR), could be  
17 induced by exogenous application of the hormone analogue dexamethasone  
18 (DEX)(10). After verification of the functionality of the ETT-GR fusion protein (SI Fig.  
19 1), inflorescence tissues of transformed plants were treated either with DEX and  
20 cycloheximide (CHX) to induce the transcriptional effects of ETT on its target genes  
21 while simultaneously blocking protein synthesis, or with CHX alone to block protein  
22 synthesis without ETT-induction. The efficiency of the CHX treatments given has been  
23 previously demonstrated (11). Apical regions of inflorescences, containing unopened  
24 flowers only, were harvested from both groups of plants two hours after induction. RNA  
25 was then extracted and processed for expression analyses on CATMA GST  
26 microarrays (12).  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 The results of three independent experiments were compared, resulting in a list of 85  
42 putative ETT targets (p-value < 0.005; SI Tab. 1). Among these, 65 genes (76.5%)  
43 were repressed, while the remaining 20 (23.5%) were activated. This predominantly  
44 repressive activity of ETT is in agreement both with previous experimental data and  
45 the presence of a glutamine-rich central region in ETT, shared with other predominantly  
46 repressive ARFs (13). Interestingly, 13 (15.3%) of the genes that were rapidly  
47 regulated following ETT-activation encoded proteins predicted to be involved in cell-  
48 wall structure and/or remodeling, including four PMEs (AT5G07420, AT3G17060,  
49 AT2G26450 and AT5G27870), three PMEIs (AT5G50030, AT2G47050 and  
50 AT1G10770), one expansin (AT1G69530, also known as Ath.EXPA1), one member of  
51 the OLE E 1 allergen and extensin family (AT1G29140), two arabinogalactan proteins  
52 (AT3G01700 and AT5G24105, also known as Ath.AGP11 and 41, respectively), and  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 two pectate lyase-like proteins (AT1G14420 and AT3G01270, also known as Ath.PLL8  
5 and 10, respectively).  
6  
7

8  
9  
10 To compare the frequencies of cell-wall-related genes between putatively ETT-  
11 regulated sequences and the entire *A. thaliana* genome, we used Functional  
12 Classification SuperViewer, based on Gene Ontology from TAIR (14). The result of this  
13 analysis (SI Fig. 2) shows a highly statistically significant, five-fold over-representation  
14 of cell-wall-related genes, among putative immediate ETT targets.  
15  
16  
17  
18

19  
20 All 13 putative ETT cell-wall targets identified here are expressed in flower buds  
21 approaching anthesis, though predominantly in anthers (15). The expansin  
22 AT1G69530 additionally shows high expression in the stamen filaments and petals of  
23 mature flowers and in siliques and germinating seeds. As all 13 genes are  
24 downregulated on induction of ETT activity, their expression in wild-type tissues in  
25 which ETT is expressed, including the gynoecium, is not necessarily to be expected.  
26 However, gene ontology terms in TAIR (<https://www.arabidopsis.org/>), based on  
27 comparisons of transcriptomic data, do indicate these 13 genes to be expressed in  
28 carpel tissues.  
29  
30  
31  
32  
33  
34  
35  
36

37  
38 To confirm the repression by ETT of the 13 cell-wall-related genes identified, we  
39 performed qRT-PCR analyses on inflorescences from Col-0 plants and from  
40 *35S:ETTm* transformants (16) in which *ETT* mRNA is stabilized by silent mutations in  
41 both of its tasiR-ARF binding sites. All 13 cell wall-related targets identified were  
42 significantly down-regulated in *35S:ETTm* (Fig. 1), consistent with the results of our  
43 microarray analyses (SI Tab. 1).  
44  
45  
46  
47  
48  
49  
50

#### 51 Comparison of datasets produced using four distinct methods discriminates between 52 possible direct and indirect targets of ETT. 53

54  
55 We scanned a genome-wide database of presumptive *A. thaliana* promoter sequences  
56 using both of the published PSSMs for ETT (5). The highest affinity sites corresponded  
57 to a weight score (17) of 8.1 in these scans. A weight-score cutoff value of 7.0 produced  
58 a list of 808 promoters containing at least one potential ETT-binding site, though only  
59 ~8% of genes identified to bind ETT in ChIP-seq analyses (9) were present in this list  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 (data not shown). We therefore reduced the weight-score cutoff to 6.0, which produced  
5 a list of 6695 genes (SI Tab. 2) that included ~37% of those previously identified by  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

(data not shown). We therefore reduced the weight-score cutoff to 6.0, which produced a list of 6695 genes (SI Tab. 2) that included ~37% of those previously identified by ChIP-seq.

We compared datasets of ETT-regulated genes produced by four distinct methods: *in vivo* induction of ETT-GR (Tab. S1), RNA-seq comparison of wild-type and *ett-3* inflorescences without exogenous indole acetic acid (IAA)(9), ChIP-seq of *ETT-GFP* transformants without exogenous IAA (9), and *in-silico* scans of *A. thaliana* promoters (SI Tab. 2), the results of which are shown in Fig. 2. Twenty three of the genes identified in the present work by the *in-vivo* activation of ETT-GR were furthermore identified as potentially direct ETT-targets by ChIP-seq and/or promoter-scanning procedures. These genes included four cell-wall-related proteins: PME1 AT5G50030, PME AT3G17060, expansin AT1G69530 and pectolyase-like protein AT1G14420. A further 26 putative immediate ETT-targets showed differential expression between wild-type and *ett-3* inflorescences, though were not predicted as potential direct targets of ETT in ChIP-seq or promoter-scanning procedures. The majority of the cell-wall proteins identified through the *in-vivo* activation of ETT-GR were present in this group, including PMEs AT2G26450 and AT5G07420, PMEIs AT1G10770 and AT2G47050, OLE E 1 protein AT1G29140, pectolyase-like protein AT3G01270, and arabinogalactan protein AT3G01700 (Fig 2). Only two cell wall proteins identified by the *in-vivo* activation of ETT-GR (SI Tab. 1) and subsequent qPCR studies (Fig. 1) failed to be identified in one or more of the other three analyses used for comparison: arabinogalactan protein AT5G24105 and PME AT5G27870.

## Discussion

Transcriptomics experiments suggest a key role for PMEIs in the regulation of tissue patterning by *ETT*.

Andres-Robin et al. (7) showed a correlation between the regulation by ETT of tissue-patterning in the *A. thaliana* gynoecium and enzymatic and biochemical characteristics that pointed to a link with the methylesterification status of cell-wall pectins. A model emerged in which ETT acts to promote PME activity in the cell wall. Here, we provide evidence that ETT acts as a negative regulator of four PMEs and three PMEIs (SI Tab.

1  
2  
3  
4 1). The regulation of all seven of these genes was confirmed by qRT-PCR (Fig. 1), and  
5 in six cases by comparisons with one or more of three further datasets (Fig. 2).  
6  
7

8  
9  
10 From the overexpression experiments in mutant and wild-type backgrounds performed  
11 by Andres-Robin et al. (7), the negative regulation by ETT of PMEIs would be  
12 consistent with the known effects of ETT on the methylesterification of pectins, cell-  
13 wall stiffness and valve length, whereas the negative regulation by ETT of PMEs would  
14 be predicted to have the opposite effect. We therefore postulate that the regulation of  
15 valve-length by ETT is controlled predominantly through the negative regulation of  
16 PMEIs. The significance of the simultaneous negative regulation by ETT of several  
17 PMEs is unclear, though may be an example of a “gas and brake mechanism” (18), in  
18 which an upstream regulator acts in the same direction on mutually antagonistic  
19 classes of downstream components. Such mechanisms are known in the control of  
20 shade-avoidance in plants and are postulated to prevent exaggerated downstream  
21 effects.  
22  
23

24  
25  
26 The above conclusions on the possible developmental significance of ETT-targets  
27 remain, however, speculative as the biochemical properties and developmental effects  
28 of these molecules have yet to be directly investigated. For example, certain PMEs are  
29 known to show distinct enzymatic activities depending on the prior pattern of pectin  
30 methylesterification encountered (19), while certain PMEIs are known to show  
31 differential interaction specificities among PMEs (20)(21). In addition, both PME  
32 activities and certain PMEI-PMI interactions are known to be pH-dependent (19)(21).  
33 We do not, therefore, know at present whether the PMEIs identified here as immediate  
34 ETT-targets would be active against the PMEs also shown to be regulated by ETT.  
35 Neither do we yet know whether those PMEs, if over-expressed in the gynoecium,  
36 would show similar effects on pectin demethylesterification and valve-length to the  
37 effects previously demonstrated for PME5 (7).  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

58 Comparison of datasets suggests the presence of rapidly acting intermediate  
59 components in the regulation of PMEs and PMEIs by ETT.  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 Though several PMEs and PMEIs, among other cell wall-related genes, appear to be  
5 rapidly downregulated as a response to ETT, it is currently unclear whether ETT  
6 interacts directly with the promoters of these genes. Comparisons between datasets  
7 confirm the regulatory interactions identified for three PMEIs and three PMEs (Fig. 2).  
8 However, the presumptive promoters of only two of these genes, PMEI AT5G50030  
9 and PME AT3G17060, contain potential ETT-binding sites, and neither gene emerged  
10 as a putative direct target of ETT in the ChIP-seq analysis of Simonini et al. (9). Taken  
11 together, the data described here suggest the involvement of PMEs and PMEIs in  
12 ETT's mode of *action* in the gynoecium, but suggest the operation of one or more  
13 rapidly-acting intermediate components, downstream of ETT, in the transcriptional  
14 regulation of these genes.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

25  
26 For example, ETT might repress one or more transcriptional activators, whose activity  
27 would thereby be reduced in the gynoecium, leading to lower transcription rates of their  
28 own direct targets, possibly including some of the likely indirect targets of ETT identified  
29 here. Our experiments (SI Tab. 1) and those of Simonini et al. (9) have indicated a  
30 number of transcription factors as possible direct targets of ETT, so that observation  
31 could represent a starting point to investigate this hypothesis further.  
32  
33  
34  
35  
36  
37  
38

## 39 **Material and methods**

### 40 Plant growth conditions.

41 Wild-type, transgenic and/or mutant *A. thaliana* lines were grown on peat-based  
42 compost at 20°C under short-day conditions (8h light/16h dark cycles) for 4 weeks  
43 before being transferred in long-day conditions (16h light/8h dark cycles) to induce  
44 flowering for genetic transformation, induction experiments and/or gene expression  
45 analyses.  
46  
47  
48  
49  
50  
51  
52

### 53 Transgene constructs, induction experiments and microarray analyses.

54 The *ETT* coding sequence was inserted into *pG0229-35S::GR* (10) between the  
55 *CaMV-35S* promoter and a sequence encoding the hormone-binding domain of the rat  
56 glucocorticoid receptor, so as to conserve the reading frame of the latter element. The  
57 resulting plasmid was transferred to *Agrobacterium tumefaciens* C58/pMP90, a culture  
58 of which was then used to transform plants of the *A. thaliana* Col-0 ecotype by standard  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 “floral dip” procedures. A homozygous individual, stably expressing the transgene from  
5 a single insertion event, was selected from the ensuing T2 generation.  
6  
7

8  
9 The ETT-GR transgene construct was then introduced into an *ett-22* mutant  
10 background by cross pollination. The functionality of the ETT-GR fusion protein was  
11 demonstrated on homozygous plants from which all flowers later than Stage 8 (22) had  
12 been removed. Inflorescences were dipped for 2 min, once per day for five days, into  
13 aqueous solutions of DEX (10 µg/ml) containing Silwet L-77 surfactant (0.01% v/v), or  
14 mock solutions lacking DEX. Gynoecia were observed three days after final treatments  
15 using a Leica MZ12 stereomicroscope coupled to a DFC320 digital camera. Fifty  
16 measurements were made using ImageJ for each treatment-group of the *ett-22*  
17 *35S:ETT-GR* line, and 30 measurements for each treatment-group of the *35S:ETT-GR*  
18 and *ett-22* lines.  
19  
20  
21  
22  
23  
24  
25  
26

27  
28 For microarray analyses, two populations, each composed of ten T3 descendants of  
29 the original T2 *35S:ETT-GR* transformant in a wild-type Col-0 genetic background,  
30 were grown to maturity. CHX and DEX treatments, RNA extraction and microarray  
31 analyses were performed as described by Reymond et al. (11), and the entire  
32 experiment was repeated twice to provide three biological replicates. Statistical  
33 analysis of microarray data was also performed as previously described (11). The  
34 complete microarray dataset is available at CATdb (<http://tools.ips2.u-psud.fr/CATdb> )  
35 as Project: GEN45-Carpel\_development. The list of putative ETT-targets obtained was  
36 functionally analyzed using Classification Superviewer (14) at <http://bar.utoronto.ca>.  
37  
38  
39  
40  
41  
42  
43  
44

#### 45 46 47 Quantitative RT-PCR. 48

49 Total RNA was extracted from inflorescences using Spectrum Plant Total RNA Kit  
50 (Sigma). Any contaminating DNA was eliminated by treatments with Turbo DNAase  
51 (Ambion) and 4-µg aliquots of total RNA were then reverse transcribed using RevertAid  
52 Reverse Transcriptase (Fermentas). Reverse transcriptase reactions of 20 µl volume  
53 were diluted to 1 ml, and 5-µl aliquots of these dilutions were subjected to qPCR using  
54 SYBR GREEN (Roche) in a StepOne Plus apparatus (Applied). Primer efficiencies  
55 were determined from standard curves using serial cDNA dilutions and PCR  
56 amplification was performed using a three-step protocol, incorporating a melting curve.  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Results were normalized to the expression of GAPDH (AT3G26650), chosen from a  
5 list of potential control genes using BESTKEEPER (23), and analyzed using the  $2^{-(\Delta\Delta Ct)}$   
6 method. Each expression value was determined from the mean of a minimum of three  
7 plants, and each experiment was repeated twice independently. The primers used are  
8 given in the SI Tab 3.  
9

#### 10 *In-silico* promoter analyses and global data comparisons.

11 A genome-wide set of *A. thaliana* promoters (<http://arabidopsis.med.ohio-state.edu/>)  
12 (24) was scanned with PSSMs for ETT (5) using the Matrix-Scan program in RSAT  
13 (17). Scans were combined and filtered on the basis of score-values. Gene lists  
14 produced using various different methods were compared in Venny 2.1  
15 (<https://bioinfogp.cnb.csic.es/tools/venny/>).  
16  
17

#### 18 **Acknowledgments**

19 We are grateful to R. Scott Poethig for seeds of *35S:ETTm* and acknowledge funding  
20 from Génoplante (G-45) and the French National Research Agency (ANR-BLAN-0211-  
21 01) to CPS. IPS2 is supported by LabEx Saclay Plant Sciences-SPS (ANR-10-LABX-  
22 0040-SPS). AMR and MCR received doctoral studentship from the French Ministry of  
23 Higher Education, Research and Innovation and from the Rhone-Alps Region,  
24 respectively.  
25  
26

#### 27 **References**

- 28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46 1. A. Sessions, *et al.*, ETTIN patterns the Arabidopsis floral meristem and reproductive  
47 organs. *Development* **124**, 4481–4491 (1997).  
48  
49  
50 2. I. Pekker, J. P. Alvarez, Y. Eshed, Auxin response factors mediate Arabidopsis organ  
51 asymmetry via modulation of KANADI activity. *Plant Cell* **17**, 2899–2910 (2005).  
52  
53 3. D. Garcia, S. A. Collier, M. E. Byrne, R. A. Martienssen, Specification of leaf polarity in  
54 Arabidopsis via the trans-acting siRNA pathway. *Curr. Biol.* **16**, 933–938 (2006).  
55  
56 4. T. J. Guilfoyle, G. Hagen, Auxin response factors. *J. Plant Growth Regul.* **20**, 281–291  
57 (2001).  
58  
59 5. J. M. Franco-Zorrilla, *et al.*, DNA-binding specificities of plant transcription factors and  
60 their potential to define target genes. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 2367–2372  
61 (2014).  
62  
63  
64  
65

- 1
- 2
- 3
- 4 6. M. L. Bulyk, P. L. F. Johnson, G. M. Church, Nucleotides of transcription factor binding
- 5 sites exert interdependent effects on the binding affinities of transcription factors.
- 6 *Nucleic Acids Res.* **30**, 1255–1261 (2002).
- 7
- 8
- 9 7. A. Andres-Robin, *et al.*, Evidence for the Regulation of Gynoecium Morphogenesis by
- 10 ETTIN via Cell Wall Dynamics. *Plant Physiol.* **178**, 1222–1232 (2018).
- 11
- 12 8. A. Peaucelle, *et al.*, Arabidopsis phyllotaxis is controlled by the methyl-esterification
- 13 status of cell-wall pectins. *Curr. Biol.* **18**, 1943–1948 (2008).
- 14
- 15 9. S. Simonini, S. Bencivenga, M. Trick, L. Ostergaard, Auxin-Induced Modulation of
- 16 ETTIN Activity Orchestrates Gene Expression in Arabidopsis. *Plant Cell* **29**, 1864–1882
- 17 (2017).
- 18
- 19
- 20 10. H. Yu, T. Ito, F. Wellmer, E. M. Meyerowitz, Repression of AGAMOUS-LIKE 24 is a
- 21 crucial step in promoting flower development. *Nat. Genet.* **36**, 157–161 (2004).
- 22
- 23 11. M. C. Reymond, *et al.*, A light-regulated genetic module was recruited to carpel
- 24 development in Arabidopsis following a structural change to SPATULA. *Plant Cell* **24**,
- 25 2812–2825 (2012).
- 26
- 27
- 28 12. M. L. Crowe, *et al.*, CATMA: a complete Arabidopsis GST database. *Nucleic Acids Res.*
- 29 **31**, 156–158 (2003).
- 30
- 31
- 32 13. S. B. Tiwari, G. Hagen, T. Guilfoyle, The roles of auxin response factor domains in
- 33 auxin-responsive transcription. *Plant Cell* **15**, 533–543 (2003).
- 34
- 35 14. N. J. Provar, T. Zhu, A browser-based functional classification SuperViewer for
- 36 Arabidopsis genomics. *Curr Comput. Mol Biol*, 271–272 (2003).
- 37
- 38 15. A. V. Klepikova, A. S. Kasianov, E. S. Gerasimov, M. D. Logacheva, A. A. Penin, A
- 39 high resolution map of the Arabidopsis thaliana developmental transcriptome based on
- 40 RNA-seq profiling. *Plant J.* **88**, 1058–1070 (2016).
- 41
- 42 16. C. Hunter, *et al.*, Trans-acting siRNA-mediated repression of ETTIN and ARF4
- 43 regulates heteroblasty in Arabidopsis. *Development* **133**, 2973–2981 (2006).
- 44
- 45 17. J.-V. Turatsinze, M. Thomas-Chollier, M. Defrance, J. van Helden, Using RSAT to scan
- 46 genome sequences for transcription factor binding sites and cis-regulatory modules. *Nat.*
- 47 *Protoc.* **3**, 1578–1588 (2008).
- 48
- 49 18. C. D. Crocco, M. Holm, M. J. Yanovsky, J. F. Botto, Function of B-BOX under shade.
- 50 *Plant Signal. Behav.* **6**, 101–104 (2011).
- 51
- 52 19. F. Senechal, *et al.*, PECTIN METHYLESTERASE INHIBITOR7 modulates the
- 53 processive activity of co-expressed PECTIN METHYLESTERASE3 in a pH-dependent
- 54 manner. *J. Biol. Chem.* **290**, 23320–23335 (2015).
- 55
- 56 20. F. Senechal, A. Mareck, P. Marcelo, P. Lerouge, J. Pelloux, Arabidopsis PME17
- 57 Activity can be Controlled by Pectin Methylesterase Inhibitor4. *Plant Signal. Behav.* **10**,
- 58 UNSP e983351 (2015).
- 59
- 60
- 61
- 62
- 63
- 64
- 65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

21. L. Hocq, *et al.*, Combined Experimental and Computational Approaches Reveal Distinct pH Dependence of Pectin Methylesterase Inhibitors. *Plant Physiol.* **173**, 1075–1093 (2017).
22. D. Smyth, J. Bowman, E. Meyerowitz, Early flower development in Arabidopsis. *Plant Cell* **2**, 755–767 (1990).
23. M. W. Pfaffl, A. Tichopad, C. Prgomet, T. P. Neuvians, Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper--Excel-based tool using pair-wise correlations. *Biotechnol. Lett.* **26**, 509–515 (2004).
24. A. Yilmaz, *et al.*, AGRIS: the Arabidopsis Gene Regulatory Information Server, an update. *Nucleic Acids Res.* **39**, D1118–D1122 (2011).

1  
2  
3  
4 **Figure and table legends**

5  
6 Figure 1. qRT-PCR analyses of 13 cell-wall-related genes.

7 All genes studied are significantly lower expressed ( $p \leq 0.05$ ) in plants overexpressing  
8 a stabilized *ETTm* transgene compared to wild-type. Standard errors of the mean are  
9 shown.  
10

11  
12  
13  
14  
15 Figure 2. Comparison of four datasets to confirm putative immediate ETT targets.

16 Total numbers of genes occurring in each combination of datasets are indicated. Ten  
17 chloroplast genes that were rapidly regulated following ETT-GR induction (SI Tab. 1)  
18 were excluded from the analysis. The positions of 13 cell-wall-related genes (SI Tab.  
19 1, Fig 1) are indicated in boxes.  
20  
21  
22  
23  
24

25  
26 SI Figure 1. Phenotypic effects of ETT-GR activation.

27 Dexamethasone treatment to activate the ETT-GR fusion protein had no effect on  
28 valve/gynoecium length-ratio in *35S:ETT-GR* transformants in a wild-type genetic  
29 background, or in untransformed *ett-22* mutants, but restored the wild-type  
30 valve/gynoecium length-ratio to *35S:ETT-GR* transformants in an *ett-22* mutant  
31 background, demonstrating the *in-vivo* functionality of the *35S:ETT-GR* construct.  
32  
33  
34  
35  
36  
37

38  
39 SI Figure 2. Functional classification of immediate ETT-targets using Classification  
40 Superviewer.

41 Cell-wall-related genes are the most significantly over-represented group that  
42 responds rapidly to ETT-GR activation. Error bars represent standard deviations for  
43 100 bootstraps of the input set. P-values  $< 10^{-2}$  (\*) and  $< 10^{-4}$  (\*\*).  
44  
45  
46  
47  
48  
49

50 SI Table 1. Genes rapidly upregulated following ETT-GR activation in transgenic  
51 plants.

52  
53  
54  
55 SI Table 2. Potential ETT-binding sites in *Arabidopsis thaliana* promoters.

56  
57  
58  
59 SI Table 3. Primers used for qRT-PCR analyses.  
60  
61  
62  
63  
64  
65

Fig 1




Fig 2

