



HAL
open science

Symmetry group of the equiangular cubed sphere

Jean-Baptiste Bellet

► **To cite this version:**

Jean-Baptiste Bellet. Symmetry group of the equiangular cubed sphere. Quarterly of Applied Mathematics, 2022, 80, pp.69-86. hal-03071135v2

HAL Id: hal-03071135

<https://hal.science/hal-03071135v2>

Submitted on 4 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SYMMETRY GROUP OF THE EQUIANGULAR CUBED SPHERE

JEAN-BAPTISTE BELLET

ABSTRACT. The equiangular cubed sphere is a spherical grid, widely used in computational physics. This paper deals with mathematical properties of this grid. We identify the symmetry group, *i.e.* the group of the orthogonal transformations that leave the cubed sphere invariant. The main result is that it coincides with the symmetry group of a cube. The proposed proof emphasizes metric properties of the cubed sphere. We study the geodesic distance on the grid, which reveals that the shortest geodesic arcs match with the vertices of a cuboctahedron. The results of this paper lay the foundation for future numerical schemes, based on rotational invariance of the cubed sphere.

1. INTRODUCTION

1.1. Cubed sphere. Various fields in computational physics, for instance climatology modelling [31], involve numerical computations on the sphere. This includes the use of spherical grids [32]. The grids obtained by radial projection of a circumscribed cube on the sphere are among the most employed. These *cubed sphere grids* have been originally introduced in [27], and further studied, for example in [13, 18–20, 22, 23, 25]. A wide variety of numerical methods have been successfully adapted to cubed sphere grids, *e.g.* in [3–10, 12, 14, 21, 26, 29, 30] and the references therein.

This paper deals with mathematical properties of a cubed sphere structured by equiangular great circles: the *equiangular cubed sphere* CS_N [3, 20], depicted in Figure 1 and in Appendix A. As can be observed, the cubed sphere is quasi-uniform, is not polarized along a specific axis, and is shaped by the cube, including discontinuities across “edges” (radial projection of the edges of the circumscribed cube). Furthermore, some symmetry properties, such as invariance under permutation of the cartesian coordinates, can be noticed.

The symmetry of the cubed sphere has been previously used in various contexts, for instance in [3, 16, 17, 25]. However, to the best of our knowledge, a systematic study of the isometries of CS_N is not available so far. This paper aims at filling this gap by identifying the symmetry group of CS_N . Such a study provides a valuable mathematical background for the applications. Indeed, knowing the symmetry group of a grid supports the design of numerical schemes, such as interpolation methods [24], quadrature rules [28], or Discrete Fourier Transforms [15].

1.2. Main results. We study the symmetries of the cubed sphere, by means of its symmetry group [1]:

Definition 1.1. The symmetry group of CS_N is the set \mathcal{G}_N of the orthogonal matrices that leave CS_N invariant:

$$\mathcal{G}_N = \{Q \in \mathbb{R}^{3 \times 3} : Q^\top Q = QQ^\top = I_3 \text{ and } Q\text{CS}_N = \{Qu, u \in \text{CS}_N\} = \text{CS}_N\}.$$

For $N = 1$, $\text{CS}_1 = \{-1/\sqrt{3}, 1/\sqrt{3}\}^3$ is a scaling of $\{-1, 1\}^3$. Hence the group \mathcal{G}_1 is exactly the symmetry group \mathcal{O} of the cube $\{-1, 1\}^3$: $\mathcal{G}_1 = \mathcal{O}$. This group is well known: it is isomorphic to the group $\mathfrak{S}_4 \times \mathbb{Z}/2\mathbb{Z}$ [1, pp. 37, 38, 55]; any symmetry of the cube is indeed identified with a permutation of the four principal diagonals, combined with a toggle for inversion of the cube, or not. We refer to Appendix B for a matricial representation of \mathcal{O} . The main result of this paper is that $\mathcal{G}_N = \mathcal{O}$, for every $N \geq 1$:

Theorem 1.2 (Symmetry group of the equiangular cubed sphere). *Let $N \geq 1$. The symmetry group \mathcal{G}_N of the cubed sphere CS_N coincides with the symmetry group \mathcal{O} of the cube $\{-1, 1\}^3$. In other words, an orthogonal matrix Q leaves CS_N invariant if, and only if, it leaves $\{-1, 1\}^3$ invariant.*

Date: October 4, 2021.

2020 Mathematics Subject Classification. 86A08, 86A10, 65M50, 20B30, 52B15.

Key words and phrases. Cubed sphere, spherical grid, symmetry, octahedral group.




FIGURE 1. Cubed sphere grid CS_N ($N = 3$). A cartesian grid C_N (black dots) is defined on the faces of the cube $[-1, 1]^3$, by means of the two-dimensional grid $\text{T}_N \times \text{T}_N$, where $\text{T}_N = \{\tan \frac{k\pi}{2N}, -\frac{N}{2} \leq k \leq \frac{N}{2}\}$ is an equiangular grid. The cubed sphere grid CS_N (white dots) is defined as the radial projection of C_N on the unit sphere: $\text{CS}_N = \rho(\text{C}_N)$, with $\rho : x \in \mathbb{R}^3 \setminus \{0\} \mapsto \frac{x}{\|x\|} \in \mathbb{S}^2$. On the faces of the cube, the grid C_N defines a mesh (dotted straight lines). The projection ρ maps this mesh onto a mesh of the sphere: the (dotted) straight lines are mapped onto equiangular arcs of great circles (white curves).

The most difficult inclusion to be proved is $\mathcal{G}_N \subset \mathcal{O}$. One possible approach deduces this result from $\mathcal{O} \subset \mathcal{G}_N$, using group theory as Appendix C. Instead of this theoretical approach, this paper focuses on a direct proof which investigates metric properties of CS_N . This direct approach has the advantage of deepening the knowledge of CS_N . More precisely, we solve the problem of the shortest geodesic arcs on CS_N :

$$\min\{\theta(u, v) := \arccos u \cdot v; u \in \text{CS}_N, v \in \text{CS}_N, u \neq v\}. \quad (\text{P})$$

One original feature of problem (P) is that u and v are allowed to belong to distinct grid lines, or even distinct panels; in particular, any spherical diagonal of the mesh is realizable. The main difficulty of problem (P) comes from this specificity. This is somehow the lock of Theorem 1.2. Any solution of (P) is given by the following theorem.

Theorem 1.3 (Shortest arcs on the equiangular cubed sphere). *Let $u, v \in \text{CS}_N$ with $u \neq v$. The arc $\{u, v\}$ is optimal for (P) if, and only if, there exists an isometry of the cube, $Q \in \mathcal{O}$, such that*

$$\begin{cases} \{u, v\} = \{Q\rho(1, -\tan \frac{\pi}{4N}, 1), Q\rho(1, \tan \frac{\pi}{4N}, 1)\}, & \text{if } N \text{ is odd,} \\ \{u, v\} = \{Q\rho(1, 0, 1), Q\rho(1, \tan \frac{\pi}{2N}, 1)\}, & \text{if } N \text{ is even.} \end{cases}$$

In other words, the minimal arcs are described by the orbit of

$$\{\rho(1, -\tan \frac{\pi}{4N}, 1), \rho(1, \tan \frac{\pi}{4N}, 1)\}, \quad \{\rho(1, 0, 1), \rho(1, \tan \frac{\pi}{2N}, 1)\},$$

for the group action $(Q, \{u, v\}) \mapsto \{Qu, Qv\}$, $Q \in \mathcal{O}$, $\{u, v\} \subset \text{CS}_N$. The matricial representation (B.1) of \mathcal{O} shows immediately the following corollary.

Corollary 1.4. (i) *If N is odd, there are precisely 12 minimal arcs on CS_N (one per edge):*

$$\{\rho(-\delta, \epsilon_1, \epsilon_2), \rho(\delta, \epsilon_1, \epsilon_2)\}, \{\rho(\epsilon_1, -\delta, \epsilon_2), \rho(\epsilon_1, \delta, \epsilon_2)\}, \{\rho(\epsilon_1, \epsilon_2, -\delta), \rho(\epsilon_1, \epsilon_2, \delta)\}, \\ \delta = \tan \frac{\pi}{4N}, \epsilon_1 = \pm 1, \epsilon_2 = \pm 1.$$

(ii) *If N is even, there are precisely 24 minimal arcs on CS_N (two per edge):*

$$\{\rho(0, \epsilon_1, \epsilon_2), \rho(\delta, \epsilon_1, \epsilon_2)\}, \{\rho(\epsilon_1, 0, \epsilon_2), \rho(\epsilon_1, \delta, \epsilon_2)\}, \{\rho(\epsilon_1, \epsilon_2, 0), \rho(\epsilon_1, \epsilon_2, \delta)\}, \\ \delta = \pm \tan \frac{\pi}{2N}, \epsilon_1 = \pm 1, \epsilon_2 = \pm 1.$$

Roughly speaking, the minimal arcs are “short arcs around the midpoints on the edges”, as displayed in Figure 2. These midpoints match with the vertices of the *cuboctahedron* (B.3). Our proof of Theorem 1.2 is mainly based on the invariance of this cuboctahedron under \mathcal{G}_N .




FIGURE 2. The bold arcs, around the midpoints on the edges, are minimal arcs on CS_N . Left: N is odd ($N = 3$); right: N is even ($N = 4$). These arcs match with the vertices of a cuboctahedron (black line).

1.3. Organization of the paper. The paper is organized as follows. In Section 2, we prove Theorem 1.2. In Section 3, we prove Theorem 1.3. The proof is based on a series of lemmas which are proved in Section 4. In Appendix A, we recall the definition of CS_N . Some geometrical and matricial properties of \mathcal{O} are given in Appendix B. In Appendix C, we give a proof of Theorem 1.2, based on group theory.

2. PROOF OF THE MAIN THEOREM

Lemma 2.1. *Any symmetry of $\{-1, 1\}^3$ leaves CS_N invariant, i.e. $\mathcal{O} \subset \mathcal{G}_N$.*

Proof. Firstly, it is clear from the definition (A.1) that CS_N is stable under the following operations:

- permutation of some coordinates;
- toggle the sign of some coordinates.

The last point is due to the symmetry of the set T_N ($u \in \text{T}_N \Rightarrow -u \in \text{T}_N$). Secondly, by (B.2), any $Q \in \mathcal{O}$ is a composition of two such operations; therefore, $Q\text{CS}_N \subset \text{CS}_N$. Since Q is orthogonal, then $Q\text{CS}_N = \text{CS}_N$ and $Q \in \mathcal{G}_N$. \square

Remark 2.2. The result can be extended for other cubed sphere grids: the equiangular property of the grid T_N does not matter; the key property is the symmetry of T_N .

Theorem 2.3. *Let Ω be the cuboctahedron (B.3). Let $Q \in \mathcal{G}_N$. The isometry Q leaves Ω invariant.*

Proof. Case 1: N is odd. We introduce the midpoints of the minimal arcs:

$$\Omega' = \left\{ \frac{1}{2}(u + v), \{u, v\} \text{ is minimal for (P)} \right\}.$$

By Corollary 1.4, Ω' is a scaling of Ω :

$$\Omega' = \left\{ (2 + \tan^2 \frac{\pi}{4N})^{-1/2} x, x \in \Omega \right\}.$$

We prove that Ω' is left invariant by Q . Fix $w = \frac{u+v}{2} \in \Omega'$ with $\{u, v\}$ minimal. Then $\{Qu, Qv\}$ is minimal due to $Q \in \mathcal{G}_N$. As a result $Qw = \frac{Qu+Qv}{2} \in \Omega'$. Thus $Q\Omega' \subset \Omega'$. Finally $Q\Omega = \Omega$.

Case 2: N is even. We prove that $\rho(\Omega) = \{2^{-1/2}x, x \in \Omega\}$ is left invariant by Q . By Corollary 1.4,

$$\rho(\Omega) = \{u \in \text{CS}_N : n(u) = 2\},$$

where $n(u)$ denotes the number of v such that $\{u, v\}$ is minimal:

$$n(u) = \#\{v \in \text{CS}_N : \{u, v\} \text{ is minimal for (P)}\}, \quad u \in \text{CS}_N.$$

By the way, the arc $\{u, v\}$ is minimal if, and only if, $\{Qu, Qv\}$ is minimal (due to $Q \in \mathcal{G}_N$). At the end, for every $u \in \text{CS}_N$, $n(u) = n(Qu)$, and

$$u \in \rho(\Omega) \Leftrightarrow n(u) = 2 \Leftrightarrow n(Qu) = 2 \Leftrightarrow Qu \in \rho(\Omega).$$

Then $Q\rho(\Omega) = \rho(\Omega)$ and $Q\Omega = \Omega$. \square

Proof of Theorem 1.2. We have proved $\mathcal{O} \subset \mathcal{G}_N$ in Lemma 2.1. The converse is a corollary to Theorems 2.3 and B.5: any $Q \in \mathcal{G}_N$ leaves the cuboctahedron Ω invariant, so it leaves the cube $\{-1, 1\}^3$ invariant. \square

3. PROOF OF THEOREM 1.3

In this section we minimize the great circle distance $\theta(u, v) = \arccos u \cdot v$ between $u \neq v \in \text{CS}_N$. Briefly, any minimal arc on CS_N is minimal on the edge $(I) \cap (V)$, up to an isometry of the cube.

Notation 3.1. On the edge $(I) \cap (V) = \{\rho(1, \tan \frac{k\pi}{2N}, 1), |k| \leq \frac{N}{2}\}$, the shortest arc-length is denoted by

$$m^* := \min\{\theta(u, v), u \neq v \in (I) \cap (V)\}. \quad (3.1)$$

Theorem 3.2. *Let $u \neq v \in \text{CS}_N$ be such that $\{u, v\}$ is optimal for (P). Then there exists $Q \in \mathcal{O}$ such that $\{Qu, Qv\}$ is optimal for (3.1), on the edge $(I) \cap (V)$.*

Proof of Theorem 3.2. By Lemma 2.1, for every $Q \in \mathcal{O}$, $\{Qu, Qv\} \subset \text{CS}_N$ is optimal: $\theta(Qu, Qv) = \theta(u, v)$. One or both of the following cases applies.

Case 1: u, v belong to the same panel. We define an isometry $Q \in \mathcal{O}$ such that $Qu, Qv \in (I)$. Then $\{Qu, Qv\}$ is optimal for (3.3), on the panel (I) . By Lemma 3.5, $\{Qu, Qv\}$ is optimal for (3.2), on the grid lines of the panel (I) . By Lemma 3.4, there exists $P \in \mathcal{O}$ such that $\{PQu, PQv\}$ is optimal for (3.1), on the edge $(I) \cap (V)$.

Case 2: u, v belong to adjacent panels. We set an isometry $Q \in \mathcal{O}$ such that $Qu \in (I)$ and $Qv \in (V)$. Then $\{Qu, Qv\}$ is optimal for (3.4), on the adjacent panels (I) and (V) ; Lemma 3.6 shows that Qu and Qv belong to the same panel. We conclude using Case 1 for $\{Qu, Qv\}$.

Case 3: u, v belong to opposite panels. We set an isometry $Q \in \mathcal{O}$ such that $Qu \in (I)$ and $Qv \in (III)$. We apply Lemma 3.7: $\{Qu, Qv\}$ is optimal for (3.5), on the opposite panels (I) and (III) , with $Qu \cdot Qv = \frac{1}{3}$. If $N = 1$, $\text{CS}_N = \{-1/\sqrt{3}, 1/\sqrt{3}\}^3$, then Qu and Qv belong to the same edge. So we conclude with Case 1 for $\{Qu, Qv\}$. If $N \geq 2$, this case is impossible. Otherwise, $\theta(u, v) = \theta(Qu, Qv) > m^*$ which contradicts the optimality of $\{u, v\}$. \square

Proof of Theorem 1.3. Problem (P) is an optimization problem, over a finite set. Therefore there exists an optimal arc: $\{u, v\} \subset \text{CS}_N$, with $u \neq v$. Then, by Theorem 3.2, there exists $Q \in \mathcal{O}$ such that $\{Qu, Qv\} \subset (I) \cap (V)$ is optimal for (3.1), on the edge $(I) \cap (V)$. This arc is one of the arcs of Lemma 3.3. Furthermore the minimal value (P) is $\theta(u, v) = \theta(Qu, Qv) = m^*$. This shows that $\{Qu, Qv\}$ is optimal for the full problem (P). Then, by Lemma 2.1, for every $P \in \mathcal{O}$, $\{PQu, PQv\}$ is optimal for (P): $\theta(PQu, PQv) = \theta(u, v)$. \square

We formulate hereafter the lemmas that are used in the proofs of Theorems 1.3 and 3.2. They are proved in Section 4.

Lemma 3.3 (Shortest arc on an edge). *(i) If N is odd, there exists a unique minimal arc for (3.1):*

$$\{u, v\} = \{\rho(1, -\tan \frac{\pi}{4N}, 1), \rho(1, \tan \frac{\pi}{4N}, 1)\}.$$

(ii) If N is even, there are precisely two minimal arcs for (3.1):

$$\{Qu, Qv\} = \{\rho(1, 0, 1), \rho(1, \pm \tan \frac{\pi}{2N}, 1)\}, \text{ with } Q = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \pm 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \in \mathcal{O}.$$

Lemma 3.4 (Shortest arc on the grid lines). *The shortest arc-length on the grid lines of the panel (I) satisfies:*

$$\min \{\theta(u, v); u = \rho(1, X, Y), v = \rho(1, Z, T), X = Z \text{ or } Y = T, (X, Y) \neq (Z, T) \in \mathbb{T}_N^2\} = m^*. \quad (3.2)$$

Moreover, the arc $\{u, v\}$ is optimal for (3.2), if, and only if, $\{Qu, Qv\}$ is optimal for (3.1), where Q is one of the following symmetry of the cube

$$I_3, Q_1 := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, Q_2 := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{bmatrix}, Q_3 := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix} \in \mathcal{O}.$$

Lemma 3.5 (Shortest arc on a panel). *The shortest arc-length between distinct points on the panel (I) satisfies:*

$$\min\{\theta(u, v); u \in (I), v \in (I), u \neq v\} = m^*; \quad (3.3)$$

moreover, the arc $\{u, v\}$ is optimal for (3.3) if, and only if, $\{u, v\}$ is optimal for (3.2).

The shortest arcs on the panel (I) are displayed in Figure 3.




FIGURE 3. Shortest arcs between distinct points in (I) (in dark). If N is odd (left), there are 4 minimal arcs, obtained by rotation of $\{\rho(1, -\tan \frac{\pi}{4N}, 1), \rho(1, \tan \frac{\pi}{4N}, 1)\} \subset (I) \cap (V)$. If N is even (right), there are 8 minimal arcs, obtained by symmetry of $\{\rho(1, 0, 1), \rho(1, \tan \frac{\pi}{2N}, 1)\} \subset (I) \cap (V)$.

Lemma 3.6 (Shortest arc between adjacent panels). *Let $u \in (I) \setminus (V)$ and $v \in (V) \setminus (I)$. The arc (u, v) is not a minimal arc between (I) and (V):*

$$\theta(u, v) > \min\{\theta(u', v'); u' \in (I), v' \in (V), u' \neq v'\}. \quad (3.4)$$

Lemma 3.7 (Shortest arc between opposite panels). *The shortest arc-length between the panels (I) and (III) satisfies:*

$$\min\{\theta(u, v); u \in (I), v \in (III), u \neq v\} = \theta(\rho(1, -1, 1), \rho(1, 1, 1)) \geq m^*, \quad (3.5)$$

with equality if, and only if, $N = 1$.

Lemmas 3.6–3.7 are illustrated in Figure 4.




FIGURE 4. The distance between points on distinct panels is dominated by the distance between some points in a panel. Left: the edge between the corners $\rho(1, 1, 1)$ and $\rho(-1, 1, 1)$ is a minimal arc between the opposite panels (I) and (III). Middle: $\rho(1, T, Z) \in (I)$ is closer to $\rho(1, X, Y) \in (I)$ than the symmetric $\rho(Z, T, 1) \in (V)$. Right: the distance between $\rho(1, X, Y)$ and $\rho(t, X, 1)$ (on the dashed line) decreases when t increases from Y to 1; $\rho(1, X, 1) \in (I) \cap (V)$ is closer to $\rho(1, X, Y)$ than $\rho(Y, X, 1) \in (V)$.

4. PROOF OF LEMMAS 3.3–3.7

In this section, we minimize the arc-length $\theta(u, v)$ over various subsets of the cubed sphere, or, equivalently, we maximize the cos-angle $\cos\theta(u, v) = u \cdot v$ (cos is decreasing on $[0, \pi]$). Most of the proofs are based on Lemma 4.1.

4.1. **Technical lemma.** This subsection is about the continuous face containing the first panel:

$$\rho(\{1\} \times [-1, 1]^2) = \{\rho(1, X, Y), (X, Y) \in [-1, 1]^2\} \supset (I).$$

We investigate various strategies that increase the cos-angle

$$F(X, Y, Z, T) := \rho(1, X, Y) \cdot \rho(1, Z, T) = \frac{1 + XZ + YT}{(1 + X^2 + Y^2)^{1/2}(1 + Z^2 + T^2)^{1/2}},$$

$$X, Y, Z, T \in [-1, 1].$$

Firstly, the cos-angle satisfies the following symmetries:

$$F(X, Y, Z, T) = F(Y, X, T, Z) = F(Z, T, X, Y) = F(T, Z, Y, X) \quad (4.1)$$

$$= F(-X, Y, -Z, T) = F(X, -Y, Z, -T) = F(-X, -Y, -Z, -T). \quad (4.2)$$

Secondly, the following inequalities are easily checked:

$$XZ \leq 0 \Rightarrow F(X, Y, Z, T) = F(-|X|, Y, |Z|, T) \leq F(|X|, Y, |Z|, T), \quad (4.3)$$

$$YT \leq 0 \Rightarrow F(X, Y, Z, T) = F(X, -|Y|, Z, |T|) \leq F(X, |Y|, Z, |T|), \quad (4.4)$$

$$F(|X|, |Y|, -|Z|, -|T|) \leq F(X, Y, Z, T) \leq F(|X|, |Y|, |Z|, |T|). \quad (4.5)$$

Lastly, we summarize various monotonicity properties:

Lemma 4.1. Let $F(X, Y, Z, T) = \rho(1, X, Y) \cdot \rho(1, Z, T)$, $X, Y, Z, T \in [-1, 1]$.

(i) Let $-1 \leq X < Z \leq 1$. The function $Y \in [0, 1] \mapsto F(X, Y, Z, Y)$ is increasing.

(ii) Let $0 < \delta \leq \frac{\pi}{4}$, and $0 \leq Y \leq 1$. The function $x \in [-\frac{\delta}{2}, \frac{\pi}{4} - \delta] \mapsto F(\tan x, Y, \tan(x + \delta), Y)$ is decreasing if $Y > 0$, and constant if $Y = 0$.

(iii) The function $X \in [0, 1] \mapsto F(-X, 1, X, 1)$ is decreasing.

(iv) Let $-1 < Z \leq 1$ and $-1 \leq Y \leq 1$. The function $X \in [-1, Z] \mapsto F(X, Y, Z, Y)$ is increasing.

(v) Let $-1 \leq X < 1$ and $-1 \leq Y \leq 1$. The function $Z \in [X, 1] \mapsto F(X, Y, Z, Y)$ is decreasing.

(vi) Let $0 \leq X < Z \leq 1$ and $0 < Y \leq 1$. The function $T \in [0, Y] \mapsto F(X, Y, Z, T)$ is increasing.

(vii) Let $0 \leq X < Z \leq 1$ and $0 < T \leq 1$. The maximum of the function $Y \in [0, T] \mapsto F(X, Y, Z, T)$ is reached only for $Y = Y^*(X, Z, T) = \frac{T(1+X^2)}{1+ZX} \in (0, T]$.

(viii) Let $0 < \delta \leq \frac{\pi}{4}$ and $0 < T \leq 1$. The function

$$x \in [0, \frac{\pi}{4} - \delta] \mapsto F(\tan x, Y^*(\tan x, \tan(x + \delta), T), \tan(x + \delta), T) \text{ is decreasing.}$$

(ix) Let $0 \leq Y \leq 1$, $-1 \leq Z, T \leq 0$. The function $X \in [0, 1] \mapsto F(X, Y, Z, T)$ is non-increasing.

(x) Let $0 \leq X, Y \leq 1$, $-1 \leq T \leq 0$. The function $Z \in [-1, 0] \mapsto F(X, Y, Z, T)$ is non-decreasing.

Before the proof, we comment the results from a geometrical point of view; see Figure 5.

(i) The length of an iso-Y arc between two fixed meridians (iso-X) decreases from the equator $Y = 0$ to the edge $Y = 1$.

(ii) In the Northern Hemisphere, the length of an iso-Y arc between two meridians, separated from a longitude δ , decreases when the meridians move towards the central position (longitudes $\pm\delta/2$). Along the equator, it is constant.

(iii) Along the edge $Y = 1$, the arc-length between the symmetric meridians $X = -X_1$ and $X = X_1$ decreases with X_1 .

(iv)-(v) The length of an iso-Y arc between two meridians decreases when the longitudes of the meridians get closer.

(vi) In the eastern part of the Northern Hemisphere, we consider a spherical quadrangle delimited by two meridians and two iso-Y arcs. The two meridians and the northernmost iso-Y arc are fixed. The length of the NorthWest-SouthEast diagonal decreases when the southernmost iso-Y arc moves towards the northernmost one.

(vii) With the same pattern than (vi), the length of the SouthWest-NorthEast diagonal reaches its minimum value for some position of the southernmost iso-Y arc ($Y = Y^*$).

(viii) The minimal SouthWest-NorthEast diagonal of (vii) decreases when the westernmost meridian moves towards the central meridian $X = 0$, while keeping a fixed longitude δ between the meridians.

(ix-x) We consider the arc between a point in the eastern part of the Northern Hemisphere and a point in the western part of the Southern Hemisphere. The arc-length does not decrease when the




FIGURE 5. Illustrations of Lemma 4.1. We plot two arcs of great circle associated with two values of F ; the bold arc is shorter (F is larger). The dashed arcs are iso- X or iso- Y , in $\{\rho(1, X, Y), (X, Y) \in [-1, 1]^2\}$.

- (i) $F(X, Y, Z, Y) < F(X, 1, Z, 1)$, $X < Z$, $0 \leq Y < 1$.
- (ii) $F(\tan x, Y, \tan(x + \delta), Y) < F(-\tan \frac{\delta}{2}, Y, \tan \frac{\delta}{2}, Y)$, $0 < \delta$, $0 < Y$, $-\frac{\delta}{2} < x$.
- (iii) $F(-X_1, 1, X_1, 1) < F(-X_2, 1, X_2, 1)$, $0 \leq X_2 < X_1$.
- (iv) $F(X_1, Y, Z, Y) < F(X_2, Y, Z, Y)$, $-1 \leq X_1 < X_2 \leq Z$.
- (vi) $F(X, Y, Z, T) < F(X, Y, Z, Y)$, $0 \leq X < Z$, $0 \leq T < Y$.
- (vii) $F(X, Y, Z, T) < F(X, Y^*(X, Z, T), Z, T)$, $0 \leq X < Z$, $0 < T \leq 1$, $0 \leq Y \leq T$, $Y \neq Y^*(X, Z, T) := \frac{T(1+X^2)}{1+ZX}$.
- (viii) $F(\tan x, Y^*(\tan x, \tan(x + \delta), T), \tan(x + \delta), T) < F(0, T, \tan \delta, T)$, $0 < \delta, T, x$.
- (ix) $F(1, Y, Z, T) \leq F(X, Y, Z, T)$, $X, Y \geq 0$, $Z, T \leq 0$.

easternmost point moves towards the East (along the iso- Y arc), or when the westernmost point moves towards the West.

Proof of Lemma 4.1. For each case, we denote by f the considered function. We give an expression that contains the sign of the derivative f' ; these expressions have been checked by symbolic computation. By the chain-rule, the derivative f' always involves the gradient ∇F , whose components are analogous expressions to:

$$\frac{\partial F(X, Y, Z, T)}{\partial X} = \frac{(1+Z^2+T^2)(Z(1+Y^2)-X(1+TY))}{(1+X^2+Y^2)^{3/2}(1+Z^2+T^2)^{3/2}}.$$

- (i) The derivative of $f(Y) = F(X, Y, Z, Y)$, $0 \leq Y \leq 1$, is $f'(Y) = \frac{\partial F}{\partial Y} + \frac{\partial F}{\partial T}$. Thus the sign of $f'(Y)$ is the sign of: $Y(Z - X)^2(1 + Y^2 - ZX)$. Then $f'(0) = 0$ and $f'(Y) > 0$ if $Y > 0$.
- (ii) We set $X = \tan x$ and $Z = \tan(x + \delta)$. The derivative of $f(x) = F(X, Y, Z, Y)$, $x \in [-\frac{\delta}{2}, \frac{\pi}{4} - \delta]$, is $f'(x) = \frac{\partial F}{\partial X} \frac{dX}{dx} + \frac{\partial F}{\partial Z} \frac{dZ}{dx}$, with $\frac{dX}{dx} = 1 + X^2$ and $\frac{dZ}{dx} = 1 + Z^2$. The sign of $f'(x)$ is the sign of: $-(1 + Y^2)(Z - X)^2 Y^2 (X + Z)$. If $Y = 0$, then $f'(x) = 0$. Otherwise, if $Y > 0$, the sign of $f'(x)$ is the sign of $-(X + Z)$ ($Z \neq X$). Since $-(X + Z)$ is a decreasing function of x (derivative: $-2 - X^2 - Z^2 < 0$) that takes the value 0 for $x = -\frac{\delta}{2}$, then $f'(-\frac{\delta}{2}) = 0$, and $f'(x) < 0$ if $x > -\frac{\delta}{2}$.
- (iii) For $f(X) = F(-X, 1, X, 1)$, $0 \leq X \leq 1$, the sign of the derivative $f'(X) = -\frac{\partial F}{\partial X} + \frac{\partial F}{\partial Z}$ is the sign of $-8X(2 + X^2)$.
- (iv) The sign of the derivative of $f(X) = F(X, 1, Z, 1)$, $-1 \leq X \leq Z$, is the sign of $(1 + Y^2)(Z - X)$.
- (v) The sign of the derivative of $f(Z) = F(X, Y, Z, Y)$, $X \leq Z \leq 1$, is the sign of $(1 + Y^2)(X - Z)$.
- (vi) The sign of the derivative of $f(T) = F(X, Y, Z, T)$, $0 \leq T \leq Y$, is the sign of $Y(1 + Z^2) - T(1 + ZX)$. Then $f'(T) > 0$ if $T < Y \frac{1+Z^2}{1+ZX}$, whereas $Y < Y \frac{1+Z^2}{1+ZX}$.
- (vii) The sign of the derivative of $f(Y) = F(X, Y, Z, T)$, $0 \leq Y \leq T$, is the sign of $T(1 + X^2) - Y(1 + ZX)$. For $Y^* = \frac{T(1+X^2)}{1+ZX} \in (0, T]$, $f'(Y) > 0$ if $Y < Y^*$, $f'(Y^*) = 0$, and $f'(Y) < 0$ if $Y > Y^*$.
- (viii) We set $X = \tan x$, $Z = \tan(x + \delta)$, $Y^* = Y^*(X, Z, T)$, $f(x) = F(X, Y^*, Z, T)$, $x \in [0, \frac{\pi}{4} - \delta]$.

Then

$$f'(x) = \frac{\partial F}{\partial X}(X, Y^*, Z, T)(1 + X^2) + \frac{\partial F}{\partial Y}(X, Y^*, Z, T)\frac{dY^*}{dx} + \frac{\partial F}{\partial Z}(X, Y^*, Z, T)(1 + Z^2).$$

By construction of Y^* , $\frac{\partial F}{\partial Y}(X, Y^*, Z, T) = 0$. Then, the sign of $(1 + ZX)^3 f'(x)$ is the sign of

$$-(1 + X^2)(Z - X)^2 Z T^2 [(1 + ZX)^2 + T^2(1 + X^2)] < 0.$$

(ix) The sign of the derivative of $f(X) = F(X, Y, Z, T)$, $0 \leq X \leq 1$, is the sign of $Z(1 + Y^2) - X(1 + TY) \leq 0$.

(x) The sign of the derivative of $f(Z) = F(X, Y, Z, T)$, $-1 \leq Z \leq 0$, is the sign of $X(1 + T^2) - Z(1 + YT) \geq 0$. \square

4.2. Proof of Lemmas 3.3–3.7.

Proof of Lemma 3.3. We solve: $\max\{F(X, 1, Z, 1), X \neq Z \in \mathbb{T}_N\}$. Let $X \neq Z \in \mathbb{T}_N$.

Case 1: N is even

Case 1.a: $XZ < 0$. Combination of (4.3) and Lemma 4.1,(iv) shows:

$$F(X, 1, Z, 1) = F(-|X|, 1, |Z|, 1) < F(0, 1, |Z|, 1), \quad 0 \neq |Z| \in \mathbb{T}_N.$$

Case 1.b: $XZ \geq 0$. By (4.1)-(4.2),

$$F(X, 1, Z, 1) = F(|X|, 1, |Z|, 1) = F(|Z|, 1, |X|, 1) = F(X', 1, Z', 1),$$

with $0 \leq X' = \min(|X|, |Z|) < Z' = \max(|X|, |Z|)$; $X', Z' \in \mathbb{T}_N$.

Write $X' = \tan x$, $Z' = \tan(x + \delta)$, with $0 \leq x < z = x + \delta \leq \frac{\pi}{4}$, and $\delta \geq \frac{\pi}{2N}$. By Lemma 4.1, (ii) and (v),

$$F(X, 1, Z, 1) = F(X', 1, Z', 1) \leq F(0, 1, \tan \delta, 1) \leq F(0, 1, \tan \frac{\pi}{2N}, 1),$$

with equalities if, and only if, $x = 0$ and $\delta = \frac{\pi}{2N}$, *i.e.* $\{|X|, |Z|\} = \{X', Z'\} = \{0, \tan \frac{\pi}{2N}\}$. This is the maximum.

Case 2: N is odd.

Case 2.a: $XZ \geq 0$. As Case 1.b,

$$F(X, 1, Z, 1) = F(\tan x, 1, \tan(x + \delta), 1), \quad 0 \leq x < z = x + \delta \leq \frac{\pi}{4}, \delta \geq \frac{\pi}{2N}.$$

Then $\tan \frac{\delta}{2} \geq \tan \frac{\pi}{4N}$ and Lemma 4.1,(ii)-(iii), shows that

$$F(X, 1, Z, 1) < F(-\tan \frac{\delta}{2}, 1, \tan \frac{\delta}{2}, 1) \leq F(-\tan \frac{\pi}{4N}, 1, \tan \frac{\pi}{4N}, 1).$$

Case 2.b: $XZ < 0$. Lemma 4.1,(iv)-(v), combined with (4.3), shows:

$$F(X, 1, Z, 1) = F(-|X|, 1, |Z|, 1) \leq F(-\tan \frac{\pi}{4N}, 1, |Z|, 1) \leq F(-\tan \frac{\pi}{4N}, 1, \tan \frac{\pi}{4N}, 1),$$

with equality if, and only if, $|X| = |Z| = \tan \frac{\pi}{4N}$. This is the maximum. \square

Proof of Lemma 3.4. We maximize $F(X, Y, Z, T)$, $(X, Y) \neq (Z, T) \in \mathbb{T}_N^2$, and, $X = Z$ or $Y = T$. Let (X, Y, Z, T) be an optimal solution. Equivalently, $\{u = \rho(1, X, Y), v = \rho(1, Z, T)\}$ is minimal for (3.2). We prove that there exists $Q \in \{I, Q_1, Q_2, Q_3\}$ such that $\{Qu, Qv\}$ is minimal for (3.1); this implies $\theta(u, v) = \theta(Qu, Qv) = m^*$, because $Q \in \mathcal{O}$.

Case 1: $X \neq Z$ and $Y = T \geq 0$. Combination of (4.1) and Lemma 4.1,(i) shows that

$$F(X, Y, Z, T) = F(Z, T, X, Y) \leq F(X, 1, Z, 1) = F(Z, 1, X, 1),$$

with equality if, and only if, $Y = 1$. By optimality of (X, Y, Z, T) , we obtain $Y = 1$. Then $\{u, v\} \subset (I) \cap (V)$ is optimal for (3.1), on the edge $(I) \cap (V)$.

Case 2: $X \neq Z$ and $Y = T < 0$. The invariance (4.2) shows that $F(X, -Y, Z, -T) = F(X, Y, Z, T)$, where $(X, -Y) \neq (Z, -T) \in \mathbb{T}_N^2$ with $-Y = -T \geq 0$. Then $(X, -Y, Z, -T)$ is optimal. We deduce from Case 1 that $\{Q_3u, Q_3v\} = \{\rho(1, X, -Y), \rho(1, Z, -T)\}$ is optimal for (3.1).

Case 3: $X = Z \geq 0$ and $Y \neq T$. The invariance (4.1) shows $F(X, Y, Z, T) = F(Y, X, T, Z)$, where $(Y, X) \neq (T, Z) \in \mathbb{T}_N^2$ with $X = Z \geq 0$. Then (Y, X, T, Z) is optimal. By Case 1, $\{Q_1u, Q_1v\} = \{\rho(1, Y, X), \rho(1, T, Z)\}$ is optimal for (3.1).

Case 4: $X = Z < 0$ and $Y \neq T$. By invariance (4.1)-(4.2), $F(X, Y, Z, T) = F(-Y, -X, -T, -Z)$, where $(-Y, -X) \neq (-T, -Z) \in \mathbb{T}_N^2$ with $-X = -Z \geq 0$. Then $(-Y, -X, -T, -Z)$ is optimal. By Case 1, $\{Q_2u, Q_2v\} = \{\rho(1, -Y, -X), \rho(1, -T, -Z)\}$ is optimal for (3.1).

Conversely, for all $Q \in \{I, Q_1, Q_2, Q_3\}$ and $\{u^*, v^*\}$ minimal for (3.1), $\{u, v\} = \{Q^\top u^*, Q^\top v^*\}$ is realizable for (3.2), and $\theta(u, v) = \theta(Q^\top u^*, Q^\top v^*) = m^*$ is the minimal value of (3.2), so $\{u, v\}$ is minimal for (3.2). \square

Proof of Lemma 3.5. Equivalently, we maximize $F(X, Y, Z, T)$ on the first panel, with $(X, Y) \neq (Z, T) \in \mathbb{T}_N^2$. Let $X \neq Z$ and $Y \neq T$ in \mathbb{T}_N . We prove that $F(X, Y, Z, T)$ is strictly dominated by a value on the grid lines: $F(X', Y', Z', T')$, with $(X', Y') \neq (Z', T') \in \mathbb{T}_N^2$ and, $X' = Z'$ or $Y' = T'$.

Case 1: $XZ \geq 0$ and $YT \geq 0$.

Case 1.a: $0 \leq X < Z \leq 1$ and $0 \leq T < Y \leq 1$. By Lemma 4.1,(vi), $F(X, Y, Z, T) < F(X, Y, Z, Y)$.

Case 1.b: $0 \leq X < Z \leq 1$ and $0 \leq Y < T \leq 1$. According to Lemma 4.1,(vii)-(viii),

$$F(X, Y, Z, T) \leq F(X, Y^*(X, Z, T), Z, T) \leq F(0, Y^*(0, \tan \delta, T), \tan \delta, T), \quad \delta = \arctan Z - \arctan X,$$

with equalities if, and only if, $Y = Y^*(X, Z, T)$ and $X = 0$. If $X \neq 0$, the last inequality is strict. Otherwise, $Y^*(X, Z, T) = Y^*(0, \tan \delta, T) = T > Y$, and the first inequality is strict. In both cases:

$$F(X, Y, Z, T) < F(0, T, \tan \delta, T), \quad \tan \delta \geq \tan \frac{\pi}{2N}.$$

Application of Lemma 4.1,(v) and (ii), shows

$$F(0, T, \tan \delta, T) \leq F(0, T, \tan \frac{\pi}{2N}, T) \leq F(-\tan \frac{\pi}{4N}, T, \tan \frac{\pi}{4N}, T).$$

If N is even, resp. odd, we conclude with the first, resp. second, inequality.

Case 1.c: $0 \leq Z < X \leq 1$. By relation (4.1), $F(X, Y, Z, T) = F(Z, T, X, Y)$. The right member is dominated using Case 1.a or Case 1.b.

Case 2: $XZ < 0$ or $YT < 0$.

Case 2.a: $X = -Z$ or $Y = -T$. Similarly to (4.3)-(4.4),

$$F(X, Y, Z, T) < \begin{cases} F(|X|, Y, |Z|, T) = F(|X|, Y, |X|, T), & \text{if } X = -Z, \\ F(X, |Y|, Z, |T|) = F(X, |Y|, Z, |Y|), & \text{if } Y = -T. \end{cases}$$

Case 2.b: $X \neq -Z$ and $Y \neq -T$. By (4.5):

$$F(X, Y, Z, T) \leq F(|X|, |Y|, |Z|, |T|).$$

The right member is dominated using Case 1. \square

Proof of Lemma 3.6. Let $u \in (I) \setminus (V)$ and $v \in (V) \setminus (I)$: $u = \rho(1, X, Y)$, and $v = \rho(Z, T, 1)$, with $(X, Y), (Z, T) \in \mathbb{T}_N^2$, such that $Y, Z < 1$. We prove that $u \cdot v < \cos m^*$.

Case 1: $(T, Z) \neq (X, Y)$, as in the middle of Figure 4. Let

$$Q = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \in \mathcal{O}, \quad w = Qv = \rho(1, T, Z) \in (I).$$

It is clear that $0 < (1 - Z)(1 - Y)$, so $Z + XT + Y < 1 + XT + YZ$, i.e. $u \cdot v < F(X, Y, T, Z) = u \cdot w$. By the way, $u \neq w \in (I)$ so $u \cdot w \leq \cos m^*$.

Case 2: $(T, Z) = (X, Y)$, as in the right of Figure 4. We prove that $u \cdot v < F(X, Y, X, 1)$ using the auxiliary function

$$f(t) = \rho(1, X, Y) \cdot \rho(t, X, 1), \quad t \in [Y, 1];$$

$f(t)$ represents the cos-angle between $\rho(1, X, Y)$ and $\rho(t, X, 1)$. The sign of $f'(t)$ is the sign of $(1 + X^2 + t^2) - (X^2 + Y + t)t = X^2(1 - t) + 1 - Yt > 0$. Thus f is increasing and $f(Y) < f(1)$, i.e. $u \cdot v < F(X, Y, X, 1)$. Due to $\rho(1, X, Y) \neq \rho(1, X, 1) \in (I)$, $u \cdot v < F(X, Y, X, 1) \leq \cos m^*$. \square

Proof of Lemma 3.7. Since $v \in (III) \Leftrightarrow -v \in (I)$, we solve $m := \min\{\theta(u, -w); u \neq w \in (I)\}$, or, equivalently, $\cos m = \max\{-u \cdot w; u \neq w \in (I)\} = -\min\{F(X, Y, Z, T); (X, Y) \neq (Z, T) \in \mathbb{T}_N^2\}$. Fix $X, Y, Z, T \in \mathbb{T}_N$. We apply: (4.5), Lemma 4.1,(ix-x), (4.1), and Lemma 4.1,(ix-x) again:

$$\begin{aligned} F(X, Y, Z, T) &\geq F(|X|, |Y|, -|Z|, -|T|) \\ &\geq F(1, |Y|, -|Z|, -|T|) \\ &\geq F(1, |Y|, -1, -|T|) = F(|Y|, 1, -|T|, -1) \\ &\geq F(1, 1, -|T|, -1) \\ &\geq F(1, 1, -1, -1). \end{aligned}$$

Thus the minimal value of F is reached on a diagonal:

$$F(1, 1, -1, -1) = \rho(1, 1, 1) \cdot \rho(1, -1, -1).$$

Its opposite value represents the length of an edge:

$$\cos m = -F(1, 1, -1, -1) = \rho(1, 1, 1) \cdot \rho(1, -1, 1).$$

By Lemma 3.3, $m > m^*$ if $N \geq 2$, and $m = m^*$ if $N = 1$. □

5. CONCLUSION AND PERSPECTIVES

In this paper we have identified the symmetry group of the equiangular cubed sphere: it is the symmetry group of the cube. Moreover, we have studied the geodesic distance between points of the grid. Our results provide some theoretical foundation for upcoming numerical computation on the cubed sphere.

The symmetry group of a grid plays a central role in several contexts. It can be used to build spherical quadrature rules which are valid for as many spherical harmonics as possible [28]. Our main result shows that the group of the cube is the suitable symmetry group for the determination of quadrature weights on the cubed sphere. This background supports a quadrature rule of ongoing works [2]. Moreover, for quadrature rules, the geometric distribution of the nodes is often examined. Our study of the geodesic distance could serve as a tool to quantify the ‘‘uniformity’’ of the cubed sphere grid.

Another subject of interest concerns the building of a discrete Fourier analysis on the cubed sphere, based on the invariance under the action of the symmetry group, in the spirit of [15]. Here again, our result is a first step in this direction, since it identifies the group to be considered.

APPENDIX A. EQUIANGULAR CUBED SPHERE

In this section we recall the definition of the equiangular cubed sphere. Let $N \geq 1$ be a fixed integer, and the one-dimensional grid

$$\mathbb{T}_N = \left\{ \tan \frac{k\pi}{2N}, -\frac{N}{2} \leq k \leq \frac{N}{2} \right\},$$

represented in Figure 6. The index k , resp. the angle $\frac{k\pi}{2N}$, takes the $N + 1$ equidistributed values from $-\frac{N}{2}$ to $\frac{N}{2}$, resp. from $-\frac{\pi}{4}$ to $\frac{\pi}{4}$. Then we define a cartesian grid on the faces of the cube $[-1, 1]^3$, based on the two-dimensional grid $\mathbb{T}_N \times \mathbb{T}_N$, as in Figure 1,

$$\mathbb{C}_N = \{(1, u, v), (-1, u, v), (u, 1, v), (u, -1, v), (u, v, 1), (u, v, -1); u, v \in \mathbb{T}_N\}.$$

To finish with, we project \mathbb{C}_N on the unit sphere $\mathbb{S}^2 = \{x \in \mathbb{R}^3 : \|x\| = 1\}$.

Definition A.1. The equiangular cubed sphere grid \mathbb{CS}_N is the radial projection of the cartesian grid \mathbb{C}_N on the sphere: $\mathbb{CS}_N = \rho(\mathbb{C}_N)$, with $\rho : x \in \mathbb{R}^3 \setminus \{0\} \mapsto \frac{x}{\|x\|} \in \mathbb{S}^2$. Equivalently,

$$\mathbb{CS}_N := \left\{ \frac{1}{r}(\pm 1, u, v), \frac{1}{r}(u, \pm 1, v), \frac{1}{r}(u, v, \pm 1); u, v \in \mathbb{T}_N, r = \sqrt{1 + u^2 + v^2} \right\}. \quad (\text{A.1})$$

The radial projection ρ can be checked to be injective on the faces of the cube $[-1, 1]^3$. Therefore, there is a one-to-one correspondence between \mathbb{CS}_N and \mathbb{C}_N ; $(u_1, u_2, u_3) \in \mathbb{C}_N$ are the so-called *gnomonic* coordinates of the point $\rho(u_1, u_2, u_3) \in \mathbb{CS}_N$. By construction, the geometry of the cubed




FIGURE 6. Left: construction of the equiangular grid $T_N = \{\tan \frac{k\pi}{2N}, -\frac{N}{2} \leq k \leq \frac{N}{2}\}$ ($N = 6$). On the unit circle $x_1^2 + x_2^2 = 1$, we consider $N + 1$ equidistributed angles: $\frac{k\pi}{2N}, -\frac{N}{2} \leq k \leq \frac{N}{2}$. The origin and the point $(1, \tan \frac{k\pi}{2N}) \in \{1\} \times T_N$ (black dot on the dotted line) are aligned with $(\cos \frac{k\pi}{2N}, \sin \frac{k\pi}{2N})$ (black dot on the arc of circle). Right: construction of a panel of the cubed sphere by intersection of two equiangular families of great circles ($N = 5$).

sphere is shaped by the geometry of the cube. In particular, we define twelve *edges*, as the projection of the edges of the cube, and six *panels*, as the projection of the grid of the faces:

$$\begin{aligned} \text{CS}_N &= \rho(C_N) = (I) \cup (II) \cup (III) \cup (IV) \cup (V) \cup (VI), \\ (I) &= \rho(\{1\} \times T_N \times T_N), \quad (II) = \rho(T_N \times \{1\} \times T_N), \quad (III) = \rho(\{-1\} \times T_N \times T_N), \\ (IV) &= \rho(T_N \times \{-1\} \times T_N), \quad (V) = \rho(T_N \times T_N \times \{1\}), \quad (VI) = \rho(T_N \times T_N \times \{-1\}). \end{aligned}$$

Two panels are said to be *adjacent*, resp. *opposite*, if they are the projection of adjacent, resp. opposite, faces.

As in Figures 1 and 6, the cubed sphere meshes the sphere with equiangular arcs of great circles: the radial projection maps the cartesian straight lines of the cube onto arcs of great circles. For instance, in panel (I),

$$\begin{aligned} \rho(\{(1, \tan \frac{k\pi}{2N}, z), |z| \leq 1\}) &\subset \{u \in \mathbb{S}^2 : u_1 \sin \frac{k\pi}{2N} - u_2 \cos \frac{k\pi}{2N} = 0\}, \quad |k| \leq \frac{N}{2}; \\ \rho(\{(1, y, \tan \frac{j\pi}{2N}), |y| \leq 1\}) &\subset \{u \in \mathbb{S}^2 : u_1 \sin \frac{j\pi}{2N} - u_3 \cos \frac{j\pi}{2N} = 0\}, \quad |j| \leq \frac{N}{2}. \end{aligned}$$

The panel (I) is obtained by intersection of two equiangular families of great circles:

$$\rho(1, \tan \frac{k\pi}{2N}, \tan \frac{j\pi}{2N}) \in \{u \in \mathbb{S}^2 : u_1 \sin \frac{k\pi}{2N} - u_2 \cos \frac{k\pi}{2N} = 0\} \cap \{u \in \mathbb{S}^2 : u_1 \sin \frac{j\pi}{2N} - u_3 \cos \frac{j\pi}{2N} = 0\};$$

the longitude $\frac{k\pi}{2N}$ and the latitude $\frac{j\pi}{2N}$ both scan a uniform grid on $[-\frac{\pi}{4}, \frac{\pi}{4}]$. Analogous properties can be derived for the other panels.

APPENDIX B. SYMMETRY GROUP OF THE CUBE

In this section we recall properties about the symmetry group of the cube. We give short direct proofs for completeness.

Theorem B.1. *Let $e_1 = (1, 0, 0)$, $e_2 = (0, 1, 0)$ and $e_3 = (0, 0, 1)$. The symmetry group \mathcal{O} of the cube $\{-1, 1\}^3$ is:*

$$\mathcal{O} = \{[\epsilon_1 e_{\sigma_1} \quad \epsilon_2 e_{\sigma_2} \quad \epsilon_3 e_{\sigma_3}], \sigma \in \mathfrak{S}_3, \epsilon \in \{-1, 1\}^3\}. \quad (\text{B.1})$$

Proof. \subset : Fix an orthogonal matrix Q such that $C_1 = \{-1, 1\}^3$ is left invariant by Q . We remark that for $x_1, x_2 \in C_1$, $\|x_1 - x_2\| = 2 \Leftrightarrow \exists \epsilon \in \{-1, 1\}, \exists 1 \leq \sigma \leq 3, x_1 - x_2 = 2\epsilon e_\sigma$ (x_1 and x_2 are the vertices of an edge); this is due to the following identity: $\|x_1 - x_2\|^2 = \sum_{j=1}^3 |e_j \cdot x_1 - e_j \cdot x_2|^2$, with $|e_j \cdot x_1 - e_j \cdot x_2| \in \{0, 2\}$. Then, the matrix Q maps $x_0 = (1, 1, 1) \in C_1$ onto $Qx_0 \in C_1$. For $1 \leq i \leq 3$, the distance between $x_0, x_0 - 2e_i \in C_1$ is 2. Then the distance between $Qx_0, Qx_0 - 2Qe_i \in C_1$ is also 2, i.e. there are $\sigma_i \in \{1, 2, 3\}$ and $\epsilon_i \in \{-1, 1\}$ such that $Qe_i = \epsilon_i e_{\sigma_i}$. To finish with, the indices $\sigma_1, \sigma_2, \sigma_3$ define a permutation; otherwise Q would contain at least two collinear columns.

\supset : Conversely, let $Q = [\epsilon_1 e_{\sigma_1} \ \epsilon_2 e_{\sigma_2} \ \epsilon_3 e_{\sigma_3}]$, with $\sigma \in \mathfrak{S}_3$ and $\epsilon \in C_1$. It is clear that $Q^\top Q = I_3$. Let $x \in C_1$. Then Qx is a permutation of $[\epsilon_1 x_1 \ \epsilon_2 x_2 \ \epsilon_3 x_3]^\top \in C_1$:

$$Qx = [e_{\sigma_1} \ e_{\sigma_2} \ e_{\sigma_3}] [\epsilon_1 x_1 \ \epsilon_2 x_2 \ \epsilon_3 x_3]^\top, \quad (\text{B.2})$$

so $Qx \in C_1$. Then $QC_1 \subset C_1$. But Q is bijective; so the finite sets QC_1 and C_1 have the same cardinal number, and $QC_1 = C_1$. \square

Remark B.2. The group \mathcal{O} contains $3! \cdot 2^3 = 48$ symmetries, determined by a permutation of \mathfrak{S}_3 and three signs.

From a geometrical point of view, any symmetry of the cube permutes the centers of the faces. The result can also be formulated as follows.

Corollary B.3. *The symmetry group \mathcal{O} of the cube $\{-1, 1\}^3$ coincides with the symmetry group of the octahedron $\{(1, 0, 0), (0, 1, 0), (0, 0, 1), (-1, 0, 0), (0, -1, 0), (0, 0, -1)\}$.*

Definition B.4. The centers of the edges of the cube $[-1, 1]^3$ define the cuboctahedron

$$\Omega := \{(0, \epsilon, \eta), (\epsilon, 0, \eta), (\epsilon, \eta, 0), \epsilon = \pm 1, \eta = \pm 1\}. \quad (\text{B.3})$$

Theorem B.5. *The symmetry group of the cuboctahedron Ω coincides with the symmetry group \mathcal{O} of the cube $\{-1, 1\}^3$.*

Proof. \subset : Let Q be in the symmetry group of Ω : Q is orthogonal and leaves Ω invariant. Then Q maps $(1, 0, 1)$ onto $x = (x_1, x_2, x_3) \in \Omega$; let j be such that $x_j = 0$, and $x_i = \pm 1, i \neq j$. The vector $(1, 0, 1)$ and $(1, 0, -1)$ are orthogonal, so Q maps $(1, 0, -1)$ onto $\bar{x} = (\bar{x}_1, \bar{x}_2, \bar{x}_3) \in \Omega$ such that $x \cdot \bar{x} = 0$. Thus $\bar{x}_j = 0$, and there are $i \neq j$, such that $\bar{x}_i = x_i$, and $\bar{x}_k = -x_k$, for $k \neq i, j$. Then, the vector $e_1 = \frac{1}{2}((1, 0, 1) + (1, 0, -1))$ is mapped onto $Qe_1 = \frac{1}{2}(x + \bar{x}) = x_i e_i$. This shows that Q maps the center of face e_1 onto the center of face $Qe_1 = \epsilon_1 e_{\sigma_1}$, with $\epsilon_1 = x_i \in \{-1, 1\}$, and $\sigma_1 = i \in \{1, 2, 3\}$. A similar procedure shows that $Qe_2 = \epsilon_2 e_{\sigma_2}$, $Qe_3 = \epsilon_3 e_{\sigma_3}$, with $\epsilon_2, \epsilon_3 \in \{-1, 1\}$, and $\sigma_2, \sigma_3 \in \{1, 2, 3\}$. To finish with, the vectors Qe_i are orthogonal so $(\sigma_1, \sigma_2, \sigma_3) \in \mathfrak{S}_3$.

\supset : Let $Q \in \mathcal{O}$: $Q = [\epsilon_1 e_{\sigma_1} \ \epsilon_2 e_{\sigma_2} \ \epsilon_3 e_{\sigma_3}]$, with $\sigma \in \mathfrak{S}_3$, $\epsilon \in C_1$. Fix $x \in \Omega$: $x = \eta_1 e_{i_1} + \eta_2 e_{i_2}$, with $\eta_1, \eta_2 = \pm 1, 1 \leq i_1 \neq i_2 \leq 3$. It is clear that $Qx = \eta_1 \epsilon_{i_1} e_{\sigma_{i_1}} + \eta_2 \epsilon_{i_2} e_{\sigma_{i_2}} \in \Omega$. Then $Q\Omega \subset \Omega$. \square

APPENDIX C. PROOF OF THE MAIN THEOREM USING GROUP THEORY

In this section, we prove shortly Theorem 1.2 using classification of subgroups of orthogonal groups. We refer to [1, Theorems 11.1, 19.1 and 19.2] for more details.

Theorem C.1 (Lagrange's Theorem). *The order of a subgroup of a finite group is always a divisor of the order of the group.*

Theorem C.2. *A finite subgroup of $O_2 = \{Q \in \mathbb{R}^{2 \times 2} : Q^\top Q = I_2\}$ is either cyclic or dihedral.*

Theorem C.3. *A finite subgroup of $SO_3 = \{Q \in \mathbb{R}^{3 \times 3} : Q^\top Q = I_3, \det Q = 1\}$ is isomorphic either to a cyclic group, a dihedral group, or the rotation group of a Platonic solid (tetrahedron, cube/octahedron, dodecahedron/icosahedron).*

Proof of Theorem 1.2 using group theory. Lemma 2.1 proves $\mathcal{O} \subset \mathcal{G}_N$. We deduce $\mathcal{O} = \mathcal{G}_N$ from Theorem C.3.

Firstly, we focus on the rotation subgroups, $\mathcal{O}^+ = \mathcal{O} \cap SO_3$ and $\mathcal{G}_N^+ = \mathcal{G}_N \cap SO_3$. Indeed, $\mathcal{O} = \mathcal{G}_N \Leftrightarrow \mathcal{O}^+ = \mathcal{G}_N^+$, because $-I_3 \in \mathcal{O} \subset \mathcal{G}_N$, so $\mathcal{O} = \mathcal{O}^+ \cup \{-Q, Q \in \mathcal{O}^+\}$, and $\mathcal{G}_N = \mathcal{G}_N^+ \cup \{-Q, Q \in \mathcal{G}_N^+\}$. Secondly, \mathcal{G}_N^+ is a finite subgroup of SO_3 , by injection into a finite permutation group. Indeed, $\text{CS}_N = \{x_j, 1 \leq j \leq M\}$ is a finite set which contains three linearly independent vectors. So $\sigma : Q \in \mathcal{G}_N^+ \mapsto \sigma[Q] \in \mathfrak{S}_M$ such that $Qx_j = x_{\sigma[Q](j)}, 1 \leq j \leq M$, is injective.

By Theorem C.3, \mathcal{G}_N^+ is isomorphic either to a cyclic group, a dihedral group, or the rotation group of a Platonic solid. Since $\mathcal{O}^+ \subset \mathcal{G}_N^+$, we eliminate all the candidates, except the rotation group of the cube. By Lagrange's Theorem, the order of \mathcal{O}^+ (24) is a divisor of the order of \mathcal{G}_N^+ . Then \mathcal{G}_N^+ is not isomorphic to the group of the tetrahedron (order 12), neither the group of the dodecahedron/icosahedron (order 60). By the way \mathcal{G}_N^+ is not cyclic, because \mathcal{O}^+ is not commutative. And to finish with, \mathcal{G}_N^+ is not

dihedral. Indeed, a dihedral group is isomorphic to a finite subgroup of O_2 [11, p. 152]. So, by Theorem C.2, the subgroups of a dihedral group are either cyclic or dihedral. But the rotation group O^+ of the cube is neither cyclic nor dihedral, by Theorem C.3. \square

Remark C.4. By Remark 2.2, the proof is still valid if T_N is replaced by any symmetric grid. Therefore, the group of any cubed sphere (with T_N symmetric) coincides with the group of the cube.

ACKNOWLEDGEMENTS

The author gratefully acknowledges the many helpful suggestions of J.-P. Croisille during the preparation of the paper. The author wishes to express his thanks to A. Pasquale for several helpful comments concerning group theory.

REFERENCES

- [1] M. Armstrong. *Groups and symmetry*. Springer, 1988.
- [2] J.-B. Bellet, M. Brachet, and J.-P. Croisille. Quadrature and symmetry on the Cubed Sphere. *Working paper <https://hal.archives-ouvertes.fr/hal-03223150>*, 2021.
- [3] M. Brachet. *Schémas compacts hermitiens sur la Sphère: applications en climatologie et océanographie numérique*. PhD thesis, 2018 (in French).
- [4] M. Brachet and J.-P. Croisille. Spherical Shallow Water simulation by a cubed sphere finite difference solver. *Quarterly Journal of the Royal Meteorological Society*, 147(735):786–800, 2021.
- [5] C. Chen and F. Xiao. Shallow water model on cubed-sphere by multi-moment finite volume method. *Journal of Computational Physics*, 227(10):5019–5044, 2008.
- [6] S. Chevrot, R. Martin, and D. Komatitsch. Optimized discrete wavelet transforms in the cubed sphere with the lifting scheme—implications for global finite-frequency tomography. *Geophysical Journal International*, 191(3):1391–1402, 2012.
- [7] M. Faham and H. Nasir. Weakly Orthogonal Spherical Harmonics in a Non-Polar Spherical Coordinates and its Application to Functions on Cubed-Sphere. *Sultan Qaboos University Journal for Science*, 17(2):200–213, 2012.
- [8] L. Ivan, H. De Sterck, S. A. Northrup, and C. P. T. Groth. Multi-dimensional finite-volume scheme for hyperbolic conservation laws on three-dimensional solution-adaptive cubed-sphere grids. *Journal of Computational Physics*, 255:205–227, 2013.
- [9] B. A. Jones, G. H. Born, and G. Beylkin. Comparison of the Cubed-Sphere Gravity Model with the Spherical Harmonics. *Journal of Guidance, Control, and Dynamics*, 33(2):415–425, 2010.
- [10] H.-G. Kang and H.-B. Cheong. An efficient implementation of a high-order filter for a cubed-sphere spectral element model. *Journal of Computational Physics*, 332:66–82, 2017.
- [11] Y. Kosmann-Schwarzbach. *Groupes et symétries : groupes finis, groupes et algèbres de Lie, représentations*. Éditions de l'École Polytechnique, 2006.
- [12] D. Lee and A. Palha. A mixed mimetic spectral element model of the rotating shallow water equations on the cubed sphere. *Journal of Computational Physics*, 375:240–262, 2018.
- [13] J. L. McGregor. Semi-Lagrangian Advection on Conformal-Cubic Grids. *Monthly Weather Review*, 124(6):1311–1322, 1996.
- [14] R. D. Nair, S. J. Thomas, and R. D. Loft. A Discontinuous Galerkin Transport Scheme on the Cubed Sphere. *Monthly Weather Review*, 133(4):814–828, 2005.
- [15] G. Peyré. *L'algèbre discrète de la transformée de Fourier*. Ellipses, 2004.
- [16] B. Portelenelle and J.-P. Croisille. An efficient quadrature rule on the Cubed Sphere. *Journal of Computational and Applied Mathematics*, 328:59–74, 2018.
- [17] R. J. Purser. Sets of optimally diversified polyhedral orientations. *Office note, National Centers for Environmental Prediction (U.S.)*, 489, 2017.
- [18] R. J. Purser. Möbius Net Cubed-Sphere Gnomonic Grids. *Office note, National Meteorological Center (U.S.)*, 496, 2018.
- [19] R. J. Purser and M. Rančić. Smooth quasi-homogeneous gridding of the sphere. *Quarterly Journal of the Royal Meteorological Society*, 124(546):637–647, 1998.
- [20] R. J. Purser and M. Tong. A minor modification of the gnomonic cubed-shaped sphere grid that offers advantages in the context of implementing moving hurricane nests. *Office note, National Centers for Environmental Prediction (U.S.)*, 486, 2017.
- [21] W. M. Putman. *Development of the finite-volume dynamical core on the cubed-sphere*. PhD thesis, 2007.
- [22] M. Rančić, R. J. Purser, and F. Mesinger. A global shallow-water model using an expanded spherical cube: Gnomonic versus conformal coordinates. *Quarterly Journal of the Royal Meteorological Society*, 122(532):959–982, 1996.
- [23] M. Rančić, R. J. Purser, D. Jović, R. Vasic, and T. Black. A Nonhydrostatic Multiscale Model on the Uniform Jacobian Cubed Sphere. *Monthly Weather Review*, 145(3):1083–1105, 2017.
- [24] E. Rodriguez Bazan and E. Hubert. Multivariate interpolation: Preserving and exploiting symmetry. *Journal of Symbolic Computation*, 107:1–22, 2021.

- [25] C. Ronchi, R. Iacono, and P. S. Paolucci. The “cubed sphere”: a new method for the solution of partial differential equations in spherical geometry. *Journal of Computational Physics*, 124(1):93–114, 1996.
- [26] J. A. Rossmanith. A wave propagation method for hyperbolic systems on the sphere. *Journal of Computational Physics*, 213(2):629–658, 2006.
- [27] R. Sadourny. Conservative finite-difference approximations of the primitive equations on quasi-uniform spherical grids. *Monthly Weather Review*, 100(2):136–144, 1972.
- [28] S. L. Sobolev. Cubature formulas on the sphere invariant under finite groups of rotations. *Dokl. Akad. Nauk SSSR*, 146:310–313, 1962.
- [29] S. J. Thomas, J. M. Dennis, H. M. Tufo, and P. F. Fischer. A Schwarz preconditioner for the cubed-sphere. *SIAM Journal on Scientific Computing*, 25(2):442–453, 2003.
- [30] P. A. Ullrich, P. H. Lauritzen, and C. Jablonowski. Some considerations for high-order ‘incremental remap’-based transport schemes: edges, reconstructions, and area integration. *International Journal for Numerical Methods in Fluids*, 71(9):1131–1151, 2013.
- [31] W. M. Washington, L. Buja, and A. Craig. The computational future for climate and earth system models: on the path to petaflop and beyond. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 367(1890):833–846, 2009.
- [32] D. L. Williamson. The evolution of dynamical cores for global atmospheric models. *Journal of the Meteorological Society of Japan. Ser. II*, 85B:241–269, 2007.

UNIVERSITÉ DE LORRAINE, CNRS, IECL, F-57000 METZ, FRANCE
Email address: `jean-baptiste.bellet@univ-lorraine.fr`