



Cross-sections and rate coefficients for rotational excitation of aluminium hydroxide by helium

Manel Naouai, Faouzi Najar, Cheikh Bop, Kamel Hammami

► To cite this version:

Manel Naouai, Faouzi Najar, Cheikh Bop, Kamel Hammami. Cross-sections and rate coefficients for rotational excitation of aluminium hydroxide by helium. Monthly Notices of the Royal Astronomical Society, 2018, 480 (4), pp.5412 - 5418. 10.1093/mnras/sty2223 . hal-03078747

HAL Id: hal-03078747

<https://hal.science/hal-03078747>

Submitted on 16 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Cross-sections and rate coefficients for rotational excitation of aluminium hydroxide by helium

Manel Naouai ¹★ Faouzi Najar,¹ Cheikh Tidiane Bop ² and Kamel Hammami¹

¹Laboratoire de Spectroscopie Atomique Moléculaire et Applications, Département de Physique, Faculté des Sciences de Tunis, Campus Universitaire Farhat Hached, 2092 Tunis El Manar, Tunisia

²Laboratoire Atomes Lasers, Département de Physique, Faculté des Sciences et Techniques, Université Cheikh Anta Diop, Dakar 5005, Senegal

Accepted 2018 August 13. Received 2018 June 30; in original form 2018 February 15

ABSTRACT

Aluminium hydroxide (AlOH) is ubiquitous in the interstellar medium. However, there are no reliable data available in the literature that can be used to interpret the observations. Therefore, in this paper, we deal with the rotational rate coefficients of AlOH induced by collision with helium. The potential energy surface of the $\text{AlOH}-\text{He}$ complex was computed within the coupled cluster *ab initio* method with single, double and perturbative triple excitation, in conjunction with the augmented-correlation consistent-polarized valence quadruple zeta Gaussian basis set. Using this potential energy surface, we computed rotational cross-sections for total energies ranging up to 1500 cm^{-1} with the close-coupling approach. By thermally averaging these cross-sections over the kinetic temperature range $5\text{--}300 \text{ K}$, we derived the AlOH rate coefficients corresponding to the first 13 rotational levels. Propensity rules have been found in favour of $|\Delta j| = 1$. The data calculated in this paper have been compared with previous results and significant differences are observed. Such a finding might greatly affect the calculation of the AlOH abundance. In addition, this might encourage experimental work to settle the matter.

Key words: molecular data – molecular processes – scattering – ISM: abundances – ISM: molecules.

1 INTRODUCTION

A large number of aluminium-containing species has been detected in the interstellar medium (ISM). Indeed, aluminium isocyanide (AlCN ; Ziurys et al. 2002), aluminium fluoride (AlF) and aluminium chloride (AlCl) were discovered toward the asymptotic giant branch star IRC+10216 in the envelopes of carbon-rich evolved stars (Ziurys 2006).

Because of the high abundance of its elemental form, aluminium is thought to be the main precursor of AlOH through the single replacement reaction (Morton et al. 1973)



This chemical equation has been the subject of several investigations, theoretical as well as experimental (Hauge, Kauffman & Margrave 1980; McClean, Nelson & Campbell 1993; Álvarez-Barcia & Flores 2011, 2016). These studies provide some additional information: the chemical reaction in equation (1) is favourable in the gas phase.

In radio astronomy, AlOH is considered to be one of the most important aluminium-bearing species because of its large electric

dipole moment (Li et al. 2003). Recently, along with several other Al-bearing molecules, AlOH was identified towards the circumstellar gas of Mira (o Ceti) by Kamiński et al. (2016). Using the Atacama Pathfinder Experiment, they found three rotational transitions of AlOH : $13 \rightarrow 12$, $15 \rightarrow 14$ and $35 \rightarrow 34$. In addition, this molecule was observed towards the envelope of the oxygen-rich star, VY Canis Majoris (VY CMa), using the Arizona Radio Observatory (ARO; Tenenbaum & Ziurys 2010). More precisely, three rotational emission lines were detected: $9 \rightarrow 8$, $7 \rightarrow 6$ and $5 \rightarrow 4$. It is worth noting that AlO , which is the second aluminium reservoir in VY CMa with a fractional abundance ~ 17 times smaller than that of AlOH , was also observed with the ARO (Tenenbaum & Ziurys 2009). Therefore, AlOH plays an important role in the chemistry of aluminium in the super-giant VY CMa star, making it crucial to determine its exact abundance in this medium. In fact, its abundance has been estimated at $\sim 1 \times 10^{-7}$ based on radiative transfer analysis. In addition, assuming the local thermodynamic equilibrium (LTE) model of Tsuji (1973), Tenenbaum & Ziurys (2010) predicted an abundance of $\sim 2 \times 10^{-6}$.

To satisfy the astrophysical precision when modelling the interstellar molecular clouds, radiative transitions and rate coefficients induced by the most abundant collision partners (such as He and H_2) are necessary. For AlOH , rate coefficients induced by collision with

* E-mail: manelnaouai6@gmail.com




Figure 1. Jacobi coordinate system (R, θ) of AlOH–He.




Figure 2. Contour plot of the interaction energy (in cm^{-1}). Negative contour lines (blue) are equally spaced by 2 cm^{-1} and positive contour lines (red) are equally spaced by 50 cm^{-1} . The zero energy contour line is black.

He have been reported by Tchakoua et al. (2015). However, these data have not been without criticism. In terms of magnitude rank, the rate coefficients of the AlOH–He system computed by Tchakoua et al. (2015) are much larger than those of similar systems such as HCP–He and AlF–He (for further discussion, see Section 4). In this regime, it is necessary to recompute the collisional rates of AlOH due to He impact, in order to provide reliable astrophysical data.

Although the AlOH emission lines observed in the ISM are rotational, theoretical calculations cannot be performed regardless of the vibrational modes. For this purpose, the AlOH vibrational frequencies should be determined first. Indeed, these have been the target of various experimental and theoretical studies. For example, Pilgrim, Robbins & Duncan (1993) measured the Al–OH stretch frequency ($\nu_{\text{Al-OH}} = 895 \text{ cm}^{-1}$) and Hauge et al. (1980) measured in a noble gas matrix the two stretch frequency modes ($\nu_{\text{Al-OH}} =$




Figure 3. Radial coefficients V_λ ($\lambda = 0, \dots, 4$) of the interaction energy as a function of the intermolecular distance R .

810 cm^{-1} and $\nu_{\text{AlO-H}} = 3790 \text{ cm}^{-1}$). These measurements have been theoretically ($\nu_{\text{Al-OH}} = 884 \text{ cm}^{-1}$ and $\nu_{\text{AlO-H}} = 4313 \text{ cm}^{-1}$) confirmed by Vacek, Deleeuw & Schaefer (1993).

This paper is organized as follows. In Section 2, we describe the new potential energy surface (PES) of AlOH–He as well as the computational details. In Section 3, we outline the inelastic cross-sections and rate coefficients of the AlOH–He complex. In Section 4, we present a comparative study between the present work and the previous work. Finally, we give concluding remarks in Section 5.

2 POTENTIAL ENERGY SURFACE

The studied system is made up of a linear molecule (AlOH) and a structureless atom (He). In the rigid rotator approximation, the Al–O and O–H bond lengths are frozen into their experimental equilibrium geometries, $r_{\text{AlO}} = 3.1796 \text{ bohr}$ and $r_{\text{OH}} = 1.6597 \text{ bohr}$ (Apponi, Barclay & Ziurys 1993). As illustrated in Fig. 1, the complex is described by the Jacobi coordinate system (R, θ), where R is the intermolecular distance and θ represents the enclosed angle between the intermolecular axis and the AlOH axis. The AlOH–He van der Waals system can be described using one single electron configuration. For this purpose, we performed preliminary calculations using the complete active space self-consistent field (CASSCF) approach to check the validity of this approximation (Knowles & Werner 1985; Werner & Knowles 1985). Indeed, the dominant configuration of the ground electronic state weighs ~ 0.96 for all geometries. Therefore, the coupled cluster method with single, double and perturbative triple excitation, hereafter CCSD(T) (Adler, Knizia & Werner 2007) is suitable to probe the interaction potential of this monoconfigurational system. This approach supplemented by the augmented-correlation consistent-polarized Valence Quadruple Zeta (aug-cc-pVQZ) basis set (Kendall, Dunning & Harrison 1992) and mid-bond functions of Cybulski & Tocylowski (1999) [3s 3p 2d2f1g] – hereafter, this level of theory will be denoted as CCSD(T)/aug-cc-pVQZ+bf – was employed as implemented in the MOLPRO-2010 quantum chemistry package (Werner et al. 2010) to determine the interaction energy. This latter was calculated from $R = 4\text{--}100 \text{ bohr}$ (4, 4.25, 4.5, 4.75, 5, 5.25, 5.5, 5.75, 6, 6.25, 6.5, 6.75, 7, 7.25, 7.5, 7.75, 8, 8.25, 8.5, 8.75, 9, 9.25, 9.5,

Table 1. MOLSCAT parameter values.

R_{\min} (bohr)	R_{\max} (bohr)	STEPS	μ (au)	B_0 (cm^{-1})	D_0 (cm^{-1})
4	50	200, 100	3.66877366	0.5250414 ^a	8.276×10^{-7a}
3	30	20, 16			

^aSee Apponi et al. (1993).




Figure 4. Rotational excitation cross-sections of AlOH by He as a function of the kinetic energy for $\Delta j = 1, 2, 3$ transitions.




Figure 5. De-excitation cross-sections at total energies $E = 100, 500$ and 1500 cm^{-1} from the rotational level $j = 12$ to $j' = 11, \dots, 0$.

9.75, 10, 10.5, 11, 11.5, 12, 12.5, 13, 13.5, 14, 14.5, 15, 15.5, 16, 16.5, 17, 17.5, 18, 18.5, 19, 19.5, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 50, 100) and θ from 0° to 180° with a step of 10° . It is worth mentioning that the basis set superposition error (BSSE) was corrected using the counterpoise procedure of Boys & Bernardi (1970),

$$V(R, \theta) = E_{\text{AlOH-He}}(R, \theta) - E_{\text{AlOH}}(R, \theta) - E_{\text{He}}(R, \theta), \quad (2)$$

where $E_{\text{AlOH-He}}$ is the energy of the complex and E_{AlOH} and E_{He} are calculated in the full basis set of the complex stand for the energies of AlOH and He, respectively.




Figure 6. Rotational de-excitation rate coefficients of AlOH by He as a function of kinetic temperature for $\Delta j = -1, -2, -3$ transitions.

Fig. 2 illustrates some contour plots of the AlOH–He PES. As can be seen, the interaction potential presents a strong anisotropy with a single well depth of -32.680 cm^{-1} located at $\theta = 0^\circ$ and $R = 8.2 \text{ bohr}$ (i.e. towards the H end of the AlOH molecule). In addition, the shape is typically identical to the PES of Tchakoua et al. (2015), which has a slightly deeper potential well (-32.995 cm^{-1}). Hence, our PES can be unambiguously used to perform the dynamical calculations.

We have expanded the PES over the Legendre polynomial functions to determine the basic input needed in the dynamic computations :

$$V(R, \theta) = \sum_{\lambda=0}^{\lambda_{\max}=18} V_\lambda(R) P_\lambda(\cos \theta). \quad (3)$$

We included 19 radial terms (V_λ) in the development and this led to an analytical PES similar to the *ab initio* PES with deviations of a few cm^{-1} . Fig. 3 illustrates the R dependence of the first five V_λ . The radial coefficients corresponding to $\lambda = 1$ strongly outweigh the others. The impact of such a predominance on the cross-sections and rate coefficients is discussed in the next section.

3 DYNAMICAL STUDY

3.1 Cross-sections

We computed the rotational energy levels of AlOH using the experimental spectroscopic constants presented in Table 1. The rotational cross-sections of AlOH in collision with He were calculated using the close coupling (CC) approach (Arthurs & Dalgarno 1960) as is implemented in the MOLSCAT quantum package (Hutson & Green 1994). It is worth mentioning that the coupled equations have been

Table 2. Rotational de-excitation rate coefficients of AlOH by He as a function of kinetic temperature (in K) for $1 \leq j \leq 12$.

j	j'	$K_{j \rightarrow j'} (\text{cm}^3 \text{s}^{-1})$								
		5	10	30	50	100	150	200	250	300
1	0	8.3164(-11)	7.4824(-11)	6.1724(-11)	5.6672(-11)	5.1772(-11)	5.0243(-11)	4.9812(-11)	4.9828(-11)	4.9999(-11)
2	0	1.8117(-11)	1.6798(-11)	1.4270(-11)	1.3975(-11)	1.4371(-11)	1.4703(-11)	1.4955(-11)	1.5172(-11)	1.5357(-11)
2	1	1.0745(-10)	1.0107(-10)	8.9409(-11)	8.5171(-11)	8.1753(-11)	8.1408(-11)	8.2009(-11)	8.2935(-11)	8.3881(-11)
3	0	8.2213(-12)	8.3295(-12)	9.3635(-12)	1.0112(-11)	1.1233(-11)	1.1960(-11)	1.2527(-11)	1.2997(-11)	1.3385(-11)
3	1	2.5947(-11)	2.4799(-11)	2.3090(-11)	2.3981(-11)	2.6391(-11)	2.7741(-11)	2.8565(-11)	2.9138(-11)	2.9551(-11)
3	2	1.1334(-10)	1.0979(-10)	9.7555(-11)	9.3263(-11)	9.0503(-11)	9.0961(-11)	9.2256(-11)	9.3756(-11)	9.5160(-11)
4	0	2.6605(-12)	2.6693(-12)	3.2133(-12)	3.8145(-12)	4.7875(-12)	5.2719(-12)	5.5335(-12)	5.6861(-12)	5.7755(-12)
4	1	1.4609(-11)	1.5643(-11)	1.7870(-11)	1.9421(-11)	2.1966(-11)	2.3764(-11)	2.5173(-11)	2.6322(-11)	2.7254(-11)
4	2	2.9401(-11)	2.9108(-11)	2.7404(-11)	2.9175(-11)	3.3523(-11)	3.5949(-11)	3.7392(-11)	3.8353(-11)	3.9016(-11)
4	3	1.0993(-10)	1.1106(-10)	1.0011(-10)	9.5608(-11)	9.2756(-11)	9.3477(-11)	9.5130(-11)	9.6974(-11)	9.8674(-11)
5	0	2.0381(-12)	2.2244(-12)	2.8649(-12)	3.3070(-12)	4.0563(-12)	4.6016(-12)	5.0189(-12)	5.3472(-12)	5.6060(-12)
5	1	5.3788(-12)	5.5877(-12)	7.0973(-12)	8.8652(-12)	1.1846(-11)	1.3354(-11)	1.4176(-11)	1.4661(-11)	1.4952(-11)
5	2	1.7537(-11)	1.9100(-11)	2.1465(-11)	2.3055(-11)	2.5893(-11)	2.8141(-11)	3.0012(-11)	3.1574(-11)	3.2854(-11)
5	3	2.9430(-11)	2.9720(-11)	2.8567(-11)	3.0843(-11)	3.6654(-11)	4.0174(-11)	4.2357(-11)	4.3831(-11)	4.4856(-11)
5	4	1.0875(-10)	1.1128(-10)	1.0133(-10)	9.6802(-11)	9.3987(-11)	9.4861(-11)	9.6719(-11)	9.8773(-11)	1.0066(-10)
6	0	8.9361(-13)	9.5684(-13)	1.5509(-12)	2.2023(-12)	3.2646(-12)	3.7905(-12)	4.0752(-12)	4.2429(-12)	4.3442(-12)
6	1	4.2695(-12)	4.3961(-12)	5.4528(-12)	6.3024(-12)	7.8398(-12)	9.0812(-12)	1.0095(-11)	1.0916(-11)	1.1574(-11)
6	2	7.0488(-12)	7.2174(-12)	8.9656(-12)	1.1278(-11)	1.5712(-11)	1.8220(-11)	1.9684(-11)	2.0595(-11)	2.1175(-11)
6	3	2.1016(-11)	2.2034(-11)	2.3963(-11)	2.5392(-11)	2.8257(-11)	3.0702(-11)	3.2818(-11)	3.4624(-11)	3.6119(-11)
6	4	2.9502(-11)	2.9520(-11)	2.8657(-11)	3.1111(-11)	3.7402(-11)	4.1477(-11)	4.4154(-11)	4.6032(-11)	4.7379(-11)
6	5	1.1116(-10)	1.1265(-10)	1.0253(-10)	9.7852(-11)	9.5074(-11)	9.6181(-11)	9.8289(-11)	1.0057(-10)	1.0264(-10)
7	1	1.9531(-12)	2.0364(-12)	3.2599(-12)	4.7713(-12)	7.5876(-12)	9.1621(-12)	1.0080(-11)	1.0653(-11)	1.1021(-11)
7	2	5.8507(-12)	5.7600(-12)	6.7505(-12)	7.7333(-12)	9.6935(-12)	1.1347(-11)	1.2738(-11)	1.3888(-11)	1.4819(-11)
7	3	8.8562(-12)	8.7511(-12)	1.0276(-11)	1.2596(-11)	1.7439(-11)	2.0454(-11)	2.2351(-11)	2.3606(-11)	2.4452(-11)
7	4	2.5475(-11)	2.5374(-11)	2.6286(-11)	2.7389(-11)	3.0178(-11)	3.2827(-11)	3.5184(-11)	3.7213(-11)	3.8899(-11)
7	5	2.9723(-11)	2.9326(-11)	2.8528(-11)	3.1094(-11)	3.7414(-11)	4.1562(-11)	4.4381(-11)	4.6424(-11)	4.7934(-11)
7	6	1.1438(-10)	1.1437(-10)	1.0367(-10)	9.8796(-11)	9.5894(-11)	9.7158(-11)	9.9504(-11)	1.0202(-10)	1.0431(-10)
8	2	2.9185(-12)	2.8984(-12)	4.2724(-12)	6.0858(-12)	9.7160(-12)	1.1951(-11)	1.3358(-11)	1.4291(-11)	1.4925(-11)
8	3	7.6550(-12)	7.1634(-12)	7.8043(-12)	8.7546(-12)	1.1002(-11)	1.3038(-11)	1.4774(-11)	1.6216(-11)	1.7385(-11)
8	4	1.1158(-11)	1.0530(-11)	1.1573(-11)	1.3728(-11)	1.8402(-11)	2.1441(-11)	2.3443(-11)	2.4829(-11)	2.5804(-11)
8	5	3.1030(-11)	2.9288(-11)	2.8631(-11)	2.9260(-11)	3.1728(-11)	3.4477(-11)	3.7044(-11)	3.9295(-11)	4.1180(-11)
8	6	3.0357(-11)	2.9395(-11)	2.8420(-11)	3.1072(-11)	3.7387(-11)	4.1471(-11)	4.4260(-11)	4.6313(-11)	4.7854(-11)
8	7	1.1890(-10)	1.1658(-10)	1.0481(-10)	9.9713(-11)	9.6556(-11)	9.7784(-11)	1.0022(-10)	1.0288(-10)	1.0531(-10)
9	3	3.7854(-12)	3.7537(-12)	5.1770(-12)	7.0746(-12)	1.0899(-11)	1.3336(-11)	1.4942(-11)	1.6058(-11)	1.6849(-11)
9	4	8.9402(-12)	8.3989(-12)	8.7447(-12)	9.5846(-12)	1.1913(-11)	1.4239(-11)	1.6293(-11)	1.8023(-11)	1.9435(-11)
9	5	1.2897(-11)	1.2184(-11)	1.2865(-11)	1.4854(-11)	1.9262(-11)	2.2145(-11)	2.4071(-11)	2.5435(-11)	2.6417(-11)
9	6	3.4635(-11)	3.2625(-11)	3.0806(-11)	3.0978(-11)	3.2994(-11)	3.5660(-11)	3.8302(-11)	4.0689(-11)	4.2719(-11)
9	7	3.0003(-11)	2.9237(-11)	2.8314(-11)	3.1061(-11)	3.7424(-11)	4.1478(-11)	4.4240(-11)	4.6279(-11)	4.7818(-11)
9	8	1.1904(-10)	1.1723(-10)	1.0556(-10)	1.0043(-10)	9.7119(-11)	9.8224(-11)	1.0061(-10)	1.0328(-10)	1.0575(-10)
10	4	4.8842(-12)	4.7540(-12)	6.1297(-12)	8.0466(-12)	1.1864(-11)	1.4287(-11)	1.5903(-11)	1.7050(-11)	1.7883(-11)
10	5	1.0464(-11)	9.7282(-12)	9.6654(-12)	1.0351(-11)	1.2584(-11)	1.5013(-11)	1.7255(-11)	1.9191(-11)	2.0797(-11)
10	6	1.5043(-11)	1.4030(-11)	1.4215(-11)	1.6020(-11)	2.0160(-11)	2.2880(-11)	2.4706(-11)	2.6008(-11)	2.6954(-11)
10	7	3.8822(-11)	3.6186(-11)	3.2990(-11)	3.2666(-11)	3.4170(-11)	3.6621(-11)	3.9185(-11)	4.1572(-11)	4.3641(-11)
10	8	3.0241(-11)	2.9402(-11)	2.8341(-11)	3.1114(-11)	3.7509(-11)	4.1557(-11)	4.4321(-11)	4.6370(-11)	4.7923(-11)
10	9	1.2098(-10)	1.1853(-10)	1.0641(-10)	1.0118(-10)	9.7713(-11)	9.8672(-11)	1.0094(-10)	1.0353(-10)	1.0594(-10)
11	5	5.9840(-12)	5.8018(-12)	7.1134(-12)	9.0277(-12)	1.2796(-11)	1.5166(-11)	1.6747(-11)	1.7876(-11)	1.8705(-11)
11	6	1.1688(-11)	1.0922(-11)	1.0526(-11)	1.1060(-11)	1.3145(-11)	1.5544(-11)	1.7828(-11)	1.9848(-11)	2.1551(-11)
11	7	1.6906(-11)	1.5768(-11)	1.5528(-11)	1.7159(-11)	2.1059(-11)	2.3639(-11)	2.5385(-11)	2.6642(-11)	2.7564(-11)
11	8	4.1796(-11)	3.9167(-11)	3.4993(-11)	3.4227(-11)	3.5259(-11)	3.7474(-11)	3.9893(-11)	4.2193(-11)	4.4210(-11)
11	9	3.0245(-11)	2.9530(-11)	2.8432(-11)	3.1202(-11)	3.7611(-11)	4.1657(-11)	4.4429(-11)	4.6498(-11)	4.8076(-11)
11	10	1.2135(-10)	1.1915(-10)	1.0705(-10)	1.0181(-10)	9.8265(-11)	9.9121(-11)	1.0128(-10)	1.0376(-10)	1.0608(-10)
12	6	7.1053(-12)	6.8826(-12)	8.1122(-12)	1.0011(-11)	1.3711(-11)	1.6020(-11)	1.7567(-11)	1.8683(-11)	1.9507(-11)
12	7	1.2724(-11)	1.1985(-11)	1.1299(-11)	1.1701(-11)	1.3658(-11)	1.5992(-11)	1.8239(-11)	2.0248(-11)	2.1959(-11)
12	8	1.8629(-11)	1.7414(-11)	1.6798(-11)	1.8264(-11)	2.1930(-11)	2.4370(-11)	2.6038(-11)	2.7257(-11)	2.8162(-11)
12	9	4.3999(-11)	4.1604(-11)	3.6739(-11)	3.5622(-11)	3.6281(-11)	3.8297(-11)	4.0568(-11)	4.2750(-11)	4.4673(-11)
12	10	3.0301(-11)	2.9692(-11)	2.8557(-11)	3.1293(-11)	3.7696(-11)	4.1739(-11)	4.4511(-11)	4.6590(-11)	4.8182(-11)
12	11	1.2129(-10)	1.1943(-10)	1.0748(-10)	1.0230(-10)	9.8777(-11)	9.9578(-11)	1.0166(-10)	1.0405(-10)	1.0627(-10)

solved using the log-derivative propagator of Manolopoulos (1986). For a best compromise between result accuracy and computational time, we carried out preliminary calculations to fix the integration parameters of the propagator. In addition, Table 3 displays


some convergence tests for the rotational basis, j_{\max} . As can be seen, the convergence is reached with $j_{\max} = 23$, for a total energy (E) of 600 cm^{-1} . Indeed, the cross-sections slightly vary (0.002 \AA^2) when passing from $j_{\max} = 21$ to 23. For the entire total en-

Table 3. Cross-sections (in \AA^2) for different values of j_{\max} for total energy $E = 600 \text{ cm}^{-1}$.

Transitions	$E (\text{cm}^{-1})$	Tchakoua et al. (2015)				j_{\max}	This work			
		17	19	21	23		17	19	21	23
0 → 1	600	75.884	75.884	75.884	75.884		8.012	8.012	8.012	8.012
0 → 2		4.423	4.423	4.423	4.423		4.180	4.180	4.180	4.180
1 → 2		77.776	77.776	77.776	77.776		7.723	7.722	7.722	7.722
2 → 3		78.666	78.666	78.666	78.666		7.498	7.498	7.498	7.498
5 → 6		79.405	79.405	79.405	79.405		7.123	7.126	7.126	7.126
5 → 7		8.290	8.290	8.290	8.290		3.866	3.867	3.866	3.866
6 → 7		79.552	79.552	79.552	79.552		7.172	7.168	7.167	7.167
8 → 10		8.527	8.527	8.527	8.527		3.592	3.513	3.512	3.512
11 → 7		–	–	–	–		1.708	1.714	1.713	1.712
11 → 8		–	–	–	–		3.071	3.058	3.054	3.054
11 → 9		–	–	–	–		3.019	2.977	2.981	2.983
11 → 10		–	–	–	–		6.545	6.528	6.528	6.527
12 → 8		–	–	–	–		1.864	1.759	1.756	1.757
12 → 9		–	–	–	–		3.090	3.038	3.051	3.052
12 → 10		–	–	–	–		3.270	3.045	2.999	2.997
12 → 11		–	–	–	–		6.608	6.575	6.550	6.548

Table 4. Comparison the AlOH–He, HCP–He and AlF–He global minima (in units of cm^{-1}).

Coordinates		This work	Hammami et al.	Gotoum et al.
$R (\text{bohr})$	$\theta (\text{°})$	AlOH–He	HCP–He	AlF–He
8.20	0	32.680	–	–
9.20	0	–	22.018	–
7.75	0	–	–	24.056

**Figure 7.** Our rotational de-excitation rate coefficients of AlOH–He in comparison with results obtained by Tchakoua et al. (2015).

ergy range, which expands up to 1500 cm^{-1} , j_{\max} was fixed at 17 for $0 \leq E \leq 100 \text{ cm}^{-1}$, 20 for $100 \leq E \leq 300 \text{ cm}^{-1}$, 23 for $300 \leq E \leq 600 \text{ cm}^{-1}$, 27 for $600 \leq E \leq 1000 \text{ cm}^{-1}$ and 33 for $1000 \leq E \leq 1500 \text{ cm}^{-1}$.

The total energy range was spanned using steps of 0.1 cm^{-1} for $0 \leq E \leq 35 \text{ cm}^{-1}$, 0.2 cm^{-1} for $35 \leq E \leq 100 \text{ cm}^{-1}$, 5 cm^{-1} for $100 \leq E \leq 300 \text{ cm}^{-1}$, 10 cm^{-1} for $300 \leq E \leq 600 \text{ cm}^{-1}$, 20 cm^{-1} for $600 \leq E \leq 1000 \text{ cm}^{-1}$ and a step of 50 cm^{-1} for $1000 \leq E \leq 1500 \text{ cm}^{-1}$.

Fig. 4 outlines the rotational excitation cross-sections for $\Delta j = 1, 2$ and 3. Several resonances are observed at low kinetic energies (E_k), more precisely at $E_k \leq 30 \text{ cm}^{-1}$. This behaviour is due to the global minimum of the interaction potential, which temporarily traps the He atom leading to the formation of quasi-bound states of the AlOH–He complex. In addition, Fig. 4 shows that the $0 \rightarrow 1$ and $1 \rightarrow 2$ transitions outweigh in the whole kinetic energy range. Although the $\Delta j = 2$ ($0 \rightarrow 2$ and $1 \rightarrow 3$) transitions predominate the $\Delta j = 3$ transitions ($0 \rightarrow 3$ and $1 \rightarrow 4$) at low kinetic energy, an inversion occurs at $E_k \geq 500 \text{ cm}^{-1}$. Therefore, we can conclude that the propensity rules are in favour of the odd Δj transitions. Indeed, this trend was predicted by the dominance of the V_1 and V_3 radial coefficients (see Fig. 3).

Fig. 5 displays the dependence of the de-excitation cross-sections with respect to the final rotational level (j') for selected kinetic energies ($E_k = 100, 500$ and 1500 cm^{-1}). The behaviour of these curves allows us to confirm that the propensity rules favour the $\Delta j = \pm 1$ transitions.

3.2 Rate coefficients

The rate coefficients ($K_{j \rightarrow j'}$), which depend on the kinetic temperature (T), were determined by averaging the cross-sections ($\sigma_{j \rightarrow j'}$) over the Maxwell–Boltzmann distribution as follows:

$$K_{j \rightarrow j'}(T) = \left(\frac{8}{\pi \mu \beta} \right)^{1/2} \beta^2 \int_0^\infty E_k \sigma_{j \rightarrow j'}(E_k) e^{-\beta E_k} dE_k. \quad (4)$$

Here, $\beta = 1/(k_B T)$, where k_B is the Boltzmann constant and μ is the reduced mass (see Table 1). Based on the total energy range, we were able to compute the AlOH rate coefficients corresponding

Table 5. Temperature variation of four sets of rate coefficients (in $\text{cm}^3 \text{s}^{-1}$) for $\Delta j = -1$ transitions.

System		Temperature (K)	$1 \rightarrow 0$	$2 \rightarrow 1$	$3 \rightarrow 2$	$4 \rightarrow 3$	$5 \rightarrow 4$
AlOH–He	Tchakoua et al.	10	1.079(−9)	7.81(−10)	4.380(−10)	8.152(−10)	2.091(−9)
		30	2.087(−10)	1.511(−10)	8.470(−11)	1.576(−10)	4.044(−10)
		150	1.870(−11)	1.354(−11)	7.590(−12)	1.413(−11)	3.624(−11)
	This work	10	7.482(−11)	1.011(−10)	1.098(−10)	1.111(−10)	1.113(−10)
		30	6.172(−11)	8.941(−11)	9.756(−11)	1.001(−10)	1.013(−10)
		150	5.667(−11)	8.517(−11)	9.326(−11)	9.561(−11)	9.680(−11)
HCP–He ^a		20	0.796(−11)	1.366(−11)	1.550(−11)	1.343(−11)	1.172(−11)
		50	1.072(−11)	1.640(−11)	1.796(−11)	1.758(−11)	1.709(−11)
		150	1.748(−11)	2.908(−11)	3.124(−11)	3.182(−11)	3.207(−11)
AlF–He ^b		10	0.9362(−10)	1.240(−10)	1.396(−10)	1.531(−10)	1.659(−10)
		30	1.155(−10)	1.498(−10)	1.630(−10)	1.733(−10)	1.821(−10)
		150	1.493(−10)	2.107(−10)	2.262(−10)	2.333(−10)	2.382(−10)

^a See Hammami et al. (2008). ^b See Gotoum et al. (2011).

to the first 13 rotational levels in the kinetic temperature range 5–300 K.

Fig. 6 illustrates the rotational de-excitation rate coefficients for $\Delta j = -1, -2$ and -3 transitions. These curves exhibit that the $\Delta j = -1$ transitions strongly predominate for all kinetic temperatures. Moreover, Table 2 presents the downward rotational rate coefficients computed in this work. These data are very important for the astrophysical community. They allow us to determine the physical conditions of interstellar clouds where AlOH was observed without assuming the LTE. Even if AlOH mainly collides with H₂, which is the most abundant species in the media mentioned above, the AlOH–H₂ rate coefficients can be roughly approximated from those of AlOH–He. As helium is a closed-shell atom with two electrons, the AlOH rate coefficients induced by collision with para-H₂ ($j = 0$) can be derived by multiplying those of AlOH—He by the mass scaling factor which is $\sim \sqrt{2}$. Such an estimation was justified in early works on heavy molecules, such as SiS (Lique et al. 2008) and SO (Lique et al. 2007). However, this approximation was not valid for many other systems, including relatively light molecules, (Gotoum et al. 2012; Lanza et al. 2014; Najar et al. 2017). As a result, the present study provides an opportunity to study the collisional de-excitation of AlOH by H₂.

4 COMPARISON STUDY

Despite the agreements between our PES and that of Tchakoua et al. (2015), significant differences are noticed in the dynamic results. In fact, our inelastic cross-sections do not exceed 250 Å² while those available in the literature achieve 600 Å². To better appreciate the disagreement, in Table 3 we give both sets of cross-sections computed at 600 cm^{−1} using different values of j_{\max} . For most of the transitions (mainly those for which $\Delta j = 1$), the old values of cross-sections are 10 to 100 times greater than the new values. We expect that these large values of cross-sections reported by Tchakoua et al. (2015) are due to convergence problems, as they did not take into account the transitions involving $j \geq 10$. The discrepancies mentioned above will persist for the rate coefficients, which are derived by thermally averaging the cross-sections. In Fig. 7, we compare the two sets of rotational rate coefficients by plotting the $5 \rightarrow 4$ and $5 \rightarrow 3$ transitions. As expected, the comparison reveals that the rate coefficients of Tchakoua et al. (2015) are ~ 100 times larger than ours, at low kinetic temperatures ($T \leq 50$ K). Due to

the lack of experimental rate coefficients for AlOH–He, we rely on the data of HCP–He (Hammami et al. 2008) and AlF–He (Gotoum et al. 2011) in order to settle the matter mentioned above. In fact, these molecules are isoelectronic to AlOH and HCP is geometrically identical to it. Moreover, AlOH–He, HCP–He and AlF–He have very similar PESs in term of shape and well depth (see Table 4). All these points support the pertinence of the comparison and predict rate coefficients in the same magnitude rank for the three systems. Therefore, in Table 5 we show the rate coefficients of the three systems for selected temperatures and transitions. The HCP (AlF) data vary from 0.8×10^{-11} to 3.3×10^{-11} (1.0×10^{-10} to 2.4×10^{-10}) and those of AlOH computed in this work (calculated by Tchakoua et al. 2015) are between 7.5×10^{-11} and 1.1×10^{-10} (7.6×10^{-12} and 1.1×10^{-9}). According to the comparison made in Table 5, the rate coefficients of AlOH presented in this paper are more reliable than those provided by Tchakoua et al. (2015).

5 CONCLUSIONS

In this paper, we have revisited the rotational excitations and de-excitations of AlOH due to impact with He. A new two-dimensional PES was computed with the CCSD(T)/aug-cc-pVQZ+bf *ab initio* method. This PES was then used for the quantum calculations of cross-sections considering total energies up to 1500 cm^{−1}. During the scattering computations, the CC quantum approach was used. By thermally averaging these data over the Maxwell–Boltzmann distribution, rate coefficients of the 13 lowest rotational levels of AlOH were determined for kinetic temperatures ranging from 5 to 300 K. The new rate coefficients of AlOH have been compared with the previous ones of Tchakoua et al. (2015) and significant discrepancies are noticed concerning the magnitude ranks. Then we compared both sets of data with those of similar molecules (HCP and AlF). This leads us to conclude that the new rate coefficients of AlOH, which are in good agreement with those of HCP and AlF, are more reliable than the previous ones. Therefore, we expect that the data presented in this paper will play a key role in the accurate modelling of the abundance of AlOH in the ISM. In addition, the discrepancies between the previous and new rates of AlOH mentioned above might encourage experimental work to confirm our findings.

ACKNOWLEDGEMENTS

The authors acknowledge Dr François Lique, Laboratoire Ondes et Milieux complexes, UMR CNRS 6294, Université Le Havre, for discussion and advice.

REFERENCES

- Álvarez-Barcia S., Flores J. R., 2016, *Phys. Chem. Chem. Phys.*, 92, 382
 Álvarez-Barcia S., Flores J. R., 2011, *Chem. Phys.*, 92, 382
 Adler T. B., Knizia G., Werner H. J., 2007, *J. Chem. Phys.*, 127, 221106
 Apponi A. J., Barclay W. L., Ziurys L. M., 1993, *ApJ*, 414, 129
 Arthurs A. M., Dalgarno A., 1960, *Proc. R. Soc. A.*, 256, 540
 Boys S. F., Bernardi F., 1970, *Mol. Phys.*, 19, 553
 Cybulski S. M., Toczyłowski R., 1999, *J. Chem. Phys.*, 111, 10520
 Gotoum N., Nkem C., Hammami K., Charfadine M. A., Owono Owono L. C., Jaidane N., 2011, *Ap&SS*, 332, 209
 Gotoum N., Nkem C., Hammami K., Owono Owono L. C., Jaidane N., 2012, *Ap&SS*, 337, 553
 Hammami K., Owono Owono L. C., Jaidane N., Ben Lakhdar Z., 2008, *J. Molecular Structure: THEOCHEM*, 860, 45
 Hauge F. H., Kauffman J. W., Margrave J. L., 1980, *J. Am. Chem. Soc.*, 102, 6005
 Hutson J. M., Green S., 1994, MOLSCAT computer code, version 14, Collaborative Computational Project No. 6. Science and Engineering Research Council, UK
 Kamiński T. et al., 2016, *A&A*, 592, A42
 Kendall R. A., Dunning T. H., Harrison R. J., 1992, *J. Chem. Phys.*, 96, 6796
 Knowles P. J., Werner H. J., 1985, *Chem. Phys. Lett.*, 115, 259
 Lanza M., Kalugina Y., Wiesenfeld L., Faure A., Lique F., 2014, *MNRAS*, 443, 3351
 Li S., Sattelmeyer K. W., Yamaguchi Y., Schaefer H. F., 2003, *J. Chem. Phys.*, 119, 12830
 Lique F., Senent M. L., Spielfiedel A., Feautrier N., 2007, *J. Chem. Phys.*, 126, 164312
 Lique F., Tobola R., Klos J., Feautrier N., Spielfiedel A., Vincent L. F. M., Chalasiński G., Alexander M. H., 2008, *A&A*, 478, 567
 McClean R. E., Nelson H. H., Campbell M. L., 1993, *J. Phys. Chem.*, 97, 9673
 Manolopoulos D. E., 1986, *J. Chem. Phys.*, 85, 6425
 Morton D. C., Drake J. F., Jenkins E. B., Rogerson J. B., Spitzer L., York D. G., 1973, *ApJ*, 181, 103
 Najar F., Naouai M., El Hanini H., Jaidane N., 2017, *MNRAS*, 472, 2919
 Pilgrim J. S., Robbins D. L., Duncan M. A., 1993, *Chem. Phys. Lett.*, 202, 203
 Tchakoua T., Pamboundom M., Nsangou M., Motapon O., 2015, *Ap&SS*, 357, 142
 Tenenbaum E. D., Ziurys L. M., 2009, *ApJ*, 694, 59
 Tenenbaum E. D., Ziurys L. M., 2010, *ApJ*, 712, 93
 Tsuji T., 1973, *A&A*, 23, 411
 Vacek G., Deleeuw B. J., Schaefer H. F., 1993, *J. Chem. Phys.*, 98, 8704
 Werner H. J., Knowles P. J., 1985, *J. Chem. Phys.*, 82, 5053
 Werner H. J. et al., 2010, MOLPRO, version 2010.1, a package of *ab initio* programs, <http://www.molpro.net>
 Ziurys L. M., 2006, *Proceedings of the National Academy of Sciences*, 103, 12 274
 Ziurys L. M., Savage C., Highberger J. L., Apponi A. J., Guélin M., Cericharo J., 2002, *ApJ*, 564, L45

This paper has been typeset from a TeX/LaTeX file prepared by the author.