



Identification of a circular code periodicity in the bacterial ribosome: origin of codon periodicity in genes?

Christian J. Michel, Julie Thompson

► To cite this version:

Christian J. Michel, Julie Thompson. Identification of a circular code periodicity in the bacterial ribosome: origin of codon periodicity in genes?. RNA Biology, 2020, 17 (4), pp.571-583. 10.1080/15476286.2020.1719311 . hal-03078903


HAL Id: hal-03078903

<https://hal.science/hal-03078903>

Submitted on 2 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Identification of a circular code periodicity in the bacterial ribosome: origin of codon periodicity in genes?

Christian J. Michel & Julie D. Thompson

To cite this article: Christian J. Michel & Julie D. Thompson (2020): Identification of a circular code periodicity in the bacterial ribosome: origin of codon periodicity in genes?, RNA Biology, DOI: [10.1080/15476286.2020.1719311](https://doi.org/10.1080/15476286.2020.1719311)

To link to this article: <https://doi.org/10.1080/15476286.2020.1719311>



Accepted author version posted online: 21 Jan 2020.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Publisher: Taylor & Francis & Informa UK Limited, trading as Taylor & Francis Group

Journal: *RNA Biology*

DOI: 10.1080/15476286.2020.1719311

**Identification of a circular code periodicity in the bacterial ribosome:
origin of codon periodicity in genes?**

Christian J. Michel *, Julie D. Thompson *

Department of Computer Science, ICube, CNRS, University of Strasbourg, Strasbourg, France

Email: c.michel@unistra.fr, thompson@unistra.fr

* To whom correspondence should be addressed; Email: c.michel@unistra.fr

Keywords: three-base periodicity; circular code periodicity; ribosome; 16S rRNA; protein-coding gene.

Abstract

Three-base periodicity (TBP), where nucleotides and higher order n -tuples are preferentially spaced by 3, 6, 9, etc. bases, is a well-known intrinsic property of protein-coding DNA sequences. However, its origins are still not fully understood. One hypothesis is that the periodicity reflects a primordial coding system that was used before the emergence of the modern standard genetic code (SGC). Recent evidence suggests that the X circular code, a set of 20 trinucleotides allowing the reading frames in genes to be retrieved locally, represents a possible ancestor of the SGC. Motifs from the X circular code have been found in the reading frame of protein-coding regions in extant organisms from bacteria to eukaryotes, in many transfer RNA (tRNA) genes and in important functional regions of the ribosomal RNA (rRNA), notably in the peptidyl transferase center and the decoding center (Dila et al., 2019b). Here, we have used a powerful correlation function to search for periodicity patterns involving the 20 trinucleotides of the X circular code in a large set of bacterial protein-coding genes, as well as in the translation machinery, including rRNA and tRNA sequences. As might be expected, we found a strong circular code periodicity 0 modulo 3 in the protein-coding genes. More surprisingly, we also identified a similar circular code periodicity in a large region of the 16S rRNA. This region includes the 3' major domain corresponding to the primordial proto-ribosome decoding center and containing numerous sites that interact with the tRNA and messenger RNA (mRNA) during translation. Furthermore, 3D structural analysis shows that the periodicity region surrounds the mRNA channel that lies between the head and the body of the SSU. Our results support the hypothesis that the X circular code may constitute an ancestral translation code involved in reading frame retrieval and maintenance, traces of which persist in modern mRNA, tRNA and rRNA despite their long evolution and adaptation to the SGC.

Introduction

The genetic code defines the set of rules needed to translate the information in DNA into proteins. Virtually all living organisms use the same standard genetic code (SGC) to determine how the 64 DNA trinucleotides (also known as codons) are translated into 20 amino acids and the stop signal. The degeneracy of the genetic code (most amino acids are coded by more than one codon) and specific codon usage bias in different organisms leads to a biased distribution of codons, and an intrinsic property of protein-coding DNA, known as three-base periodicity (TBP), defined as the preferential spacing of nucleotides and other n -tuples such as trinucleotides by distances of 3, 6, 9, etc. bases (Shepherd, 1981a,b), i.e. a periodicity 0 modulo 3.

This periodic phenomenon has intrigued biologists for decades (e.g. Arquès and Michel, 1990a,b; Gutiérrez et al., 1994; Trifonov, 1998). For example, it led to the proposal that the ancestral forms of present-day genes might have been coded by the primitive comma-free codes RRY and RNY ($R=\{A,G\}$, $Y=\{C,T\}$, N being any base) (Crick et al., 1976; Eigen and Winkler-Oswatitsch 1981; Eskesen et al., 2004). To illustrate the notion of TBP, for a RRY code, a word $RRY|RRY|RRY|...$ implies that any letter Y

is distant from another letter Y by a multiple of 3 letters (3, 6, etc.), and any trinucleotide RRY is also distant from another trinucleotide RRY by a multiple of 3 letters (0, 3, 6, etc.), etc. Obviously, in real genetic sequences, the preferential occurrence of some codons in genes (such as the X circular code, defined below or the codon usage biases observed in different genomes) implies a modulo 3 periodic signal which is very noisy but which can be identified by sensitive statistical-signal analysis functions. It was also suggested that TBP may have a structural or functional role, for example to maintain the reading frame (Trifonov, 1987) or to regulate gene expression in some way (Shabalina et al., 2006; Ding et al., 2014). Furthermore, powerful modern algorithms utilize the TBP to predict coding regions in unannotated genomes (Shepherd 1981b; Tiwara et al., 1997; Guigó et al., 2000; Yin and Yau, 2005, 2007; Marhon and Kramer, 2011; Chen and Ji, 2011). Unraveling the origins of TBP may help understand the forces that shaped the code during the early evolution of life on Earth. It has been suggested that TBP is simply due to species-specific amino acid or codon usage bias (Ohno, 1988; Tsonis et al., 1991; Eskesen et al., 2004), although recently this has been shown to be insufficient to explain TBP in modern genes (Howe and Song, 2012). It has also been proposed that TBP additionally reflects a tendency for trinucleotides to cluster in the same phase (Sánchez and López-Villaseñor, 2006). In addition to TBP in protein-coding genes, a small number of studies have also identified periodicities in homologous regions of transfer RNA (tRNA) and ribosomal RNA (rRNA) genes (Block et al., 1985; Nazarea et al., 1985; Johnson and Wang, 2010), and some authors have concluded that this might reflect a primal pre-translational code (Rodin et al., 2011).

One potential primordial translation code is the X circular code (Arquès and Michel, 1996). According to coding theory, circular codes are a weaker version of comma-free codes, where any word written on a circle (the last letter becoming the first in the circle) has a unique decomposition into trinucleotides of the circular code. The mathematical formalisms (definitions, theorems, properties, enumeration, etc.) of codes, circular codes and a special class of circular codes, known as comma-free codes, can be found in two reviews (Michel, 2008; Fimmel and Strüngmann, 2018), and are summarized here in Figure 1. A circular code with words of 3 letters (e.g. trinucleotides) on a 4-letter alphabet (e.g. the genetic alphabet) is said to be maximal when it contains 20 words (Arquès and Michel 1996). Indeed, there is no circular code with trinucleotides on the genetic alphabet that has a strictly larger size than 20 words. There are 12,964,440 maximal circular codes (Arquès and Michel 1996). Remarkably, one of the maximal circular codes, called the X circular code, was found to be overrepresented in the reading frame of protein coding genes from bacteria, archaea, eukaryotes, plasmids and viruses (Michel 2017, 2015; Arquès and Michel 1996). The X circular code consists of 20 trinucleotides

$$X = \{AAC, AAT, ACC, ATC, ATT, CAG, CTC, CTG, GAA, GAC, \\ GAG, GAT, GCC, GGC, GGT, GTA, GTC, GTT, TAC, TTC\} \quad (1)$$

and codes the 12 following amino acids (three and one letter notation)

$$\mathcal{X} = \{Ala, Asn, Asp, Gln, Glu, Gly, Ile, Leu, Phe, Thr, Tyr, Val\} \\ = \{A, N, D, Q, E, G, I, L, F, T, Y, V\}. \quad (2)$$

The trinucleotide set X has several strong mathematical properties. In particular, it is self-complementary, i.e. 10 trinucleotides of X are complementary to the other 10 trinucleotides of X , e.g. $AAC \in X$ is complementary to $GTT \in X$. Moreover, the $+1/-2$ and $+2/-1$ circular permutations of X , denoted X_1 and X_2 respectively, are also maximal circular codes (C^3) and are complementary to each other (Arquès and Michel 1996). The class of circular codes, like comma-free codes, also have the property of synchronizability, i.e. they are hypothesized to retrieve and maintain the reading frame by using an appropriate window of nucleotides. In any sequence generated by a trinucleotide comma-free code, the reading frame can be determined in a window length of at most 3 consecutive nucleotides, while for the X circular code, at most 13 consecutive nucleotides (i.e. at most 4 trinucleotides) are enough to always retrieve the reading frame. In other words, a sequence 'motif' containing several consecutive X trinucleotides is sufficient to determine the correct reading frame (see Figure 1 for examples).




Figure 1. Hierarchy of a set of words, a code, a circular code and a comma-free code. The symbol "|" denotes the decomposition on a line for a code or non-code, and on a circle for a code which is circular or non-circular. The set $\{AAA, AAC\}$ is a code as there is an unique decomposition on a line for any words of the code (an example of an unique decomposition on a line is given with the word $AACAAA$). The genetic code $\{AAA, \dots, TTT\}$ is a code from a mathematical point of view. The frame 0 is the reading frame. Words belonging to the code are shown in green, and other words are in red. The code $\{ACT, GGT, GTA, GTC\}$ is circular, since only the frame 0 (reading frame) can be read by words of the code. There is an unique decomposition on a circle for any words of the circular code (an example of an unique decomposition on a circle is given with the word $GTAGGTACTGTC$; green symbol "|"). The

code {*GGT, GTG, TGT*} is not circular as several frames (here frames 0 and 1) can be read by words of the code (several decompositions on a circle; green and red symbols "|"). The code {*GGT, GTA, GTC*} is comma-free, since words of the code appear only in the reading frame.

The hypothesis of the *X* circular code as a primordial coding system is supported by evidence from several statistical analyses of modern genomes. For example, it was shown in a large scale study of 138 eukaryotic genomes (El Soufi and Michel, 2016) that *X* motifs are found preferentially in protein-coding genes compared to non-coding regions with a ratio of ~8 times more *X* motifs located in genes. More detailed studies of the complete gene sets of yeast and mammal genomes (Michel et al., 2017; Dila et al., 2019a) confirmed the strong enrichment of *X* motifs in genes and further demonstrated a statistically significant enrichment in the reading frame compared to frames 1 and 2 (p -value $< 10^{-10}$). In addition, it was shown that most of the mRNA sequences from these organisms (e.g. 98% of experimentally verified genes in *S. cerevisiae*) contain *X* motifs.

In addition to mRNA sequences, conserved *X* motifs have also been found in many tRNA genes (Michel, 2013), as well as many important functional regions of the ribosomal RNA, notably the decoding center (Michel, 2012; El Soufi and Michel, 2014; El Soufi and Michel, 2015; Dila et al., 2019b), which suggest their involvement in universal gene translation mechanisms. Intriguingly, the theoretical minimal RNA rings, short RNAs designed to code for all coding signals without coding redundancy among frames, are also biased for codons from the *X* circular code (Demongeot and Seligmann, 2019a). These RNA rings, despite being designed based on coding constraints, attempt to mimic primitive tRNAs and potentially reflect ancient translation machineries (Demongeot and Moreira, 2007; Demongeot and Seligmann, 2019b). Based on the combined results of these previous studies, we hypothesized that the *X* circular code was an ancestor code of the SGC, which would have been used to code a smaller set of amino acids and with the additional ability to identify and maintain the reading frame. This primordial circular code would have existed before the emergence of complex start/stop codon recognition systems (see the model proposed in Figure 8 in Dila et al., 2019b), although the molecular mechanisms underlying this process are not known. It is also unknown whether circular codes continue to contribute to frame recognition in extant organisms that use the standard genetic code to code for highly complex proteins.

In this paper, we extend our study of the ribosome (Dila et al., 2019b) and investigate whether the *X* circular code, like the SGC, presents a periodicity property. To achieve this, we used a powerful circular code correlation function to search for periodicity patterns involving the 20 trinucleotides of the *X* circular code in coding sequences and in the translation machinery, including rRNA and tRNA, from bacteria. As might be expected, we found a strong circular code periodicity in the coding regions. More surprisingly, we also identified a statistically significant circular code periodicity of *X* trinucleotides in a large region of the 16S rRNA for a large set of >100 sequences from diverse bacteria.

Accepted Manuscript

Materials and methods

We define here a circular code correlation function which gives exact probabilities (with the exception of numerical approximations). This approach is particularly adapted to identifying periodicities in short and noisy sequences, such as the ribosomal and transfer RNAs.

Circular code correlation function

A language F , e.g. a genome or a set of genomes, consists of $|F|$ words, e.g. protein coding genes, ribosomal RNA (rRNA) genes, etc., on the alphabet $B = \{A, C, G, T\}$ (F is a finite subset of all words over B). Let $w = l_1 l_2 \dots l_{|w|}$ be a word of F of length $|w|$ letters (nucleotides), $l_i \in B$ for $i \in \{1, \dots, |w|\}$. Let m and m' be 2 motifs of respective lengths $|m|$ et $|m'|$ on B . Then, the word correlation function $A_{m,m'}(i, w)$ in w is defined by

$$A_{m,m'}(i, w) = \frac{1}{l(w)} \sum_{p=1}^{l(w)} \delta_m(p) \cdot \delta_{m'}(p + |m| + i), \quad i = 0, \dots, imax \quad (3)$$

with

$$\delta_m(p) = \begin{cases} 1 & \text{if the motif in position } p..p + |m| - 1 \text{ is } m \\ 0 & \text{otherwise} \end{cases}$$

and $l(w) = |w| - (|mm'| + imax) + 1$ with the length $|mm'| = |m| + |m'|$.

Note that when $i = 0$, the motif m in position $p..p + |m| - 1$ and the motif m' in position $p + |m| + i..p + |m| + i + |m'| - 1$, i.e. $p + |m|..p + |mm'| - 1$, are consecutive.

This definition of $A_{m,m'}(i, w)$ can also be understood as follows. Let an i -motif $mN^i m'$ be 2 motifs m and m' separated by i , $i \in \{0, \dots, imax\}$, any letters $N \in B$. In order to count the occurrences of $mN^i m'$ in a word w of F under the same conditions for all $i \in \{0, \dots, imax\}$, i.e. without probability bias, only the $l(w) = |w| - (|mm'| + imax) + 1$ first letters of w are analysed (a few i -motifs at the end of the sequence are thus not considered, since $l(w)$ is a function of $imax$ and not of i). Indeed, when $p = l(w)$ and $i = imax$, then the motif m' in position $p + |m| + i = |w| - |m'| + 1$ has its last letter $l_{|m'|}$ in the last position of w .

Remark 1. The definition of $A_{m,m'}(i, w)$ is a generalization of the classical letter correlation function used in signal analysis when the motifs m and m' are letters, i.e. when $m = l$ and $m' = l'$ (see Appendix A).

Remark 2. As a consequence of Equation (3), the word correlation function $A_{m,m'}(i, w)$ gives exact probabilities (with the exception of numerical approximations) which can be retrieved mathematically when the word w has a basic structure or a combination of basic structures, e.g. l^n , $(l_1 l_2)^n$, $(l_1 l_2)^n (l_1 l_2 l_3)^m$, etc. (see Appendix A and in particular, the example computations in A.3). However, only a computed function $A_{m,m'}(i, w)$ can be used in real genetic sequences.

Remark 3. As consequences of Remark 1 and Remark 2, the function $A_{l,l'}(i, w)$ (particular case when $m = l$ and $m' = l'$) is similar but not identical to the classical correlation function which is in bijection with the Fourier transform. Indeed, the classical correlation function does not correct the side effect

induced by the finite length of the word w (see Appendix A and in particular, the example computations in A.3).

In order to study the correlation function of the circular code X based on 20 trinucleotides, we choose $|m| = |m'| = 3$ and extend Equation (3) to a set of motifs. Let $B^3 = \{AAA, \dots, TTT\}$ be the set of the 64 trinucleotides with the following partition into 2 classes $C_2 = \{X, \bar{X} : X \cap \bar{X} = \emptyset, X \cup \bar{X} = B^3\}$ by recalling

$X = \{AAC, AAT, ACC, ATC, ATT, CAG, CTC, CTG, GAA, GAC, GAG, GAT, GCC, GGC, GGT, GTA, GTC, GTT, TAC, TTC\}$ which was given in (1). Then, the circular code autocorrelation function $A_{X,X}(i, w)$ in w is defined by

$$A_{X,X}(i, w) = \sum_{m \in X} \sum_{m' \in X} A_{m,m'}(i, w), \quad i = 0, \dots, imax \quad (4)$$

with $A_{m,m'}(i, w)$ defined in Equation (3).

Equation (4) is easily extended to a language. Thus, the circular code autocorrelation function $A_{X,X}(i, F)$ in F is defined by

$$A_{X,X}(i, F) = \frac{1}{|F|} \sum_{w \in F} A_{X,X}(i, w), \quad i = 0, \dots, imax \quad (5)$$

with $A_{X,X}(i, w)$ defined in Equation (4).

The function $i \rightarrow A_{X,X}(i, F)$, which gives the occurrence probability that the circular code X appears any i letters N after X in the language F , is called the circular code autocorrelation function XN^iX (associated with the i -motif XN^iX based on the circular code X). It is represented by a curve with:

- on the abscissa, the number i of letters N between X and itself (i.e. X and X), i varying from 0 to $imax$, which is chosen to be equal to 20 in the described results.
- on the ordinate, the occurrence probability $A_{X,X}(i, F)$ of XN^iX in F .

Remark 4. $\sum_{m \in \{X, \bar{X}\}} \sum_{m' \in \{X, \bar{X}\}} A_{m,m'}(i, F) = 1$ for all i letters N^i , $i \in \{0, \dots, imax\}$, and any F . The curve $A_{X,X}(i, F)$ is a horizontal line of value 1.

Remark 5. $A_{X,X}(i, F) = \frac{20 \cdot 20}{64 \cdot 64} = \frac{25}{256} \approx 0.0977$ for all i letters N^i , $i \in \{0, \dots, imax\}$, in a random language F , and in particular in a random word (sequence) w (case $|F| = 1$). The curve $A_{X,X}(i, F)$ is a horizontal line of value 0.0977. $A_{X,X}(i, F) = \frac{20 \cdot 20}{61 \cdot 61} \approx 0.107$ for all i letters N^i , $i \in \{0, \dots, imax\}$, in a random language F without the three stop codons.

Remark 5 is particularly interesting as any correlation curve without horizontal line can be associated with a non-random language F or a non-random word (sequence) w .

In Appendix A, we compare the method developed here with the two classical correlation functions used in signal analysis.

Protein-coding genes

Bacterial protein-coding genes were obtained from the GenBank database (<http://www.ncbi.nlm.nih.gov/genome/browse/>). Only one genome for each species was selected. Genes without initiation codons, without stop codons, with nucleotides different from *B* and with lengths non-modulo 3 were excluded. This resulted in a set of bacterial genes $F = \text{Genes}_{\text{Bac}}$ containing 465,762 genes with a total length of 2,339,752,707 trinucleotides.

Ribosomal RNA sequences and structure

Multiple sequence alignments for 16S small subunit (SSU) rRNAs and 23S large subunit (LSU) rRNAs were obtained from the Comparative RNA Web (CRW) site at <http://www.rna.icmb.utexas.edu/DAT/3C/Alignment>. In order to obtain a broad but sparse sampling of the bacterial domain, we used the seed alignment containing complete sequences for rRNAs from bacteria, and selected 1 representative sequence from each subgroup. This resulted in two alignments, each containing 103 sequences from the organisms provided in the Appendix B. Each alignment was then divided into two equal parts, corresponding to the 5' and 3' regions of the ribosome sequences. For the 16S rRNA alignment, the 3' region corresponds to nucleotides 1-765 (*E. coli* numbering) and the 5' region corresponds to nucleotides 766-1530 (*E. coli* numbering). For the 23S rRNA alignment, the 5' region corresponds to nucleotides 1-1447 (*E. coli* numbering) and the 3' region corresponds to nucleotides 1448-2895 (*E. coli* numbering).

The secondary structures of the SSU rRNA for *E. coli* were downloaded from <http://apollo.chemistry.gatech.edu/RibosomeGallery/>. Mapping of information on to secondary structures was performed with RiboVision (apollo.chemistry.gatech.edu/RiboVision) (Bernier et al., 2014). Coordinates of the high-resolution crystal structure of the *T. thermophilus* ribosome (PDB entry 4W2F) were obtained from the PDB database (<https://www.rcsb.org/>). This was chosen because it contains mRNA nucleotides and three deacylated tRNAs in the A, P and E sites. Numbering of the *T. thermophilus* SSU rRNA is the same as for *E. coli*. Visualization and analysis of the three-dimensional structures, as well as image preparation were performed with PyMOL (The PyMOL Molecular Graphics System, Version 1.2r3pre, Schrödinger, LLC).

Transfer RNA sequences

Transfer RNA (tRNA) sequences were downloaded from the tRNAdb database at <http://trna.bioinf.uni-leipzig.de>. All bacterial sequences were selected and grouped according to the corresponding amino acids. This resulted in 20 sets of tRNA sequences corresponding to each amino acid, with the number of sequences in each set given in the Appendix B.

Results

Circular code periodicity in bacterial genes

We first applied the circular code autocorrelation method to a large set of bacterial genes $F = \text{Genes}_{\text{Bac}}$ (see Method). As shown in Figure 2, the values of the function $A_{X,X}(i, \text{Genes}_{\text{Bac}})$ are higher for multiples $(3i)$ where $i \in \{0,1,2 \dots\}$ than for multiples of $(3i + 1)$ or $(3i + 2)$, indicating that a circular code periodicity 0 modulo 3 is present in the genes for the circular code X . Note that the average values are around 0.1 as expected by Remark 5. While this result is new for a circular code, the observation of three base periodicity in genes of eukaryotes, bacteria, viruses, chloroplasts and mitochondria is classical and has been described in the past by several authors using different methods, in particular at the sequence level by Shepherd (1981a,b) and at the population level by Fickett (1982), Michel (1986, Fig. 1) and Arquès and Michel (1987a,b, 1990a,b).




Figure 2. Circular code periodicity 0 modulo 3 identified by the circular code autocorrelation function $A_{X,X}(i, \text{Genes}_{\text{Bac}})$ in bacterial genes. The abscissa represents the number i of letters N between X and itself (i.e. X and X), i varying from 0 to $i_{\text{max}} = 20$. The ordinate gives the occurrence probability $A_{X,X}(i, \text{Genes}_{\text{Bac}})$ (Equation (5)) of XN^iX in $\text{Genes}_{\text{Bac}}$.


Circular code periodicity in bacterial ribosomes

Next, we applied the circular code autocorrelation method to the 23S and 16S rRNA sequences from 103 bacterial organisms. In an initial study, we used the full length rRNA sequences, however no periodicity was observed (data not shown). Therefore, we divided the sequences into two parts corresponding to the 5' and 3' regions of each sequence, and calculated the circular code correlation functions for each region independently (Figure 3). Although no periodicity is identified in the 5' and 3' regions of the 23S rRNA (Figure 3A,B respectively) or the 5' region of the 16S rRNA (Figure 3C), we report these negative results in order to highlight the unicity of the circular code periodicity in the 3' region of the 16S rRNA (Figure 3D).


For the first time, the circular code correlation function $A_{X,X}(i, 16\text{S rRNA}_{766-1530})$ identifies a circular code periodicity 0 modulo 3 (up to $i = 15$) in the 3' region (766-1530) of bacterial 16S rRNA (Figure 3D). Obviously, since the 3' regions of 16S rRNA are relatively short, this modulo 3 periodicity is not regular compared to the one observed in genes (compared to Figure 2). However, an elementary calculus proves that this periodicity 0 modulo 3 is significant. Indeed, the probability that

$A_{X,X}(0, 16S \text{ rRNA}_{766-1530}) > A_{X,X}(1, 16S \text{ rRNA}_{766-1530})$ is equal to $1/2$. The probability that $A_{X,X}(i, 16S \text{ rRNA}_{766-1530}) > A_{X,X}(i-1, 16S \text{ rRNA}_{766-1530})$ and $A_{X,X}(i, 16S \text{ rRNA}_{766-1530}) > A_{X,X}(i+1, 16S \text{ rRNA}_{766-1530})$ with $i \equiv 0 \pmod{3}$ and $i > 0$ is equal to $1/3$. By assuming independence between the events, the probability of a periodicity 0 modulo 3 until $i = 15$ is equal to $P = \frac{1}{2} \cdot \left(\frac{1}{3}\right)^5 \approx 0.002$.


A. 23S rRNA, 5' region



B. 23S rRNA, 3' region



C. 16S rRNA, 5' region



D. 16S rRNA, 3' region




Figure 3. Circular code autocorrelation functions for bacterial 23S and 16S rRNA. The abscissa represents the number i of letters N between X and itself (i.e. X and X), i varying from 0 to $i_{max} = 20$. The ordinate gives the occurrence probability $A_{X,X}(i, F)$ (Equation (5)) of XN^iX in F . **A.** Circular code autocorrelation function $A_{X,X}(i, 23S \text{ rRNA}_{1-1447})$ in the 5' region (1-1447) of 23S bacterial rRNA. **B.** Circular code autocorrelation function $A_{X,X}(i, 23S \text{ rRNA}_{1448-2895})$ in the 3' region (1448-2895) of 23S rRNA. **C.** Circular code autocorrelation function $A_{X,X}(i, 16S \text{ rRNA}_{1-765})$ in the 5' region (1-765) of 16S rRNA. **D.** Circular code autocorrelation function $A_{X,X}(i, 16S \text{ rRNA}_{766-1530})$ in the 3' region (766-1530) of 16S rRNA, revealing the circular code periodicity 0 modulo 3.

The circular code periodicity 0 modulo 3 identified in the 3' region of 16S rRNA leads to a direct biological conclusion: a unit of genetic information based on trinucleotides exists in the 3' region of 16S rRNA, similarly to the protein coding genes.

Circular code periodicity in the bacterial tRNA of alanine

We also applied the same correlation function in a large set of bacterial tRNA genes corresponding to the 20 amino acids (see Method). For the first time, the circular code autocorrelation function

$A_{X,X}(i, \text{tRNA}_{\text{Ala}})$ identifies a circular code periodicity 0 modulo 3 (up to $i = 12$) in the tRNA of alanine (Figure 4). Obviously, this modulo 3 periodicity is noisy as the tRNA has a short length and constrained 2D and 3D structures. However, the calculus developed in the previous section proves that this periodicity 0 modulo 3 is significant and equal to $P = \frac{1}{2} \cdot \left(\frac{1}{3}\right)^4 \approx 0.006$.




Figure 4. Circular code periodicity 0 modulo 3 identified by the circular code autocorrelation function $A_{X,X}(i, \text{tRNA}_{\text{Ala}})$ in the bacterial tRNA of alanine. The abscissa represents the number i of letters N between X and itself (i.e. X and X), i varying from 0 to $imax = 20$. The ordinate gives the occurrence probability $A_{X,X}(i, \text{tRNA}_{\text{Ala}})$ (Equation (5)) of XN^iX in tRNA_{Ala} .

No modulo 3 periodicity is observed in the 19 remaining tRNAs. To date, we have no explanation for the absence of this signal property in the other tRNAs and additional studies will have to be considered in the future.

Structural and functional analysis of circular code periodicity in 16S rRNA

Modern ribosomes are highly sophisticated molecular machines, consisting of two subunits that come together during the initiation of protein synthesis, remain together as individual amino acids are added to the growing peptide, and finally separate again in conjunction with the release of the finished protein (Opron and Burton, 2018). Each subunit is a large nucleoprotein complex. In bacteria, the large subunit (LSU) contains the 23S rRNA and 5S rRNA, whereas the 16S rRNA makes up the bulk of the small subunit (SSU). The 16S rRNA is important for subunit association and translational accuracy. It consists of 1542 bases and the structural arrangement creates a 5' domain, central domain, 3' major domain, and 3' minor domain.

As illustrated in Figure 3, the observed circular code periodicity (0 modulo 3) in the 3' region (766-1530) of bacterial 16S rRNA covers part of the central domain (helices h24-h27), all of the 3' major domain (helices h28-h43) and part of the 3' minor domain (helix h44). Notably, the 3' major domain contains the decoding center and interacts with both tRNA and mRNA. The decoding center is widely accepted to be an essential building block of the primeval 'proto-ribosome' that was already present in

the Last Universal Common Ancestor (LUCA) (Petrov et al., 2015; Agmon, 2018), where it may have simply been a location to bind RNAs in an open structure configuration (de Farias et al., 2019).




Figure 5. Schema of the 2D structure of the bacterial 16S rRNA (*E. coli*), showing the classical division into 4 domains: 5' domain, central domain (C), 3' major (M) and 3' minor (m) domains. Interaction sites with tRNA are indicated by colored boxes, and numbering is according to *E. coli* sequence. The region shown in red corresponds to the sequence segment with the circular code periodicity (0 modulo 3).

The spatial organization of the circular code periodicity is shown in Figure 6. The periodicity is mainly localized in the SSU close to the interaction sites with the mRNA and tRNAs, but also extends into the body, along the interface with the LSU formed by helix 44 in the 3' minor domain (Figure 6A). No periodicity was observed in the LSU. Within the SSU, the periodicity region covers all of the head

(formed by the 3' major domain), and part of the central domain that forms the platform (Figure 6B). The mRNA and tRNAs lie across the neck (h28) of the SSU between the platform and the head. Finally, Figure 6C,D show the nucleotides close to the mRNA (<15 Å) and highlight the close packing of the periodicity around the decoding center, with 102 out of 134 (76%) 16S nucleotides within the periodicity region.




Figure 6. 3D structure of the bacterial ribosome (*E. coli*). A. rRNA of the LSU (beige) and SSU (gray), the region (red) with the circular code periodicity (0 modulo 3), the mRNA segment (green), and the A-site (cyan), P-site (teal) and E-site tRNAs (light teal). B. SSU rRNA (gray) with the periodicity region (red) and the mRNA segment (green). The mRNA lies across the neck of the SSU between the platform and the head. C. Nucleotides close to the mRNA (<15 Å) with the tRNAs (colored as in A), SSU RNA (gray) and periodicity region (red). D has been rotated 90° with respect to C.

Discussion

We have developed a new method that allows us to calculate exact probabilities of observing circular code autocorrelation, even for sequences with short lengths (as demonstrated in Appendix A). Using this method, we confirmed that the 20 trinucleotides of the *X* circular code present a periodicity 0 modulo 3 in the protein-coding regions of bacterial genomes. Furthermore, for the first time, we identify three base periodicity in ribosomal RNA sequences. This would imply that the trinucleotide may be a fundamental unit of information within rRNA sequences, and is line with numerous previous studies showing that some rRNA sequences contain protein-coding genes (reviewed in Root-Bernstein and Root-Bernstein, 2016).

Importantly, the observed periodicity is restricted to the 3' region of the 16S rRNA, which contains the decoding center and other interaction sites with the mRNA and tRNAs in A, P and E sites. Our 3D structural analysis shows that the periodicity region surrounds the mRNA channel between the head and the body of the SSU. Previously, we showed that this region contains a number of *X* motifs that are universally conserved in bacteria, but are also present in archaea and eukaryotic rRNA sequences (Dila et al., 2019b). This leads us to the question: do the *X* motifs in the 16S rRNA interact somehow with *X* motifs in the mRNA of protein-coding genes to regulate translation? Other mRNA-rRNA interactions are known to affect translation efficiency or quality. For example, hybridization between the Shine-Dalgarno sequence in the 5' UTR of bacterial mRNA and the anti-Shine-Dalgarno region of the 16S rRNA directs the ribosome to the start codon of the mRNA (Amin et al., 2018). Additional examples of mRNA-rRNA interactions include non-Shine-Dalgarno ribosome binding sites in the 5' UTR (Barendt et al., 2013), internal Shine-Dalgarno sequences (O'Connor et al., 2013) or recoding signals that direct ribosomal frameshifting (Atkins et al., 2016).

The periodicity property in the 16S 3' region largely corresponds to the 'proto-SSU' that has been proposed to represent the primordial ribosomal SSU (Harish and Caetano-Anollés, 2012; Caetano-Anollés, 2015; Petrov et al., 2015; Agmon, 2018). Thus, our results provide additional support for the hypothesis that the primordial coding system was RNA-based, and this RNA translation template then evolved to form the modern tRNA, mRNA and rRNA sequences (Chatterjee and Jadav, 2019; Root-Bernstein and Root-Bernstein, 2019). According to this theory, the initial replicator whose biomolecular activity initiated Darwinian evolution on Earth (Szathmáry, 2006; Yarus, 2011), consisted of short RNA oligonucleotides and was probably stabilized by small peptides containing amino acids such as glycine, alanine, aspartic acid or valine (Fournier et al., 2010; Maier et al., 2013; Attwater et al., 2018). Necessary features of such an RNA translation template include some level of specificity between nucleotide triplets and the amino acids (Kunnev and Gospodinov, 2018), and self-complementarity between nucleotides to allow replication (Banwell et al., 2018). The mathematical properties of the *X* circular code meet these requirements: (i) it provides a mapping between trinucleotides and the early amino acids, (ii) it is circular and has the capacity to detect the reading

frame, and (iii) it is self-complementary. Therefore, it is tempting to speculate that the TBP in protein-coding genes arose from the periodicity property of the *X* circular code in the primordial ribosome. Finally, the circular code autocorrelation function also allowed us to identify a circular code periodicity in the tRNA of alanine. The link between tRNAs and rRNAs has been highlighted by other groups and it is widely believed that rRNAs may have evolved by concatenation of tRNA-like molecules (e.g. Harish and Caetano-Anollés, 2012; de Farias et al., 2019). It is noteworthy that alanine, along with glycine, is generally predicted to be one of the most ancient amino acids to be included in the genetic code (Koonin, 2017; Demongeot and Seligmann, 2019c). Furthermore, we previously proposed that the comma-free code {GGC, GCC} was used initially to code Ala and Gly, and that this code quickly evolved to circular codes that included more and more amino acids (Dila et al., 2019b). A more in depth study of TBP in tRNA sequences is planned in the near future to determine whether a weaker periodicity property remains to be found in the tRNAs coding for other amino acids.

Acknowledgements

This work was supported by Institute funds from the French Centre National de la Recherche Scientifique and the University of Strasbourg. The authors would like to thank the BISTRO and BICS Bioinformatics Platforms for their assistance. This work was supported by the ANR under Grant Elixir-Excelebrate: GA-676559 and under RAinRARE: ANR-18-RAR3-0006-02.

Disclosure of interest

The authors report no conflict of interest.

References

- Agmon I. 2018. Hypothesis: Spontaneous advent of the prebiotic translation system via the accumulation of L-shaped RNA elements. *International Journal of Molecular Sciences* 19, E4021.
- Amin M.R., Yurovsky A., Chen Y., Skiena S., Futcher B. 2018. Re-annotation of 12,495 prokaryotic 16S rRNA 3' ends and analysis of Shine-Dalgarno and anti-Shine-Dalgarno sequences. *PLoS One* 13(8):e0202767.
- Arquès D.G., Michel C.J. 1987a. Study of a perturbation in the coding periodicity. *Mathematical Biosciences* 86, 1-14.
- Arquès D.G., Michel C.J. 1987b. A purine-pyrimidine motif verifying an identical presence in almost all gene taxonomic groups. *Journal of Theoretical Biology* 128, 457-461.
- Arquès D.G., Michel C.J. 1990a. Periodicities in coding and noncoding regions of the genes. *Journal of Theoretical Biology* 143, 307-318.
- Arquès D.G., Michel C.J. 1990b. A model of DNA sequence evolution. Part 1: Statistical features and classification of gene populations, 743-753. Part 2: Simulation model, 753-766. Part 3: Return of the model to the reality, 766-770. *Bulletin of Mathematical Biology* 52, 741-772.
- Arquès D.G., Michel C.J. 1996. A complementary circular code in the protein coding genes. *Journal of Theoretical Biology* 182, 45-58.
- Atkins J.F., Loughran G., Bhatt P.R., Firth A.E., Baranov P.V. 2016. Ribosomal frameshifting and transcriptional slippage: From genetic steganography and cryptography to adventitious use. *Nucleic Acids Research* 44, 7007-7078.
- Attwater J., Raguram A., Morgunov A.S., Gianni E., Holliger P. 2018. Ribozyme-catalysed RNA synthesis using triplet building blocks. *Elife*. 7, e35255.
- Banwell E.F., Piette B.M.A.G., Taormina A., Heddle J.G. 2018. Reciprocal nucleopeptides as the ancestral Darwinian self-replicator. *Molecular Biology and Evolution* 35, 404-416.
- Barendt P.A., Shah N.A., Barendt G.A., Kothari P.A., Sarkar C.A. 2013. Evidence for context-dependent complementarity of non-Shine-Dalgarno ribosome binding sites to Escherichia coli rRNA. *ACS Chemical Biology* 8, 958-966.
- Bernier C.R., Petrov A.S., Waterbury C.C., Jett J., Li F., Freil L.E., Xiong X., Wang L., Migliozzi B.L., HersHKovits E., Xue Y., Hsiao C., Bowman J.C., Harvey S.C., Grover M.A., Wartell Z.J., Williams L.D. 2014. RiboVision suite for visualization and analysis of ribosomes. *Faraday Discussions* 169, 195-207.
- Bloch D.P., McArthur B., Mirrop S. 1985. tRNA-rRNA sequence homologies: evidence for an ancient modular format shared by tRNAs and rRNAs. *Biosystems* 17, 209-225.
- Caetano-Anollés G. 2015. Ancestral insertions and expansions of rRNA do not support an origin of the ribosome in its peptidyl transferase center. *Journal Molecular Evolution* 80, 162-165.
- Chatterjee S., Yadav S. 2019. The origin of prebiotic information system in the peptide/RNA world: A simulation model of the evolution of translation and the genetic code. *Life* 9, E25.

- Chen B., Ji P. 2011. Visualization of the protein-coding regions with a self adaptive spectral rotation approach. *Nucleic Acids Research* 39, e3.
- Crick F.H., Brenner S., Klug A., Pieczek G. 1976. A speculation on the origin of protein synthesis. *Origins Life* 7, 389-397.
- de Farias S.T., Rêgo T.G., José M.V. 2019. Origin of the 16S ribosomal molecule from ancestor tRNAs. *Sci* 1, 8.
- Demongeot J., Moreira A. 2007. A possible circular RNA at the origin of life. *Journal of Theoretical Biology* 249, 314-324.
- Demongeot J., Seligmann H. 2019a. Spontaneous evolution of circular codes in theoretical minimal RNA rings. *Gene* 705, 95-102.
- Demongeot J., Seligmann H. 2019b. The uroboros theory of life's origin: 22-nucleotide theoretical minimal RNA rings reflect evolution of genetic code and tRNA-rRNA translation machineries. *Acta Biotheoretica* 67, 273-297.
- Demongeot J., Seligmann H. 2019c. Evolution of tRNA into rRNA secondary structures. *Gene Reports* 17, 100483.
- Dila G., Michel C.J., Poch O., Ripp R., Thompson J.D. 2019a. Evolutionary conservation and functional implications of circular code motifs in eukaryotic genomes. *Biosystems* 175, 57-74.
- Dila G., Ripp R., Mayer C., Poch O., Michel C.J., Thompson J.D. 2019b. Circular code motifs in the ribosome: a missing link in the evolution of translation? *RNA* 25, 1714-1730.
- Ding Y., Tang Y., Kwok C.K., Zhang Y., Bevilacqua P.C., Assmann S.M. 2014. In vivo genome-wide profiling of RNA secondary structure reveals novel regulatory features. *Nature* 505, 696-700.
- Eigen M., Winkler-Oswatitsch R. 1981 Transfer-RNA, an early gene? *Naturwissenschaften* 68, 282-292.
- El Soufi K., Michel C.J. 2014. Circular code motifs in the ribosome decoding center. *Computational Biology and Chemistry* 52, 9-17.
- El Soufi K., Michel C.J. 2015. Circular code motifs near the ribosome decoding center. *Computational Biology and Chemistry* 59, 158-176.
- El Soufi K., Michel C.J. 2016. Circular code motifs in genomes of eukaryotes. *Journal of Theoretical Biology* 408, 198-212.
- Esken S.T., Esken F.N., Kinghorn B., Ruvinsky A. 2004. Periodicity of DNA in exons. *BMC Molecular Biology* 5, 12.
- Fickett J.W. 1982. Recognition of protein coding regions in DNA sequences. *Nucleic Acids Research* 10, 5303-5318.
- Fimmel E., Strüngmann L. 2018. Mathematical fundamentals for the noise immunity of the genetic code. *Biosystems* 164, 186-198.
- Fournier G.P., Neumann J.E., Gogarten J.P. 2010. Inferring the ancient history of the translation machinery and genetic code via recapitulation of ribosomal subunit assembly orders. *PLoS One* 5, e9437.

- Guigó R., Agarwal P., Abril J.F., Burset M., Fickett J.W. 2000. An assessment of gene prediction accuracy in large DNA sequences. *Genome Research* 10, 1631-1642.
- Gutiérrez G., Oliver J.L., Marin A. 1994. On the origin of the periodicity of three in protein coding DNA sequences. *Journal of Theoretical Biology* 167, 413-414.
- Harish A., Caetano-Anollés G. 2012. Ribosomal history reveals origins of modern protein synthesis. *PLoS One* 7, e32776.
- Howe E.D., Song J.S. 2012. Categorical spectral analysis of periodicity in human and viral genomes *Biosystems* 107, 142-144.
- Johnson D.B., Wang L. 2010. Imprints of the genetic code in the ribosome. *Proceedings of the National Academy of Sciences U.S.A.* 107, 8298-8303.
- Koonin E.V. 2017. Frozen accident pushing 50: stereochemistry, expansion, and chance in the evolution of the genetic code. *Life* 7, 22.
- Kunnev D., Gospodinov A. 2018. Possible emergence of sequence specific RNA aminoacylation via peptide intermediary to initiate Darwinian evolution and code through origin of life. *Life* 8, E44.
- Maier U.G., Zauner S., Woehle C., Bolte K., Hempel F., Allen J.F., Martin W.F. 2013. Massively convergent evolution for ribosomal protein gene content in plastid and mitochondrial genomes. *Genome Biology and Evolution* 5, 2318-2329.
- Marhon S.A., Kremer S.C. 2011. Gene prediction based on DNA spectral analysis: a literature review. *Journal Computational Biology* 18, 639-676.
- Michel C.J. 1986. New statistical approach to discriminate between protein coding and non-coding regions in DNA sequences and its evaluation. *Journal of Theoretical Biology* 120, 223-236.
- Michel C.J. 2008. A 2006 review of circular codes in genes. *Computer and Mathematics with Applications* 55, 984-988.
- Michel C.J. 2012. Circular code motifs in transfer and 16S ribosomal RNAs: a possible translation code in genes. *Computational Biology and Chemistry* 37, 24-37.
- Michel C.J. 2013. Circular code motifs in transfer RNAs. *Computational Biology and Chemistry* 45, 17-29.
- Michel C.J. 2015. The maximal C^3 self-complementary trinucleotide circular code X in genes of bacteria, eukaryotes, plasmids and viruses. *Journal of Theoretical Biology* 380, 156-177.
- Michel C.J. 2017. The maximal C^3 self-complementary trinucleotide circular code X in genes of bacteria, archaea, eukaryotes, plasmids and viruses. *Life* 7, 20, 1-16.
- Michel C.J., Nguefack Ngoune V., Poch O., Ripp R., Thompson J.D. 2017. Enrichment of circular code motifs in the genes of the yeast *Saccharomyces cerevisiae*. *Life* 7, 52, 1-20.
- Nazarea A.D., Bloch D.P., Semrau A.C. 1985. Detection of a fundamental modular format common to transfer and ribosomal RNAs: second-order spectral analysis. *Proceedings of the National Academy of Sciences U.S.A.* 82, 5337-5341.

- O'Connor P.B., Li G.W., Weissman J.S., Atkins J.F., Baranov P.V. 2013. rRNA:mRNA pairing alters the length and the symmetry of mRNA-protected fragments in ribosome profiling experiments. *Bioinformatics*. 29, 1488-1491.
- Ohno S. 1988. Codon preference is but an illusion created by the construction principle of coding sequences. *Proceedings of the National Academy of Sciences U.S.A.* 85, 4378-4382.
- Opron K., Burton Z.F. 2018. Ribosome structure, function, and early evolution. *International Journal of Molecular Sciences* 20, E40.
- Petrov A.S., Gulen B., Norris A.M., Kovacs N.A., Bernier C.R., Lanier K.A., Fox G.E., Harvey S.C., Wartell R.M., Hud N.V., Williams L.D. 2015. History of the ribosome and the origin of translation. *Proceedings of the National Academy of Sciences U.S.A.* 112, 15396-15401.
- Rodin A.S., Szathmáry E., Rodin S.N. 2011. On origin of genetic code and tRNA before translation. *Biology Direct* 22, 6-14.
- Root-Bernstein R., Root-Bernstein M. 2016. The ribosome as a missing link in prebiotic evolution II: Ribosomes encode ribosomal proteins that bind to common regions of their own mRNAs and rRNAs. *Journal of Theoretical Biology* 397, 115-127.
- Root-Bernstein R., Root-Bernstein M. 2019. The ribosome as a missing link in prebiotic evolution III: Over-representation of tRNA- and rRNA-like sequences and plieofunctionality of ribosome-related molecules argues for the evolution of primitive genomes from ribosomal RNA modules. *International Journal Molecular Sciences* 20, E140.
- Sánchez J., López-Villaseñor I. 2006. A simple model to explain three-base periodicity in coding DNA. *FEBS Letters* 580, 6413-6422.
- Shabalina S.A., Ogurtsov A.Y., Spiridonov N.A. 2006. A periodic pattern of mRNA secondary structure created by the genetic code. *Nucleic Acids Research* 34, 2428-2437.
- Shepherd J.C.W. 1981a. Periodic correlations in DNA sequences and evidence suggesting their evolutionary origin in a comma-less genetic code. *Journal Molecular Evolution* 17, 94-102.
- Shepherd J.C.W. 1981b. Method to determine the reading frame of a protein from the purine/pyrimidine genome sequence and its possible evolutionary justification. *Proceedings of the National Academy of Sciences U.S.A.* 78, 1596-1600.
- Szathmáry E. 2006. The origin of replicators and reproducers. *Philosophical Transactions of the Royal Society B* 361, 1761-1776.
- Tiwari S., Ramachandran S., Bhattacharya S., Ramaswamy R. 1997. Prediction of probable genes by Fourier analysis of genomic sequences. *Computer Applications in the Biosciences* 263-270.
- Trifonov E.N. 1987. Translation framing code and frame-monitoring mechanism as suggested by the analysis of mRNA and 16S rRNA nucleotide sequences. *Journal Molecular Biology* 194, 643-652.
- Trifonov E.N. 1998. 3-, 10.5-, and 400-base periodicities in genome sequences. *Physica A* 249, 511-516.
- Tsonis A.A., Elsner J.B., Tsonis P.A. 1991. Periodicity in DNA coding sequences: implications in gene evolution. *Journal of Theoretical Biology* 151, 323-331.

- Yarus M. 2011. Getting Past the RNA World: The initial Darwinian ancestor. *Cold Spring Harbor Perspectives Biology*, a003590.
- Yin C., Yau S. 2005. A Fourier characteristic of coding sequences: origins and a non-Fourier approximation. *Journal Computational Biology* 12, 1153.
- Yin C., Yau S. 2007. Prediction of protein coding regions by the 3-base periodicity analysis of a DNA sequence. *Journal of Theoretical Biology* 247, 687-694.

Accepted Manuscript

APPENDIX A

We compare the circular code autocorrelation method developed here (Equations (3) and (5)) with the two classical correlation methods used in Fourier analysis. After recalling these classical formulas, we extend them to an i -motif $mN^i m'$, i.e. 2 motifs m and m' separated by i , $i \in \{0, \dots, imax\}$, any letters $N \in B$.

A.1. Classical correlation methods

The power spectral density is the Fourier transform of the correlation function which is classically estimated in a discrete signal on a word $w = l_1 l_2 \dots l_{|w|}$ according to

$$\hat{A}_{l,l}(i, w) = \frac{1}{|w|} \sum_{p=1}^{|w|-i} \delta_l(p) \cdot \delta_l(p+i), \quad i = 0, \dots, |w| - 1 \quad (6)$$

where $\delta_l(p) = \begin{cases} 1 & \text{if the letter in position } p \text{ is } l \\ 0 & \text{otherwise} \end{cases}$.

This estimate $\hat{A}_{l,l}(i, w)$ is so called "biased", because when the correlation lag i approaches the length $|w|$, it differs from the exact probability calculus. The estimate $\hat{A}_{l,l}(i, w)$ has some drastic effects with short words (see the examples in Section A.3). Thus, another estimate is also proposed by normalizing the denominator

$$\hat{A}'_{l,l}(i, w) = \frac{1}{|w| - i} \sum_{p=1}^{|w|-i} \delta_l(p) \cdot \delta_l(p+i), \quad i = 0, \dots, |w| - 1 \quad (7)$$

where $\delta_l(p)$ is defined in Equation (6).

While this estimate $\hat{A}'_{l,l}(i, w)$ gives exact probability calculus with long words, it becomes less accurate with short words (see the examples in Section A.3).

A.2. Extension of classical correlation methods to an i -motif

In order to compare Equation (6) with Equation (3) associated with the i -motif $XN^i X$, we omit the Fourier case $i = 0$ and we trivially extend Equation (6) to an i -motif $mN^i m'$ separated by i any letters $N \in B$

$$\hat{B}_{m,m'}(i, w) = \frac{1}{|w| - |mm'| + 1} \sum_{p=1}^{|w|-i-|mm'|+1} \delta_m(p) \cdot \delta_{m'}(p + |m| + i), \quad i = 0, \dots, |w| - |mm'| \quad (8)$$

where $\delta_m(p)$ and $|mm'|$ are defined in Equation (3).

Note that the case $i = 0$ does not have the same meaning for the Equations (6) and (8). Similarly, Equation (7) is extended to an i -motif $XN^i X$ as follows

$$\hat{B}'_{m,m'}(i, w) = \frac{1}{|w| - i - |mm'| + 1} \sum_{p=1}^{|w|-i-|mm'|+1} \delta_m(p) \cdot \delta_{m'}(p + |m| + i), \quad i = 0, \dots, |w| - |mm'| \quad (9)$$

where $\delta_m(p)$ and $|mm'|$ are defined in Equation (3).

Equation (8) (not shown for Equation (9)) easily extends to a sequence population F as follows

$$\hat{B}_{m,m'}(i, F) = \frac{1}{|F|} \sum_{w \in F} \hat{B}_{m,m'}(i, w), \quad i = 0, \dots, |w| - 1 \quad (10)$$

A.3. Application examples

We give computation examples of the correlation function $A_{m,m'}(i, w)$ on the sequences $(RRY)^+$ and $(RNY)^+$ by choosing, for sake of simplicity, the letters $m = m' = R$ on the 2-letter alphabet $B' = \{R, Y\}$ ($N = \{R, Y\}$).

A.3.1. Sequence $w = (RRY)^+$

In this first example, we apply the correlation function $A_{R,R}(i, w)$ on the sequence $w = (RRY)^+ = RRYRRY \dots$

(i) Exact calculus of $A_{R,R}(i, (RRY)^+)$ leads trivially to the following solution

$$A_{R,R}(i, (RRY)^+) = \begin{cases} \frac{1}{3} \approx 0.3333 & \text{for } i \equiv 0 \pmod{3} \\ \frac{1}{3} \approx 0.3333 & \text{for } i \equiv 1 \pmod{3} \\ \frac{2}{3} \approx 0.6667 & \text{for } i \equiv 2 \pmod{3} \end{cases}$$

(ii) The computation of $A_{R,R}(i, (RRY)^n)$ (Equation (3)) in a simulated sequence of $n = 33$ consecutive trinucleotides RRY is associated with the exact probabilities (Table 1 and Figure 7A).

Table 1. Correlation function $A_{R,R}(i, (RRY)^n)$ in a simulated sequence $(RRY)^n$ of trinucleotide length $n \in \{33, 333, 3333\}$ where i represents the number of letters N between R and itself, i varying from 0 to $imax = 7$, and $A_{R,R}(i, (RRY)^n)$ of RN^iR in $(RRY)^n$ is computed according to $A_{R,R}(i, (RRY)^{33})$ (Equation (3)) and $\hat{B}'_{m,m'}(i, (RRY)^n)$ (Equation (9)).

i	Equation (3) with $(RRY)^{33}$	Equation (9) with $(RRY)^{33}$	Equation (9) with $(RRY)^{333}$	Equation (9) with $(RRY)^{3333}$
0	0.3333	0.3366	0.3337	0.3334
1	0.3333	0.3300	0.3330	0.3333
2	0.6667	0.6667	0.6667	0.6667
3	0.3333	0.3367	0.3337	0.3334
4	0.3333	0.3299	0.3330	0.3333
5	0.6667	0.6667	0.6667	0.6667
6	0.3333	0.3368	0.3337	0.3334
7	0.3333	0.3298	0.3330	0.3333




Figure 7. Correlation function $A_{R,R}(i, (RRY)^n)$ in a simulated sequence $(RRY)^n$. The abscissa represents the number i of letters N between R and itself (i.e. R and R), i varying from 0 to $imax = 20$. The ordinate gives the occurrence probability $A_{R,R}(i, (RRY)^n)$ of RN^iR in $(RRY)^n$ computed according to **A**: $A_{R,R}(i, (RRY)^{33})$ (Equation (3)) and **B**: $\hat{B}'_{m,m'}(i, (RRY)^{33})$ (Equation (8)).

(iii) The computation of $\hat{B}_{m,m'}(i, (RRY)^n)$ (Equation (8) with $m = m' = R$ and $|mm'| = 2$) in a simulated sequence $(RRY)^n$ of trinucleotide length $n = 33$ (Figure 7B) strongly differs from the exact probabilities (Table 1 and Figure 7A).

(iv) As expected, the computation of $\hat{B}_{m,m'}(i, (RRY)^n)$ (Equation (9) with $m = m' = R$ and $|mm'| = 2$) of $(RRY)^n$ leads to values close to the exact probabilities with a simulated sequence of short length ($n = 33$ trinucleotides) and to the exact probabilities with a simulated sequence of large length ($n = 3333$ trinucleotides) (Table 1).

In summary, only Equation (3) allows to compute exact probabilities with a sequence of a short length, i.e. about 100 nucleotides which is the length of a tRNA for example.

A.3.2. Sequence $w = (RNY)^+$

In this second example, we show that even Equation (3) is not enough to retrieve exact probabilities in a noisy sequence of short length. However, Equation (5) extending Equation (3) to a sequence population, retrieves the exact probabilities. We will apply the correlation function $A_{R,R}(i, w)$ on the sequence $w = (RNY)^+ = RNYRNY \dots$, N being randomly chosen between R and Y with equiprobability ($1/2$) for sake of simplicity, in order to introduce (basic) noise and evaluate the behavior of the computed correlation functions.

(i) Exact calculus of $A_{R,R}(i, (RNY)^+)$ leads trivially to the following solution

$$A_{R,R}(i, (RNY)^+) = \begin{cases} \frac{1}{6} \approx 0.1667 & \text{for } i \equiv 0 \pmod{3} \\ \frac{1}{6} \approx 0.1667 & \text{for } i \equiv 1 \pmod{3} \\ \frac{5}{12} \approx 0.4167 & \text{for } i \equiv 2 \pmod{3} \end{cases}$$

(ii) The computation of $A_{R,R}(i, (RNY)^n)$ (Equation (3)) in a simulated sequence of $n = 33$ consecutive trinucleotides RNY is close to the exact probabilities (Figure 8A).

(iii) The computation of $\hat{B}_{m,m'}(i, (RNY)^{33})$ (Equation (8) with $m = m' = R$ and $|mm'| = 2$) in a simulated sequence $(RNY)^n$ of trinucleotide length $n = 33$ (Figure 8B) again differs from the exact probabilities.




Figure 8. Correlation function $A_{R,R}(i, (RNY)^n)$ in a simulated sequence $(RNY)^n$. The abscissa represents the number i of letters N between R and itself, i varying from 0 to $imax = 20$. The ordinate gives the occurrence probability $A_{R,R}(i, (RNY)^n)$ of RN^iR in $(RNY)^n$ computed according to **A**: $A_{R,R}(i, (RNY)^{33})$ (Equation (3)) and **B**: $\hat{B}_{m,m'}(i, (RNY)^{33})$ (Equation (8)).

We continue the example by showing the importance of a sequence population when the sequences of short lengths are noisy. As an illustration example, we chose a population with $|F| = 100$ sequences of $n = 33$ consecutive trinucleotides RNY , noted $(RNY)_F^n = (RNY)_{100}^{33}$.

(iii) The computation of $A_{R,R}(i, (RNY)_F^n)$ (Equation (5)) in simulated sequences $(RNY)_{100}^{33}$ retrieves the exact probabilities (Figure 9A).

(iv) The computation of $\hat{B}_{m,m'}(i, (RNY)_F^n)$ (Equation (10) with $m = m' = R$ and $|mm'| = 2$) in simulated sequences $(RNY)_{100}^{33}$ (Figure 9B) again differs from the exact probabilities significantly.




Figure 9. Correlation function $A_{R,R}(i, (RNY)_F^n)$ in simulated sequences $(RNY)_{100}^{33}$ with $|F| = 100$ sequences of $n = 33$ consecutive trinucleotides RNY . The abscissa represents the number i of letters N between R and itself, i varying from 0 to $imax = 20$. The ordinate gives the occurrence probability $A_{R,R}(i, (RNY)_F^n)$ of RN^iR in $(RNY)_F^n$ according to **A**: $A_{R,R}(i, (RNY)_{100}^{33})$ (Equation (5)) and **B**: $\hat{B}_{m,m'}(i, (RNY)_{100}^{33})$ (Equation (10)).

In conclusion, the correlation method developed in Section Materials and methods allows to retrieve exact probabilities with noisy sequences of short lengths, and thus is well adapted to study rRNAs and tRNAs.

APPENDIX B

Table 1. Bacterial organisms used in the ribosomal RNA multiple sequence alignments.

<i>Actinoplanes utahensis</i>	<i>Myxococcus xanthus</i>
<i>Aeromonas ichthiosmia</i>	<i>Neisseria meningitidis</i>
<i>Agrobacterium tumefaciens</i>	<i>Nitrospira moscoviensis</i>
<i>Aquifex aeolicus</i>	<i>Paracoccus denitrificans</i>
<i>Bacillus cereus</i>	<i>Pelobacter acetylenicus</i>
<i>Bacillus globisporus</i>	<i>Pirellula marina</i>
<i>Bacillus halodurans</i>	<i>Piscirickettsia salmonis</i>
<i>Bacillus licheniformis</i>	<i>Polynucleobacter necessarius</i>
<i>Bacteroides fragilis</i>	<i>Propionigenium modestum</i>
<i>Bartonella quintana</i>	<i>Proteus vulgaris</i>
<i>Bifidobacterium bifidum</i>	<i>Pseudomonas aeruginosa</i>
<i>Brevundimonas diminuta</i>	<i>Pseudomonas fluorescens</i>
<i>Buchnera aphidicola</i>	<i>Psychrobacter pacificensis</i>
<i>Caedibacter caryophila</i>	<i>Rahnella aquatilis</i>
<i>Caloramator indicus</i>	<i>Rhizobium</i> sp.
<i>Chlorobium vibrioforme</i>	<i>Rhizobium tropici</i>
<i>Chlorogloeopsis</i> sp	<i>Rhodopseudomonas palustris</i>
<i>Clavibacter xyli</i>	<i>Rhodospirillum rubrum</i>
clone CS981 (X81184)	<i>Rhodothermus marinus</i>
clone SAR (U34043)	<i>Rice yellow dwarf phytoplasma</i>
<i>Clostridium ghoni</i>	<i>Rubrobacter radiotolerans</i>
<i>Clostridium hastiforme</i>	<i>Ruminobacter amylophilus</i>
<i>Clostridium sphenoides</i>	<i>Saccharococcus thermophilus</i>
<i>Coprothermobacter proteolyticus</i>	<i>Salinicoccus roseus</i>
<i>Deferribacter thermophilus</i>	<i>Sargasso Sea (X52169)</i>
<i>Desulfacinum infernum</i>	<i>Serratia marcescens</i>
<i>Desulfobacterium frappieri</i>	<i>Shewanella algae</i>
<i>Desulfofustis glycolicus</i>	<i>Simkania negevensis</i>
<i>Desulfogalobium retbaense</i>	<i>Sinorhizobium meliloti</i>
<i>Desulfotalea psychrophila</i>	<i>Spirochaeta</i> sp.
<i>Desulfotomaculum thermosapovorans</i>	<i>Spirulina platensis</i>
<i>Desulfurella acetivorans</i>	<i>Sporobacter termitidis</i>
<i>Dichelobacter nodosus</i>	<i>Staphylococcus condimenti</i>
endosymbiont of L29265	<i>Streptococcus macedonicus</i>
epibiont of L35522	<i>Streptococcus pyogenes</i>
<i>Escherichia coli</i>	<i>Streptomyces acidiscabies</i>
<i>Frankia</i> sp.	<i>Streptomyces sampsonii</i>
<i>Geotoga subterranea</i>	<i>Sulfobacillus thermosulfidooxidans</i>
<i>Glycaspis brimblecombei</i> (AF263561)	symbiont S (M27040)
<i>Haloanaerobium lacuroseus</i>	<i>Synechocystis</i> PCC6803
<i>Halomonas</i> sp. NIBH P1H25	<i>Syntrophus buswellii</i>
<i>Helicobacter pylori</i>	<i>Thermomonospora chromogena</i>
<i>Kineococcus like bacterium</i> AS2960	<i>Thermotoga maritima</i>
<i>Lactobacillus acidophilus</i>	uncultured bacterium (AY212656)
<i>Lactococcus lactis</i>	uncultured <i>Pseudomonas</i> sp (DQ234150)
<i>Lactosphaera pasteurii</i>	<i>Ureaplasma urealyticum</i>
<i>Legionella lytica</i>	<i>Vibrio vulnificus</i>
<i>Magnetobacterium bavaricum</i>	<i>Xylella fastidiosa</i>
<i>Mesorhizobium loti</i>	<i>Zoogloea ramigera</i>
<i>Moraxella lacunata</i>	<i>Zoogloea ramigera</i>
<i>Mycobacterium leprae</i>	<i>Zymomonas mobilis</i>
<i>Mycoplasma capricolum</i>	

Table 2. Bacterial tRNA sequences used in the analysis.

Amino acid	No. of sequences	Amino acid	No. of sequences	Amino acid	No. of sequences
Ala	361	Gly	406	Pro	317
Arg	329	His	158	Ser	707
Asn	197	Ile	204	Thr	427
Asp	181	Leu	688	Trp	163
Cys	150	Lys	260	Tyr	172
Gln	229	Met	511	Val	337
Glu	237	Phe	173		

Accepted Manuscript