



**HAL**  
open science

# Neutrosophic Linear Diophantine Equations With Two Variables

Hasan Sankari, Mohammad Abobala

► **To cite this version:**

Hasan Sankari, Mohammad Abobala. Neutrosophic Linear Diophantine Equations With Two Variables. Neutrosophic Sets and Systems, 2020. <hal-03081126>

**HAL Id: hal-03081126**

**<https://hal.science/hal-03081126v1>**

Submitted on 18 Dec 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



# Neutrosophic Linear Diophantine Equations With Two Variables

Hasan Sankari<sup>1</sup> and Mohammad Abobala<sup>2</sup>

<sup>1</sup> Tishreen University, Faculty of Science, Department of Mathematics, Lattakia, Syria

<sup>2</sup> Tishreen University, Faculty of Science, Department of Mathematics, Lattakia, Syria

<sup>1</sup>e-mail: Hasan.Sankari2@gmail.com

<sup>2</sup>e-mail: mohammadabobala777@gmail.com

**Abstract:** This paper is devoted to study for the first time the neutrosophic linear Diophantine equations with two variables in the neutrosophic ring of integers  $Z(I)$ , and refined neutrosophic ring of integers  $Z(I_1, I_2)$ . This work introduces an algorithm to solve the linear Diophantine equation  $AX + BY = C$  in  $Z(I), Z(I_1, I_2)$ .

**Keywords:** Neutrosophic ring, refined neutrosophic ring, neutrosophic linear Diophantine equation, refined neutrosophic linear Diophantine equation.

---

## 1. Introduction

Neutrosophy is a new kind of logic founded by F. Smarandache to deal with the indeterminacy in nature, mathematics and reality. It plays an interesting role in the progression of algebraic studies. Many neutrosophical algebraic structures were introduced and handled such as neutrosophic groups, neutrosophic rings, refined neutrosophic rings, and n-refined neutrosophic rings. See [1,2,3,4,5,6,8,10,11]. On the other hand neutrosophic sets were used to deal with health care [12], decision making [13], financial goals [14], computer science, and industry [15,16,17,18,20]. Recently, the interesting in neutrosophic number theory has increased. Relationships between neutrosophic rings and refined neutrosophic rings were studied in [1]. Also, some number theoretical concepts were presented in the neutrosophic ring of integers  $Z(I)$  such as division, primes and factors [7]. The theory of neutrosophic numbers is concerning with properties of neutrosophic integers, by the same, refined neutrosophic number theory is dealing with the properties of refined neutrosophic

integers. One of the most important number theoretical concepts is the concept of linear Diophantine equations, these equations were solved in the case of classical integers [9]. Through this paper, we aim to find an algorithm to solve such equations in the case of neutrosophic integers and refined neutrosophic integers by using classical number theoretical methods, where a relationship between neutrosophic equations and classical equations is described.

This work continues the efforts of establishing neutrosophical number theory. It studies the concept of linear Diophantine equations with two variables with respect to neutrosophic integers and refined neutrosophic integers. We determine the sufficient condition for the solvability of these equations and introduce an algorithm which gives the solution in easy way.

## 2. Preliminaries

### Theorem 2.1: [9]

Let  $AX + BY = C$  be a linear Diophantine equation, where  $A, B, C \in \mathbb{Z}$ . Then it is solvable if and only if  $\gcd(A, B) | C$ . To check the solution's form of classical linear Diophantine equation based on Euclidean division theorem in the ring of integers  $\mathbb{Z}$ , see [9].

### Definition 2.2:[6]

Let  $(R, +, \times)$  be a ring,  $R(I) = \{a + bI \mid a, b \in R\}$  is called the neutrosophic ring where  $I$  is a neutrosophic element with condition  $I^2 = I$ .

If  $R = \mathbb{Z}$ , then  $R(I)$  is called the neutrosophic ring of integers.

### Definition 2.3:[4]

Let  $(R, +, \times)$  be a ring,  $(R(I_1, I_2), +, \times)$  is called a refined neutrosophic ring generated by  $R, I_1, I_2$ .

If  $R = \mathbb{Z}$ , then  $(R(I_1, I_2), +, \times)$  is called the refined neutrosophic ring of integers.

### Definition 2.4: [5]

Let  $(G, *)$  be a group. Then the neutrosophic group is generated by  $G$  and  $I$  under  $*$  denoted by  $N(G) = \langle G \cup I, * \rangle$ .

$I$  is called the indeterminate (neutrosophic element) with the property  $I^2 = I$ .

### 3. Main results

#### Definition 3.1:

Let  $Z(I) = \{a + bI; a, b \in \mathbb{Z}\}$  be the neutrosophic ring of integers. The neutrosophic linear Diophantine equation with two variables is defined as follows:

$$AX + BY = C; A, B, C \in Z(I).$$

#### Theorem 3.2:

Let  $Z(I) = \{a + bI; a, b \in \mathbb{Z}\}$  be the neutrosophic ring of integers. The neutrosophic linear

Diophantine equation  $AX + BY = C$  with two variables  $X = x_1 + x_2I, Y = y_1 + y_2I$ , where

$A = a_1 + a_2I, B = b_1 + b_2I$  is equivalent to the following two classical Diophantine equations:

$$(1) a_1x_1 + b_1y_1 = c_1.$$

$$(2) (a_1 + a_2)(x_1 + x_2) + (b_1 + b_2)(y_1 + y_2) = c_1 + c_2.$$

Proof:

It is sufficient to show that  $AX + BY = C$  implies (1) and (2).

$AX + BY = C$  is equivalent to:

$$(a_1 + a_2I)(x_1 + x_2I) + (b_1 + b_2I)(y_1 + y_2I) = c_1 + c_2I, \text{ by easy computing we find}$$

$$[a_1x_1 + b_1y_1] + [a_1x_2 + a_2x_1 + a_2x_2 + b_1y_2 + b_2y_1 + b_2y_2]I = c_1 + c_2I, \text{ hence}$$

$a_1x_1 + b_1y_1 = c_1$ , and  $a_1x_2 + a_2x_1 + a_2x_2 + b_1y_2 + b_2y_1 + b_2y_2 = c_2$ . We can see that we get equation

(1). For equation (2) we take

$$a_1x_2 + a_2x_1 + a_2x_2 + b_1y_2 + b_2y_1 + b_2y_2 = c_2, \text{ by adding equation (1) to the two sides we obtain}$$

$a_1x_1 + b_1y_1 + a_1x_2 + a_2x_1 + a_2x_2 + b_1y_2 + b_2y_1 + b_2y_2 = c_1 + c_2$ , which implies equation (2)

$$(a_1 + a_2)(x_1 + x_2) + (b_1 + b_2)(y_1 + y_2) = c_1 + c_2.$$

The following theorem determines the criteria for the solvability of neutrosophic linear Diophantine equation.

**Theorem 3.3:**

Let  $Z(I) = \{a + bI; a, b \in \mathbb{Z}\}$  be the neutrosophic ring of integers. The neutrosophic linear

Diophantine equation  $AX + BY = C$  with two variables  $X = x_1 + x_2I, Y = y_1 + y_2I$  and

$A = a_1 + a_2I, B = b_1 + b_2I$  is solvable if and only if  $\gcd(a_1, b_1) | c_1, \gcd(a_1 + a_2, b_1 + b_2) | c_1 + c_2$ .

Proof:

By Theorem 2.1, we can solve the neutrosophic linear Diophantine equation by solving (1) and (2).

Equation (1) is solvable if and only if  $\gcd(a_1, b_1) | c_1$  according to Theorem 2.1.

Equation (2) is solvable if and only if  $\gcd(a_1 + a_2, b_1 + b_2) | c_1 + c_2$  according to Theorem 2.1.

Thus our proof is complete.

**Example 3.4:**

(a) The neutrosophic Diophantine equation  $(2 + 2I)X + (3 + 4I)Y = 5 + 5I$  is solvable, that is because

$$\gcd(2,3) | 5, \text{ and } \gcd(4,7) | 10.$$

(b) The neutrosophic Diophantine equation  $(2 + 3I)X + (4 + 5I)Y = 5 + I$  is not solvable, since  $\gcd(2,4) = 2$  does not divide 5.

Now, we describe an algorithm to solve a neutrosophic linear Diophantine equation  $AX + BY = C$ .

**Remark 3.5:**

Let  $Z(I) = \{a + bI; a, b \in \mathbb{Z}\}$  be the neutrosophic ring of integers. Consider a neutrosophic linear

Diophantine equation  $AX + BY = C$  with two variables  $X = x_1 + x_2I, Y = y_1 + y_2I$  and

$A = a_1 + a_2I, B = b_1 + b_2I$ . To solve this equation follow these steps:

(a) Check the solvability of  $AX + BY = C$  by Theorem 3.3.

(b) Solve  $a_1x_1 + b_1y_1 = c_1$ .

(c) Solve  $(a_1 + a_2)(x_1 + x_2) + (b_1 + b_2)(y_1 + y_2) = c_1 + c_2$ .

(d) Compute  $x_2, y_2$ .

**Example 3.6:**

The neutrosophic Diophantine equation  $(2 + 2I)X + (3 + 4I)Y = 5 + 5I$  is solvable according to Example 3.4 .

$2x_1 + 3y_1 = 5$  is a classical linear Diophantine equation. It has a solution  $x_1 = 4, y_1 = -1$ .

$(2 + 2)(x_1 + x_2) + (3 + 4)(y_1 + y_2) = 5 + 5$ , i.e  $4M + 7N = 10; M = x_1 + x_2, N = y_1 + y_2$ . It is a

classical linear Diophantine equation with  $M, N$  as variables. It has a solution  $M = -1, N = 2$ .

$x_2 = M - x_1 = -5, y_2 = N - y_1 = 3$ , thus the equation  $(2 + 2I)X + (3 + 4I)Y = 5 + 5I$  has a solution

$X = 4 - 5I, Y = -1 + 3I$ .

**Definition 3.7:**

Let  $Z(I_1, I_2) = \{(a, bI_1, cI_2); a, b, c \in Z\}$  be the refined neutrosophic ring of integers. The refined neutrosophic linear Diophantine equation with two variables is defined as follows:

$$AX + BY = C; A, B, C \in Z(I_1, I_2).$$

**Theorem 3.8:**

Let  $Z(I_1, I_2) = \{(a, bI_1, cI_2); a, b, c \in Z\}$  be the refined neutrosophic ring of integers,

$AX + BY = C; A, B, C \in Z(I_1, I_2)$  be a refined neutrosophic linear Diophantine equation, where

$$X = (x_0, x_1I_1, x_2I_2), Y = (y_0, y_1I_1, y_2I_2), A = (a_0, a_1I_1, a_2I_2),$$

$B = (b_0, b_1I_1, b_2I_2), C = (c_0, c_1I_1, c_2I_2)$ . Then  $AX + BY = C$  is equivalent to the following three

Diophantine equations:

(1)  $a_0x_0 + b_0y_0 = c_0$ .

$$(2) (a_0 + a_2)(x_0 + x_2) + (b_0 + b_2)(y_0 + y_2) = c_0 + c_2.$$

$$(3) (a_0 + a_1 + a_2)(x_0 + x_1 + x_2) + (b_0 + b_1 + b_2)(y_0 + y_1 + y_2) = c_0 + c_1 + c_2.$$

Proof:

By replacing  $A, B, C, X, Y$  we find

$$AX = (a_0, a_1 I_1, a_2 I_2)(x_0, x_1 I_1, x_2 I_2) =$$

$$(a_0 x_0, [a_0 x_1 + a_1 x_0 + a_1 x_1 + a_1 x_2 + a_2 x_1] I_1, [a_0 x_2 + a_2 x_0 + a_2 x_2] I_2),$$

$$BY = (b_0, b_1 I_1, b_2 I_2)(y_0, y_1 I_1, y_2 I_2) =$$

$$(b_0 y_0, [b_0 y_1 + b_1 y_0 + b_1 y_1 + b_1 y_2 + b_2 y_1] I_1, [b_0 y_2 + b_2 y_0 + b_2 y_2] I_2), \text{ thus the equation}$$

$AX + BY = C$  implies

$$(*) a_0 x_0 + b_0 y_0 = c_0. \text{ (Equation (1)).}$$

$$(**) a_0 x_2 + a_2 x_0 + a_2 x_2 + b_0 y_2 + b_2 y_0 + b_2 y_2 = c_2.$$

$$(***) a_0 x_1 + a_1 x_0 + a_1 x_1 + a_1 x_2 + a_2 x_1 + b_0 y_1 + b_1 y_0 + b_1 y_1 + b_1 y_2 + b_2 y_1 = c_1.$$

By adding (\*) to (\*\*) we get  $(a_0 + a_2)(x_0 + x_2) + (b_0 + b_2)(y_0 + y_2) = c_0 + c_2$ . (Equation (2)).

By adding (2) to (\*\*\*) we get

$$(a_0 + a_1 + a_2)(x_0 + x_1 + x_2) + (b_0 + b_1 + b_2)(y_0 + y_1 + y_2) = c_0 + c_1 + c_2. \text{ (Equation (3)).}$$

### Theorem 3.9:

Let  $Z(I_1, I_2) = \{(a, b I_1, c I_2); a, b, c \in Z\}$  be the refined neutrosophic ring of integers,

$AX + BY = C; A, B, C \in Z(I_1, I_2)$  be a refined neutrosophic linear Diophantine equation, where

$$X = (x_0, x_1 I_1, x_2 I_2), Y = (y_0, y_1 I_1, y_2 I_2), A = (a_0, a_1 I_1, a_2 I_2),$$

$B = (b_0, b_1 I_1, b_2 I_2), C = (c_0, c_1 I_1, c_2 I_2)$ . Then  $AX + BY = C$  is solvable if and only if:

$$(a) \gcd(a_0, b_0) | c_0.$$

$$(b) \gcd(a_0 + a_2, b_0 + b_2) | c_0 + c_2.$$

$$(c) \gcd(a_0 + a_1 + a_2, b_0 + b_1 + b_2) | c_0 + c_1 + c_2.$$

The proof is similar to that of Theorem 3.3.

**Example 3.10:**

(a) Consider the refined neutrosophic linear Diophantine equation

$$(1, 2I_1, 3I_2).X + (3, 3I_1, 8I_2)Y = (2, 4I_1, I_2), \text{ we have}$$

$$\gcd(1, 3) = 1 | 2, \gcd(1 + 3, 3 + 8) = \gcd(4, 11) = 1 | (2 + 1 = 3),$$

$\gcd(1 + 2 + 3, 3 + 3 + 8) = \gcd(6, 14) = 2$  which does not divide  $2 + 4 + 1 = 7$ , thus it is not solvable.

(b) Consider the refined neutrosophic linear Diophantine equation

$$(1, 2I_1, 3I_2).X + (3, 3I_1, 8I_2)Y = (2, 4I_1, 2I_2), \text{ we have}$$

$$\gcd(1, 3) = 1 | 2, \gcd(1 + 3, 3 + 8) = \gcd(4, 11) = 1 | (2 + 2 = 4),$$

$\gcd(1 + 2 + 3, 3 + 3 + 8) = \gcd(6, 14) = 2 | (2 + 4 + 2 = 8)$ . Thus it is solvable.

**Remark 3.11:**

Let  $Z(I_1, I_2) = \{(a, bI_1, cI_2); a, b, c \in \mathbb{Z}\}$  be the refined neutrosophic ring of integers,

$AX + BY = C; A, B, C \in Z(I_1, I_2)$  be a refined neutrosophic linear Diophantine equation, where

$$X = (x_0, x_1I_1, x_2I_2), Y = (y_0, y_1I_1, y_2I_2), A = (a_0, a_1I_1, a_2I_2),$$

$B = (b_0, b_1I_1, b_2I_2), C = (c_0, c_1I_1, c_2I_2)$ , we summarize the algorithm of solution as follows:

(a) Check the solvability condition.

$$(b) \text{ Solve } a_0x_0 + b_0y_0 = c_0.$$

$$(c) \text{ Solve } (a_0 + a_2)(x_0 + x_2) + (b_0 + b_2)(y_0 + y_2) = c_0 + c_2.$$

(d) Compute  $x_2, y_2$ .

(e) Solve  $(a_0 + a_1 + a_2)(x_0 + x_1 + x_2) + (b_0 + b_1 + b_2)(y_0 + y_1 + y_2) = c_0 + c_1 + c_2$ .

(f) Compute  $x_1, y_1$ .

### Example 3.12:

According to Example 3.10, we found that  $(1, 2I_1, 3I_2).X + (3, 3I_1, 8I_2)Y = (2, 4I_1, 2I_2)$  is solvable.

We consider  $x_0 + 3y_0 = 2$ . It has a solution  $x_0 = -1, y_0 = 1$ . We take

$(1 + 3)(x_0 + x_2) + (3 + 8)(y_0 + y_2) = 2 + 2$ , i.e.  $4M + 11N = 4; M = x_0 + x_2, \text{ and } N = y_0 + y_2$ , it has

a solution  $M = 1, N = 0$ , thus  $x_2 = M - x_0 = 2, y_2 = N - y_0 = -1$ . The third equation is

$(1 + 2 + 3)(x_0 + x_1 + x_2) + (3 + 3 + 8)(y_0 + y_1 + y_2) = 2 + 4 + 2$ , i.e.

$6S + 14T = 8; S = x_0 + x_1 + x_2, T = y_0 + y_1 + y_2$ . It has a solution  $S = -1, T = 1$ , thus

$x_1 = S - x_0 - x_2 = -2, y_1 = T - y_0 - y_2 = 1$ . The solution of

$(1, 2I_1, 3I_2).X + (3, 3I_1, 8I_2)Y = (2, 4I_1, 2I_2)$  is  $X = (-1, -2I_1, 2I_2), Y = (1, I_1, -I_2)$ .

## 5. Conclusion

In this paper, we have determined the criteria for the solvability of linear Diophantine equation in the neutrosophic ring of integers and refined neutrosophic rings of integers by finding the relationship between neutrosophic equations and classical equations. Also, we have presented an algorithm which gives a solution of these equations, and constructed some examples to clarify the validity of this work.

**Funding:** This research received no external funding

**Conflicts of Interest:** The authors declare no conflict of interest

## References

[1]. Abobala, M., "Classical Homomorphisms Between Refined Neutrosophic Rings and Neutrosophic Rings", International Journal of Neutrosophic Science, Vol. 5, pp.72-75, 2020.

[2]. Abobala, M., "On Some Special Substructures of Neutrosophic Rings and Their Properties", International Journal of Neutrosophic Science", Vol. 4, pp.72-81, 2020.

- [3].Abobala, M., "On Some Special Substructures of Refined Neutrosophic Rings", International Journal of Neutrosophic Science, Vol. 5, pp.59-66, 2020.
- [4].Adeleke, E.O., Agboola, A.A.A., and Smarandache, F., "Refined Neutrosophic Rings I", International Journal of Neutrosophic Science, Vol 2 , pp 77-81, 2020.
- [5].Agboola, A.A.A., Akwu, A.D., and Oyebo, Y.T., "Neutrosophic Groups and Subgroups", International J .Math. Combin, Vol. 3, pp. 1-9, 2012.
- [6].Agboola, A.A.A., Akinola, A.D., and Oyebola, O.Y.," Neutrosophic Rings I ", International J. Math combin, Vol 4,pp 1-14, 2011.
- [7]. Ceven, Y., and Tekin, S., " Some Properties of Neutrosophic Integers", Kırklareli University Journal of Engineering and Science, Vol. 6, pp.50-59, 2020.
- [8].Kandasamy, V.W.B., and Smarandache, F., "Some Neutrosophic Algebraic Structures and Neutrosophic N-Algebraic Structures", Hexis, Phonex, Arizona 2006.
- [9]. Sankari, H., " Number Theory", Tishreen University Press, pp. 123-201, 2017.
- [10].Smarandache, F., and Abobala, M., "n-Refined Neutrosophic Rings", International Journal of Neutrosophic Science, Vol. 6, pp.83-90 , 2020.
- [11] Smarandache, F., and Abobala, M., "n-Refined neutrosophic Vector Spaces", International Journal of Neutrosophic Science, Vol. 7, pp. 47-54, 2020.
- [12] Abdel-Basst, Mohamed, Rehab Mohamed, and Mohamed Elhoseny. "<? covid19?> A model for the effective COVID-19 identification in uncertainty environment using primary symptoms and CT scans." Health Informatics Journal (2020): 1460458220952918.
- [13] Abdel-Basset, M., Gamal, A., Son, L. H., & Smarandache, F. (2020). A Bipolar Neutrosophic Multi Criteria Decision Making Framework for Professional Selection. Applied Sciences, 10(4), 1202.
- [14] Abdel-Basset, M., Gamal, A., Chakraborty, R. K., & Ryan, M. A new hybrid multi-criteria decision-making approach for location selection of sustainable offshore wind energy stations: A case study. Journal of Cleaner Production, 280, 124462.
- [15] Abdel-Basst, M., Mohamed, R., & Elhoseny, M. (2020). A novel framework to evaluate innovation value proposition for smart product-service systems. Environmental Technology & Innovation, 101036.

- [16] Abdel-Basset, M., Mohamed, R., Zaied, A. E. N. H., Gamal, A., & Smarandache, F. (2020). Solving the supply chain problem using the best-worst method based on a novel Plithogenic model. In *Optimization Theory Based on Neutrosophic and Plithogenic Sets* (pp. 1-19). Academic Press.
- [17] K. Hamza Alhasan and F. Smarandache, "Neutrosophic Weibull distribution and Neutrosophic Family Weibull Distribution," *Neutrosophic Sets and Systems*, vol. 28, pp. 191-199, 2019.
- [18] S. K. Patro and F. Smarandache, "THE NEUTROSOPHIC STATISTICAL DISTRIBUTION, MORE PROBLEMS, MORE SOLUTIONS," *Neutrosophic Sets and Systems*, vol. 12, pp. 73-79, 2016.
- [19] M. Sahin, N. Olgun, V. Uluçay, A. Kargin and F. Smarandache, "A New Similarity Measure Based on Falsity Value between Single Valued Neutrosophic Sets Based on the Centroid Points of Transformed Single Valued Neutrosophic Numbers with Applications to Pattern Recognition," *Neutrosophic Sets and Systems*, vol. 15, pp. 31-48, 2017.
- [20] M. M. Lotfy, S. ELhafeez, M. Eisa and A. A. Salama, "Review of Recommender Systems Algorithms Utilized in Social Networks based e-Learning Systems & Neutrosophic System," *Neutrosophic Sets and Systems*, vol. 8, pp. 32-41, 2015.

Received: June 18, 2020. Accepted: Nov 20, 2020