



HAL
open science

Comment on “Meta-analysis reveals declines in terrestrial but increases in freshwater insect abundances”

Marion Desquilbet, Laurence Gaume, Manuela Grippa, Régis Cereghino, Jean-François Humbert, Jean-Marc Bonmatin, Pierre-André Cornillon, Dirk Maes, Hans van Dyck, David Goulson

► To cite this version:

Marion Desquilbet, Laurence Gaume, Manuela Grippa, Régis Cereghino, Jean-François Humbert, et al.. Comment on “Meta-analysis reveals declines in terrestrial but increases in freshwater insect abundances”. *Science*, 2020, 370 (6523), 10.1126/science.abd8947. hal-03082926

HAL Id: hal-03082926

<https://hal.science/hal-03082926>

Submitted on 10 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comment on “Meta-analysis reveals declines in terrestrial but increases in freshwater insect abundances”

Authors: Marion Desquilbet^{1*†}, Laurence Gaume^{2†}, Manuela Grippa³, Régis Céréghino⁴, Jean-François Humbert⁵, Jean-Marc Bonmatin⁶, Pierre-André Cornillon⁷, Dirk Maes⁸, Hans Van Dyck⁹, David Goulson¹⁰

Affiliations:

¹ Toulouse School of Economics, INRAE, University of Toulouse Capitole, Toulouse, France.

² AMAP, University of Montpellier, CNRS, CIRAD, INRAE, IRD, Montpellier, France.

³ GET, University of Toulouse 3, CNRS, IRD, Toulouse, France.

⁴ Laboratoire Ecologie Fonctionnelle et Environnement, Université de Toulouse, CNRS, Toulouse, France.

⁵ iEES Paris, INRAE, CNRS, IRD, UPEC, Université de Paris, Sorbonne Université, Paris France.

⁶ Centre de Biophysique Moléculaire, CNRS, 45071 Orléans, France.

⁷ Univ Rennes, CNRS, IRMAR - UMR 6625, F-35000 Rennes, France.

⁸ Res Inst Nat & Forest INBO, Species Divers Grp, Brussels, Belgium.

⁹ Behavioural Ecology & Conservation Group, Earth & Life Institute, UCLouvain, Louvain-la-Neuve, Belgium.

¹⁰ Evolution, Behaviour, and Environment, School of Life Sciences, University of Sussex, Falmer, East Sussex, BN1 9QG, UK.

*Correspondence to: marion.desquilbet@inrae.fr.

†These authors contributed equally to this work.

Abstract:

Van Klink *et al.* (Reports, 24 April 2020, p. 1459) argue for a more nuanced view of insect decline, and human responsibility for this decline, than previously suggested. However, shortcomings in data selection and methodology raise questions over their conclusions on trends and drivers. We call for more rigorous methodology to be applied in meta-analyses of ecological data.

Main Text:

Recent evidence has emerged on insect decline and on the loss of their crucial ecological services (1, 2), but the magnitude and causes of this decline are debated (3, 4). In their meta-analysis, Van Klink *et al.* estimated a 9% decline in abundance per decade for terrestrial insects, and an 11% increase for freshwater insects. We argue that problems in data selection and in the methodology of the meta-analysis undermine the main conclusions of their article. In particular, we argue that the evidence for freshwater insect increases is seriously flawed.

We thoroughly examined the 166 studies included in their analysis, paying particular attention to the 14 studies identified as outliers, on which their estimated positive trend for freshwater insects was based. We identify problems in the data processing of 113 studies, including all outliers.

Fifty-seven studies, including five outliers (identified in the meta-analysis by numbers 313, 1261, 1364, 1408, 1427), were field or natural experiments (5), examining experimental conditions susceptible to impact insect communities. However, the effect of these experiments was not considered. Outlier 1364 included 172 experimental plots in a seven-hectare field.

Outlier 1427 compared streams affected, or not, by a wildfire in a natural experiment. Crop experiments carried out in each plot, and the wildfire, both introduced strong site heterogeneity that was not considered in the meta-analysis.

No attempt was made to weight studies according to their representativeness in terms of geographic location, anthropogenic impact including farming methods and pesticide use, protected status or insect assemblages. Twelve studies dealt with exceptional circumstances that cannot be extrapolated, such as the area around Chernobyl or the creation of a polder or a reservoir. They included outliers that addressed restoration activities in freshwaters designed to

mitigate the impacts of historical mining (1423) and salinity (1503), and an outlier that considered high-flying migrant insects without accounting for the major impact of wind on their local abundance (1493).

Studies were also heterogeneous in terms of number of years with data (only two years for 21 studies), site sizes (ranging from one tree branch to a whole country) and distances among sites within a study (from 3.5 m to 1760 km). While the identification of differences across geographical regions was a main purpose of the meta-analysis, the geographical coverage of the database was very uneven, with 76% of studies focused on the US and Europe. A single dataset from Sweden accounted for 43% of freshwater sites.

Heterogeneous taxonomic levels were considered in the dataset. Among the 63 freshwater assemblages, 27 contained crustaceans, molluscs or worms, including outliers 1423, 1427 and 1503. As insects represented a highly variable proportion of these assemblages, insect trends cannot be inferred from overall invertebrate trends. In 90 of the 166 studies, only one taxonomic order was considered. Stress-tolerant or invasive taxa were overrepresented compared to those known to be susceptible to environmental change. For example, in the 13 studies on dipterans, nine pertained to mosquitoes, often invasive in temperate countries (6), or chironomids, often stress-tolerant (7). By contrast, bees, particularly threatened by intensive agricultural practices, are represented in only two of nine studies pertaining to hymenopterans. For wider assemblages, increases in abundance and biomass sometimes reflected the proliferation of such opportunistic taxa while hiding diversity losses. Finally, five times, the meta-analysis considered fewer insect taxa than available in the original datasets.

Other types of problems were found for 37 studies. Outlier 70 presented a temporal skew due to undermapping of common species in an earlier period (8). The original dataset of outlier 1006 included 81 missing values coded as “101”, mistaken as abundances of 101. Outliers 1476, 1477 and 1478 were overweighted by separating their common dataset into three studies and including more sites than in the original publications. Ten studies, including outlier 1409, inconsistently included five to eight abundance or biomass data per month while the model considered the week as the finest time resolution.

Overall, we identified data processing problems in approximately two thirds of terrestrial studies and three quarters of freshwater studies (Figure 1 and Table 1).

[Figure and Table]

As for statistical methods, the model included random site effects assuming that site effects were independently and identically distributed, but this assumption was not met, notably because the inclusion of experimental sites implied non-random effects and there was a geographical dependency among sites in some studies. Moreover, the meta-analysis did not follow the usual standards, consisting in selecting the best model among several models of abundance or biomass trends and checking its validity, considering also within-year temporal autocorrelation, regional and scale effects, taxa effects, differences in abundance and biomass trends, non-linear dynamics and changepoints.

The analysis of drivers also shows methodological shortcomings. Possible drivers of insect trends were analysed by matching study data with external databases based on geographical coordinates, but this methodology is prone to numerous errors. The ESA CCI database has a 300 m × 300 m resolution only allowing a rough assessment of local land use, and may easily

misclassify croplands and grasslands when processing satellite images because of their similar spectral signatures (9, 10). The meta-analysis considered that 48 terrestrial studies had at least one site with some local cropland cover. Actually, from corresponding publications or satellite images, we find that cropland cover was inadequately assessed in 31 of these studies.

5 In freshwaters, pollutants are carried by water throughout the watershed. Therefore, drivers of insect population change should examine land use upstream (11) and not only around the sampling site. The trophic status of water, reported in almost all studies, would have allowed a better assessment of anthropogenic pressures.

10 These issues call into question the conclusion that insect abundance trends were positively associated with crop cover at the local scale in both terrestrial and freshwater ecosystems.

Overall, we argue that these methodological problems invalidate the estimates of the meta-analysis of van Klink *et al.* To avoid replicating such issues, ecologists must apply rigorous standards for systematic literature reviews and meta-analyses (12), as is becoming the rule for example in environmental health (13).

15 **References and Notes:**

1. S. Seibold *et al.*, Arthropod decline in grasslands and forests is associated with landscape-level drivers. *Nature* **574**, 671-674 (2019).
2. B. C. Lister, A. Garcia, Climate-driven declines in arthropod abundance restructure a rainforest food web. *Proc. Natl. Acad. Sci. U.S.A.* **115**, E10397-E10406 (2018).
- 20 3. G. A. Montgomery *et al.*, Is the insect apocalypse upon us? How to find out. *Biol. Conserv.* **241**, 108327 (2020).

4. P. Cardoso *et al.*, Scientists' warning to humanity on insect extinctions. *Biol. Conserv.* **242**, (2020).
5. J. M. Diamond, Ecology: Laboratory, field and natural experiments. *Nature* **304**, 586-587 (1983).
- 5 6. J. M. Medlock *et al.*, A review of the invasive mosquitoes in Europe: Ecology, public health risks, and control options. *Vector-Borne and Zoonotic Dis.* **12**, 435-447 (2012).
7. C. Lindegaard, "Classification of water-bodies and pollution" in *The Chironomidae: Biology and ecology of non-biting midges*, P. D. Armitage, L. C. Pinder, P. Cranston, Eds. (Springer, Netherlands, 1995), pp. 385-404.
- 10 8. D. Maes, H. Van Dyck, Butterfly diversity loss in Flanders (north Belgium): Europe's worst case scenario? *Biol. Conserv.* **99**, 263–276 (2001).
9. W. Li *et al.*, Gross and net land cover changes in the main plant functional types derived from the annual ESA CCI land cover maps (1992–2015). *Earth Syst. Sci. Data* **10**, 219-234 (2018).
- 15 10. L. Vilar, J. Garrido, P. Echavarría, J. Martínez-Vega, M. P. Martín, Comparative analysis of CORINE and climate change initiative land cover maps in Europe: Implications for wildfire occurrence estimation at regional and local scales. *Int. J. Appl. Earth Obs. Geoinf.* **78**, 102-117 (2019).
11. J. D. Allan, Landscapes and Riverscapes: The Influence of Land Use on Stream Ecosystems. *Annu. Rev. Ecol. Evol. Syst.* **35**, 257-284 (2004).
- 20 12. J. Koricheva, J. Gurevitch, K. Mengersen, *Handbook of Meta-analysis in Ecology and Evolution*. (Princeton University Press, 2013).

13. J. McPartland, J. Lam, C. Lanier-Christensen, A Valuable Contribution toward Adopting Systematic Review in Environmental Health. *Environ. Health Perspect.* **122**, A10-A10 (2014).

Acknowledgments: M.D. and L.G. thank M. Elias and G. Le Moguédec for their helpful comments on the manuscript. M. Elias and A. Toporov are acknowledged for their help in translating the Russian articles, as well as F. Kjellberg for his help in translating the Swedish database.

Funding: This work was publicly funded by the French National Research Agency (grants ANR-17-EURE-0010 to M. D. and ANR-16-IDEX-0006 to L. G. under the "Investissements d'avenir" program and grant ANR-15-CE21-0006 to M.D.).

Author contributions: M.D. and L.G. performed both the detailed and overall analysis of the meta-analysis and wrote the original draft. M.G. contributed to the analysis of data based on geographical coordinates and the comparison with satellite images, and processed the map. R.C. contributed to the analysis of databases on freshwater insects and, like J.F.H., he provided methodological comments on the analysis of freshwater ecosystems. P.A.C. reviewed the statistical analysis of the meta-analysis. D.M. and H.V.D. participated in the analysis of some Lepidoptera databases. J.M.B and D.G. extensively edited the manuscript. All authors contributed to the general comment and reviewed the manuscript.

Competing interests: Authors declare no competing interests.

Data and materials availability: There are no new data in this comment.

Fig. 1. Geographical distribution of studies giving rise to data processing concerns




Table 1. Types of problems encountered

Problem types	Number of studies	Study identifiers
Field or natural experiment, experimental conditions not considered	57	63, 294, 300, 301, 313 , 1261 , 1335, 1339, 1357, 1364 , 1365, 1367, 1376, 1384, 1385, 1387, 1388, 1391, 1393, 1396, 1398, 1407, 1408 , 1410, 1411, 1413, 1415, 1417, 1419, 1421, 1424, 1426, 1427 , 1430, 1431, 1433, 1437, 1439, 1441, 1465, 1468, 1473, 1479, 1484, 1485, 1487, 1494, 1497, 1502, 1504, 1505, 1506, 1513, 1516, 1519, 1521, 1527
Non-insects (and non-arachnids*) included in the assemblage	31	1395, 1396, 1402, 1421, 1423 , 1424, 1425, 1427 , 1428, 1432, 1435, 1448, 1449, 1451, 1452, 1454, 1455, 1456, 1458, 1466, 1473, 1498, 1500, 1503 , 1504, 1506, 1507, 1509, 1511, 1513, 1521
Only two years with data	21	1335, 1365, 1376, 1384, 1385, 1402, 1411, 1415, 1419, 1421, 1425, 1435, 1439, 1465, 1468, 1480, 1502, 1508, 1513, 1515, 1521
Split studies (several studies from one dataset or one geographical location)	18	79, 380, 1006 , 1263, 1266, 1267, 1345, 1347, 1353, 1357, 1424, 1476 , 1477 , 1478 , 1485, 1487, 1495, 1496
Exceptional situation	12	478, 1324, 1395, 1422, 1423 , 1451, 1456, 1464, 1493 , 1498, 1503 , 1517
More than four data per month	10	249, 294, 301, 1263, 1351, 1409 , 1476 , 1477 , 1481, 1501
Change in site numbers	9	70 , 1006 , 1102, 1263, 1266, 1267, 1388, 1476 , 1477
Error in insect count	6	1006 , 1393, 1431, 1448, 1457, 1509
Data exists for other taxa	5	70 , 502, 1393, 1480, 1494
Data exists for a longer period	3	502, 1263, 1494
Time bias in data	2	70 , 465
Very unbalanced data	1	1475
No identified problem	53	375, 1310, 1312, 1328, 1340, 1346, 1349, 1361, 1377, 1378, 1379, 1381, 1382, 1392, 1394, 1397, 1400, 1401, 1403, 1404, 1405, 1406, 1412, 1414, 1416, 1418, 1429, 1434, 1440, 1444, 1445, 1446, 1453, 1459, 1460, 1461, 1462, 1467, 1470, 1471, 1472, 1474, 1488, 1491, 1510, 1512, 1518, 1520, 1524, 1525, 1526

Note: The references of the studies numbered with the identifiers indicated in the Table are available in the supplementary materials of van Klink *et al.*, in Table S1 and in the section “references and notes”. Outliers are in bold. *Arachnids were considered as insects in the meta-analysis.