



HAL
open science

Second Coordination Sphere Effects in an Evolved Ru Complex Based on Highly Adaptable Ligand Results in Rapid Water Oxidation Catalysis

Nataliia Vereshchuk, Roc Matheu, Jordi Benet-Buchholz, Muriel Pipelier, Jacques Lebreton, Didier Dubreuil, Arnaud Tessier, Carolina Gimbert-Suriñach, Mehmed Ertem, Antoni Llobet

► To cite this version:

Nataliia Vereshchuk, Roc Matheu, Jordi Benet-Buchholz, Muriel Pipelier, Jacques Lebreton, et al.. Second Coordination Sphere Effects in an Evolved Ru Complex Based on Highly Adaptable Ligand Results in Rapid Water Oxidation Catalysis. *Journal of the American Chemical Society*, 2020, 142 (11), pp.5068-5077. 10.1021/jacs.9b11935 . hal-03085843

HAL Id: hal-03085843

<https://hal.science/hal-03085843>

Submitted on 22 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

Second Coordination Sphere Effects in an Evolved Ru Complex Based on a Highly Adaptable Ligand Results in Rapid Water Oxidation Catalysis

Journal:	<i>Journal of the American Chemical Society</i>
Manuscript ID	Draft
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Vereshchuk, Natalii; The Barcelona Institute of Science and Technology, Inst of Chem. Res. of Catalonia (ICIQ) Matheu, Roc; University of California Berkeley, Chemistry Benet-Buchholz, Jordi; Universitat Rovira i Virgili Institut Catala d'Investigacio Quimica, Chemistry Pipelier, Muriel; Universite de Nantes Faculte de Droit et des Sciences Politiques, UMR CNRS 6230 Lebreton, Jacques; Universite de Nantes, Dubreuil, Didier; Chimie et Interdisciplinarite Synthese Analyse Modelisation, Tessier, Arnaud; Laboratoire CEISAM, Faculté des Sciences et Techniques - CNRS UMR 6230 Gimbert Suriñach, Carolina; Universitat Rovira i Virgili Institut Catala d'Investigacio Quimica, Ertem, Mehmed; Brookhaven National Laboratory, Chemistry Llobet, Antoni; The Barcelona Institute of Science and Technology, Inst of Chem. Res. of Catalonia (ICIQ)

SCHOLARONE™
Manuscripts

Second Coordination Sphere Effects in an Evolved Ru Complex Based on a Highly Adaptable Ligand Results in Rapid Water Oxidation Catalysis

Nataliia Vereshchuk^{1,2}, Roc Matheu¹, Jordi Benet-Buchholz¹, Muriel Pipelier³, Jacques Lebreton³, Didier Dubreuil³, Arnaud Tessier^{*,3}, Carolina Gimbert-Suriñach¹, Mehmed Z. Ertem^{*,4} and Antoni Llobet^{*,1,5}

¹ Institute of Chemical Research of Catalonia (ICIQ), Barcelona Institute of Science and Technology (BIST), Avda. Països Catalans 16, 43007 Tarragona, Spain.

² Departament de Química Física i Inorgànica, Universitat Rovira i Virgili, Marcel·lí Domingo s/n, 43007 Tarragona, Spain.

³ Université de Nantes, CNRS, CEISAM, UMR 6230, Faculté des Sciences et des Techniques, 2 rue de la Houssinière, BP 92208, 44322 Nantes, France.

⁴ Chemistry Division, Energy & Photon Sciences Directorate, Brookhaven National Laboratory, Upton, New York, 11973-5000, USA.

⁵ Departament de Química, Universitat Autònoma de Barcelona, Cerdanyola del Vallès, 08193 Barcelona, Spain.

Corresponding authors: arnaud.tessier@univ-nantes.fr; mzertem@bnl.gov and allobet@iciq.es

Abstract

A new Ru complex containing the deprotonated 2,2':6',2''-terpyridine,6,6''-diphosphonic acid (H_4tPa) and pyridine (py) of general formula, $[Ru^II(H_3tPa-\kappa-N^3O)(py)_2]^+$, **2**⁺, has been prepared and thoroughly characterized by means of spectroscopic, electrochemical techniques, X-ray diffraction analysis and with DFT calculations. Complex **2**⁺, presents a dynamic behavior in solution that involves the synchronous coordination and the decoordination of the dangling phosphonic groups of the tPa^{4-} ligand. However, at oxidation state IV complex **2**⁺, becomes seven coordinated with the two phosphonic groups now bonded to the metal center. Further at this oxidation state at neutral and basic pH, the Ru complex undergoes coordination of an exogenous OH^- group from the solvent that leads to an intramolecular aromatic O-atom insertion into the CH bond of one of the pyridyl groups forming the corresponding pheoxo-phosphonate Ru complex $[Ru^{III}(tPaO-\kappa-N^2O_pO_c)(py)_2]^{2-}$, **4**²⁻, where $tPaO^{5-}$ is 3-hydroxo-[2,2':6',2''-terpyridine]-6,6''-diyl)bis(phosphonate) ligand. This new in situ generated Ru complex, **4**²⁻, has been isolated and spectroscopically and electrochemically characterized. In addition, a crystal structure has been also obtained using single crystal X-ray diffraction techniques. Complex **4**²⁻, turns out to be an exceptional water oxidation catalyst achieving record high TOF_{max} in the of $16,000\ s^{-1}$. A mechanistic analysis complemented with DFT calculations has also been carried out showing the critical role of intramolecular second coordination sphere effects exerted by the phosphonate groups in lowering the activation energy at the rate determining step.

Keywords

Water oxidation catalysis, Ru complexes, water splitting, redox properties transition metal complexes,

1. Introduction

Molecular water oxidation catalysis is a field that has been rapidly developing over the last decade mainly due to their potential application in new energy conversion schemes based on water splitting with sunlight.¹⁻⁷ In particular, Ru complexes have been leading the field and today the amount of information extracted from these complexes has generated a deep understanding of the different parameters involved in the catalytic cycle.^{8,9} This has been possible thanks to the spectroscopic characterization of the reaction intermediates as well as due to the kinetic characterization of the different steps involved and complemented with a thorough computational analysis.¹⁰⁻¹³ This wealth of information based on Ru complexes as water oxidation catalysts (WOCs) has also been extended to other transition metals as well as oxides reported as WOCs for their design and mechanistic proposals.¹⁴⁻¹⁷

The best WOCs known today are based on the Flexible Adaptative Multidentate Equatorial (FAME) ligands¹⁸ containing polypyridyl carboxylate groups such as H₂tda¹⁹ and H₂bda²⁰⁻²² (see Chart I for the drawings of the ligands) and their related derivatives where the carboxylate groups have been partially or totally substituted by phosphonate groups.²³⁻²⁶ In particular, the complex [Ru^{IV}(OH)(tda-κ-N³O)(py)₂]⁺ (py for pyridine), **1**⁺, has been recently shown to achieve TOF_{max} of 8,000 s⁻¹ at pH 7.0.¹⁹ Further, the analysis of its performance shows a delicate balance among multiple equilibria at different oxidation states with diverse coordination number and degree of hydrogen bonding that are responsible for its performance as a catalyst.^{19,27} Because of these delicate equilibria, the replacement of carboxylate groups in H₂tda by phosphonate groups leading to the H₄tPa ligand (Chart I), is expected to generate significant differences in its catalytic behavior. The electronic nature and geometry of the phosphonate with regard to those of the carboxylate group as well as an increase in steric will be responsible for the different catalytic behavior of the new Ru complexes based on H₄tPa.²⁸ In addition, the increased number of acidic protons with substantially different pK_a's are expected to not only facilitate remote proton-coupled electron transfer (PCET) events during redox leveling,²⁹⁻³⁵ but also give rise to beneficial second sphere coordination effects such as hydrogen bonding interactions. Further, the phosphonate groups are expected to act as intramolecular proton acceptor, at the O-O bond formation step, significantly reducing its activation energies.

Here on we present a new family of Ru complexes derived from the H₄tPa ligand together with a thorough characterization of their redox and water oxidation catalytic properties.

2. Results

2.1 Synthesis and structure of the precursor complex **2**⁺

The synthesis of the H₄tPa ligand was carried out in three steps using 2,2':6',2''-terpyridine (trpy) as the starting material as described in the supporting information. Initially, the trpy ligand is oxidized to 2,2':6',2''-terpyridine-1,1''-dioxide using metachloroperbenzoic acid. The treatment of the latter with POEt₃ forms tetraethyl 2,2':6',2''-terpyridine-6,6''-diphosphonate that is hydrolyzed with trimethylbromosilane yielding the H₄tPa ligand. The synthetic strategy for the preparation of the complexes described in this work is outlined in Figure 1. Complex [Ru^{II}(H₃tPa-κ-N³O)(py)₂]⁺, **2**⁺, is prepared by refluxing stoichiometric amounts of H₄tPa and [Ru(dmsO)₄Cl₂] in *n*-BuOH, followed by addition of excess of pyridine and further refluxing overnight. Proper column chromatography in silica generates pure **2**Cl in moderate yields where the equatorial H₃tPa⁻ ligand acts in a monoanionic fashion. Crystals of **2**PF₆·3H₂O sufficiently good for X-ray analysis were obtained by counteranion exchange crystallization from an acidic aqueous solution of **2**Cl and KPF₆. The ORTEP view of its solvated cationic moiety is presented in Figure 1. The Ru(II) center features an octahedral type of geometry where the H₃tPa⁻ ligand coordinates in the equatorial plane in a κ-N³O fashion leaving a dangling non-coordinated phosphonic acid group. The octahedral geometry is completed with two pyridyl groups in the axial position. The bonding parameters are unremarkable³⁶ except for a distortion due to the geometrical constraints of the HtPa³⁻ ligand at the equatorial zone, that produces a N-Ru-O angle of 115°; 25° larger than the ideal O_h geometry. The same angle for the [Ru^{II}(tda)(py)₂] is 125°,¹⁹ that is 10° higher which is a consequence of the different bonding parameters associated with the phosphonate and carboxylate moieties.^{28,37} Finally, there are three H₂O molecules in the unit cell, interacting with the phosphonic groups by hydrogen bonding as can be observed in Figure 1. Density functional theory (DFT) calculations at the M06 level of theory³⁸ in conjunction with SMD aqueous continuum solvation model³⁹ (see computational methods for further details) have been carried out in order to complement the experimental work described here for **2**⁺. The computed structure for **2**⁺, shows a very good agreement with the experimental structure in terms of bonding parameters (Figure S62) revealing its good degree of reliability.

Addition of two equivalents of Ce(IV) generates the corresponding seven coordinated Ru(IV) complex [Ru^{IV}(tPa-κ-N³O²)(py)₂], **3**, where the tPa⁴⁻ ligand now acts in a pentadentate manner and the metal center possesses a pentagonal bipyramidal geometry as can be observed in the computed

1
2
3 structure shown in Figure 1. This complex is diamagnetic at room temperature (RT) with a high field
4 low spin d^4 electronic configuration, $(e_1'')^4(e_2')^0(a_1')^0$, analogous to related Ru(IV) complexes.^{19,40,41}
5 Its ^1H , ^{13}C and ^{31}P NMR are presented in the SI see Figures S27-29.
6
7

8 9 2.2 Spectroscopy and dynamic behavior of **2**



10
11 The spectroscopic properties were investigated by means of UV-vis and NMR spectroscopy. The UV-
12 vis spectra of $[\text{Ru}^{\text{II}}(\text{H}_2\text{tPa-}\kappa\text{-N}^3\text{O})(\text{py})_2]$, **2**, and its one and two electron oxidized species $[\text{Ru}^{\text{III}}(\text{H}_2\text{tPa-}$
13 $\kappa\text{-N}^3\text{O})(\text{py})_2]^+$ and $[\text{Ru}^{\text{IV}}(\text{HtPa-}\kappa\text{-N}^3\text{O}^2)(\text{py})_2]^+$ were generated via redox titration using Ce(IV) as the
14 oxidant at pH 1.0 in aqueous triflic acid (Figures S45-S46). Complex **2** is a diamagnetic low spin d^6
15 ion with a typical $(t_{2g})^6(e_g)^0$ electronic configuration and its NMR spectra and assignment are
16 presented in the Figure 2 and the supporting information (Figures S12-S26 and Figures S35-S36). At
17 room temperature, the NMR spectrum displays eight resonances as if the complex had C_{2v}
18 symmetry. However, at low temperature the resonances of the protons H_a , H_b , H_c and H_e split
19 indicating the presence of a dynamic behavior (Figure 2). The dynamic behavior is associated with
20 the synchronic coordination and decoordination of the dangling phosphonate that has an activation
21 energy of 13.3 kcal/mol based on Eyring plots generated from VT-NMR (Scheme S2). By
22 computational methods, an estimate activation energy of 7.9 kcal/mol (see Figure S37) was
23 obtained for this dynamic behavior in line with the experimental observations. The value of the
24 activation energy is similar to that measured for Ru-bda complexes involving coordination and
25 decoordination of the dangling carboxylates.⁴²
26
27
28
29
30
31
32
33
34
35
36

37 2.3 Redox properties of **2**

38
39 The redox properties of **2** were investigated by electrochemical techniques in a three-electrode cell
40 using glassy carbon as a working electrode and all potentials were measured using a $\text{Hg}/\text{Hg}_2\text{SO}_4$ as
41 a reference electrode and converted to vs NHE by adding 0.65 V to the measured potential. Cyclic
42 Voltammetry of **2** were performed in aqueous solution as a function of pH that produces a different
43 degree of protonation at the phosphonate groups. A cyclic voltammetry at pH 7.0 is shown in
44 Figure 3 (red trace) where two chemically reversible and electrochemically quasireversible redox
45 processes can be observed. The first redox process at $E_{1/2} = 0.83$ V is assigned to a PCET process to
46 generate $[\text{Ru}^{\text{III}}(\text{tPa-}\kappa\text{-N}^3\text{O}^2)(\text{py})_2]^-$ whereas the second wave at $E_{1/2} = 0.92$ V is due to a one electron
47 oxidation process to form $[\text{Ru}^{\text{IV}}(\text{tPa-}\kappa\text{-N}^3\text{O}^2)(\text{py})_2]$. The one electron nature of the two processes is
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 corroborated by the spectrophotometric redox titration (Figure S45-S46), and the degree of
4 protonation is deduced from the Pourbaix diagram shown in Figure 3 right.

5
6
7 The four protons potentially available from the two phosphonic groups in **2**, allow to obtain a rich
8 family of complexes depending on the degree of oxidation and protonation as shown in the Pourbaix
9 diagram. The pK_a 's coincide with the dashed vertical lines and are deduced from the change of slope
10 in the E^o vs. pH plot. The first and second pK_a 's due to the monodeprotonation of each phosphonic
11 group are expected to occur below pH 1.0.⁴³⁻⁴⁵ Further, the pK_a for the third and fourth proton loss
12 are labeled, pK_a and $pK_{a'}$, and are defined in equations 1 and 2 respectively,
13
14
15
16
17
18
19



22
23
24
25
26
27
28 The latter have been independently calculated via spectrophotometric acid-base titration as shown
29 in Figure S47. It is interesting to see how due to the different proton content of the species involved
30 at a given pH, the difference between the IV/III-III/II redox couples changes considerably as a
31 function of pH as can be graphically observed in Figure S49A. Indeed, at pH 7.0 this difference is only
32 100 mV but it increases up to 330 mV at pH 13.0 (Figure S49B).
33
34
35
36
37

38 *2.4 Generation of the active Ru species via O-atom insertion*

39
40 The synthesis of the active catalyst $[Ru^{III}(tPaO-\kappa-N^2O_pO_c)(py)_2]^{2-}$, **4²⁻**, (see a drawing of the H₅tPaO
41 ligand in Chart 1; O_p refers to oxygen atom coordinated to the Ru center via the oxygen atom from
42 the phosphonic group and O_c from the pyridyl group) was carried out electrochemically using
43 $[Ru^{II}(H_3tPa-\kappa-N^3O)(py)_2]^+$, **2⁺**, (**2⁺** is predicted to convert to **2** at pH 7.7) as a precursor as outlined in
44 Figure 1.
45
46
47
48

49 A sample of **2⁺** is dissolved in a pH 7.7 aqueous phosphate buffer solution and placed in a two-
50 compartment electrochemical cell using a Pt mesh as a working electrode. Then a potential of 1.30
51 V is applied for 110 minutes. During this process, the initial **2⁺** precursor complex is quantitatively
52 transformed into the final complex **4²⁻**, where an O atom insertion between the Ru center and one
53
54
55
56
57
58
59
60

1
2
3 of the pyridyl groups takes place. One electron reduction of $\mathbf{4}^{2-}$ yields the diamagnetic $\mathbf{4}^{3-}$ complex
4 that was characterized by ^1H , ^{13}C and ^{31}P NMR spectroscopy and whose spectra are shown in the
5 (Figure S39-S44). Electrochemically only the initial and final species are detected (Figure S50).
6
7

8
9 Single crystals of $\mathbf{4}^{3-}$ were obtained by adding excess Cs^+ to a solution of $\mathbf{4}^{2-}$ in MeOH. A crystal
10 structure of $\mathbf{4}^{3-}$ is displayed in Figure 1 showing the phenoxide bonding and the decoordination of
11 the initial pyridyl-phosphonic group. The complex was further characterized by UV-vis spectroscopy
12 using an Optically Transparent Thin Layer Electrochemical (OTTLE) cell (Figure S48).
13
14
15


16
17 In order to get further insight into the potential sequence of reactions that generate $\mathbf{4}^{2-}$ from $\mathbf{2}^+$ a
18 computational analysis was carried out and the results obtained are summarized in Scheme 1. As
19 the current flows at an applied potential of 1.30 V, the initial Ru(II) complex $\mathbf{2}^+$ is initially converted
20 to seven coordinate (CN7) Ru(IV) species, $[\text{Ru}^{\text{IV}}(\text{tPa-}\kappa\text{-N}^3\text{O}^2)(\text{py})_2]$, $\mathbf{3}$, in agreement with its redox
21 properties described in the previous section. This complex then reacts with a water molecule with
22 concomitant proton release inducing the decoordination of one of the pyridyl phosphonate groups
23 to generate $[\text{Ru}^{\text{IV}}(\text{O})(\text{HtPa-}\kappa\text{-N}^2\text{O})(\text{py})_2]$. The latter then undergoes a PCET step to form $[\text{Ru}^{\text{V}}(\text{O})(\text{tPa-}$
24 $\kappa\text{-N}^2\text{O})(\text{py})_2]$. This species is responsible for the intramolecular hydroxylation of the dangling pyridyl
25 group that is highly favored from a thermodynamic perspective ($\Delta G = -9.0$ kcal/mol) and that finally
26 leads to pure $\mathbf{4}^{2-}$ following a deprotonation event ($\Delta G = -54.6$ kcal/mol). A transition state for the
27 electrophilic oxygen insertion into the CH bond has also been located with a low free energy of
28 activation (ΔG^\ddagger) of 14 kcal/mol.
29
30
31
32
33
34
35
36
37

38 The redox and electrocatalytic properties of $\mathbf{4}^{2-}$ were investigated by CV and Coulombimetric
39 techniques. Figure 3 (black trace, left) shows the CV of $\mathbf{4}^{2-}$ at pH 7.0 phosphate buffer solution where
40 an electrochemically and chemically reversible wave appears at $E_{1/2} = 0.35$ V that is assigned to the
41 Ru(III)/Ru(II) couple. At higher anodic potentials a very large electrocatalytic current is observed at
42 approximately 1.4 V that is due to the oxidation of water to dioxygen. This second wave is associated
43 with a two electron process and oxido coordination, reaching the Ru(V)=O species that is responsible
44 for the reaction with solvent water and O-O bond formation. A two electron process is proposed
45 since no other waves are observed besides the III/II wave and the corresponding Ru(IV)=O species
46 are not sufficiently reactive for fast water oxidation and further supported by DFT.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2.5 Water oxidation catalysis and mechanisms

A kinetic characterization of the electrocatalytic water oxidation reaction was carried out for $\mathbf{4}^{2-}$, based on Foot of the Wave Analysis (FOWA)⁴⁶⁻⁴⁸ at pH 7.2 as shown in Figure S54. A $TOF_{max} = 1.6 \cdot 10^4 (\pm 0.2 \cdot 10^4) s^{-1}$ is obtained that is found to be independent of catalyst concentration between 0.5 and 3.0 mM. This first order kinetics behavior with regard to catalyst concentration points towards a water nucleophilic attack (WNA)⁴⁹ type of mechanism where the O-O bond formation between the Ru(V)=O species and the solvent is the rate determining step (rds). This is further supported by theoretical calculations as will be shown in the next section. The performance of $\mathbf{4}^{2-}$ as WOC was further tested by a bulk electrolysis experiments described in detail in the supporting information (Figure S53). More than 42 million turnovers with Faradaic efficiencies above 93% were obtained using Saveant's methodology that takes into account the concentration of the catalyst close to the electrode surface and manifest the remarkable stability of this catalyst.²¹⁻²³ It is also interesting to point out here that catalyst $\mathbf{4}^{2-}$ is also active at lower pH albeit with lower efficiency (Figure S51).

A thorough mechanistic analysis was also carried out based on theoretical calculations and is summarized in Figure 4 top, where the different reaction intermediates and the computed ΔG s and redox potentials associated with each step are presented (see computational methods in the SI for further details). We propose that once the Ru(III) complex, $\mathbf{4}^{2-}$, is generated, water coordination together with a two electron oxidation process proceeds to form Ru^V=O species, $[Ru^V(O)(HtPaO-\kappa-N^3O_p)(py)_2]^-$, which will in turn react with a solvent H₂O to generate the corresponding hydroxoperoxido derivative, as indicated in the equation below,



The optimized TS for WNA reaction is presented in Figure 4 bottom. It is interesting to realize here that the coordination number (CN) for the Ru(V) species is 6 with the tPaO⁵⁻ ligand acting in a $\kappa-N^2O_c$ fashion and thus with a dangling phosphonate group. This dangling group is strategically situated so that it can act as an intramolecular proton acceptor to facilitate the WNA TS and to reduce the activation free energy ($\Delta G^\ddagger = 22.1$ kcal/mol). Once Ru(III)-OOH is formed then it suffers a PCET that

1
2
3 forms the Ru(IV)-OO that spontaneously evolves dioxygen and generates the initial Ru(II) species
4 completing the catalytic cycle.
5
6
7
8
9

10 **3. Discussion**

11
12 Detailed understanding of water oxidation catalysis is challenging due to the large number of
13 reactions and intermediates involved in this complex process. The electronic demands imposed by
14 the reaction intermediates accessed at each oxidation state at the different stages of the catalytic
15 cycle involve a change in metal coordination number. Along this line, high oxidation states will prefer
16 hard base ligands while lower oxidation states will prefer softer ones. Additional complexity arises
17 because the variety of reactions handled by the catalyst including outer sphere electron transfers
18 (OSET), PCET and chemical reactions that might not necessarily involve ET such as proton transfer
19 or O-O bond formation. Moreover, non-desired pathways that derail these reactions from the
20 catalytic cycle towards non-productive or decomposition products also need to be understood,⁵⁰
21 to be able to avoid them and to obtain robust water oxidation catalysts. To control all the parameters
22 indicated above, each step should occur with an activation barrier as low as possible so that efficient
23 water oxidation catalysis can take place.
24
25
26
27
28
29
30
31
32

33 *3.1 Geometrical and electronic consequences due to the presence of the phosphonate groups*

34
35 Complex $[\text{Ru}^{\text{IV}}(\text{O})(\text{tda})(\text{py})_2]$, **1**, is an example of a catalyst that complies with the above
36 requirements and up to now appears as the fastest water oxidation catalyst ever reported.⁵¹ The
37 structural consequences of changing carboxylate by phosphonates can be graphically observed by
38 comparing the structures of the diamagnetic seven coordinated catalyst precursors $[\text{Ru}^{\text{IV}}(\text{tda}-\kappa\text{-}$
39 $\text{N}^3\text{O}^2)(\text{py})_2]^{2+}$ and $[\text{Ru}^{\text{IV}}(\text{tPa}-\kappa\text{-N}^3\text{O}^2)(\text{py})_2]$ (Figures S61 and S63). While the X-ray structure of
40 $[\text{Ru}^{\text{IV}}(\text{tda}-\kappa\text{-N}^3\text{O}^2)(\text{py})_2]^{2+}$ shows all the coordinating atoms of the tda²⁻ ligand nearly on the equatorial
41 plane, the coordinating oxygen atoms of the phosphonate in $[\text{Ru}^{\text{IV}}(\text{tPa}-\kappa\text{-N}^3\text{O}^2)(\text{py})_2]$ complex are
42 significantly off the equatorial plane as observed in the computed structure shown in Figure 1.
43 Furthermore, this distortion also breaks the planarity of the trpy moiety of the tPa⁴⁻ ligand
44 diminishing the aromatic π delocalization. This distortion is due mainly to the different geometric
45 parameters of the phosphonic acid group (pseudo T_d) as compared to the carboxylate (pseudo C_{2v}),
46 and results in weaker Ru-O bonds for the phosphonic complex. From an electronic perspective the
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 phosphonate group acts as a stronger sigma donor than the carboxylate, leading in general to lower
4 redox potentials and higher pK_a (pK_a 's are 3 and 11 for $[Ru^{II}(H_2tPa)(py)_2]$ and 6.4 for
5 $[Ru^{III}(HtPa)(py)_2]$).²⁷ The existence of a range of pK_a values obtained at different oxidation states
6 from the auxiliary ligands is interesting because it can promote remote PCET, that is a PCET where
7 the proton is not associated with the Ru-OH₂ or Ru-OH group, but from auxiliary ligands such as the
8 phosphonic ones. This is important because for typical polypyridyl Ru-aqua complexes such as
9 $[Ru(trpy)(bpy)(H_2O)]^{2+}$,⁵² the complex can access Ru^{IV}=O species by two consecutive PCET processes
10 from its aqua derivatives but access to reactive Ru(V) occurs as an ET only process and therefore
11 requires high energy input and overpotential (1.8 V for $[Ru(trpy)(bpy)(H_2O)]^{2+}$).⁵³⁻⁵⁵ Thus, the
12 possibility to reach Ru(V) via three PCET processes can open up the access to a highly active Ru^V=O
13 species at much lower overpotentials.
14
15
16
17
18
19
20
21

22 *3.2 The importance of the second coordination sphere effects*

23
24
25 The two electron oxidation of the octahedral $[Ru^{II}(H_2tPa-\kappa-N^3O)(py)_2]$, **2**, complex generates the CN7
26 diamagnetic $[Ru^{IV}(tPa-\kappa-N^3O^2)(py)_2]$. Here the importance of ligand flexibility and design is
27 manifested by leaving a non-coordinated dangling phosphonic acid group at oxidation state II where
28 the Ru complex needs to be CN6. The ligand flexibility also allows for a dynamic behavior where the
29 phosphonic groups synchronically coordinate and decoordinate fast at room temperature. Upon
30 reaching Ru(IV) the dangling group immediately coordinates thanks to the flexibility of the ligands
31 achieving CN7. At oxidation state IV in basic solution, $[Ru^{IV}(tPa-\kappa-N^3O^2)(py)_2]$ undergoes OH⁻
32 substitution to form either $[Ru^{IV}(OH)(tPa-\kappa-N^2O)(py)_2]^-$ or $[Ru^{IV}(O)(HtPa-\kappa-N^2O)(py)_2]^-$. Further
33 oxidation leads to the formation of highly reactive $[Ru^V(O)(tPa-\kappa-N^2O)(py)_2]^-$ species. The generation
34 of the Ru^{IV}-OH (or Ru^{IV}=O) occurs simultaneously with the breaking of both the Ru-N and Ru-O bonds
35 of one of the pyridyl-phosphonato arms of the tPa⁴⁻ ligand (Scheme 1). This is due to the steric
36 congestion generated by the phosphonate group that precludes the formation of the CN7
37 complexes $[Ru^{IV}(O)(HtPa-\kappa-N^3O)(py)_2]^-$ and $[Ru^V(O)(tPa-\kappa-N^3O)(py)_2]^-$. Once the highly reactive
38 Ru^V=O species is generated then the dangling pyridyl-phosphonato arm has the perfect geometry
39 to undergo intramolecular oxygen atom transfer to the non-coordinated pyridyl ring. This
40 intramolecular O-atom insertion into the CH bond generates the catalytically active complex
41 precursor $[Ru^{III}(tPaO-\kappa-N^2O_pO_c)(py)_2]^{2-}$, **4**²⁻.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Two electron oxidation of $\mathbf{4}^{2-}$ forms the $\text{Ru}^{\text{V}}=\text{O}$ species that performs O-O bond formation via WNA and generates the $\text{Ru}^{\text{III}}\text{-OOH}$ intermediate. This step is the rate determining step (rds) of the catalytic cycle and is favored by the intramolecular proton transfer from the incoming water molecule to the phosphonate group that again has the right geometry to promote this step reducing the activation free energy. This renders $\mathbf{4}^{2-}$ as an extremely powerful catalyst giving a TOF_{max} of $16,000 \text{ s}^{-1}$ with an overpotential of 530 mV at pH 7.0. It gives a higher TOF_{max} ($8,000 \text{ s}^{-1}$) than $\mathbf{1}$ at lower overpotential at (75 mV.)

In summary, the dangling phosphonate group is responsible for low energy pathway of the two key reactions: (i) The formation of active catalyst precursor $[\text{Ru}^{\text{III}}(\text{tPaO-}\kappa\text{-N}^2\text{O}_p\text{O}_c)(\text{py})_2]^{2-}$, $\mathbf{4}^{2-}$, via oxygen insertion and (ii) the intramolecular proton transfer from the incoming water molecule for the O-O bond formation step that is the rds in the WNA mechanism. These two key reactions occur intramolecularly thanks to the right positioning of the $\text{Ru}^{\text{V}}=\text{O}$ groups versus the dangling phosphonate group and thus are entropically highly favored.

The present work highlights the importance of designing catalysts with the right *second coordination sphere environment* in the field of redox catalysis and in particular in the catalytic oxidation of water to dioxygen and the fate of the catalysts during turnover.


4. Supporting Information. Additional experimental details, electrochemical and spectroscopic data. CIF files for complexes $\{[\text{Ru}^{\text{II}}(\text{H}_3\text{tPa-}\kappa\text{-N}^3\text{O})(\text{py})_2](\text{H}_2\text{O})_3\}(\text{PF}_6)$ and $\{[\text{Ru}^{\text{II}}(\text{tPaO-}\kappa\text{-N}^2\text{O}_p\text{O}_c)(\text{py})_2](\text{H}_2\text{O})_{8.5}(\text{CH}_3\text{OH})\}\text{Cs}_3$, with CCDC numbers 1890782 and 1954776 respectively are available at <https://www.ccdc.cam.ac.uk/>.

5. Acknowledgements


Support from MINECO, FEDER, "La Caixa" and AGAUR are gratefully acknowledged (CTQ2016-80058-R, CTQ2015-73028-EXP, SEV 2013-0319, ENE2016-82025-REDT, CTQ2016-81923-REDC, and 2017-SGR-1631). The work at Brookhaven National Laboratory (BNL; M.Z.E) was carried out under contract DE-SC0012704 with the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, and utilized computational resources at the Center for Functional Nanomaterials, which is

1
2
3 a U.S. DOE Office of Science Facility, and the Scientific Data and Computing Center, a component of
4 the Computational Science Initiative, at BNL under Contract No. DE-SC0012704.
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


Chart 1. Drawing and labeling of ligands discussed in this work.




H₂tda



H₂bda



H₄tPa



H₅tPaO

Scheme 1. Computed reaction pathway at pH 7.0 for the generation of the catalytically active species $[\text{Ru}^{\text{III}}(\text{tPaO-}\kappa\text{-N}^2\text{O}_p\text{O}_C)(\text{py})_2]^{2-}$, $\mathbf{4}^{2-}$, from the precursor complex $[\text{Ru}^{\text{II}}(\text{H}_2\text{tPa-}\kappa\text{-N}^3\text{O})(\text{py})_2]$, $\mathbf{2}$. Redox potentials (E) in units of volts (V) vs NHE, ΔG s and ΔG^\ddagger in units of kcal/mol. Axial pyridyl ligands are omitted for clarity.




Figure 1. Synthetic scheme and structures of the complexes described in this work. X-ray structure ORTEP views for $\{[\text{Ru}^{\text{II}}(\text{H}_3\text{tPa}-\kappa\text{-N}^3\text{O})(\text{py})_2](\text{H}_2\text{O})_3\}^+$, $2^+ \cdot 3\text{H}_2\text{O}$, showing the 3 hydrogen bonded water molecules and for $[\text{Ru}^{\text{III}}(\text{tPaO}-\kappa\text{-N}^2\text{O}_p\text{O}_c)(\text{py})_2]^{2-}$, 4^{2-} . Calculated structure for $[\text{Ru}^{\text{IV}}(\text{tPa}-\kappa\text{-N}^3\text{O}^2)(\text{py})_2]$, **3**. Color code: Ru, cyan; P, pink; N, blue; O, red; C, black; H, white.




Figure 2. VT $^1\text{H-NMR}$ of $[\text{Ru}^{\text{II}}(\text{H}_2\text{tPa-}\kappa\text{-N}^3\text{O})(\text{py})_2]$, **2** in CD_3OD together with labeling.




Figure 3. Left, cyclic voltammetry of 0.6 mM $[\text{Ru}^{\text{II}}(\text{HtPa}-\kappa\text{-N}^3\text{O})(\text{py})_2]^-$, **2**⁻, at pH 7.0 in a 0.1 M phosphate buffer aqueous solution (red trace) and at pH = 1.0 in 0.1 M triflic acid (purple trace). Solid line black trace, cyclic voltammetry of 0.6 mM $[\text{Ru}^{\text{III}}(\text{tPaO}-\kappa\text{-N}^2\text{O}_p\text{O}_c)(\text{py})_2]^{2-}$, **4**²⁻, at pH 7.0 in a 0.1 M phosphate buffer aqueous solution, generated from the precursor **2**⁻, by a 3h bulk electrolysis at 1.30 V. Right, Pourbaix diagram for **2**. The dominant species as a function of pH and potential are indicated with the labels such as $[\text{Ru}^{\text{II}}(\text{H}_2\text{tPa})]$ where the oxidation state is indicated as well as the degree of protonation of the H_4tPa ligand. The dashed vertical lines indicate the pK_a of the species involved. The axial pyridyl ligands are omitted for clarity.






Figure 4. Top, calculated catalytic cycle for catalyst $[\text{Ru}^{\text{V}}(\text{O})(\text{HtPaO}-\kappa\text{-N}^2\text{O}_\text{C})(\text{py})_2]^-$ at pH 7.0 Redox potentials (E) (experimental in green and calculated in blue) in units of volts (V) vs NHE, ΔG s and ΔG^\ddagger in units of kcal/mol. Axial pyridyl ligands are omitted for clarity. Bottom, optimized TS for the O-O bond formation step. Color code: Ru, cyan; P, orange; N, blue; O, red; C, black; H, white.



- 1
- 2
- 3 (1) Berardi, S.; Drouet, S.; Francàs, L.; Gimbert-Suriñach, C.; Guttentag, M.; Richmond, C.; Stoll,
- 4 T.; Llobet, A. Molecular Artificial Photosynthesis. *Chem. Soc. Rev.* **2014**, *43* (22), 7501–7519.
- 5 (2) Lewis, N. S. Research Opportunities to Advance Solar Energy Utilization. *Science* (80-.). **2016**,
- 6 *351* (6271), aad1920.
- 7 (3) Alstrum-Acevedo, J. H.; Brennaman, M. K.; Meyer, T. J. Chemical Approaches to Artificial
- 8 Photosynthesis. 2. *Inorg. Chem.* **2005**, *44* (20), 6802–6827.
- 9 (4) Kärkäs, M. D.; Johnston, E. V.; Verho, O.; Åkermark, B. Artificial Photosynthesis: From
- 10 Nanosecond Electron Transfer to Catalytic Water Oxidation. *Acc. Chem. Res.* **2014**, *47* (1),
- 11 100–111.
- 12 (5) Cook, T. R.; Dogutan, D. K.; Reece, S. Y.; Surendranath, Y.; Teets, T. S.; Nocera, D. G. Solar
- 13 Energy Supply and Storage for the Legacy and Nonlegacy Worlds. *Chem. Rev.* **2010**, *110* (11),
- 14 6474–6502.
- 15 (6) Walter, M. G.; Warren, E. L.; McKone, J. R.; Boettcher, S. W.; Mi, Q.; Santori, E. A.; Lewis, N.
- 16 S. Solar Water Splitting Cells. *Chem. Rev.* **2010**, *110* (11), 6446–6473.
- 17 (7) De Luna, P.; Hahn, C.; Higgins, D.; Jaffer, S. A.; Jaramillo, T. F.; Sargent, E. H. What Would It
- 18 Take for Renewably Powered Electrosynthesis to Displace Petrochemical Processes? *Science*
- 19 (80-.). **2019**, *364* (6438), eaav3506.
- 20 (8) Garrido-Barros, P.; Gimbert-Suriñach, C.; Matheu, R.; Sala, X.; Llobet, A. How to Make an
- 21 Efficient and Robust Molecular Catalyst for Water Oxidation. *Chem. Soc. Rev.* **2017**, *46* (20),
- 22 6088–6098.
- 23 (9) Matheu, R.; Garrido-Barros, P.; Gil-Sepulcre, M.; Ertem, M. Z.; Sala, X.; Gimbert-Suriñach, C.;
- 24 Llobet, A. The Development of Molecular Water Oxidation Catalysts. *Nat. Rev. Chem.* **2019**,
- 25 *3* (5), 331–341.
- 26 (10) Sala, X.; Maji, S.; Bofill, R.; García-Antón, J.; Escriche, L.; Llobet, A. Molecular Water Oxidation
- 27 Mechanisms Followed by Transition Metals: State of the Art. *Acc. Chem. Res.* **2014**, *47* (2),
- 28 504–516.
- 29 (11) Gimbert-Suriñach, C.; Moonshiram, D.; Francàs, L.; Planas, N.; Bernales, V.; Bozoglian, F.;
- 30 Guda, A.; Mognon, L.; López, I.; Hoque, M. A.; et al. Structural and Spectroscopic
- 31 Characterization of Reaction Intermediates Involved in a Dinuclear Co-Hbpp Water Oxidation
- 32 Catalyst. *J. Am. Chem. Soc.* **2016**, *138* (47), 15291–15294.
- 33 (12) Romain, S.; Bozoglian, F.; Sala, X.; Llobet, A. Oxygen-Oxygen Bond Formation by the Ru-Hbpp
- 34 Water Oxidation Catalyst Occurssolely via an Intramolecular Reaction Pathway. *J. Am. Chem.*
- 35 *Soc.* **2009**, *131* (8), 2768–2769.
- 36 (13) Shaffer, D. W.; Xie, Y.; Concepcion, J. J. O–O Bond Formation in Ruthenium-Catalyzed Water
- 37 Oxidation: Single-Site Nucleophilic Attack vs. O–O Radical Coupling. *Chem. Soc. Rev.* **2017**, *46*
- 38 (20), 6170–6193.
- 39 (14) Wang, L. P.; Van Voorhis, T. Direct-Coupling O2 Bond Forming a Pathway in Cobalt Oxide
- 40 Water Oxidation Catalysts. *J. Phys. Chem. Lett.* **2011**, *2* (17), 2200–2204.
- 41 (15) McCrory, C. C. L.; Jung, S.; Ferrer, I. M.; Chatman, S. M.; Peters, J. C.; Jaramillo, T. F.
- 42 Benchmarking Hydrogen Evolving Reaction and Oxygen Evolving Reaction Electrocatalysts
- 43 for Solar Water Splitting Devices. *J. Am. Chem. Soc.* **2015**, *137* (13), 4347–4357.
- 44 (16) Hong, W. T.; Stoerzinger, K. A.; Lee, Y. L.; Giordano, L.; Grimaud, A.; Johnson, A. M.; Hwang,
- 45 J.; Crumlin, E. J.; Yang, W.; Shao-Horn, Y. Charge-Transfer-Energy-Dependent Oxygen
- 46 Evolution Reaction Mechanisms for Perovskite Oxides. *Energy Environ. Sci.* **2017**, *10* (10),
- 47 2190–2200.
- 48 (17) Surendranath, Y.; Kanan, M. W.; Nocera, D. G. Mechanistic Studies of the Oxygen Evolution
- 49 Reaction by a Cobalt-Phosphate Catalyst at Neutral PH. *J. Am. Chem. Soc.* **2010**, *132* (46),
- 50 16501–16509.
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1
2
3 (18) Matheu, R.; Ertem, M. Z.; Gimbert-Suriñach, C.; Sala, X.; Llobet, A. Seven Coordinated
4 Molecular Ruthenium–Water Oxidation Catalysts: A Coordination Chemistry Journey. *Chem.*
5 *Rev.* **2019**, *119* (6), 3453–3471.
6 (19) Matheu, R.; Ertem, M. Z.; Benet-Buchholz, J.; Coronado, E.; Batista, V. S.; Sala, X.; Llobet, A.
7 Intramolecular Proton Transfer Boosts Water Oxidation Catalyzed by a Ru Complex. *J. Am.*
8 *Chem. Soc.* **2015**, *137* (33), 10786–10795.
9 (20) Richmond, C. J.; Matheu, R.; Poater, A.; Falivene, L.; Benet-Buchholz, J.; Sala, X.; Cavallo, L.;
10 Llobet, A. Supramolecular Water Oxidation with Rubda-Based Catalysts. *Chem. - A Eur. J.*
11 **2014**, *20* (52), 17282–17286.
12 (21) Duan, L.; Bozoglian, F.; Mandal, S.; Stewart, B.; Privalov, T.; Llobet, A.; Sun, L. A Molecular
13 Ruthenium Catalyst with Water-Oxidation Activity Comparable to That of Photosystem II.
14 *Nat. Chem.* **2012**, *4* (5), 418–423.
15 (22) Song, N.; Concepcion, J. J.; Binstead, R. A.; Rudd, J. A.; Vannucci, A. K.; Dares, C. J.; Coggins,
16 M. K.; Meyer, T. J. Base-Enhanced Catalytic Water Oxidation by a Carboxylate–Bipyridine
17 Ru(II) Complex. *Proc. Natl. Acad. Sci.* **2015**, *112* (16), 4935–4940.
18 (23) Xie, Y.; Shaffer, D. W.; Lewandowska-Andralojc, A.; Szalda, D. J.; Concepcion, J. J. Water
19 Oxidation by Ruthenium Complexes Incorporating Multifunctional Bipyridyl Diphosphonate
20 Ligands. *Angew. Chemie - Int. Ed.* **2016**, *55* (28), 8067–8071.
21 (24) Kamdar, J. M.; Marelius, D. C.; Moore, C. E.; Rheingold, A. L.; Smith, D. K.; Grotjahn, D. B.
22 Ruthenium Complexes of 2,2'-Bipyridine-6,6'-Diphosphonate Ligands for Water Oxidation.
23 *ChemCatChem* **2016**, *8* (19), 3045–3049.
24 (25) Shaffer, D. W.; Xie, Y.; Szalda, D. J.; Concepcion, J. J. Lability and Basicity of Bipyridine-
25 Carboxylate-Phosphonate Ligand Accelerate Single-Site Water Oxidation by Ruthenium-
26 Based Molecular Catalysts. *J. Am. Chem. Soc.* **2017**, *139* (43), 15347–15355.
27 (26) Surendranath, Y.; Dincă, M.; Nocera, D. G. Electrolyte-Dependent Electrosynthesis and
28 Activity of Cobalt-Based Water Oxidation Catalysts. *J. Am. Chem. Soc.* **2009**, *131* (7), 2615–
29 2620.
30 (27) Matheu, R.; Ertem, M. Z.; Gimbert-Suriñach, C.; Benet-Buchholz, J.; Sala, X.; Llobet, A.
31 Hydrogen Bonding Rescues Overpotential in Seven-Coordinated Ru Water Oxidation
32 Catalysts. *ACS Catal.* **2017**, *7* (10), 6525–6532.
33 (28) Albrecht-Schmitt, T.; Bujoli, B.; Cahill, C.; Murugavel, R.; Rocha, J.; Hix, G.; Shimizu, G.;
34 Zubietta, J.; Zon, J.; Brunet, E.; et al. *Metal Phosphonate Chemistry*; The Royal Society of
35 Chemistry, 2011.
36 (29) Hammes-Schiffer, S. Catalysts by Design: The Power of Theory. *Acc. Chem. Res.* **2017**, *50* (3),
37 561–566.
38 (30) Meyer, T. J.; Sheridan, M. V.; Sherman, B. D. Mechanisms of Molecular Water Oxidation in
39 Solution and on Oxide Surfaces. *Chem. Soc. Rev.* **2017**, *46* (20), 6148–6169.
40 (31) Huynh, M. H. V.; Meyer, T. J. Proton-Coupled Electron Transfer. *Chem. Rev.* **2007**, *107* (11),
41 5004–5064.
42 (32) Weinberg, D. R.; Gagliardi, C. J.; Hull, J. F.; Murphy, C. F.; Kent, C. A.; Westlake, B. C.; Paul, A.;
43 Ess, D. H.; McCafferty, D. G.; Meyer, T. J. Proton-Coupled Electron Transfer. *Chem. Rev.* **2012**,
44 *112* (7), 4016–4093.
45 (33) Warren, J. J.; Tronic, T. A.; Mayer, J. M. Thermochemistry of Proton-Coupled Electron
46 Transfer Reagents and Its Implications. *Chem. Rev.* **2010**, *110* (12), 6961–7001.
47 (34) Shaffer, D. W.; Xie, Y.; Concepcion, J. J. O–O Bond Formation in Ruthenium-Catalyzed Water
48 Oxidation: Single-Site Nucleophilic Attack: Vs. O–O Radical Coupling. *Chem. Soc. Rev.* **2017**,
49 *46* (20), 6170–6193.
50 (35) Hammes-Schiffer, S.; Soudackov, A. V. Proton-Coupled Electron Transfer in Solution,
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Proteins, and Electrochemistry. *J. Phys. Chem. B* **2008**, *112* (45), 14108–14123.
- 4 (36) Laurent, F.; Plantalech, E.; Donnadiou, B.; Jiménez, A.; Hernández, F.; Martínez-Ripoll, M.;
5 Biner, M.; Llobet, A. Synthesis, Structure and Redox Properties of Ruthenium Complexes
6 Containing the Tpm Facial and the Trpy Meridional Tridentate Ligands: Crystal Structures of
7 [RuCl₃(Trpy)] and [Ru(Tpm)(Py)₃](PF₆)₂. *Polyhedron* **1999**, *18* (25), 3321–3331.
- 8 (37) Kuppuraj, G.; Dudev, M.; Lim, C. Factors Governing Metal–Ligand Distances and Coordination
9 Geometries of Metal Complexes. *J. Phys. Chem. B* **2009**, *113* (9), 2952–2960.
- 10 (38) Zhao, Y.; Truhlar, D. G. The M06 Suite of Density Functionals for Main Group
11 Thermochemistry, Thermochemical Kinetics, Noncovalent Interactions, Excited States, and
12 Transition Elements: Two New Functionals and Systematic Testing of Four M06-Class
13 Functionals and 12 Other Function. *Theor. Chem. Acc.* **2008**, *120* (1), 215–241.
- 14 (39) Marenich, A. V.; Cramer, C. J.; Truhlar, D. G. Supporting Information (PART I) Universal
15 Solvation Model Based on Solute Electron Density and on a Continuum Model of the Solvent
16 Defined by the Bulk Dielectric Constant and Atomic Surface Tensions Contents : *J. Phys.*
17 *Chem. B* **2009**, *113*, 6378.
- 18 (40) Hoque, M. A.; Benet-Buchholz, J.; Llobet, A.; Gimbert-Suriñach, C. Catalytic Oxidation of
19 Water to Dioxygen by Mononuclear Ru Complexes Bearing a 2,6-Pyridinedicarboxylato
20 Ligand. *ChemSusChem* **2019**, *12* (9), 1949–1957.
- 21 (41) Matheu, R.; Ertem, M. Z.; Pipelier, M.; Lebreton, J.; Dubreuil, D.; Benet-Buchholz, J.; Sala, X.;
22 Tessier, A.; Llobet, A. The Role of Seven-Coordination in Ru-Catalyzed Water Oxidation. *ACS*
23 *Catal.* **2018**, *8* (3), 2039–2048.
- 24 (42) Matheu, R.; Ghaderian, A.; Francàs, L.; Chernev, P.; Ertem, M. Z.; Benet-Buchholz, J.; Batista,
25 V. S.; Haumann, M.; Gimbert-Suriñach, C.; Sala, X.; et al. Behavior of Ru–Bda Water-Oxidation
26 Catalysts in Low Oxidation States. *Chem. – A Eur. J.* **2018**, *24* (49), 12838–12847.
- 27 (43) Brauman, J. I.; Bryson, J. A.; Kahl, D. C.; Nelson, N. J. Equilibrium Acidities in Dimethyl
28 Sulfoxide. *J. Am. Chem. Soc.* **1970**, *92* (22), 6679–6680.
- 29 (44) Nazeeruddin, M. K.; Zakeeruddin, S. M.; Humphry-Baker, R.; Kaden, T. A.; Grätzel, M.
30 Determination of PKa Values of 4-Phosphonato-2,2':6',2''-Terpyridine and Its Ruthenium(II)-
31 Based Photosensitizer by NMR, Potentiometric, and Spectrophotometric Methods. *Inorg.*
32 *Chem.* **2000**, *39* (20), 4542–4547.
- 33 (45) Ordóñez, M.; Viveros-Ceballos, J. L.; Sayago, F. J.; Cativiela, C. Stereoselective Synthesis of α -
34 Amino- H -Phosphinic Acids and Derivatives. *Synth.* **2017**, *49* (5), 987–997.
- 35 (46) Costentin, C.; Drouet, S.; Robert, M.; Savéant, J.-M. Turnover Numbers, Turnover
36 Frequencies, and Overpotential in Molecular Catalysis of Electrochemical Reactions. Cyclic
37 Voltammetry and Preparative-Scale Electrolysis. *J. Am. Chem. Soc.* **2012**, *134* (27), 11235–
38 11242.
- 39 (47) Rountree, E. S.; McCarthy, B. D.; Eisenhart, T. T.; Dempsey, J. L. Evaluation of Homogeneous
40 Electrocatalysts by Cyclic Voltammetry. *Inorg. Chem.* **2014**, *53* (19), 9983–10002.
- 41 (48) Matheu, R.; Neudeck, S.; Meyer, F.; Sala, X.; Llobet, A. Foot of the Wave Analysis for
42 Mechanistic Elucidation and Benchmarking Applications in Molecular Water Oxidation
43 Catalysis. *ChemSusChem* **2016**, *9* (23), 3361–3369.
- 44 (49) Wasylenko, D. J.; Ganesamoorthy, C.; Henderson, M. A.; Koivisto, B. D.; Osthoff, H. D.;
45 Berlinguette, C. P. Electronic Modification of the [Ru(II)(Tpy)(Bpy)(OH₂)]²⁺ Scaffold: Effects on
46 Catalytic Water Oxidation. *J. Am. Chem. Soc.* **2010**, *132* (45), 16094–16106.
- 47 (50) Wang, J. W.; Sahoo, P.; Lu, T. B. Reinvestigation of Water Oxidation Catalyzed by a Dinuclear
48 Cobalt Polypyridine Complex: Identification of CoOxas a Real Heterogeneous Catalyst. *ACS*
49 *Catal.* **2016**, *6* (8), 5062–5068.
- 50 (51) Matheu, R.; Ertem, M. Z.; Gimbert-Suriñach, C.; Sala, X.; Llobet, A. Seven Coordinated
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Molecular Ruthenium-Water Oxidation Catalysts: A Coordination Chemistry Journey. *Chem. Rev.* **2019**, *119* (6), 3453–3471.
- 4
5 (52) Moyer, B. A.; Thompson, M. S.; Meyer, T. J. Chemically Catalyzed Net Electrochemical
6 Oxidation of Alcohols, Aldehydes, and Unsaturated Hydrocarbons Using the System
7 (Trpy)(Bpy)Ru(OH₂)₂²⁺/(Trpy)(Bpy)RuO₂⁺. *J. Am. Chem. Soc.* **1980**, *102* (7), 2310–2312.
- 8
9 (53) Wasylenko, D. J.; Ganesamoorthy, C.; Henderson, M. A.; Koivisto, B. D.; Osthoff, H. D.;
10 Berlinguette, C. P. Supporting Information: Electronic Modification of the [Ru II
11 (Tpy)(Bpy)(OH₂)₂]²⁺ Scaffold: Effects on Catalytic Water Oxidation. *J. Am. Chem. Soc.* **2010**,
12 *132* (45), 16094–16106.
- 13
14 (54) López, I.; Maji, S.; Benet-Buchholz, J.; Llobet, A. Oxo-Bridge Scenario behind Single-Site
15 Water-Oxidation Catalysts. *Inorg. Chem.* **2015**, *54* (2), 658–666.
- 16
17 (55) Takeuchi, K. J.; Thompson, M. S.; Pipes, D. W.; Meyer, T. J. Redox and Spectral Properties of
18 Monooxo Polypyridyl Complexes of Ruthenium and Osmium in Aqueous Media. *Inorg. Chem.*
19 **1984**, *23* (13), 1845–1851.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TOC

