## APODICTIC DISCOURSE IN ANCIENT AND MODERN MATHEMATICS

#### Satyanad KICHENASSAMY

Laboratoire de Mathématiques de Reims (CNRS UMR 9008, Université de Reims Champagne-Ardenne, France)

Tank delivered at the International Workshop "Srinivas Ramanujan: The Man Beyond Infinity", Dec. 22, 2020, organized by the Central University of Himachal Pradesh, India.

### ISSUE

- Many important Indian authors, in particular S. Ramanujan, appear not to have justified their results. If it is more that an induction, there must be an implied derivation.
- We show that, in some cases, the derivation was encoded through the discursive structure itself. This is what we call an apodictic discourse.
- Examples: Brahmagupta, Ramanujan, Hardy, Gauss, Tartaglia...

## **MATHEMATICAL DISCOURSE**

...may belong to all categories but

- research mathematicians prefer apodictic discourse...
- ...because it enables imparting motivation, history and derivation,...
- ...allows reading at different levels...
- ...and makes the continuity of knowledge clear.
- For Ramanujan : may be useful to scrutinize sequences of propositions even when they are not formally proved.

#### ARGUMENT

A. Mathematics made public is a discourse

- B. Case study: Brahmagupta's "triquadrilateral". [How do you analyze apodictic discourse.]
- C. Ramanujan's discursive strategies (examples).

For Ramanujan, almost all sources are now available: Berndt (*Notebooks*), Hardy, R.'s Collected papers. But R.'s results have often been verified, seldom derived.

#### **A. MATHEMATICS AS DISCOURSE**

## **ARGUMENTATION IS DISCURSIVE**

- Mathematical activity is in part non-discursive (Hadamard), but precise communication is a discourse.
- Non-rigorous discourse (including comments on tables, diagrams etc.) may have heuristic value; can also be misleading.
- Ex.: a figure may suggest an argument, it is not an argument.
- Ex.: Lists of rules? Procedures? Tables, memory aids? None of these is in itself a discourse.

## **ARGUMENTATION IS DISCURSIVE**

- Mathematical discourse implies an *audience* and a structured *argumentation*:
  - Is the audience oneself (math'l diaries, private notes, etc.)?
  - Is it rigorous [essential arguments present] and if so,
    - Is it aimed at imparting some content to a group to which the author does not belong (*dogmatic discourse*)?
    - Or does it indicate motivation and steps of derivations for scholars with about the same level of knowledge (*apodictic discourse*)?

### FORMS OF MATHEMATICAL DISCOURSE

- 1. Rigorous : Apodictic or dogmatic discourse
- 2. Suggestive, but non-rigorous (possibly misleading) : heuristic d., memory aids etc.
- **3.** Inconclusive: arguments based on sense perception ("cas de figure"), analogies,...
- 4. Communication in M. is through discourses.
- 5. Analysis of earlier work is a discourse analysis.

What binds mathematicians is the experience that rigorous discourse is possible.

### DOGMATIC AND APODICTIC DISCOURSES ARE COMPLEMENTARY

- Archimedes seems (Wallis, quoted by Heath, HGM II, 20) "as it were of set purpose to have covered up the traces of his investigation, as if he had grudged posterity the secret of his method of inquiry, while he wished to extort from them assent to his results". "...not only Archimedes but nearly all the ancients so hid from posterity their method of Analysis... that more modern mathematicians found it easier to invent a new Analysis than to seek out the old." Dogmatic discourse insufficient.
- Tartaglia's discourse in the *Quesiti* not understood by Cardano. Apodictic discourse requires an audience with the same background as the author.
- Neither type of discourse is "universal" since either may fail to be understood by some. But apodictic discourse may be explicated by internal analysis, dogmatic discourse may be verified. (There are intermediate types.)

### **EXAMPLES**

Refs.: https://www.normalesup.org/~kichenassamy

(How Brahmagupta obtained his results on the cyclic quadrilateral)

- *Historia Mathematica*, <u>37(</u>1) (2010) 28-61.
- Historia Mathematica, <u>39(4)</u>, (2012) 387-404.

(How Brahmagupta obtained his results on congruences)

• Gaņita Bhāratī, to appear

(How Tartaglia obtained his results on cubic equations)

• Historia Mathematica, 2015, <u>42</u> (4), 407-435.

(How Baudhāyana obtained his approx. quadrature of the circle)

• Historia Mathematica, <u>33</u>, 2006, pp.149-183.

#### **REFERENCES : DISCURSIVITY IN MATHEMATICS AND PHILOSOPHY**

- Comptes-rendus des séances de l'Académie des inscriptions et belles-lettres, Paris : Durand, <u>2012-II</u> (4), pp.781-796.
- Gaņita Bhāratī, (2011).
- Journal Asiatique, 2018, 306 (1), 85-99.
- « L'emploi métonymique de l'arbre kallāl dans la philosophie médiévale en pays tamoul. » P.-S. Filliozat et M. Zink (eds.). L'Arbre en Asie, De Boccard, Paris, 2018, pp.279-299.
- « L'irruption de l'infini : la légende de la colonne de lumière » (= lingodbhava), Comptes-Rendus des Séances de l'Académie des Inscriptions et Belles-Lettres, <u>2018</u> (4), to appear.

## **DISCOURSE ANALYSIS**

How to analyze an apodictic discourse :

- Need to know it is not merely heuristic or borrowed etc.
- Need to have an idea of the author's previous knowledge...
- ... and standard practices.
- If so, perform close reading (every element of the text, including its form, must be significant).

Upshot :

- You solve old problems in the History of Mathematics.
- You tend to read every mathematical work more closely...
- ... and may obtain new results by following the argument implied by earlier papers.

#### **B. CASE STUDY: BRAHMAGUPTA'S TRIQUADRILATERAL**

## BRAHMAGUPTA'S DISCOURSE IN B.S.S. XII.21-38

- **Objective** : given a cyclic quadrilateral,
  - express in terms of its sides all the lines generated by projection, extension and intersection of already constructed segments, (XII.21-32) and
  - parameterize all possible sets of sides (XII.33-38).
- XII.21 states the area formula; XII.21-27 give elements of derivation; XII.28 the diagonals, and XII.29-32, all perpendiculars, projections etc.
- Derivation of area formula : the triquadrilateral is naturally split into two triangles with the same circumcenter and the same circumradius.

## QUESTIONS

- XII.24 expresses the theorem that the "trilateral in a semi-circle" is right.
- XII.25 gives a general result on similar halfoblongs.
- XII.30-31 ("Brahmagupta's theorem") unknown outside India (Chasles, 1837).

This proves that mathematics is not cumulative.

**Questions:** How is the derivation motivated? Where did he specify that the quadrilateral was cyclic [= inscribed in a circle of unspecified radius] ?

## **BRAHMAGUPTA'S ARGUMENT**

The cyclic character is indicated by a neologism "*tricaturbhuja*", *lit.* triquadrilateral (XII.21), whose meaning is fixed by XII.27.

#### **Proof:**

- Term occurs only twice: XII.21, XII.27 : specific to Brahmagupta and to this situation.
- XII.27 tells how to compute the common circumradius of a trilateral and a triquadrilateral: it is inscribed, it is not a trilateral, and it contains a distinguished triangle.

**Conclusion** : the triquadrilateral is a quadrilateral obtained by completing a trilateral by a point arbitrarily taken on its circumcircle.





#### 12.21

- 12.21 sthūlaphalam tricaturbhujabāhupratibāhuyogadalaghātaķ bhujayogārdhacatuṣṭayabhujonaghātāt padam sūkṣmam
- A crude value [indeed] of the area of a triquadrilateral
- Is the product of the half-sums of opposite sides ;
- Of a group consisting of four half-sums of the sides, from which
- The sides have been subtracted [in turn], the root of the product is the refined [value].

#### **AREA OF A TRIQUADRILATERAL**

a, b, c, d : sides, s = (a + b + c + d)/2 $A \cong \frac{a+c}{2} \times \frac{b+d}{2}$  [(a, c), (b, d) pairs of opposite sides]

 $A=\sqrt{(s-a)(s-b)(s-c)(s-d)}$   $s-d = \frac{a+b+c-d}{2} = \frac{a+c}{2} + \frac{b-d}{2}$   $s-b = \frac{a-b+c+d}{2} = \frac{a+c}{2} - \frac{b-d}{2}$   $(s-b)(s-d) = \left[\left(\frac{a+c}{2}\right)^2 - \left(\frac{b-d}{2}\right)^2\right], \dots$ 

Hence : gross formula ≥ exact value

#### XII.21 : THE ORDER OF THE SIDES IS IMMATERIAL

Quadrilateral = cercle with four regions called « pradhi »s in the Śulva-s



## XII.27 : CIRCUMCIRCLE

tribhujasya vadho bhujayor dviguņita-lamboddhrto hrdayarajjuķ sā dviguņā tricaturbhujakoņa-sprg-vrtta-viṣkambhaḥ

In the trilateral, the product of the sides divided by the perpendicular multiplied by two, Is the cord of the heart, Which, times two, is, in the triquadrilateral, the diameter of the cercle that touches [its] corners.

- 12.21 Triquadrilateral is a closed figure (has an area) that has four sides.
- 12.27 It has a circumcircle determined by the height of a triangle : the triquadrilateral is obtained by completing a triangle by adding a fourth point on its circumcircle.
- NB : 12.26 describes two special cases, in which the fourth point is obtained from one of the vertices by symmetry.

#### TRIQUADRILATERAL



To find the perpendicular in XII.27, the triquadrilateral must contain a distinguished triangle.

## OTHER REMARKABLE PASSAGES

- XII.24 expresses the theorem that the "trilateral in a semicircle" is right.
- XII.25 gives a general result on similar half-oblongs.
- XII.30-31 ("Brahmagupta's theorem") unknown outside India (Chasles, 1837).
- The last part of XII gives a proof of "Gauss' Lemma" (if a | bc and a/b irreducible, then a | c)
- XVIII extends operations from rational numbers to negatives, constructible quadratic irrationals and "nonmanifest" quantities (theoretical basis of algebra, developed in his theory of equations).

Mathematics not cumulative. What is forgotten can only be recovered by historical investigation (in the modern sense).

# **C. RAMANUJAN'S DISCURSIVE STRATEGIES**

## ORIENTATION

- Compare S. Ramanujan's discursive strategies to that of other mathematicians.
- His approach is not unlike those of others, but he was not part of an extensive community of scholars [did not undergo regular training], except possibly at the end of his life.
- Upshot: better understanding of his intended meaning and possibly carry it further.
- NB : Part of R.'s work is not discursive & may not have been meant for communication: more complicated than the case of Brahmagupta.

#### **DISCURSIVE STRATEGIES**

- Ramanujan: "a rational human being who happened to be a great mathematician" (H. p. 5), "not in the least disposed to make a mystery of his achievements" (p. 11).
- Defensive: give only results. « If I had given you my methods of proof I am sure you will follow the London Professor » [who was convinced he did not understand the notion of convergence] (R to H, 27 Feb. 1913),
- Heuristic (for himself first and foremost): not always definitive results. L on R, March 1913 : « I imagine that he is satisfied if he can convince himself that his results are correct »; H on R (p. 27) "He had a "proof", a definite and very ingenious train of reasoning." We want to remove the quotes !
- Hardy regretted not having asked R. about his reasoning ("I hardly asked him a single question of this kind" p. 11)
- Carr's Synopsis (1880-6) as a model ?

## **CARR'S SYNOPSIS (PREFACE)**

"I have, in many cases, merely indicated the salient points of a demonstration, or merely referred to the theorems by which the proposition is proved. I am convinced that it is more beneficial to the student to recall demonstrations with such aids, than to read and re-read them."

#### (i.e.: conducive to active learning)

But: hybrid format (idiosyncratic, preparation for Tripos)

- Structured like an Indian text (!) beginning with units (for physical quantities), putting algebra foremost etc.
- But includes a list of propositions from Books of Euclid for reference.
- Lacks motivation: (a) why is this interesting ? (b) how does one reach this result? (c) why choose this proof ?

## « ON HIGHLY COMPOSITE NUMBERS »

- R. adopts the dogmatic style (PLMS 2, XIV, 1915, 347-409 = Paper 15, p. 78)
- H. explains the motivation: (ch. III "Round numbers") [round nb. = "the product of a considerably large number of comparatively small factors"]...

... "It is a matter of common observation that *round numbers are very rare;* the fact may be verified by anyone who will make a habit of factorising numbers, such as numbers of motor cars of railway carriages, which are presented to his attention in a random manner. Both Ramanujan and I had observed this phenomenon, which seems at first a little paradoxical." + footnote: "Half the numbers are divisible by 2, ...one-sixth by both 2 and 3. Surely then we may them expect numbers to have a *large* number of factors? But the facts seem to show the opposite. (H., ibid, p. 55)

## FROM THE NOTEBOOKS: ENTRIES 20(II-III)

Notebooks, vol. III (Berndt, 1991, p. 199)

- Entry 20(i): construction of  $\frac{355}{113} \approx \pi$ ; Cor. (i) [equilateral tr. inscribed in a circle:  $\sqrt[3]{31} = 3.14138 \dots \approx \pi$ ) and (ii)  $\sqrt[4]{97\frac{1}{2}-\frac{1}{11}} \approx \pi$
- "The appearance below of Entries 20(ii) and 20(iii) is enigmatic indeed; there does not seem to be any connection between these entries and any other result in Chapter 18."
- Entry 20(ii) : "Parametric solutions of the equation  $A^3 + B^3 = C^2$  are given by  $A = 3n^3 + 6n^2 n$ ,  $B = -3n^3 + 6n^2 + n$  and  $C = 6n^2(3n^2 + 1)$  where *n* is arbitrary

## FROM THE NOTEBOOKS: ENTRY 20(III)

Entry 20(ii) : "Parametric solutions of the equation  $A^3 + B^3 = C^2$  are given by  $A = 3n^3 + 6n^2 - n$ ,  $B = -3n^3 + 6n^2 + n$  and  $C = 6n^2(3n^2 + 1)$  where *n* is arbitrary

Entry 20(iii) : "Parametric solutions of the equation  $A^3 + B^3 + C^3 = D^3$  are given by

$$A = m^{7} - 3(p + l)m^{4} + (3p^{2} + 6p + 2)m,$$
  

$$B = 2m^{6} - 3(2p + 1)m^{3} + (3p^{2} + 3p + 1),$$
  

$$C = m^{6} - (3p^{2} + 3p + 1)$$
  

$$D = m^{7} - 3pm^{4} + (3p^{2} - 1)m,$$

where *m* and *p* denote arbitrary numbers."

**Berndt provides verification + many references.** 

## FROM THE NOTEBOOKS: ENTRY 20(II-III)

- Entry 20(ii) : "Parametric solutions of the equation  $A^3 + B^3 = C^2$  are given by  $A = 3n^3 + 6n^2 - n$ ,  $B = -3n^3 + 6n^2 + n$  and  $C = 6n^2(3n^2 + 1)$  where *n* is arbitrary
- Entry 20(iii) : "Parametric solutions of the equation  $A^3 + B^3 + C^3 = D^3$  are given by

$$A = m^{7} - 3(p + l)m^{4} + (3p^{2} + 6p + 2)m,$$
  

$$B = 2m^{6} - 3(2p + 1)m^{3} + (3p^{2} + 3p + 1),$$
  

$$C = m^{6} - (3p^{2} + 3p + 1),$$
  

$$D = m^{7} - 3pm^{4} + (3p^{2} - 1)m,$$

where m and p denote arbitrary numbers." + List of examples Parallel formulation : same method ?

## FROM THE NOTEBOOKS: A LIST OF SOLUTIONS IN 20(III)

- Entry 20(ii) : "Parametric solutions of the equation  $A^3 + B^3 = C^2$  are given by  $A = 3n^3 + 6n^2 n$ ,  $B = -3n^3 + 6n^2 + n$  and  $C = 6n^2(3n^2 + 1)$  where *n* is arbitrary
- A and B have the form  $an^3 + bn^2 n$  and  $-an^3 + bn^2 + n$
- ...optimized to kill odd powers in sum of cubes., hence

• 
$$C^2 = 6bn^4 \left[ 1 + \frac{1}{3} (b^2 - 6a)n^2 + (an^2)^2 \right]$$

- Need:  $6b = q^2$ ,  $b^2 6a = 6a$  hence  $12a = b^2 = \frac{q^4}{36}$
- Therefore q = 6r,  $b = 6r^2$  ...
- 1. Possibly, similar approach for 20(iii)
- 2. List of solutions of 20(iii) : last example has different status

## CONCLUSION

- 1. Mathematics, when communicated, is a discourse.
- 2. Rigorous results need not be made explicit (when talking to scholars)
- 3. Important texts (Brahmagupta, Tartaglia, S. Ramanujan,...) appear non-rigorous because they are not fully explicit.

<u>Conclusion</u>: 1) There are several types of rigorous discourse.

2) Textual analysis as practiced by modern historians is an essential part of mathematical activity and leads to new results.

# THANK YOU FOR YOUR ATTENTION