



HAL
open science

Delta 1-activated Notch inhibits muscle differentiation without affecting Myf5 and Pax3 expression in chick limb myogenesis

Marie-Claire Delfini, Estelle Hirsinger, Olivier Pourquié, Delphine Duprez

► **To cite this version:**

Marie-Claire Delfini, Estelle Hirsinger, Olivier Pourquié, Delphine Duprez. Delta 1-activated Notch inhibits muscle differentiation without affecting Myf5 and Pax3 expression in chick limb myogenesis. *Development* (Cambridge, England), 2000, 127 (23), pp.5213-5224. 10.1242/dev.127.23.5213 . hal-03091061

HAL Id: hal-03091061

<https://hal.science/hal-03091061v1>

Submitted on 15 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Delta 1-activated Notch inhibits muscle differentiation without affecting *Myf5* and *Pax3* expression in chick limb myogenesis

Marie-Claire Delfini¹, Estelle Hirsinger², Olivier Pourquié² and Delphine Duprez^{1,*}

¹Institut d'Embryologie Cellulaire et Moléculaire du CNRS (FRE2160) et du Collège de France, 49 bis avenue de la Belle Gabrielle, 94736 Nogent sur Marne Cedex, France

²Laboratoire de Génétique et de Physiologie du Développement (LGPD), Developmental Biology Institute of Marseille (IBDM) Campus de Luminy. Case 907, 13288 Marseille Cedex 09, France

*Author for correspondence (e-mail: duprez@infobiogen.fr)

Accepted 29 September; published on WWW 2 November 2000

SUMMARY

The myogenic basic helix-loop-helix (bHLH) transcription factors, *Myf5*, *MyoD*, myogenin and *MRF4*, are unique in their ability to direct a program of specific gene transcription leading to skeletal muscle phenotype. The observation that *Myf5* and *MyoD* can force myogenic conversion in non-muscle cells in vitro does not imply that they are equivalent. In this paper, we show that *Myf5* transcripts are detected before those of *MyoD* during chick limb development. The *Myf5* expression domain resembles that of *Pax3* and is larger than that of *MyoD*. Moreover, *Myf5* and *Pax3* expression is correlated with myoblast proliferation, while *MyoD* is detected in post-mitotic myoblasts. These data indicate that *Myf5* and *MyoD* are involved in different steps during chick limb bud myogenesis, *Myf5* acting upstream of *MyoD*. The progression of myoblasts through the differentiation steps must be carefully controlled to ensure myogenesis at the right place and time during wing development. Because

Notch signalling is known to prevent differentiation in different systems and species, we sought to determine whether these molecules regulate the steps occurring during chick limb myogenesis. *Notch1* transcripts are associated with immature myoblasts, while cells expressing the ligands *Delta1* and *Serrate2* are more advanced in myogenesis. Misexpression of *Delta1* using a replication-competent retrovirus activates the Notch pathway. After activation of this pathway, myoblasts still express *Myf5* and *Pax3* but have downregulated *MyoD*, resulting in inhibition of terminal muscle differentiation. We conclude that activation of Notch signalling during chick limb myogenesis prevents *Myf5*-expressing myoblasts from progressing to the *MyoD*-expressing stage.

Key words: *Myf5*, *MyoD*, Notch, Delta, Chick, Limb bud, Myogenesis

INTRODUCTION

In vertebrates, all the myogenic cells that form the striated skeletal muscles of the limb and trunk originate from the somites. The medial halves of the somites give rise to back and intercostal muscles or the epaxial musculature (Ordahl and Le Douarin, 1992). Cells derived from lateral dermomyotomes migrate lateroventrally to produce the muscles of the body wall and the limbs, forming the hypaxial musculature (Ordahl and Le Douarin, 1992; Christ and Ordahl, 1995). Formation of limb skeletal muscle during vertebrate embryogenesis involves cellular commitment, migration, proliferation, growth arrest and differentiation. Each step involves the expression and activity of a specific panel of factors (Olson, 1992). The myogenic bHLH transcription factors, *Myf5*, *MyoD* (also known as *MyoD1*), *Mrf4* (also known as *Myf6*) and myogenin, which are also called myogenic regulatory factors (MRFs), have been shown to initiate the skeletal muscle differentiation program.

Ectopic expression of these MRFs is able to convert several non-muscle cell types into skeletal muscle in tissue culture (Weintraub et al., 1991), in transgenic mice (Miner et al., 1992; Santerre et al., 1993) and in *Xenopus* (Ludolph et al., 1994). Conversely, knockout of these genes leads to various muscle defects (Rudnicki et al., 1992, 1993; Braun et al., 1992a,b; Hasty et al., 1993; Nabeshima et al., 1993). Moreover, cells deprived of *Myf5* or *MyoD* assume a non-muscle fate (Tajbakhsh et al., 1996; Kablar et al., 1999). These properties have led to the notion that MRFs trigger the successive events leading to skeletal muscle formation. Gene targeting has clearly defined a hierarchy among the MRF family members. Mice lacking *Myf5* and *MyoD* (Rudnicki et al., 1993) do not form myoblasts or skeletal muscle. In contrast, in myogenin-null mice, myoblasts do form, as assayed by *Myf5* and *MyoD* expression, but do not differentiate into muscle fibres (Hasty et al., 1993; Nabeshima et al., 1993). *Myf5* and *MyoD* therefore appear to lie in a genetic pathway upstream of myogenin, the latter

having a role in activating muscle cell terminal differentiation (Hasty et al., 1993; Nabeshima et al., 1993). In mice, there are numerous and consistent studies concerning the sequential expression of myogenic factors during somite and limb bud development, *Myf5* being detected before *MyoD* in somites and limbs (Ontell et al., 1995; Tajbakhsh and Buckingham, 1999). In birds, studies of the timing of the expression of these factors in axial regions have led to the conclusion that, in contrast to the situation in mice, *MyoD* expression occurs before that of *Myf5* by a few hours (Pownall and Emerson, 1992; Boricky et al., 1997; Denetclaw and Ordahl, 2000). However, Hacker and Guthrie (1998) found that *Myf5* transcripts were expressed first, followed by those of *MyoD*. In the limb, studies using in situ hybridisation of tissue sections (Williams and Ordahl, 1994), RT-PCR (Lin-Jones and Hauschka, 1996) and whole-mount in situ hybridisation (Hacker and Guthrie, 1998) have led to conflicting results concerning the timing of appearance of *MyoD* and *Myf5* transcripts.

The progression through discrete developmental steps has been studied in muscle cell lines. The presence of *Myf5* and *MyoD* is not itself sufficient to trigger differentiation in cell culture, since myoblasts exposed to growth factors continue to proliferate and to express *Myf5* and/or *MyoD* (Yutzey et al., 1990). One well-described general mechanism influencing differentiation events during development is the Notch signalling pathway (reviewed in Artavanis-Tsakonas et al., 1999). The Notch pathway has been shown to operate at different steps during *Drosophila* myogenesis (Baylies et al., 1998). However, although mice bearing null mutations in the different Notch signalling components exhibit defects of somite formation, they do not display any muscle defects (Swiatek et al., 1994; Conlon et al., 1995; de Angelis et al., 1997). The lack of effect probably reflects the functional overlap among the Notch family members. In mammals, the only direct evidence of Notch involvement in myogenesis comes from in vitro studies where activated Notch or ligand-induced Notch signalling suppresses muscle differentiation in various mouse cell lines (Kopan et al., 1994; Nye et al., 1994; Lindsell et al., 1995; Shawber et al., 1996; Jarriault et al., 1998; Nofziger et al., 1999; Kuroda et al., 1999). No such evidence has been obtained in vivo.

Knockouts and studies on cell lines have proved very powerful in determining the genetic hierarchy of MRFs and giving clues about their functions. However, the exact functions of the proteins coded by *Myf5* and *MyoD* during development are still not fully understood. Based on in vitro studies, *Myf5* and *MyoD* are widely considered to have overlapping function associated with myoblast proliferation (Lassar et al., 1994; Molkentin and Olson, 1996). In order to gain insight into the respective roles of *Myf5* and *MyoD* in avian limb myogenesis, we have characterised the cellular expression patterns of these genes. We found that in the chick limb, *Myf5* can be detected at stage 20 with an expression domain similar to that of *Pax3*, *MyoD* being detected a few hours later (stage 22) in a more restricted domain. Activation of the Notch pathway in vivo led to a downregulation of *MyoD* expression, without affecting *Pax3* and *Myf5* expression, followed by an inhibition of terminal differentiation. Together, these results suggest that *Myf5* acts upstream of *MyoD*, and that the Notch pathway is involved in the progression from the

Myf5-expressing stage to the *MyoD*-expressing stage, during chick limb bud myogenesis.

MATERIALS AND METHODS

Chick embryos

Fertilised White Leghorn eggs (HAAS, Strasbourg, France) were incubated at 37°C. All grafting experiments were performed in ovo. Young embryos were staged according to Hamburger and Hamilton (HH) (1951), while old embryos were staged according to embryonic days in ovo. To facilitate comparisons, we report both staging for young embryos.

Production of control/RCAS- or Delta-expressing cells

Infectious Delta/RCAS (Henrique et al., 1997) and control/RCAS viruses were produced in Chick Embryo Fibroblasts (CEF) as described by Duprez et al. (1998). Briefly, CEF were isolated from E10 O-line embryos (BBSRC, Institute for Animal Health, Compton, Berkshire, UK) and grown in DMEM (Gibco, BRL) containing 8% (v/v) fetal calf serum and 2% (v/v) chick serum supplemented with antibiotics. CEF were transfected transiently with retroviral recombinant DNA using Transfectam (Gibco, BRL) according to the manufacturer's instructions.

Grafting of retrovirus-infected cells

Retrovirus-expressing cells were prepared for grafting as described by Duprez et al. (1998). Pellets of approximately 50 to 100 µm in diameter were grafted into the limb field of White Leghorn embryos around stage HH 16 of development (E2.5). Embryos were harvested at different times after grafting and processed for in situ hybridisation of whole mounts or tissue sections. Embryos grafted with control/RCAS-expressing cells did not exhibit any change in morphology (see also Duprez et al., 1996) or gene expression (data not shown). The numbers of embryos processed for in situ hybridisation of whole mounts are given in the text or in Table 1. In each experiment, two to six specimens were used for in situ hybridisation of tissue sections.

Bromodeoxyuridine (BrdU) labelling in ovo

E3 embryos were injected in the amnios (near the heart and wing) with 200 µl of 10 mM BrdU (Amersham, Life Science), and were re-incubated for another 15 minutes. 1 µl of 10 mM BrdU was directly injected in the circulation of E7 embryos and fixed 1 hour after. The embryos were then fixed and processed for in situ hybridisation of sections.

In situ hybridisation of whole mounts and tissue sections

Embryos were fixed in 4% (v/v) formaldehyde and processed as previously described for in situ hybridisation of whole mounts and paraffin sections (Duprez et al., 1998, 1999). Antisense digoxigenin- and fluorescein-labelled RNA probes were prepared as follows: *Myf5* (Saitoh et al., 1993); *Pax3* and *MyoD* (Duprez et al., 1998), *Delta1* (Henrique et al., 1997), *Serrate2* and *Notch1* (kind gift from Domingos Henrique). For double in situ hybridisation, the fluorescein probe was revealed with NBT/BCIP reagents (Roche) first, then the digoxigenin probe with INT/BCIP (Roche).

Immunohistochemistry

Differentiated muscle cells were detected on sections and in cultures using a monoclonal antibody against sarcomeric myosin heavy chain, MF20 (Developmental Hybridoma Bank, University of Iowa, Iowa City). Proliferating cells were detected using a monoclonal antibody against BrdU (Amersham). In situ followed by immunohistochemistry was performed using successively the probes (*Notch1*, *Serrate2*, *Delta1*, *Pax3*, *MyoD* and *Myf5*) and the monoclonal antibody against BrdU or MF20 antibody.

RESULTS

Myf5 transcripts can be detected before those of *MyoD* during limb development

In order to clear up the controversy concerning the timing of appearance of *Myf5* and *MyoD*, we performed in situ hybridisation on serial transverse limb sections (Fig. 1). *Myf5* and *MyoD* transcripts were not detected during the migration of the muscle progenitors from the somites to the limb bud (data not shown). This is in agreement with chick (Williams and Ordahl, 1994; Lin-Jones and Hauschka, 1996) and mouse data (Tajbakhsh and Buckingham, 1994; Ontell et al., 1995). In contrast, as soon as the *Pax3*-expressing myoblasts had reached their destination (Fig. 1A), defined as stage 20 (Chevalier et al., 1977; Christ et al., 1977), we were able to detect *Myf5* transcripts (Fig. 1B), but no *MyoD* mRNA was observed (Fig. 1C). At that stage the ventral and dorsal muscle masses had not yet separated, as visualised by *Pax3* expression (Fig. 1A). Using this in situ hybridisation technique, *MyoD* expression was first detected unambiguously at stage 22/23 (see Fig. 2F and Duprez et al., 1998).

Myf5 and *MyoD* show different expression domains during limb development

Whole-mount in situ hybridisation at stage 22/23 (E4) showed the *Myf5* expression domain was larger than that of *MyoD* and closely resembled that of *Pax3* (Fig. 2A-C). The *Myf5* expression domain matched that of *Pax3* throughout development (Fig. 1, Fig. 2 and data not shown). *MyoD* transcripts (Fig. 2F,G) were located in a subregion of the ventral and dorsal muscle masses expressing *Myf5* mRNA (Fig. 2D,E). The *Myf5* expression domain extended to near the ectoderm, while *MyoD* mRNA was located more centrally within the limb. In order to understand whether *Myf5* and *MyoD* transcripts were located in the same cells where their expression domains overlapped, we performed double in situ hybridisation at stage 23. These experiments on transverse wing sections confirmed that the expression domain of *MyoD* mRNA (orange) was more restricted than and contained within that of *Myf5* (purple) (Fig. 2H,I). All three possible expression combinations were observed (Fig. 2I): (1) *Myf5*⁺/*MyoD*⁻ cells (black arrows) were preferentially located near the ectoderm, (2) *Myf5*⁻/*MyoD*⁺ cells were preferentially found near the centre of the limb (black arrowheads) and (3) cells expressing both genes were found at the interface between these regions (white arrowhead). After stage 23, the *MyoD* expression domain spread to include all muscle masses, while *Myf5* and *Pax3* transcripts were progressively downregulated (see Figs 6, 8 and Duprez et al., 1998).

Myf5 expression is associated with myoblast proliferation whereas *MyoD* transcripts are detected in postmitotic myoblasts

The *Myf5* mRNA expression domain appeared identical to that of *Pax3* (Fig. 1A,B; Fig. 2A,B). The similarity in the expression patterns of *Pax3* and *Myf5* transcripts suggested that their expression might be linked to the proliferative state of the cell. Since *Pax3* expression has already been linked with proliferation (Epstein et al., 1995; Amthor et al., 1998), we studied the proliferative/differentiation state of *Myf5*-positive cells by performing BrdU incorporation experiments at stage





Fig. 1. *Myf5* transcripts are detected before those of *MyoD* in chick limb bud. Adjacent transverse wing sections from stage 20/21 (E3) embryos were hybridised with digoxigenin-labelled antisense probes for *Pax3* (A), *Myf5* (B) and *MyoD* (C). Scale bar, 160 μ m. D, dorsal; V, ventral.

23 (see Materials and Methods). These experiments showed that some *Myf5*-expressing cells had indeed incorporated BrdU, similar to *Pax3*-expressing cells (Fig. 3A-D, arrows). In contrast, most of the cells expressing *MyoD* transcripts did not incorporate BrdU (Fig. 3E,F, arrowheads).

Location of Delta/Notch pathway components during chick limb myogenesis

One known mechanism involved in differentiation processes in many systems and species is the Notch pathway (Artavanis-Tsakonas et al., 1999). We set out to investigate whether this signalling pathway is involved in myogenesis. We first analysed the endogenous cellular expression pattern of the Notch pathway components. From the literature it appeared that the Notch receptor might be ubiquitously expressed, the specificity of its action being determined by its ligands (Artavanis-Tsakonas et al., 1999). Indeed, whole-mount in situ hybridisation showed that the ligands *Delta1* (Fig. 4A) and *Serrate2* (Fig. 4B) were located in the muscle areas of the chick limbs at E5, while *Notch1* transcripts were more uniformly distributed (Fig. 4C; Vargesson et al., 1998; Beckers et al., 1999). BrdU incorporation experiments indicated that high levels of the ligands *Delta1* (Fig. 4G, arrows) and *Serrate2* (Fig. 4D,E, arrows) were expressed in scattered cells that did not incorporate BrdU within the muscle masses. At E7, the ligand *Serrate2* was only detected in MF20-positive cells (Fig.

Fig. 2. *Myf5* expression domain differs from that of *MyoD*. Distribution of *Pax3* (A), *Myf5* (B) and *MyoD* (C) transcripts in whole-mount preparations from stage 22/23 embryos (E4). A-C are dorsal views focused on the wings. Consecutive transverse sections through the forelimb from a stage 23 embryo were hybridised with digoxigenin-labelled antisense probes for *Myf5* (D,E) and *MyoD* (F,G). On transverse sections through the forelimb from a stage 23 embryo, double in situ hybridisation was performed using the *Myf5* digoxigenin-labelled-probe (purple) and the *MyoD* fluo-labelled probe (orange) (H,I). Black arrows indicate the *Myf5*-positive cells. Arrowheads point to *MyoD*-positive cells. White arrowheads mark the double *Myf5*- and *MyoD*-positive cells. E,G,I show higher magnification of D,F,H, respectively. Scale bars: 1.2mm in A-C; 180 μ m in D,F,H; 45 μ m in E,G,I. D, dorsal; V, ventral.



4H). *Delta1* was weakly detected at that stage (data not shown). The expression domain of the receptor *Notch1* was larger than that of the ligand *Serrate2* (compare the adjacent sections of Fig. 4E and 4F), suggesting that *Notch1* was also expressed in non-myogenic cells. We could detect *Notch1* mRNA in BrdU-positive cells (Fig. 4F, arrowheads), although most of the *Notch1*-positive cells were BrdU negative (Fig. 4F). In addition, *Notch1* transcripts were clearly detected in mononucleated cells around the muscle fibres (Fig. 4I). These results indicate that the receptor *Notch1* is expressed in immature myoblasts, while the cells expressing the ligands *Delta1/Serrate2* are more advanced in myogenesis (postmitotic myoblasts and muscle fibres).

Overexpression of *Delta1* affects *MyoD* expression without affecting *Myf5* expression

In order to understand the role of Notch signalling during the different steps of myogenesis, we activated the Notch pathway by over-expressing *Delta1* using the RCAS retrovirus. The Delta/RCAS construct has been shown to be effective in retinal (Henrique et al., 1997), cartilage (Crowe et al., 1999), feather bud (Crowe et al., 1998; Viallet et al., 1998) and scale (Crowe and Niswander, 1998) formation. Based on described functions of Notch signalling in different systems and species, we hypothesised that constitutive activation of Notch signalling in muscle cells would lead to an inhibition of terminal muscle differentiation. Aggregates of Delta/RCAS-transfected cells (see methods) were grafted to stage 16 (E2.5) wing buds. In

situ hybridisation of *Delta1* transcripts in whole mounts showed the degree of virus spread 48 hours (Fig. 5A,B; $n=5$ out of 6) and 72 hours (Fig. 5C,D; $n=2$ out of 2) after grafting. In order to visualise the activation of Notch signalling, we looked for *Notch1* expression after grafting, since it has been shown that activation of Notch signalling enhances *Notch* expression (Lewis, 1996). Overexpression of *Delta1* in the limb bud led to an extension of the *Notch1* expression domain (Fig. 5I,J; $n=4$ out of 5), reflecting an activation of Notch signalling (Micchelli et al., 1997; Franklin et al., 1999). In such grafted embryos, the *MyoD* expression domain appeared reduced in the region where ectopic *Delta1* was detected (Fig.

Table 1. Muscle gene expression in whole-mount embryos following grafts of Delta1/RCAS transfected cells to the wing region at E2.5

Time after grafting	Gene transcripts		
	<i>Pax3</i>	<i>Myf5</i>	<i>MyoD</i>
24 hours	No change (2/2)	No change (5/6)*	No change (4/6)‡
48 hours	No change (7/8)§	No change (3/3)	Decrease (4/5)¶
72 hours	n.d.	No change (3/3)	Decrease (5/7)¶

Numbers in brackets indicate the number of cases giving the results out of the number of embryos examined.


*In one case *Myf5* expression appears downregulated.

‡*MyoD* expression was not detected in the control limb in four cases.

§In one case *Pax3* expression was upregulated.

¶The other cases were unchanged. n.d., not determined.

Fig. 3. *Myf5* and *Pax3* transcripts are associated with proliferative myoblasts while *MyoD* is detected in postmitotic cells. Transverse wing sections from stage 23 embryos incubated with BrdU 15 minutes before fixation were hybridised with *Pax3* (A,B), *Myf5* (C,D) and *MyoD* (E,F) probes and then incubated with the anti-BrdU antibody. B,D,F show higher magnification of the dorsal (B) and ventral (D,F) muscle masses of the sections shown in A,C,E, respectively. Arrows indicate the *Myf5*- (B) and the *Pax3*- (D) positive cells that are also BrdU positive. Arrowheads point to the *MyoD*-positive cells, which are BrdU negative. Scale bar: 160 μ m in A,C,E; 40 μ m in B,D,F. D, dorsal; V, ventral.



5B,D), compared with the control wing, after 48 hours (Fig. 5E,F; Table 1) and 72 hours (Fig. 5G,H; Table 1). In contrast, the *Myf5* (Fig. 5K,L; Table 1) and *Pax3* (Table 1) expression domains were unchanged compared to the control wing. This demonstrates that the Notch pathway operates between the *Myf5/Pax3*-expressing and the *MyoD*-expressing stages.

Delta-activated Notch inhibits terminal muscle differentiation, despite the presence of *Myf5* and *Pax3* transcripts

Seventy-two hours after the Delta/RCAS-expressing cells were grafted into the wing bud, the control and operated wings were cut transversely through the forelimb region and hybridised with the *Delta1* probe, revealing the extent of the spread of the virus (Fig. 6A,B,I,J). Adjacent sections hybridised with *MyoD* (Fig. 6C,D,K,L), *Myf5* (Fig. 6E,F,M,N) and *Pax3* (Fig. 6G,H,O,P) probes showed the normal and modified muscle pattern in the control and operated wings, respectively. The Delta/RCAS-infected right wing exhibited a downregulation of *MyoD* transcripts (compare Fig. 6C,D with 6K,L), while *Myf5* and *Pax3* expression appeared unaffected (compare Fig. 6E-H with 6M-P). Analysis of myosin expression using the MF20 antibody showed that there was a clear diminution in the number of differentiated muscle cells in the treated wing (Fig. 6O,P), compared with the control limb (Fig. 6G,H, in brown). This demonstrated that terminal differentiation is affected despite the presence of *Myf5* and *Pax3* transcripts.

Delta-activated Notch leads to disorganised muscles

We then examined the phenotype of the Delta/RCAS-infected limbs at E9.5 (7 days after grafting; $n=4$). Transverse sections were cut at the same level along the proximo-distal axis from the control (Fig. 7A,B) and manipulated (Fig. 7C,D) wings. In the limb shown in Fig. 7, only the posterior muscles were affected (Fig. 7D), although all muscles could be affected (data not shown). Examination of *MyoD* and myosin expression showed that the posterior muscles were reduced in size and disorganised (Fig. 7D), compared with the normal pattern (Fig. 7B). Moreover, the FDP (flexor digitorum profundus) muscle was absent in the *Delta1*-infected wing (Fig. 7D) but was clearly present in the control wing (Fig. 7B). Only certain

muscles were affected (Fig. 7D) in this way by Delta1/RCAS infection despite the broad distribution of ectopic *Delta1* transcripts (Fig. 7C). We interpreted this result as Delta-activated Notch acting only within a specific time window between the *Myf5* and *MyoD* expression steps (see above). This time window is probably brief, so ectopic *Delta1* would have to infect myoblasts at a very specific time in order to affect their further differentiation. High magnifications of the *Delta1*-infected muscle FCU (flexor carpi ulnaris) and control FCU (Fig. 8) confirmed the absence of myosin in the *MyoD*-negative region (Fig. 8C,D) of the infected FCU, while myosin expression could be detected in the *MyoD*-positive region (Fig. 8D; arrow). In the control muscles, we could detect the *MyoD* mRNAs (purple) and myosin (brown) in most of the cells (Fig. 8A,B).

DISCUSSION

Distinct expression patterns of *Myf5* and *MyoD* in the chick wing indicate different roles during myogenesis, *Myf5* acting upstream of *MyoD*

In situ studies of *Myf5* and *MyoD* expression during development in the chick have been apparently contradictory (see Introduction). We have shown that *Myf5* transcripts can be unambiguously detected before those of *MyoD* in the chick limb bud. A similar situation occurs in the chick somite (Hirsinger et al., 2001). These results are consistent with those obtained in mice. The expression domains of *Myf5* and *Pax3* essentially overlap during development. In contrast, *MyoD* mRNAs initially show




Fig. 4. Endogenous expression of Notch signalling components in the chick limb bud. Distribution of *Delta1* (A), *Serrate2* (B) and *Notch1* (C) transcripts in whole-mount preparations from stage 26/27 embryos (E5). (A,B,C) Dorsal views focused on the wings. Consecutive transverse wing sections from stage 29 embryos (E6.5) injected with BrdU 1 hour before fixation were hybridised with *Serrate2* (D,E) and *Notch1* (F) probes (blue), and then incubated with the anti-BrdU antibody (brown). (G) Posterior part of the ventral muscle mass indicates that the *Delta*-positive cells are BrdU-negative (arrows). Transverse wing sections from stage 31/32 (E7/E7.5) embryos were hybridised with *Serrate2* (H) and *Notch1* (I) probes (blue) and then incubated with the MF20 antibody (brown). Scale bars: 160 μ m in D; 40 μ m in E-I.

a more restricted pattern, in the centre of the limb. During subsequent limb development, the *MyoD* expression domain spreads to include all muscle masses, while *Pax3* and *Myf5* are downregulated, although a low level of *Pax3* transcripts can still be detected quite late (E10) (Duprez et al., 1998). Despite the absence of clear segregation between proliferative and postmitotic myoblasts in the limb, we observe a gradient of maturation from the ectoderm, where *Myf5*- and *Pax3*-expressing myoblasts proliferate (less differentiated), to the centre of the muscle masses, where myoblasts express *MyoD* and become postmitotic (more differentiated). There is probably an intermediate phase where the cells are *Myf5*⁺/*MyoD*⁺. This sequence of myogenic factor expression is shown in Fig. 9A.

In mice, *Myf5* and *MyoD* have been described as being activated in a mutually exclusive manner in the musculature, *Myf5* transcripts being first detected in the epaxial myotome and *MyoD* in the hypaxial myotome (Braun and Arnold, 1996; Cossu et al., 1996a, Tajbakhsh and Buckingham, 1999). Nevertheless, both genes are later co-expressed in the majority of cells with myogenic potential both in vivo and in vitro (Cossu et al., 1996b; Tajbakhsh and Buckingham, 1999). Moreover, *Myf5*-deficient embryos exhibit a 2-day delay in development of axial muscles, but normal formation of the limb musculature. Conversely, in *MyoD* mutant embryos, there is delayed limb muscle development and normal axial

musculature formation (Kablar et al., 1997). This has been interpreted, in mice, as showing that *Myf5* has a primary function in the regulation of axial muscles whereas *MyoD* is involved in limb muscle formation. Our chick expression data (see Results) provide no evidence for this dichotomy of function (*Myf5*/axial muscles versus *MyoD*/limb muscles). Instead, our results suggest involvement at different steps during myogenesis, *Myf5* acting before *MyoD*. This has been already suggested by gene targeting analysis: in the absence of *Myf5* and *Pax3*, mice do not express *MyoD* and fail to develop body skeletal muscles (Tajbakhsh et al., 1997). Moreover, it appears that *Myf5* is activated first in both epaxial and hypaxial domains of mouse somites (Tajbakhsh and Buckingham, 1999). In addition, mouse *Myf5* expression matches the main sources of myotomal precursors (Venters et al., 1999). These findings indicate that *Myf5* initiates the body skeletal muscle differentiation program in both chick and mouse. The absence of muscle phenotype in the *Myf5* knockout mice (Braun et al., 1992a,b) could be explained by *Pax3* replacing the absent *Myf5* and activating *MyoD*. Indeed, *Pax3* appears to be sufficient, in some cellular contexts, to activate *MyoD* expression and thus initiate the myogenic program in vitro (Maroto et al., 1997) and in vivo (Tajbakhsh et al., 1997; Bendall et al., 1999). Alternatively, *paraxis* could be another candidate to assume the role of *Myf5*, since the double mutation *paraxis*⁻/*Myf5*⁻ shows muscle losses not observed in the single mutations (Wilson-

Fig. 5. Ectopic *Delta1* expression downregulates *MyoD* without affecting *Myf5* expression. Viral transcripts were detected in whole-mount preparations by in situ hybridisation with a probe against *Delta1*, 48 hours (A,B) and 72 hours (C,D) after grafts to the wing at E2.5 of Delta/RCAS-expressing cells. *MyoD* transcripts are downregulated in the grafted wings 48 hours (F) and 72 hours (H) after similar grafts compared with the respective control limbs (E,G). *Notch1* transcripts are upregulated in the grafted wings (right) 24 hours (I) and 48 hours (J) after similar grafts compared with the control limbs (left). Distribution of *Myf5* transcripts in whole-mount preparations is unchanged 72 hours (K,L) after similar grafts. Arrows indicate the ectopic *Delta1* expression (B,D), the downregulation of *MyoD* (F,H), the upregulation of *Notch1* (I,J) and the unchanged *Myf5* (L) domain in the manipulated wings. Scale bars: 500 μ m in A-G,K,L; 350 μ m in I; 1mm in J.





Fig. 6. Overexpression of *Delta1* inhibits myogenesis despite the presence of *Myf5* and *Pax3* transcripts. Adjacent transverse sections of the control (A-H) and infected wings (I-P) from the same embryo 72 hours after grafting Delta/RCAS-expressing cells in E2.5 limbs were hybridised with *Delta1* (A,B,I,J), *MyoD* (C,D,K,L), *Myf5* (E,F,M,N) and *Pax3* (G,H,O,P) probes. The *Pax3* in situ hybridisation was followed by an incubation with the MF20 antibody (G,H,O,P). All the pictures are orientated similarly: dorsal towards the top, ventral towards the bottom, posterior towards the left and anterior towards the right. (B,D,F,H,J,L,N,P) High magnifications of the ventral muscle masses from the control limb (A,C,E,G) and infected limb (I,K,M,O). Scale bars 240 μ m in A,C,E,G,I,K,M,O; 60 μ m in B,D,F,H,J,L,N,P.




Fig. 7. Ectopic *Delta1* results in disorganised muscles. Adjacent transverse sections of the control (A,B) and infected (C,D) wings from the same embryo were hybridised with the RNA probes specific for *Delta1* (A,C) or *MyoD* (B,D) and then incubated with the MF20 antibody (B,D). The muscles in the posterior regions are disorganised or absent. a, anterior; D, dorsal; FCU, flexor carpi ulnaris; Δ FCU, the remains of the FCU; FDP, flexor digitorum profundus; p, posterior; r, radius; u, ulna; V, ventral;. Scale bar: 320 μ m.

Rawls et al., 1999a). The presence of muscle in the absence of *MyoD* (Rudnicki et al., 1992) has been interpreted as showing that it can be replaced by *Myf5* (Rudnicki et al., 1993; Tajbakhsh and Cossu, 1997). However, *Myf5* alone is insufficient to activate the myogenic program in the absence of the other three myogenic factors (Valdez et al., 2000), suggesting rather an overlap in the functions of myogenin, *MyoD* and *Mrf4*. It has already been shown that *Mrf4* and *MyoD* can compensate for each other's absence in muscle differentiation in mice, since the *Mrf4*⁻/*MyoD*⁻ double mutant displays a severe muscle deficiency, whereas mice lacking either *Mrf4* or *MyoD* do not show defects in muscle development (Rawls et al., 1998). The absence of axial muscle defects in *MyoD*⁻ mice could be explained if *Mrf4* were able to support muscle development. The transient expression of *Mrf4* in myotome before its expression in late embryogenesis

and postnatal muscles (Ontell et al., 1995) is consistent with this notion.

Delta-activated Notch signalling inhibits myogenesis in vivo

We have shown that Delta-activated Notch signalling in vivo downregulates *MyoD* expression and then inhibits terminal differentiation in the chick limb bud. This is the first demonstration in vivo of the involvement of Notch signalling in chick limb myogenesis. The cell-surface receptor Notch mediates communication between cells expressing Notch and cells expressing membrane-bound ligands such as *Delta1* and *Serrate2*. Our examination of *Notch1* and *Delta1/Serrate2* expression shows that high levels of the ligands are detected in postmitotic cells and muscle fibres but that *Notch1* is associated with mononucleated cells surrounding the fibres. These results




Fig. 8. Higher magnifications focused on FCU muscles from the control (A,B) and manipulated wing (C-F) from Fig. 7, hybridised with *MyoD* probe (purple) and then incubated with the MF20 antibody revealed in brown (A-D), or hybridised with the *Delta1* probe (E,F). The arrow in D indicates a myosin-positive cell in the *MyoD*-positive area. Scale bars: 80 μ m in A,C,E; 40 μ m in B,D,F.

are illustrated in Fig. 9B. Our overexpression data coupled with the in situ analysis can be interpreted as Notch signalling playing a role in maintaining the myoblasts in an undifferentiated state until myoblasts are correctly positioned to pursue their differentiation. This result is consistent with the known functions of Notch in other systems, e.g. retina, where progenitor retinal cells exposed to *Delta1* are prevented from undergoing neuronal differentiation (Henrique et al., 1997; Dorsky et al., 1995); cartilage, where misexpression of *Delta1* blocks chondrocyte maturation (Crowe et al., 1999); and feather, where overexpression of *Delta1* inhibits feather development (Crowe et al., 1998; Viallet et al., 1998). Moreover, during development of adult indirect flight muscles in *Drosophila*, Notch activation causes failure of differentiation (Anant et al., 1998).

The use of the dominant-negative form of *Delta1* that blocks Notch signalling (Henrique et al., 1997) failed to give any muscle phenotype when grafted into the limb (data not shown). This means that definitive proof of a physiological role for Notch in limb myogenesis is still lacking. However, successful block of Notch signalling using the dominant-negative form of *Delta1* has only been reported for one system, the retina (Henrique et al., 1997). An alternative explanation is that *Serrate2*, another Notch ligand that is expressed in differentiated myotubes (see Results), compensates for the lack of *Delta1* activity. *Serrate* has been indeed shown to compensate for the loss of function of *Delta* in *Drosophila* (Gu et al., 1995). Moreover, overexpression of *Serrate2* can inhibit the differentiation of C2C12 myoblast cells (Lindsell et al., 1995). *Serrate2* is expressed at the right time and place to play a role in limb myogenesis. However, it is not clear whether *Delta1* and *Serrate2* have identical functions or whether they use the same receptors during limb myogenesis.

Delta-activated Notch acts between the *Myf5* and *MyoD* steps

Our results show that neither *Myf5* nor *Pax3* is affected by ectopic activation of Notch signalling, indicating that Notch signalling acts after the *Pax3/Myf5* step (Fig. 9A). In addition, our results show that *Myf5* and *Pax3* are insufficient to allow further muscle differentiation in the absence of *MyoD*. This contrasts with the normal muscle phenotype in *MyoD* knockout mice (Rudnicki et al., 1992; Kablar et al., 1997). Our misexpression experiments do not allow us to exclude the hypothesis that *Delta1*-activated Notch acts on the *MyoD*-expressing lineage in the chick limb, leaving intact the *Myf5*-expressing pathway. But in that case we would have expected to observe normal terminal differentiation in our experimental limbs, as in the *MyoD* knockout mice (Rudnicki et al., 1992; Kablar et al., 1997). The *Myf5* (and *Pax3*) pathway is insufficient, in our experimental context, to rescue terminal muscle differentiation in the absence of *MyoD*, since we observe fewer myosin-positive cells in *Delta1*-infected limbs. This reinforces the idea that myogenic factor(s) other than *Myf5* compensate for the absence of *MyoD* in *MyoD*^{-/-} mice (Rawls et al., 1998; Wilson-Rawls et al., 1999b; Valdez et al., 2000). Alternatively, there might be a genuine difference between chick and mouse.

It is not clear whether activated-Notch acts on the transition (1) between *Myf5*-proliferative and *MyoD*-postmitotic cells or (2) between *Myf5*-postmitotic to *MyoD*-postmitotic cells. By




Fig. 9. Involvement of Notch signalling between the *Myf5*-step and the *MyoD*-step. (A) The sequence of expression of the myogenic factors during myogenesis is represented. *Pax3* and *Myf5* are associated with myoblast proliferation. *Myf5* remains expressed in post-mitotic myoblasts, which then express *MyoD* followed by the myosin proteins. Activated Notch inhibits the progression from the *Myf5* step to the *MyoD* step. (B) The myoblasts expressing the receptor *Notch1* remain in an undifferentiated state while the myoblasts expressing the ligands (*Delta1* or *Serrate2*) pursue their differentiation.

analogy with the situation in the retina, where forced expression of *Delta1* maintains proliferating neuroepithelial precursors (Henrique et al., 1997), we would have expected an extension of the *Pax3* and *Myf5* domains concomitant with an increase of the BrdU incorporation. However, the fact that we do not observe any change in the *Pax3* and *Myf5* domains (Figs 5, 6) or of BrdU labelling (data not shown) after *Delta* misexpression favours an action of Notch on *Myf5* postmitotic cells. Whatever the situation, it is clear that *Delta*-activated Notch blocks further differentiation of *Myf5*-expressing cells.

Relationship between *MyoD* and Notch signalling components

We found a decrease of *MyoD* transcripts after ectopic activation of Notch signalling. We cannot conclude from our experiments whether the downregulation of *MyoD* transcripts is the result of an inhibition of gene activation or a defect in the maintenance of *MyoD* expression. However, studies on transfected cell lines revealed that activated Notch is able to

inhibit *MyoD* transcription (Kuroda et al., 1999). Activated Notch also interferes with the muscle-inducing activity of the MyoD protein (Kopan et al., 1994). This interference has recently been shown to occur through a direct protein interaction between the ankyrin repeat region of Notch and MEF2C, an essential cofactor of MyoD, that blocks DNA binding (Wilson-Rawls et al., 1999b). Since MyoD and MEF2 participate in regulatory circuits involving positive transcriptional feedback loops (Thayer et al., 1989; Braun et al., 1989; Molkenin et al., 1995), the downregulation of *MyoD* expression we observed could also be the consequence of the inhibition of MEF2 activity.

It has been shown recently that MyoD is a direct, positive regulator of *Xenopus Delta1* (Wittenberger et al., 1999). The detection of *MyoD* transcripts before those of *Delta1* in the *Xenopus gastrula* indicates that MyoD triggers Notch signalling in this species (Wittenberger et al., 1999). From our results it is not clear whether MyoD induces *Delta1*, which would then trigger Notch signalling in adjacent myoblasts, or if Notch signalling is activated before the onset of *MyoD* expression. Chick *MyoD* and *Delta1* expression seem to occur together in the limb (data not shown). However, *Serrate2* transcripts are detected before those of *MyoD* in the limb (data not shown). Thus, we favour the hypothesis that high levels of ligand (*Delta1*, *Serrate2*) expression in a few cells would activate Notch signalling in adjacent cells. The ligand-positive cells would then differentiate by activating *MyoD*. The existence of a positive feedback loop (*MyoD* towards *Delta1*) would enhance and lock the differentiation process.

Many arguments converge to suggest that MyoD expression is linked to cell-growth arrest

In vitro studies have generally concluded that *MyoD* is expressed in proliferative myoblasts (Lassar et al., 1994; Molkenin and Olson 1996), although this does not fit with the endogenous expression of *MyoD* in postmitotic cells of mouse and chick myotomes (Ontell et al., 1995; Brand-Saberi et al., 1996; Amthor et al., 1998; Hirsinger et al., 2001). In the chick limb, where the distinction between proliferative and postmitotic myoblasts is not obvious (in contrast to the situation in somites), we have shown that the majority of *MyoD*-expressing cells do not incorporate BrdU, indicating that they have withdrawn from the cell cycle and are postmitotic myoblasts. Moreover, experimental evidence from in vitro studies shows that MyoD induces withdrawal from the cell cycle independently of muscle differentiation (Davies et al., 1987; Crescenzi et al., 1990; Sorrentino et al., 1990; Trough et al., 1993). No such effects have been reported for Myf5. In addition, Myf5 and MyoD show different expression profiles during the cell cycle in C2 cells. MyoD expression is maximal upon cell cycle exit (Kitzmann et al., 1998). A recent study of satellite cells reached the same conclusion that MyoD could be necessary for the transition from proliferation to differentiation (Yablonka-Reuveni et al., 1999). Interestingly, MyoD is able to activate myogenin (Hollenberg et al., 1993) and a cyclin inhibitor, p21, without any new protein synthesis (Otten et al., 1997; Cenciarelli et al., 1999). p21 activation is linked to cell-cycle exit (reviewed in Walsh and Perlman, 1997).

In conclusion, in vitro studies and genetic analysis, coupled with our in situ and in vivo experiments on chick limb, indicate

that *Myf5* initiates the myogenic program and that *MyoD* expression is the manifestation of subsequent differentiation. The in vivo signal regulating the transition from the *Myf5* step to the *MyoD* step in the chick limb may involve Notch signalling.

We thank Marie-Aimée Teillet and Heather Etchevers for critical reading of the manuscript. We are grateful to Francis Beaujean, Michel Fromaget and Sophie Gournet for the illustrations. This work was supported by the Association Française contre les Myopathies (AFM) and the Centre National de la Recherche Scientifique (CNRS).

REFERENCES

- Amthor, H., Christ, B., Weil, M. and Patel, K. (1998). The importance of timing during limb muscle development. *Curr. Biol.* **8**, 642-652.
- Anant, S., Roy S. and Vijayraghavan, K. (1998). Twist and Notch negatively regulate adult muscle differentiation in *Drosophila*. *Development* **125**, 1361-1369.
- Artavanis-Tsakonas, S., Rand, M. D. and Lake, R. J. (1999). Notch signaling: cell fate control and signal integration in development. *Science* **284**, 770-776.
- Baylies, M. K., Bate, M. and Ruiz Gomez, M. (1998). Myogenesis: a view from *Drosophila*. *Cell* **93**, 921-927.
- Beckers, J., Clark, A., Wünsch, K., De Angelis M.H. and Gossler, A. (1999). Expression of the mouse *Delta1* gene during organogenesis and foetal development. *Mech. Dev.* **84**, 165-168.
- Bendall, A. J., Ding, J., Hu G., Shern, M. M. and Abate-Shen, D. (1999). Msx1 antagonizes the myogenic activity of Pax3 in migrating limb muscle precursors. *Development* **126**, 4965-4976.
- Borycki, A. G., Strunk, K.E, Savary, R. and Emerson, C. (1997). Distinct signal/response mechanisms regulate Pax-1 and MyoD activation in sclerotomal and myotomal lineages of quail somites. *Dev. Biol.* **185**, 185-200.
- Brand-Saberi, B., Wilting, J., Ebensperger, C. and Christ, B. (1996). The formation of somite compartment in the avian embryo. *Int. J. Dev. Biol.* **40**, 411-420.
- Braun, T., Bober E., Buschhausen, D. G., Kohtz, S., Grzeschik, K. H. and Arnold, H. H. (1989). Differential expression of myogenic expression determination cells: possible autoactivation by the Myf5 gene product. *EMBO J.* **8**, 3617-3625.
- Braun, T., Rudnicki, M. A., Arnold, H. H. and Jaenish, R. (1992a). Targeted inactivation of the muscle regulatory gene Myf5 results in abnormal rib development and perinatal death. *Cell* **71**, 369-382.
- Braun, T., Rudnicki, M. A., Arnold, H. H. and Jaenish, R. (1992b). Targeted inactivation of the muscle regulatory gene Myf5 results in abnormal rib development and perinatal death. *Cell* **71**, 369-382.
- Braun, T. and Arnold, H. H. (1996). Myf5 and MyoD genes are activated in distinct mesenchymal stem cells and determine different skeletal muscle cell lineages. *Cell* **15**, 310-318.
- Cenciarelli, C., De Santa, F., Mattei, E., Ricci, L., Bucci, F., Felsani, A., Caruso, M. (1999). Critical role played by cyclin D3 in the MyoD-mediated arrest of cell cycle during myoblast differentiation. *Mol. Cell Biol.* **19**, 5203-5217.
- Chevalier, A., Kieny, M. and Mauger, A. (1977). Limb-somite relationship: origin of the limb musculature. *J. Embryol. Exp. Morphol.* **41**, 245-258.
- Christ, B., Jacob, M. and Jacob, H. (1977). Experimental analysis of the origin of the wing musculature in avian embryos. *Anat. Embryol.* **150**, 171-186.
- Christ, B. and Ordahl, C.P. (1995). Early stages of somite development. *Anat. Embryol.* **191**, 381-396.
- Conlon, R. A., Reaume, A. G. and Rossant, J. (1995). Notch1 is required for the coordinate segmentation of somites. *Development* **121**, 1533-1545.
- Cossu, G., Kelly R., Tajbakshs S., Di Donna S., Vivarelli E. and Buckingham M. (1996a). Activation of different myogenic pathways: Myf5 is induced by the neural tube and myoD by the dorsal ectoderm in mouse paraxial mesoderm. *Development* **122**, 429-437.
- Cossu, G., Tajbakshs, S. and Buckingham, M., (1996b). How is myogenesis initiated in the embryo? *Trends Genet.* **12**, 218-222.
- Crescenzi, M, Fleming, T. P., Lassar, A. B., Weintraub, H. and Aaronson, S. A. (1990). MyoD induces growth arrest-independent of differentiation in normal and transformed cells. *Proc. Natl. Acad. Sci. USA* **87**, 8442-8446.

- Crowe, R. and Niswander, L. (1998). Disruption of scale development by Delta1 misexpression. *Dev. Biol.* **195**, 70-74.
- Crowe, R., Henrique, D., Ish-Horowitz, D. and Niswander, L. (1998). A new role for Notch and Delta in cell fate decisions: patterning the feather array. *Development* **125**, 767-775.
- Crowe, R., Zikherman, J. and Niswander, L. (1999). Delta1 negatively regulates the transition from prehypertrophic to hypertrophic chondrocytes during cartilage formation. *Development* **126**, 987-998.
- Davies, R.L., Weintraub, H. and Lassar, A. B. (1987). Expression of a single transfected cDNA converts fibroblasts to myoblasts. *Cell* **51**, 987-1000.
- de Angelis, M., McIntyre I. I. and Gossler, A. (1997). Maintenance of somite borders in mice requires the Delta homologue Dll1. *Nature* **386**, 717-721.
- Denetclaw, W. F. and Ordahl, C. P. (2000). The growth of the dermomyotome and formation of early myotome lineages in thoracolumbar somites of chicken embryos. *Development* **127**, 893-905.
- Dorsky, R. I., Rapaport, D. H. and Harris, W.H. (1995). Xotch inhibits cell differentiation in the *Xenopus* retina. *Neuron* **14**, 487-496.
- Duprez, D., Bell, E., Richardson, M. K., Archer, C. W., Wolpert, L., Brickell, P. M. and Francis-West, P. (1996). Overexpression of BMP-2 and BMP-4 alters the size and shape of developing skeletal elements in the chick limb. *Mech. Dev.* **57**, 145-157.
- Duprez, D., Fournier-Thibault, C. and Le Douarin, N. (1998). SHH induces proliferation of committed skeletal muscle cells in the chick limb. *Development* **125**, 495-505.
- Duprez, D., Lapointe, F., Edom-Vovard, F., Kostakopoulou, K. and Robson, L. (1999). Sonic Hedgehog (SHH) specifies muscle pattern at tissue and cellular level, in the chick limb bud. *Mech. Dev.* **82**, 151-163.
- Epstein, J. A., Lam, L., Japeal, L., Maas, R. L. and Shapiro, D. (1995). Pax3 inhibits myogenic differentiation of cultured myoblast cells. *J. Biol. Biochem.* **20**, 11719-11722.
- Fizman, M. Y. and Fush, P. (1975). Temperature-sensitive expression of differentiation in transformed myoblasts. *Nature* **254**, 429-431.
- Franklin, J. L., Berechid, B. E., Cutting, F. B., Presente, A., Chambers, C. B., Folts, D. R., Ferreira, A., Nye, J. S. (1999). Autonomous and non-autonomous regulation of mammalian neurite development by Notch1 and Delta1. *Curr. Biol.* **9**, 1448-1457.
- Gu, Y., Hukriede, N. A. and Fleming R. J. (1995). Serrate expression can functionally replace Delta-activity during neuroblast segregation in the *Drosophila* embryo. *Development* **121**, 855-865.
- Hacker, A. and Guthrie, S. (1998). A distinct developmental programme for the cranial paraxial mesoderm in the chick embryo. *Development* **125**, 3461-3472.
- Hamburger, V. and Hamilton, H. (1951). A series of normal stages in the development of the chick embryo. *J. Morphol.* **88**, 49-92.
- Hasty, P., Bradley, A., Morris, J. H., Edmondson, D. G., Venuti, J. M., Olson, E. and Klein, W. H. (1993). Muscle deficiency and neonatal death in mice with a targeted mutation in the myogenin gene. *Nature* **364**, 501-506.
- Henrique, D., Hirsinger, E., Adam, J., Le Roux, I., Pourquie, O., Ish-Horowitz, D. and Lewis, J. (1997). Maintenance of neuroepithelial progenitor cells by Delta-Notch signalling in the embryonic chick retina. *Curr. Biol.* **7**, 661-670.
- Hirsinger, E., Malapert, P., Dubrulle, J., Delfini, M.-C., Duprez, D., Henrique, D., Ish-Horowitz, D. and Pourquie, O. (2001). Notch signalling acts in post-mitotic avian myogenic cells to control MyoD activation. *Development* (in press).
- Hollenberg, S. M., Cheng, P. F. and Weintraub, H. (1993). Use of a conditional MyoD transcription factor in studies of MyoD trans-activation and muscle differentiation. *Proc. Natl. Acad. Sci. USA* **90**, 8028-8032.
- Jarriault, S., Hirsinger, E., Pourquie, O., Logeat, F., Strong, C. F., Brou, C., Seidah, N. G. and Israel, A. (1998). Delta-1 activation of Notch1 signalling results in HES-1 trans-activation. *Mol. Cell Biol.* **18**, 7423-7431.
- Kablar, B., Krastel, K., Ying, C., Asakura, A., Tapscott, S. J. and Rudnicki, M. A. (1997). MyoD and Myf5 differentially regulate the development of limb versus trunk skeletal muscle. *Development* **124**, 4729-4738.
- Kablar, B., Krastel, K., Ying, C., Tapscott, S. J., Goldhamer, D. J. and Rudnicki, A. (1999). Myogenic determination occurs independently in somites and limb buds. *Dev. Biol.* **206**, 219-231.
- Kitzmann, M., Carnac, G., Vandromme, M., Primig, M. and Lamb, N. J. C. and Fernandez, A. (1998). The muscle regulatory factors MyoD and Myf5 undergo distinct cell cycle-specific expression in muscle cells. *J. Cell Biol.* **142**, 1447-1459.
- Kopan, R., Nye, J. S. and Weintraub, H. (1994). The intracellular domain of mouse Notch: a constitutively activated repressor of myogenesis directed at the basic helix-loop-helix region of MyoD. *Development* **120**, 2385-2396.
- Kuroda, K., Tani, S., Tamura, K., Minogoshi, S., Kurooka, H. and Honjon, T. (1999). Delta-induced Notch signaling mediated by RBP-J inhibits MyoD expression and myogenesis. *J. Biol. Chem.* **11**, 7238-7244.
- Lassar, A., Skapek, S. X. and Novitsch, B. (1994). Regulatory mechanisms that co-ordinate skeletal muscle differentiation and cell cycle withdrawal. *Curr. Opin. Cell Biol.* **6**, 788-794.
- Lewis, J. (1996). Neurogenic genes and vertebrate neurogenesis. *Curr. Opin. Neurobiol.* **6**, 3-10.
- Lindsell, C. E., Shawber, C. J., Boulter, J. and Weinmaster, G. (1995). Jagged, a mammalian ligand that activates Notch1. *Cell* **80**, 909-917.
- Lin-Jones, J. and Hauschka, S. D. (1996). Myogenic determination factor expression in the developing avian limb bud: an RT-PCR analysis. *Dev. Biol.* **174**, 407-422.
- Ludolph, D. C., Neff, A. W., Mescher, A. L., Malacinski, G. M., Parker, M. A. and Smith, R. C. (1994). Overexpression of XmyoD or XMyf5 in *Xenopus* embryos induces the formation of enlarged myotomes through recruitment of cells of non somitic lineage. *Dev. Biol.* **166**, 18-33.
- Maroto, M., Reshef, R., Münsterberg, A. E., Koester, S., Goulding, M. and Lassar, A. B. (1997). Ectopic Pax3 activates MyoD and Myf5 expression in embryonic mesoderm and neural tissue. *Cell* **89**, 139-148.
- Micelli, C. A., Rulifson, E. J. and Blair, S. S. (1997). The function and regulation of cut expression on the wing margin of *Drosophila*: Notch, Wingless and a dominant negative role for Delta and Serrate. *Development* **124**, 1485-1495.
- Miner, J. H., Miller, J. B. and Wold, B. (1992). Skeletal phenotypes initiated by ectopic myoD in transgenic mouse heart. *Development* **114**, 853-860.
- Molkentin, J. D., Black, B., Martin, J. F. and Olson, E. N. (1995). Cooperative activation of muscle gene expression by Mef2 and myogenic bHLH proteins. *Cell* **83**, 1125-1136.
- Molkentin, J. D. and Olson, E. (1996). Defining the regulatory networks for muscle development. *Curr. Biol.* **6**, 445-453.
- Nabeshima, Y., Hanaoka, K., Hayasaka, M., Esumi, E., Li S., Nonaka, I. and Nabeshima, Y. (1993). Myogenin gene disruption results in perinatal lethality because of severe muscle defect. *Nature* **364**, 532-534.
- Nofziger, D., Miyamoto, A., Lyons, K. M. and Weinmaster, G. (1999). Notch signaling imposes two distinct blocks in the differentiation of C2C12. *Development* **126**, 1689-1702.
- Nye, J. S., Kopan, R. and Axel, R. (1994). An activated Notch suppresses neurogenesis and myogenesis but not gangliogenesis in mammalian cells. *Development* **120**, 2421-2430.
- Olson, E. N. (1992). Interplay between proliferation and differentiation within the myogenic lineage. *Dev. Biol.* **154**, 261-272.
- Ordahl, C. P. and Le Douarin, N. (1992). Two myogenic lineages within the developing somite. *Development* **114**, 339-353.
- Ontell, M., Ontell, M. P. and Buckingham, M. (1995). Muscle-specific gene expression during myogenesis in the mouse. *Microsc. Res. Tech.* **30**, 354-354.
- Otten, A. D., Firpo, E. J., Gerber, A. N., Brody, L. L., Roberts, J. M., Tapscott, S. J. (1997). Inactivation of MyoD-mediated expression of p21 in tumor cell line. *Cell Growth Diff.* **8**, 1151-1160.
- Pownall, M. E. and Emerson, C. (1992). Sequential activation of three myogenic regulatory genes during somite morphogenesis in quail embryos. *Dev. Biol.* **151**, 67-79.
- Rawls, A., Valdez, M. R., Zhang, W., Richardson, J., Klein, W. H. and Olson, E. (1998). Overlapping functions of the myogenic bHLH genes MRF4 and MyoD revealed in double mutant mice. *Development* **125**, 2349-2358.
- Rudnicki, M. A., Braun, T., Hinuma, S. and Jaenish, R. (1992). Inactivation of MyoD in mice leads to up-regulation of the myogenic HLH gene Myf5 and results in apparently normal muscle development. *Cell* **71**, 383-390.
- Rudnicki, M. A., Schnegelsberg, P. N. J., Stead, R. H., Braun, T., Arnold, H.H. and Jaenish, R. (1993). MyoD or Myf5 is required for the formation of skeletal muscle. *Cell* **75**, 151-159.
- Saitoh, O., Fujisawa-Sehera, A., Nabeshima, Y. and Periasamy, M. (1993). Expression of myogenic factors in denervated chicken breast muscle: isolation of the chicken Myf5 gene. *Nucleic Acid Res.* **21**, 2503-2509.
- Santerre, R. F., Bales, K. R., Janney, M. J., Hannon, K., Fisher, F., Bailey, C. S., Morris, J., Ivarie, R. and Smith, C. K. (1993). Expression of bovine Myf5 induces ectopic skeletal muscle formation in transgenic mice. *Mol. Cell Biol.* **13**, 6044-6050.
- Shawber, C., Nofziger, D., Hsieh, J. D., Lindsell, C., Bögl, O., Hayward,

- D. and Weinmaster, G.** (1996). Notch signaling inhibits muscle cell differentiation through a CBF1-independent pathway. *Development* **122**, 3765-3773.
- Sorrentino, V., Pepperkok, R., Davis, R. L., Ansoorge, W. and Philipson, L.** (1990). Cell proliferation inhibited by MyoD1 independently of myogenic differentiation. *Nature* **345**, 813-815.
- Swiatek, P. J., Lindsell, C. E., Del Amo, F. F., Weinmaster, G., Gridley, T.** (1994). Notch1 is essential for postimplantation development in mice. *Genes Dev.* **8** 707-719.
- Tajbakhsh, S. and Buckingham, M. E.** (1994). Mouse limb muscle is determined in the absence of the earliest myogenic factor Myf5. *Proc. Natl. Acad. Sci. USA* **91**, 747-751.
- Tajbakhsh, S., Rocancourt, D. and Buckingham, M.** (1996). Muscle progenitor cells failing to respond to positional cues adopt non-myogenic fates in Myf5 null mice. *Nature* **384**, 266-270.
- Tajbakhsh, S., Rocancourt, D., Cossu, G. and Buckingham, M.** (1997). Redefining the genetic hierarchies controlling skeletal myogenesis: Pax3 and Myf5 act upstream of MyoD. *Cell* **89**, 127-138.
- Tajbakhsh, S. and Cossu, G.** (1997). Establishing myogenic identity during somitogenesis. *Curr. Opin. Genet. Dev.* **7**, 634-641.
- Tajbakhsh, S. and Buckingham, M.** (1999). The birth of muscle progenitor cells in the mouse spatiotemporal considerations. *Curr. Top. Dev. Biol.* **48**, 225-268.
- Thayer, M. J., Tapscott, S. J., Davis, R. L., Wright, W. E., Lassar, A. and Weintraub, H.** (1989). Positive auto regulation of the myogenic determination gene MyoD1. *Cell* **58**, 241-248.
- Trouch, D., Grigoriev, M., Lenormand, J-L., Robin, P., Leibovitch, SA, Sassone-Corsi, P., Harel-Bellan, A.** (1993). Repression of the c-fos promoter by MyoD on muscle cell differentiation. *Nature* **363**, 79-82.
- Valdez, M. R., Richardson J. A., Klein, H. K. and Olson E.** (2000). Failure of Myf5 to support myogenic differentiation without myogenin, MyoD and MRF4. *Dev. Biol.* **219**, 287-298.
- Vargesson, N., Patel, K. Lewis, J. and Tickle, C.** (1998). Expression patterns of Notch1, Serrate1, Serrate2 and Delta1 in tissues of the developing chick limb. *Mech. Dev.* **77**, 197-200.
- Venters, S. J., Thorsteinsdottir, S. and Duxson, M.** (1999). Early development of the myotome in the mouse. *Dev. Dyn.* **216**, 219-232.
- Viallet, J. P., Prin, F., Olivera-Martinez, I., Hirsinger, E., Pourquie, O. and Dhouailly D.** (1998). Chick *Delta1* gene expression and the formation of the feather primordia. *Mech. Dev.* **72**, 159-168.
- Walsh, K. and Perlman, H.** (1997). Cell cycle exit upon myogenic differentiation. *Curr. Opin. Gen. Dev.* **7**, 597-602.
- Weintraub, H., Davis, R., Tapscott, S., Thayer, M., Krause, M., Benezra, R., Blackwell, T. K., Turner, D., Rupp, R., Hollenberg, S., Zhuang, Y. and Lassar, A.** (1991). The myoD gene family: nodal point during specification of the muscle cell lineage. *Science* **251**, 761-766.
- Williams, B. A. and Ordahl, P. O.** (1994). Pax-3 expression in segmental mesoderm marks early stages in myogenic cell specification. *Development* **120**, 785-796.
- Wilson-Rawls, J., Hurt, C. R., Parsons, S. M. and Rawls, A.** (1999a). Differential regulation of epaxial and hypaxial muscle development by Paraxis. *Development* **126**, 5217-5229.
- Wilson-Rawls, J., Molkentin, J., Black, B. L. and Olson, E.** (1999b). Activated Notch inhibits myogenic activity of the MADS-Box transcription factor myocyte enhancer factor 2C. *Mol. Cell. Biol.* **19**, 2853-2862.
- Wittenberg, T., Steinbach O. C., Authaler, A., Kopan, R. and Rupp R. A. W.** (1999). MyoD stimulates Delta-1 transcription and triggers Notch signaling in the *Xenopus* gastrula. *EMBO J.* **18**, 1915-1922.
- Yablonka-Reuvini, Z., Rudnicki, M. A., Rvera A. J., Primig M., Abderson J. E. and Natanson, P.** (1999). The transition from proliferation to differentiation is delayed in satellite cells from mice lacking MyoD. *Dev. Biol.* **210**, 440-455.
- Yutzey, K. E., Rhodes, S. J. and Konieczny, S. F.** (1990). Differential trans-activation associated with the muscle regulatory factors MyoD1, myogenin and MRF4. *Mol. Cell. Biol.* **10**, 3934-3944.