



HAL
open science

Modelling Esker Formation on Mars

Neil Arnold, Frances Butcher, Colman Gallagher, Matthew Balme, Susan J. Conway

► **To cite this version:**

Neil Arnold, Frances Butcher, Colman Gallagher, Matthew Balme, Susan J. Conway. Modelling Esker Formation on Mars. 22nd EGU General Assembly, May 2020, held online, Unknown Region. 10.5194/egusphere-egu2020-3463 . hal-03091553

HAL Id: hal-03091553

<https://hal.science/hal-03091553>

Submitted on 3 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Open Research Online

The Open University's repository of research publications and other research outputs

Modelling Esker Formation on Mars

Conference or Workshop Item

How to cite:

Arnold, Neil S.; Butcher, Frances E. G.; Gallagher, Colman; Balme, Matthew R. and Conway, Susan J. (2020). Modelling Esker Formation on Mars. In: EGU General Assembly 2020: Sharing Geoscience Online, 4-8 May 2020, Online.

For guidance on citations see [FAQs](#).

© 2020 The Authors



<https://creativecommons.org/licenses/by/4.0/>

Version: Version of Record

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.5194/egusphere-egu2020-3463>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's [data policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

EGU2020-3463

<https://doi.org/10.5194/egusphere-egu2020-3463>

EGU General Assembly 2020

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



Modelling Esker Formation on Mars

Neil Arnold¹, Frances Butcher², Colman Gallagher³, Matthew Balme⁴, and Susan Conway⁵

¹University of Cambridge, Scott Polar Research Institute, United Kingdom of Great Britain and Northern Ireland (nsa12@cam.ac.uk)

²Department of Geography, University of Sheffield, UK.

³UCD School of Geography and UCD Earth Institute, University College Dublin

⁴School of Physical Sciences, The Open University, Milton Keynes, UK.

⁵CNRS Laboratoire de Planétologie et Géodynamique de Nantes, France

Introduction: Eskers are sinuous sedimentary ridges that are widespread across formerly glaciated landscapes on Earth. They form when sediment in subglacial tunnels is deposited by meltwater. Some sinuous ridges on Mars have been identified as eskers; whilst some are thought to have formed early in Mars' history beneath extensive ice sheets, smaller, younger systems associated with extant glaciers in Mars' mid latitudes have also been identified. Elevated geothermal heating and formation during periods with more extensive glaciation have been suggested as possible prerequisites for recent Martian esker deposition.

Here, we adapt a model of esker formation with g and other constants altered to Martian values, using it initially to investigate the impact of Martian conditions on subglacial tunnel systems, before investigating the effect of varying water discharge on esker deposition.

Methods: To investigate the effect of these values on the operation of subglacial tunnel systems we first conduct a series of model experiments with steady water discharge, varying the assumed liquid density (ρ_w) from 1000 kgm^{-3} to 1980 kgm^{-3} (the density of saturated perchlorate brine) and ice hardness (A) from $2.4 \times 10^{-24} \text{ Pa}^{-3}\text{s}^{-1}$ to $5 \times 10^{-27} \text{ Pa}^{-3}\text{s}^{-1}$ (a temperature range of 0°C to -50°C). We then investigate the impact of variable water discharge on esker formation to simulate very simply a possible release of meltwater from an assumed geothermal event beneath a Martian glacier or ice cap.

Results and Discussion: A key aspect of model behaviour is the decrease in sediment carrying capacity towards the ice margin due to increased tunnel size as ice thins. Our results suggest that Martian parameters emphasise this effect, making deposition more likely over a greater length of the conduit. Lower gravity has the largest impact; it reduces the modeled closure rate of subglacial tunnels markedly as this varies with overburden stress (and hence g) cubed. Frictional heating from flowing water also drops, but much less sensitively. Thus, for a given discharge, the tunnels tend to be larger, leading to lower water pressure and a reduction in flow power. This effect is amplified for harder ice. Higher inferred fluid density raises the flow power, but by a smaller amount.

These effects are clearly seen in the variable discharge experiments. Sediment is deposited on the falling limb of the hydrograph, when the tunnels are larger than the equivalent steady-state water discharge would produce. Sediment deposition occurs much further upglacier from the glacier snout, and occurs earlier on the falling limb leading to longer periods in which deposition occurs.

Conclusions: Our results suggest that esker formation within a subglacial meltwater tunnel would be more likely on Mars than Earth, primarily because subglacial tunnels tend to be larger for equivalent water discharges, with consequent lower water flow velocities. This allows sediment deposition over longer lengths of tunnel, and to greater depths, than for terrestrial systems. Future work will use measured bed topography of a mid-latitude esker to assess the impact of topography on deposition patterns and esker morphology, and we will expand the range of discharges and sediment supply regimes investigated.