



HAL
open science

Solutions to the NLS equation : differential relations and their different representations

Pierre Gaillard

► **To cite this version:**

Pierre Gaillard. Solutions to the NLS equation : differential relations and their different representations. 2020. hal-03091896

HAL Id: hal-03091896

<https://hal.science/hal-03091896>

Preprint submitted on 31 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Solutions to the NLS equation : differential relations and their different representations

⁺Pierre Gaillard, ⁺ Université de Bourgogne, Dijon, France :
e-mail: Pierre.Gaillard@u-bourgogne.fr,

Abstract

Solutions to the focusing nonlinear Schrödinger equation (NLS) of order N depending on $2N - 2$ real parameters in terms of wronskians and Fredholm determinants are given. These solutions give families of quasi-rational solutions to the NLS equation denoted by v_N and have been explicitly constructed until order $N = 13$. These solutions appear as deformations of the Peregrine breather P_N as they can be obtained when all parameters are equal to 0. These quasi rational solutions can be expressed as a quotient of two polynomials of degree $N(N + 1)$ in the variables x and t and the maximum of the modulus of the Peregrine breather of order N is equal to $2N + 1$.

Here we give some relations between solutions to this equation. In particular, we present a connection between the modulus of these solutions and the denominator part of their rational expressions. Some relations between numerator and denominator of the Peregrine breather are presented.

2010 AMS : 35B05, 35C99, 35Q55, 35L05, 76M99, 78M99.

Keywords : Fredholm determinants, NLS equation, Peregrine breathers, rogue waves, wronskians.

1 Introduction

We consider the one dimensional focusing nonlinear Schrödinger equation (NLS) which can be written in the form

$$iv_t + v_{xx} + 2|v|^2v = 0, \tag{1}$$

The first results concerning the NLS equation date from the works of Zakharov and Shabat in 1972 who solved it using the inverse scattering method [1, 2]. Its and Kotlyarov first constructed periodic and almost periodic algebro-geometric solutions to the focusing NLS equation in 1976 [3, 4]. Ma found in 1979 the first

breather type solution of the NLS equation [5]. In 1983, the first quasi rational solutions of NLS equation were constructed by Peregrine [6]. In 1986, Eleon-ski, Akhmediev and Kulagin obtained the two-phase almost periodic solution to the NLS equation and got the first higher order analogue of the Peregrine breather[7, 8, 9]. Other analogues of the Peregrine breathers of order 3 and 4 were constructed using Darboux transformations, in a series of articles by Akhmediev et al. [10, 11, 12, 13].

Recently, many works about NLS equation have been published using different methods. We can quote the works of Matveev et al. [14, 15] in 2010 for the representation of the solutions in terms of wronskians; those of Gaillard [16, 17, 18] for the solutions given in terms of wronskians and Fredholm determinants, and their quasi-rational solutions limit of order N depending on $2N - 2$ parameters. Akhmediev gave quasi rational solutions using Darboux transformation in several papers [19, 20, 21]. Guo, Ling and Liu in 2012 gave an other representation of the solutions as a ratio of two determinants [22] using generalized Darboux transformation. A new approach has been done by Ohta and Yang in [23] using Hirota bilinear method. Smirnov [24] gave solutions with an algebro-geometric approach. Other types of solutions were given by Zhao et al. in [25].

We give some relations between the modulus of these solutions and the denominator part of their rational expression. Some relations between numerator and denominator of the rational solutions are given.

2 Different representations of solutions to the NLS equation

2.1 Solutions of the NLS equation in terms of of Fredholm determinant

We have to define the following notations.

The terms $\kappa_\nu, \delta_\nu, \gamma_\nu$ and $x_{r,\nu}$ are functions of the parameters $\lambda_\nu, 1 \leq \nu \leq 2N$; they are defined by the formulas :

$$\begin{aligned} \kappa_\nu &= 2\sqrt{1 - \lambda_\nu^2}, & \delta_\nu &= \kappa_\nu \lambda_\nu, & \gamma_\nu &= \sqrt{\frac{1 - \lambda_\nu}{1 + \lambda_\nu}}; \\ x_{r,\nu} &= (r - 1) \ln \frac{\gamma_\nu - i}{\gamma_\nu + i}, & r &= 1, 3. \end{aligned} \quad (2)$$

The parameters $-1 < \lambda_\nu < 1, \nu = 1, \dots, 2N$, are real numbers such that

$$\begin{aligned} -1 < \lambda_{N+1} < \lambda_{N+2} < \dots < \lambda_{2N} < 0 < \lambda_N < \lambda_{N-1} < \dots < \lambda_1 < 1 \\ \lambda_{N+j} &= -\lambda_j, \quad j = 1, \dots, N. \end{aligned} \quad (3)$$

The condition (3) implies that

$$\kappa_{j+N} = \kappa_j, \quad \delta_{j+N} = -\delta_{j+N}, \quad \gamma_{j+N} = \gamma_j^{-1}, \quad x_{r,j+N} = x_{r,j}, \quad j = 1, \dots, N. \quad (4)$$

Complex numbers e_ν $1 \leq \nu \leq 2N$ are defined in the following way :

$$\begin{aligned} e_j &= i \sum_{l=1}^{N-1} a_l(j\epsilon)^{2l+1} - \sum_{l=1}^{N-1} b_l(j\epsilon)^{2l+1}, \\ e_{j+N} &= i \sum_{l=1}^{N-1} a_l(j\epsilon)^{2l+1} + \sum_{l=1}^{N-1} b_l(j\epsilon)^{2l+1}, \\ 1 \leq j &\leq N-1. \end{aligned} \quad (5)$$

$\epsilon, a_\nu, b_\nu, \nu = 1 \dots 2N$ are arbitrary real numbers.
Let I be the unit matrix, and

$$\epsilon_j = j \quad 1 \leq j \leq N, \quad \epsilon_j = N + j, \quad N + 1 \leq j \leq 2N. \quad (6)$$

Let's consider the matrix $D_r = (d_{jk}^{(r)})_{1 \leq j, k \leq 2N}$ defined by :

$$d_{\nu\mu}^{(r)} = (-1)^{\epsilon_\nu} \prod_{\eta \neq \mu} \left| \frac{\gamma_\eta + \gamma_\nu}{\gamma_\eta - \gamma_\mu} \right| \exp(i\kappa_\nu x - 2\delta_\nu t + x_{r,\nu} + e_\nu). \quad (7)$$

With these notations, the solution to the NLS equation takes the form [16, 17, 18] :

Theorem 2.1 *The function v defined by*

$$v(x, t) = \frac{\det(I + D_3(x, t))}{\det(I + D_1(x, t))} e^{2it - i\varphi}. \quad (8)$$

is a solution to the focusing NLS equation depending on $2N - 1$ real parameters $a_j, b_j, \epsilon, 1 \leq j \leq N - 1$ with the matrix $D_r = (d_{jk}^{(r)})_{1 \leq j, k \leq 2N}$ defined by

$$d_{\nu\mu}^{(r)} = (-1)^{\epsilon_\nu} \prod_{\eta \neq \mu} \left| \frac{\gamma_\eta + \gamma_\nu}{\gamma_\eta - \gamma_\mu} \right| \exp(i\kappa_\nu x - 2\delta_\nu t + x_{r,\nu} + e_\nu).$$

where $\kappa_\nu, \delta_\nu, x_{r,\nu}, \gamma_\nu, e_\nu$ being defined in(2), (3) and (5).

2.2 Wronskian representation

For this, we need to define the following notations :

$$\phi_{r,\nu} = \sin \Theta_{r,\nu}, \quad 1 \leq \nu \leq N, \quad \phi_{r,\nu} = \cos \Theta_{r,\nu}, \quad N + 1 \leq \nu \leq 2N, \quad r = 1, 3, \quad (9)$$

with the arguments

$$\Theta_{r,\nu} = \kappa_\nu x / 2 + i\delta_\nu t - ix_{r,\nu} / 2 + \gamma_\nu y - ie_\nu / 2, \quad 1 \leq \nu \leq 2N. \quad (10)$$

The functions $\phi_{r,\nu}$ are defined by

$$\phi_{r,\nu} = \sin \Theta_{r,\nu}, \quad 1 \leq \nu \leq N, \quad \phi_{r,\nu} = \cos \Theta_{r,\nu}, \quad N + 1 \leq \nu \leq 2N, \quad r = 1, 3, \quad (11)$$

We denote $W_r(y)$ the wronskian of the functions $\phi_{r,1}, \dots, \phi_{r,2N}$ defined by

$$W_r(y) = \det[(\partial_y^{\mu-1} \phi_{r,\nu})_{\nu, \mu \in [1, \dots, 2N]}]. \quad (12)$$

We consider the matrix $D_r = (d_{\nu\mu})_{\nu, \mu \in [1, \dots, 2N]}$ defined in (7). Then we have the following statement [17] :

Theorem 2.2

$$\det(I + D_r) = k_r(0) \times W_r(\phi_{r,1}, \dots, \phi_{r,2N})(0), \quad (13)$$

where

$$k_r(y) = \frac{2^{2N} \exp(i \sum_{\nu=1}^{2N} \Theta_{r,\nu})}{\prod_{\nu=2}^{2N} \prod_{\mu=1}^{\nu-1} (\gamma_\nu - \gamma_\mu)}.$$

With these notations, we have the following result [17] :

Theorem 2.3 *The function v defined by*

$$v(x, t) = \frac{W_3(\phi_{3,1}, \dots, \phi_{3,2N})(0)}{W_1(\phi_{1,1}, \dots, \phi_{1,2N})(0)} e^{2it - i\varphi}.$$

is a solution to the focusing NLS equation depending on $2N - 1$ real parameters $a_j, b_j, \epsilon, 1 \leq j \leq N - 1$ with ϕ_ν^r defined in (11)

$$\begin{aligned} \phi_{r,\nu} &= \sin(\kappa_\nu x/2 + i\delta_\nu t - ix_{r,\nu}/2 + \gamma_\nu y - ie_\nu/2), \quad 1 \leq \nu \leq N, \\ \phi_{r,\nu} &= \cos(\kappa_\nu x/2 + i\delta_\nu t - ix_{r,\nu}/2 + \gamma_\nu y - ie_\nu/2), \quad N + 1 \leq \nu \leq 2N, \quad r = 1, 3, \end{aligned}$$

$\kappa_\nu, \delta_\nu, x_{r,\nu}, \gamma_\nu, e_\nu$ being defined in(2), (3) and (5).

We can give another representation of the solutions to the NLS equation depending only on terms $\gamma_\nu, 1 \leq \nu \leq 2N$. From the relations (2), we can express the terms κ_ν, δ_ν and $x_{r,\nu}$ in function of γ_ν , for $1 \leq \nu \leq 2N$ and we obtain :

$$\begin{aligned} \kappa_j &= \frac{4\gamma_j}{(1 + \gamma_j^2)}, \quad \delta_j = \frac{4\gamma_j(1 - \gamma_j^2)}{(1 + \gamma_j^2)^2}, \quad x_{r,j} = (r - 1) \ln \frac{\gamma_j - i}{\gamma_j + i}, \quad 1 \leq j \leq N, \\ \kappa_j &= \frac{4\gamma_j}{(1 + \gamma_j^2)}, \quad \delta_j = -\frac{4\gamma_j(1 - \gamma_j^2)}{(1 + \gamma_j^2)^2}, \quad x_{r,j} = (r - 1) \ln \frac{\gamma_j + i}{\gamma_j - i}, \quad N + 1 \leq j \leq 2N. \end{aligned} \quad (14)$$

We have the following new representation [17, 27] :

Theorem 2.4 *The function v defined by*

$$v(x, t) = \frac{\det[(\partial_y^{\mu-1} \tilde{\phi}_{3,\nu}(0))_{\nu, \mu \in [1, \dots, 2N]}]}{\det[(\partial_y^{\mu-1} \tilde{\phi}_{1,\nu}(0))_{\nu, \mu \in [1, \dots, 2N]}]} e^{2it - i\varphi} \quad (15)$$

is a solution to the NLS equation (1) depending on $2N - 1$ real parameters $a_j, b_j, \epsilon, 1 \leq j \leq N - 1$. The functions $\tilde{\phi}_{r,\nu}$ are defined by

$$\begin{aligned} \tilde{\phi}_{r,j}(y) &= \sin \left(\frac{2\gamma_j}{(1 + \gamma_j^2)} x + i \frac{4\gamma_j(1 - \gamma_j^2)}{(1 + \gamma_j^2)^2} t - i \frac{(r-1)}{2} \ln \frac{\gamma_j - i}{\gamma_j + i} + \gamma_j y - ie_j \right), \\ \tilde{\phi}_{r,N+j}(y) &= \cos \left(\frac{2\gamma_j}{(1 + \gamma_j^2)} x - i \frac{4\gamma_j(1 - \gamma_j^2)}{(1 + \gamma_j^2)^2} t + i \frac{(r-1)}{2} \ln \frac{\gamma_j - i}{\gamma_j + i} + \frac{1}{\gamma_j} y - ie_{N+j} \right), \end{aligned} \quad (16)$$

where $\gamma_j = \sqrt{\frac{1 - \lambda_j}{1 + \lambda_j}}, 1 \leq j \leq N$.

λ_j is an arbitrary real parameter such that $0 < \lambda_j < 1, \lambda_{N+j} = -\lambda_j, 1 \leq j \leq N$.

The terms e_ν are defined by (5),

where a_j and b_j are arbitrary real numbers, $1 \leq j \leq N - 1$.

Remark 2.1 In the formula (15), the determinants $\det[(\partial_y^{\mu-1} f_\nu(0))_{\nu, \mu \in [1, \dots, 2N]}]$ are the wronskians of the functions f_1, \dots, f_{2N} evaluated in $y = 0$. In particular $\partial_y^0 f_\nu$ means f_ν .

2.3 Families of quasi-rational solutions of NLS equation in terms of a quotient of two determinants

The following notations are used :

$$X_\nu = \kappa_\nu x/2 + i\delta_\nu t - ix_{3,\nu}/2 - ie_\nu/2,$$

$$Y_\nu = \kappa_\nu x/2 + i\delta_\nu t - ix_{1,\nu}/2 - ie_\nu/2,$$

for $1 \leq \nu \leq 2N$, with $\kappa_\nu, \delta_\nu, x_{r,\nu}$ defined in (2).

Parameters e_ν are defined by (5).

Below the following functions are used :

$$\begin{aligned} \varphi_{4j+1,k} &= \gamma_k^{4j-1} \sin X_k, & \varphi_{4j+2,k} &= \gamma_k^{4j} \cos X_k, \\ \varphi_{4j+3,k} &= -\gamma_k^{4j+1} \sin X_k, & \varphi_{4j+4,k} &= -\gamma_k^{4j+2} \cos X_k, \end{aligned} \quad (17)$$

for $1 \leq k \leq N$, and

$$\begin{aligned} \varphi_{4j+1,N+k} &= \gamma_k^{2N-4j-2} \cos X_{N+k}, & \varphi_{4j+2,N+k} &= -\gamma_k^{2N-4j-3} \sin X_{N+k}, \\ \varphi_{4j+3,N+k} &= -\gamma_k^{2N-4j-4} \cos X_{N+k}, & \varphi_{4j+4,N+k} &= \gamma_k^{2N-4j-5} \sin X_{N+k}, \end{aligned} \quad (18)$$

for $1 \leq k \leq N$.

We define the functions $\psi_{j,k}$ for $1 \leq j \leq 2N, 1 \leq k \leq 2N$ in the same way, the term X_k is only replaced by Y_k .

$$\begin{aligned} \psi_{4j+1,k} &= \gamma_k^{4j-1} \sin Y_k, & \psi_{4j+2,k} &= \gamma_k^{4j} \cos Y_k, \\ \psi_{4j+3,k} &= -\gamma_k^{4j+1} \sin Y_k, & \psi_{4j+4,k} &= -\gamma_k^{4j+2} \cos Y_k, \end{aligned} \quad (19)$$

for $1 \leq k \leq N$, and

$$\begin{aligned} \psi_{4j+1,N+k} &= \gamma_k^{2N-4j-2} \cos Y_{N+k}, & \psi_{4j+2,N+k} &= -\gamma_k^{2N-4j-3} \sin Y_{N+k}, \\ \psi_{4j+3,N+k} &= -\gamma_k^{2N-4j-4} \cos Y_{N+k}, & \psi_{4j+4,N+k} &= \gamma_k^{2N-4j-5} \sin Y_{N+k}, \end{aligned} \quad (20)$$

for $1 \leq k \leq N$.

Then we get the following result [27] :

Theorem 2.5 *The function v defined by*

$$v(x, t) = \frac{\det((n_{jk})_{j,k \in [1, 2N]})}{\det((d_{jk})_{j,k \in [1, 2N]})} e^{2it - i\varphi} \quad (21)$$

is a quasi-rational solution of the NLS equation (1) depending on $2N - 2$ real parameters $a_j, b_j, 1 \leq j \leq N - 1$, where

$$\begin{aligned} n_{j1} &= \varphi_{j,1}(x, t, 0), 1 \leq j \leq 2N & n_{jk} &= \frac{\partial^{2k-2} \varphi_{j,1}}{\partial \epsilon^{2k-2}}(x, t, 0), \\ n_{jN+1} &= \varphi_{j,N+1}(x, t, 0), 1 \leq j \leq 2N & n_{jN+k} &= \frac{\partial^{2k-2} \varphi_{j,N+1}}{\partial \epsilon^{2k-2}}(x, t, 0), \\ d_{j1} &= \psi_{j,1}(x, t, 0), 1 \leq j \leq 2N & d_{jk} &= \frac{\partial^{2k-2} \psi_{j,1}}{\partial \epsilon^{2k-2}}(x, t, 0), \\ d_{jN+1} &= \psi_{j,N+1}(x, t, 0), 1 \leq j \leq 2N & d_{jN+k} &= \frac{\partial^{2k-2} \psi_{j,N+1}}{\partial \epsilon^{2k-2}}(x, t, 0), \\ & & & 2 \leq k \leq N, 1 \leq j \leq 2N \end{aligned}$$

The functions φ and ψ are defined in (17), (18), (19), (20).

3 Structure of the multi-parametric solutions to the NLS equation of order N depending on $2N - 2$ parameters

3.1 The quotient of two polynomials of degree $(N(N + 1))$ in x and t by an exponential depending on t

Here we present a result which states the structure of the quasi-rational solutions of the NLS equation. It was only conjectured in preceding works [16, 18]. Moreover we obtain here families of deformations of the N th Peregrine breather depending on $2N - 2$ parameters.

In this section we use the notations defined in the previous sections. The functions φ and ψ are defined in (17), (18), (19), (20).

The structure of the quasi rational solutions to the NLS equation is given by [28] :

Theorem 3.1 *The function v defined by*

$$v(x, t) = \frac{\det((n_{jk})_{j,k \in [1, 2N]})}{\det((d_{jk})_{j,k \in [1, 2N]})} e^{2it - i\varphi} \quad (22)$$

is a quasi-rational solution of the NLS equation (1) quotient of two polynomials $R(x, t)$ and $S(x, t)$ depending on $2N - 2$ real parameters a_j and $b_j, 1 \leq j \leq N - 1$. $R(x, t)$ and $S(x, t)$ are polynomials of degrees $N(N + 1)$ in x and t .

Remark 3.1 *The polynomials $R(x, t)$ and $S(x, t)$ have the same coefficients of degrees $N(N + 1)$ in $2x$ and $4t$ equal to 1. The polynomial $B(x, t)$ does not have any real root.*

3.2 The structure of the Peregrine breather of order n

There is any freedom to choose γ_j in such a way that the conditions on λ_j are checked. We know from previous works [16, 18] that the (analogue) Peregrine breathers are obtained when all the parameters a_j and b_j are equal to 0. In order to get the more simple expressions in the determinants, we choose particular solutions in the previous families.

Here we choose $\gamma_j = j\epsilon$ as simple as possible in order to have the conditions on λ_j checked, and we have [27, 28] :

Theorem 3.2 *The function v_0 defined by*

$$v_{n,0}(x, t) = \left(\frac{\det((n_{jk})_{j,k \in [1,2N]})}{\det((d_{jk})_{j,k \in [1,2N]})} e^{2it - i\varphi} \right)_{(a_j = b_j = 0, 1 \leq j \leq N-1)} \quad (23)$$

is the Peregrine breather of order N solution of the NLS equation (1) whose highest amplitude in modulus is equal to $2N + 1$.

Remark 3.2 *The previous result is given in the frame where the limit of the modulus of the solution when x or t tend to infinity is equal to 1. We know that if $v(x, t)$ is a solution to the NLS equation then $u(x, t) = av(ax, a^2t)$ is also a solution to the NLS equation, for any arbitrary real a .*

Remark 3.3 *In (23), the matrices $(n_{jk})_{j,k \in [1,2N]}$ and $(d_{jk})_{j,k \in [1,2N]}$ are defined in (22).*

We have seen in previous section that solutions of NLS equation given by (16) can be written in function uniquely of terms γ . We recall that the terms γ_j are

given by $\gamma_j = \sqrt{\frac{1 - \lambda_j}{1 + \lambda_j}}$, $1 \leq j \leq N$; λ_j is an arbitrary real parameter such that $0 < \lambda_j < 1$, $\lambda_{N+j} = -\lambda_j$, $1 \leq j \leq N$.

We can rewrite the result given in (16) in a simplest formulation as follows [27, 28] :

Theorem 3.3 *The function v defined by*

$$v(x, t) = \frac{\det((f_{jk}^{(3)})_{j,k \in [1,2N]})}{\det((f_{jk}^{(1)})_{j,k \in [1,2N]})} e^{2it - i\varphi} \quad (24)$$

is a quasi-rational solution of the NLS equation (1) depending on $2N - 2$ real parameters a_j, b_j , $1 \leq j \leq N - 1$ where

$$\begin{aligned} f_{jk}^{(r)} &= \frac{\partial^{2(k-1)}}{\partial \epsilon^{2(k-1)}} \left(\gamma^{4j-1} \sin \left[\frac{2\gamma}{1+\gamma^2} x + 4i \frac{\gamma(1-\gamma^2)}{(1+\gamma^2)^2} t - i \frac{r-1}{2} \ln \frac{\gamma-i}{\gamma+i} + \sum_{l=1}^{N-1} (a_l + ib_l) \epsilon^{2l+1} + (j-1) \frac{\pi}{2} \right] \right)_{(\epsilon=0)}, \\ f_{jN+k}^{(r)} &= \frac{\partial^{2(k-1)}}{\partial \epsilon^{2(k-1)}} \left(\gamma^{2N-4j-1} \cos \left[\frac{2\gamma}{1+\gamma^2} x - 4i \frac{\gamma(1-\gamma^2)}{(1+\gamma^2)^2} t + i \frac{r-1}{2} \ln \frac{\gamma-i}{\gamma+i} + \sum_{l=1}^{N-1} (a_l - ib_l) \epsilon^{2l+1} + (j-1) \frac{\pi}{2} \right] \right)_{(\epsilon=0)}, \\ 1 \leq k \leq N, \quad 1 \leq j \leq 2N, \quad r \in \{1, 3\}, \quad \epsilon \in]0; 1[, \quad \gamma = \epsilon(1 - \epsilon^2)^{1/2}. \end{aligned}$$

Remark 3.4 In the previous theorem, the expression $\frac{\partial^0}{\partial \epsilon^0} f(x)$ means $f(x)$.

The solution to the NLS equation can be written in the form

$$v_N(x, t) = \frac{R_N(x, t)}{S_N(x, t)} e^{2it} = \left(1 + \frac{A_N(x, t)}{B_N(x, t)} \right) e^{2it} \quad (25)$$

and the Peregrine breather in the form

$$v_{N,0}(x, t) = \frac{T_N(x, t)}{U_N(x, t)} e^{2it} = \left(1 + \frac{P_N(x, t)}{Q_N(x, t)} \right) e^{2it} \quad (26)$$

where the index 0 means that all the parameters are equal to 0.

4 Differential relation for the NLS equation

In previous works [27, 28], we have proven that the solutions v_N to the NLS equation can be written in the form

$$v_N(x, t) = \left(1 + \frac{A_N(x, t)}{B_N(x, t)} \right) e^{2it}. \quad (27)$$

We have a very simple relation between the square of the modulus of v_N and the denominator part B_N . This relation appears in a paper of Ling and Zhao [25] where the solutions to the NLS equation are given in the frame of the generalized Darboux transformation. Here this result and its proof are given in a general frame by the following theorem :

Theorem 4.1 *The solutions $v_N(x, t) = \left(1 + \frac{A_N(x, t)}{B_N(x, t)} \right) e^{2it}$ to the NLS equation verify the following relation*

$$|v_N(x, t)|^2 = 1 + (\ln B_N(x, t))_{xx}, \quad (28)$$

where the subscript $_{xx}$ means the double derivation with respect to x .

5 Relations between rational part of the solutions to the NLS equation

With the preceding notations, we get the following statement

Theorem 5.1 *The polynomials of the solutions v_N to the NLS equation defined by (25) $v_N(x, t) = \frac{R_N(x, t)}{S_N(x, t)} e^{2it}$ verify the following relations*

$$\begin{aligned} & (i(R_N)_t + (R_N)_{xx} - 2R_N)S_N^2 - ((S_N)_{xx} + i(S_N)_t)R_N S_N \\ & - 2(R_N)_x(S_N)_x S_N + 2((S_N)_x^2 + R_N R_N)R_N = 0. \end{aligned} \quad (29)$$

Proposition 5.1 *The coordinates of extrema (x_0, t_0) of solutions v_N to the NLS equation defined by (25) $v_N(x, t) = \frac{R_N(x, t)}{S_N(x, t)} e^{2it}$ verify the the following relations*

$$(R_N)_x(x_0, t_0) \overline{R_N}(x_0, t_0) S_N(x_0, t_0) + (\overline{R_N})_x(x_0, t_0) R_N(x_0, t_0) S_N(x_0, t_0) - 2(S_N)_x(x_0, t_0) R_N(x_0, t_0) \overline{R_N}(x_0, t_0) = 0, \quad (30)$$

$$(R_N)_t(x_0, t_0) \overline{R_N}(x_0, t_0) S_N(x_0, t_0) + (\overline{R_N})_t(x_0, t_0) R_N(x_0, t_0) S_N(x_0, t_0) - 2(S_N)_t(x_0, t_0) R_N(x_0, t_0) \overline{R_N}(x_0, t_0) = 0. \quad (31)$$

$$(R_N)_x(x_0, t_0) S_N(x_0, t_0) - (S_N)_x(x_0, t_0) R_N(x_0, t_0) = 0. \quad (32)$$

$$(R_N)_t(x_0, t_0) S_N(x_0, t_0) - (S_N)_t(x_0, t_0) R_N(x_0, t_0) + 2i S_N(x_0, t_0) R_N(x_0, t_0) = 0. \quad (33)$$

where \bar{a} means the complex conjugate of a .

Remark 5.1 *As a consequence of the result on the highest modulus of the P_N breather defined by (26) $v_{N,0}(x, t) = \frac{T_N(x, t)}{U_N(x, t)} e^{2it}$, we get*

$$T_N(0, 0) = (2N + 1)U_N(0, 0). \quad (34)$$

6 Conclusion

Different representations of the solutions to the NLS equation have been summarized in this paper, as well as the structure of the quasi rational solutions. Some differential relations have been given in this text for the NLS equation. From different studies realized by the author, [26, 27, 28, 29, 30, 31, 32], it seems that the maximums of the modulus of the solutions to the NLS equation are in connection with the zeros of the Yablonski-Vorob'ev polynomials [33, 34].

It would be relevant to study this conjecture.

It would be also relevant to search other types of equations verified by the polynomials (P_N, Q_N) , (R_N, S_N) , (A_N, B_N) or (T_N, U_N) .

References

- [1] V. E. Zakharov, Stability of periodic waves of finite amplitude on a surface of a deep fluid, J. Appl. Tech. Phys, V. **9**, 86-94, (1968)
- [2] V. E. Zakharov, A.B. Shabat Exact theory of two dimensional self focusing and one dimensional self modulation of waves in nonlinear media, Sov. Phys. JETP, V. **34**, 62-69, (1972)

- [3] A.R. Its, A.V. Rybin, M.A. Salle, Exact integration of nonlinear Schrödinger equation, *Teore. i Mat. Fiz.*, V. **74.**, N. 1, 29-45, (1988).
- [4] A.R. Its, V.P. Kotlyarov, Explicit expressions for the solutions of nonlinear Schrödinger equation, *Dokl. Akad. Nauk. SSSR, S. A*, V. **965.**, N. 11, (1976).
- [5] Y.C. Ma, The perturbed plane-wave solutions of the cubic nonlinear Schrödinger equation, *Stud. Appl. Math.* V. **60**, 43-58 (1979).
- [6] D.H. Peregrine, Water waves, nonlinear Schrödinger equations and their solutions, *J. Austral. Math. Soc. Ser. B*, V. **25**, 16-43, (1983).
- [7] N. Akhmediev, V. Eleonskii, N. Kulagin, Exact first order solutions of the nonlinear Schrödinger equation, *Th. Math. Phys.*, V. **72**, N. 2, 183-196, (1987).
- [8] N. Akhmediev, V. Eleonsky, N. Kulagin, Generation of periodic trains of picosecond pulses in an optical fiber : exact solutions, *Sov. Phys. J.E.T.P.*, V. **62**, 894-899, (1985).
- [9] V. Eleonskii, I. Krichever, N. Kulagin, Rational multisoliton solutions to the NLS equation, *Soviet Doklady 1986 sect. Math. Phys.*, V. **287**, 606-610, (1986).
- [10] N. Akhmediev, A. Ankiewicz, J.M. Soto-Crespo, Rogue waves and rational solutions of nonlinear Schrödinger equation, *Physical Review E*, V. **80**, N. 026601, (2009).
- [11] N. Akhmediev, A. Ankiewicz, P.A. Clarkson, Rogue waves, rational solutions, the patterns of their zeros and integral relations, *J. Phys. A : Math. Theor.*, V. **43**, 122002, 1-9, (2010).
- [12] N. Akhmediev, A. Ankiewicz, D. J. Kedziora, Circular rogue wave clusters, *Phys. Review E*, V. **84**, 1-7, (2011)
- [13] A. Chabchoub, H. Hoffmann, M. Onorato, N. Akhmediev, Super rogue waves : observation of a higher-order breather in water waves, *Phys. Review X*, V. **2**, 1-6, (2012).
- [14] P. Dubard, P. Gaillard, C. Klein, V.B. Matveev, On multi-rogue waves solutions of the NLS equation and positon solutions of the KdV equation, *Eur. Phys. J. Special Topics*, V. **185**, 247-258, (2010).
- [15] P. Dubard, V.B. Matveev, Multi-rogue waves solutions : from the NLS to the KP-I equation, *Nonlinearity*, V. **26**, 93-125, 2013
- [16] P. Gaillard, Families of quasi-rational solutions of the NLS equation and multi-rogue waves, *J. Phys. A : Meth. Theor.*, V. **44**, 1-15, (2011)

- [17] P. Gaillard, Wronskian representation of solutions of the NLS equation and higher Peregrine breathers, *Jour. Of Math. Sciences : Advances and Applications*, V. **13**, N. 2, 71-153, (2012)
- [18] P. Gaillard, Degenerate determinant representation of solution of the NLS equation, higher Peregrine breathers and multi-rogue waves, *Jour. Of Math. Phys.*, V. **54**, 013504-1-32, (2013)
- [19] D.J. Kedziora, A. Ankiewicz, N. Akhmediev, Triangular rogue wave cascades, *Phys. Rev. E*, V. **86**, 056602, 1-9, (2012).
- [20] D. J. Kedziora, A. Ankiewicz, N. Akhmediev, Circular rogue wave clusters, *Phys. Review E*, V. 84, 056611-1-7, 2011
- [21] D. J. Kedziora, A. Ankiewicz, N. Akhmediev, Classifying the hierarchy of the nonlinear Schrödinger equation rogue waves solutions, *Phys. Review E*, V. 88, 013207-1-12, 2013
- [22] B. Guo, L. Ling, Q.P. Liu, Nonlinear Schrodinger equation: Generalized Darboux transformation and rogue wave solutions, *Phys. Rev. E*, V. **85**, 026607, (2012).
- [23] Y. Ohta, J. Yang, General high-order rogue waves and their dynamics in the nonlinear Schrödinger equation, *Pro. R. Soc. A*, V. 468, 1716-1740, 2012
- [24] A.O. Smirnov, Solution of a nonlinear Schrödinger equation in the form of two phase freak waves, *Theor. Math. Phys.*, V. 173, 1403-1416, 2012
- [25] L. Ling, L.C. Zhao, Simple determinant representation for rogue waves of the nonlinear Schrödinger equation, *Phys. Rev. E*, V; 88, 043201-1-9, 2013
- [26] P. Gaillard, Deformations of third order Peregrine breather solutions of the NLS equation with four parameters, *Phys. Rev. E*, V. **88**, 042903-1-9, 2013
- [27] P. Gaillard, Other $2N-2$ parameters solutions to the NLS equation and $2N+1$ highest amplitude of the modulus of the N -th order AP breather, *J. Phys. A: Math. Theor.*, V. **48**, 145203-1-23, 2015
- [28] P. Gaillard, Multi-parametric deformations of the Peregrine breather of order N solutions to the NLS equation and multi-rogue waves, *Adv. Res.*, V. **4**, 346-364, 2015
- [29] P. Gaillard, M. Gastineau Twenty parameters families of solutions to the NLS equation and the eleventh Peregrine breather, *Commun. Theor. Phys*, V. **65**, 136-144, 2016
- [30] P. Gaillard, M. Gastineau Twenty two parameters deformations of the twelfth Peregrine breather solutions to the NLS equation, *Adv. Res.*, V. **10**, 83-89, 2016

- [31] P. Gaillard, Towards a classification of the quasi rational solutions to the NLS equation, *Theor. And Math. Phys.*, V. **189**, 1440-1449, 2016
- [32] P. Gaillard, M. Gastineau Families of deformations of the thirteenth Peregrine breather solutions to the NLS equation depending on twenty four parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. **21**, N. 3, 130-139, 2017
- [33] A.P. Vorob'ev, On the rational solutions of the second Painlevé equation, *Differ. Uravn.*, V. 1, N.1, 79-81, 1965
- [34] A.I. Yablonskii, On rational solutions of the second Painlevé equation, *Vesti AN BSSR, Ser. Fiz.-Tech. Nauk*, N. 3, 3035, 1959
- [35] P. Gaillard, V.B. Matveev, Wronskian addition formula and its applications, *Max-Planck-Institut für Mathematik, MPI 02-31*, V. **161**, 2002
- [36] P. Gaillard, A new family of deformations of Darboux-Pöschl-Teller potentials, *Lett. Math. Phys.*, V. **68**, 77-90, 2004
- [37] P. Gaillard, V.B. Matveev, New formulas for the eigenfunctions of the two-particle Calogero-Moser system, *Lett. Math. Phys.*, V. **89**, 1-12, 2009
- [38] P. Gaillard, V.B. Matveev, Wronskian and Casorai determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, *RIMS Kyoto*, N. 1653, 1-19, 2009
- [39] P. Gaillard, V.B. Matveev, Wronskian and Casorai determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, *J. Phys A : Math. Theor.*, V. **42**, 1-16, 2009
- [40] P. Gaillard, From finite-gap solutions of KdV in terms of theta functions to solitons and positons, *halshs-00466159*, 2010
- [41] P. Gaillard, Wronskian representation of solutions of NLS equation and seventh order rogue waves, *J. Mod. Phys.*, V. **4**, N. 4, 246-266, 2013
- [42] P. Gaillard, V.B. Matveev, Wronskian addition formula and Darboux-Pöschl-Teller potentials, *J. Math.*, V. **2013**, ID 645752, 1-10, 2013
- [43] P. Gaillard, Two parameters deformations of ninth Peregrine breather solution of the NLS equation and multi rogue waves, *J. Math.*, V. **2013**, 1-111, 2013
- [44] P. Gaillard, Two-parameters determinant representation of seventh order rogue waves solutions of the NLS equation, *J. Theor. Appl. Phys.*, V. **7**, N. **45**, 1-6, 2013
- [45] P. Gaillard, Six-parameters deformations of fourth order Peregrine breather solutions of the NLS equation, *J. Math. Phys.*, V. **54**, 073519-1-22, 2013

- [46] P. Gaillard, Ten parameters deformations of the sixth order Peregrine breather solutions of the NLS equation, *Phys. Scripta*, V. **89**, 015004-1-7, 2014
- [47] P. Gaillard, The fifth order Peregrine breather and its eight-parameters deformations solutions of the NLS equation, *Commun. Theor. Phys.*, V. **61**, 365-369, 2014
- [48] P. Gaillard, Higher order Peregrine breathers, their deformations and multi-rogue waves, *J. Of Phys. : Conf. Ser.*, V. **482**, 012016-1-7, 2014
- [49] P. Gaillard, M. Gastineau, Eighteen parameter deformations of the Peregrine breather of order ten solutions of the NLS equation, *Int. J. Mod. Phys. C*, V. **26**, N. 2, 1550016-1-14, 2014
- [50] P. Gaillard, Two parameters wronskian representation of solutions of nonlinear Schrödinger equation, eight Peregrine breather and multi-rogue waves, *J. Math. Phys.*, V. **5**, 093506-1-12, 2014
- [51] P. Gaillard, Hierarchy of solutions to the NLS equation and multi-rogue waves, *J. Phys. : Conf. Ser.*, V. **574**, 012031-1-5, 2015
- [52] P. Gaillard, Tenth Peregrine breather solution of the NLS, *Ann. Phys.*, V. **355**, 293-298, 2015
- [53] P. Gaillard, M. Gastineau, The Peregrine breather of order nine and its deformations with sixteen parameters solutions of the NLS equation, *Phys. Lett. A.*, V. **379**, 13091313, 2015
- [54] P. Gaillard, Higher order Peregrine breathers solutions to the NLS equation, *Jour. Phys. : Conf. Ser.*, V. **633**, 012106-1-6, 2016
- [55] P. Gaillard, M. Gastineau Patterns of deformations of Peregrine breather of order 3 and 4, solutions to the NLS equation with multi-parameters, *Journal of Theoretical and Applied Physics*, V. **10**,1-7, 2016
- [56] P. Gaillard, Rational solutions to the KPI equation and multi rogue waves, *Annals Of Physics*, V. **367**, 1-5, 2016
- [57] P. Gaillard, Fredholm and Wronskian representations of solutions to the KPI equation and multi-rogue waves, *Jour. of Math. Phys.*, V. **57**, 063505-1-13, doi: 10.1063/1.4953383, 2016
- [58] P. Gaillard, From Fredholm and Wronskian representations to rational solutions to the KPI equation depending on $2N^2$ parameters, *Int. Jour. of Appl. Sci. And Math.*, V. **4**, N. 3, 60-70, 2017
- [59] P. Gaillard, Families of Rational Solutions of Order 5 to the KPI Equation Depending on 8 Parameters, *New Hor. in Math. Phys.*, V. 1, N. 1, 26-31, 2017

- [60] P. Gaillard, 6-th order rational solutions to the KPI Equation depending on 10 parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. **21**, N. 2, 92-98, 2017
- [61] P. Gaillard, N -Order rational solutions to the Johnson equation depending on $2N - 2$ parameters, *Int. Jour. of Adv. Res. in Phys. Sci.*, V. **4**, N. 9, 19-37, 2017
- [62] P. Gaillard, Families of rational solutions to the KPI equation of order 7 depending on 12 parameters, *Int. Jour. of Adv. Res. in Phys. Sci.*, V. **4**, N. 11, 24-30, 2017
- [63] P. Gaillard, Rational solutions to the Johnson equation and rogue waves, *Int. Jour. of Inn. In Sci. and Math.*, V. **6**, N. 1, 14-19, 2018
- [64] P. Gaillard, Multiparametric families of solutions of the KPI equation, the structure of their rational representations and multi-rogue waves, *Theo. And Mat. Phys.*, V. **196**, N. 2, 1174-1199, 2018
- [65] P. Gaillard, The Johnson Equation, Fredholm and Wronskian representations of solutions and the case of order three, *Adv. In Math. Phys.*, V. **2018**, 1-18, 2018
- [66] P. Gaillard, Families of Solutions of Order 5 to the Johnson Equation Depending on 8 Parameters, *NHIMP*, V. **2**, N. 4, 53-61, 2018
- [67] P. Gaillard, Multiparametric families of solutions to the Johnson equation, *J. Phys. : Conf. Series*, V. **1141**, 012102-1-10, 2018
- [68] P. Gaillard, Rational solutions to the Boussinesq equation, *Fund. Jour. Of Math. And Appl.*, V. , 109-112, 2019