



1 H, 13 C and 15 N assignments of human Grb2 free of ligands

Louise Pinet, Ying-Hui Wang, Anaïs Vogel, Françoise Guerlesquin, Nadine Assrir, Carine van Heijenoort

► To cite this version:

Louise Pinet, Ying-Hui Wang, Anaïs Vogel, Françoise Guerlesquin, Nadine Assrir, et al.. 1 H, 13 C and 15 N assignments of human Grb2 free of ligands. *Biomolecular NMR Assignments*, 2020, 14 (2), pp.323 - 327. 10.1007/s12104-020-09970-7 . hal-03095478

HAL Id: hal-03095478

<https://hal.science/hal-03095478>

Submitted on 6 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

${}^1\text{H}$, ${}^{13}\text{C}$ and ${}^{15}\text{N}$ assignments of human Grb2 free of ligands

Louise Pinet, Ying-Hui Wang, Anaïs Vogel, Françoise Guerlesquin, Nadine Assrir, et al.

Biomolecular NMR Assignments

ISSN 1874-2718

Biomol NMR Assign

DOI 10.1007/s12104-020-09970-7



Your article is published under the Creative Commons Attribution license which allows users to read, copy, distribute and make derivative works, as long as the author of the original work is cited. You may self-archive this article on your own website, an institutional repository or funder's repository and make it publicly available immediately.



^1H , ^{13}C and ^{15}N assignments of human Grb2 free of ligands

Louise Pinet^{1,2} · Ying-Hui Wang^{1,3} · Anaïs Vogel^{1,4} · Françoise Guerlesquin⁵ · Nadine Assrir¹ · Carine van Heijenoort¹

Received: 7 May 2020 / Accepted: 31 July 2020
© The Author(s) 2020

Abstract

Growth factor receptor-bound 2 (Grb2) is an important link in the receptor tyrosine kinase signaling cascades. It is involved in crucial processes, both physiological (mainly embryogenesis) and pathological (different types of cancer). Several binding partners of all three domains (SH3–SH2–SH3) of this adaptor protein are well described, such as ErbB family members for the SH2 domain and Sos for the SH3 domains. How the different domains interact with each other, both structurally and functionally, is still unclear. These interactions could be essential for regulation processes, and therefore are of great interest. Although a lot of structural data on Grb2 exist, they describe either individual domains, ligand-bound conformations, or frozen pictures of the protein captured by crystallography. Here we report the assignment of backbone and of $^{13}\text{C}_\beta$ chemical shifts of full-length, apo-Grb2 in solution. In addition to the assigned conformation corresponding to three well-folded domains, a set of peaks compatible with the presence of an unfolded conformation of the N-terminal SH3 domain is observed. This assignment paves the way for future studies of inter-domain interactions and dynamics that have to be taken into account when studying the regulation of Grb2 interactions and signaling pathways.

Keywords Growth factor receptor-bound 2 (Grb2) · Adaptor protein · SH2 · SH3 · RTK signaling

Biological context

Growth factor receptor-bound 2 (Grb2) is an adaptor protein essential for early steps of embryogenesis, especially differentiation (Cheng et al. 1998), but it was also shown to be upregulated in a subset of breast cancers (Daly et al. 1994). It was originally discovered in humans as the missing link between receptor tyrosine kinases (RTKs) and the Ras/MAPK pathway (Lowenstein et al. 1992). Grb2 is 217 residues long (with a molecular weight of 25 kDa) and is made

up of three domains: a central SH2 domain, a well-described type of phosphotyrosine-binding domain, is flanked by two SH3 domains (called NSH3 at the N-terminus and CSH3 at the C-terminus) that bind to proline-rich motifs. Grb2 binds to phosphorylated RTKs via its SH2 domain and to the Son of sevenless guanine nucleotide exchange factor (Sos) via its SH3 domains, triggering Ras/MAPK pathway signaling (Lowenstein et al. 1992; Gale et al. 1993; Buday and Downward 1993), but is also involved in other signaling pathways (Gu and Neel 2003; Ravid et al. 2004; Yamazaki et al. 2002).

✉ Carine van Heijenoort
Carine.van-Heijenoort@cnrs.fr

Louise Pinet
l.pinet@bioc.uzh.ch

Ying-Hui Wang
ying-hui.wang@sgs.com

Anaïs Vogel
vogel.anais95@gmail.com

Françoise Guerlesquin
guerlesq@imm.cnrs.fr

Nadine Assrir
Nadine.Assrir@cnrs.fr

¹ Department of Analytical and Structural Chemistry and Biology, Institut de Chimie des Substances Naturelles, CNRS UPR2301, Université Paris-Saclay, 1, av. de la terrasse, 91190 Gif-sur-Yvette, France

² Present Address: Department of Biochemistry, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

³ Present Address: SGS Taiwan LTD, No.38, Wu Chyuan 7th Rd., New Taipei Industrial Park, Wu Ku District, New Taipei City 24890, Taiwan

⁴ Present Address: NG Biotech, ZI Courbouton, 35480 Guipry, France

⁵ LISM, Institut de Microbiologie de la Méditerranée, CNRS and Aix-Marseille University, Marseille, France

The structures of the NSH3 domain in complex with Sos proline-rich peptides (PDB 1GBR) and of the CSH3 free domain (PDB 1GFC) were solved by NMR soon after Grb2 identification (Goudreau et al. 1994; Wittekind et al. 1994; Kohda et al. 1994). The SH2 domain structure was solved shortly after by X-ray crystallography (PDB1GHU) (Thornton et al. 1996). In the meantime, the 3.1 Å resolution structure of full-length Grb2 was also solved by X-ray crystallography (PDB 1GRI) (Maignan et al. 1995). In this structure Grb2 is a dimer, but the oligomeric state of Grb2 in solution remains largely controversial, just like the presence and nature of inter-domain interactions. The only detailed analysis of full-length Grb2 in solution was performed by NMR by Yuzawa et al. (2001), but with both SH2 and SH3 ligand peptides present. The behavior of apo-Grb2 in solution is therefore still poorly defined, although it is essential to decipher the interdependence of its domains and of their binding to different motifs.

Methods and experiments

Sample preparation

The gene coding for full-length Grb2 (synthesized by MWG) was inserted into a pETM11 vector also containing the gene conferring resistance to kanamycin. The construct contains a N-terminal His6 tag followed by a TEV protease cleavage site for protein purification. Grb2 plasmid was transformed into BL21 star(DE3) *E. coli* cells. A colony from the transformation was seeded into 2xYT medium, and bacteria were then successively grown at 37 °C and diluted into unlabeled M9 minimal medium, ^{15}N - ^{13}C M9, and finally ^{15}N - ^{13}C deuterated M9, for 80% deuteration of the final sample. In the M9 media, NH_4Cl concentration was 1 g/L independently of its isotopic labeling, while glucose concentration was 4 or 3 g/L for unlabeled or ^{13}C labeled glucose, respectively. Induction with isopropyl β -D-1-thiogalactopyranoside (IPTG) was performed and bacteria were harvested after overnight protein expression at 25 °C.

Cell lysis was obtained by disruption (Constant Systems Ltd.), the lysate was ultracentrifuged and the supernatant filtered before loading on a HisTrap FF crude column (GE Healthcare Life Sciences). After elution, the His6 tag was removed by TEV cleavage, leaving two additional residues G and A at the N-terminus compared to the native Grb2 sequence. The final protein product therefore had 219 residues. Excess of TEV protease, free tag and uncleaved protein was removed with a Ni-NTA resin (Thermo Scientific), and purification was completed with gel filtration on a Superdex-75 column (GE Healthcare Life Sciences). Despite deuteration during expression and given that the purification was performed in $^1\text{H}_2\text{O}$, amide groups were all

back-exchanged from 2 to ^1H before NMR measurements without need for any additional step. The final yield of [^{15}N ; ^{13}C ; U-80% ^2H] Grb2 production was about 40 mg/L culture. 200–400 μM NMR samples were prepared in PIPES buffer at pH 7.2 (40 mM PIPES, 150 mM NaCl, 2 mM TCEP).

NMR spectroscopy

Spectra were recorded at 308K with Bruker Avance NMR spectrometers of proton frequencies 600, 800 and 950 MHz equipped with TCI cryoprobes. Backbone and $^{13}\text{C}_\beta$ assignments were obtained using a common combination of 3D experiments: HNCO, HN(CA)CO, HNCA, HNCACB and HN(CO)CACB. Assignments of Grb2 isolated SH2, NSH3 and CSH3 domains were also obtained and were used to clear up a few ambiguities in the assignment of the full-length protein. The data were processed using Topspin3.5 (Bruker) and analyzed using CCPNMR (Vranken et al. 2005).

Secondary structure verification

Overall structural integrity of each domain (SH2, CSH3 and the main form of the NSH3 domain) was assessed using chemical shifts and TALOS+ (Shen et al. 2009), and compared with secondary structures in Grb2 crystal structure (PDB 1GRI).

Extent of assignments and data deposition

Assignment of backbone ^{15}N - ^1H Resonance assignment of full-length Grb2 has been obtained using a series of heteronuclear experiments. The backbone sequential assignment of the ^1H - ^{15}N HSQC spectrum is presented in Fig. 1. The unassigned residues mainly belong to the unstructured edges of the three domains, or to the linkers between them, which indicate specific dynamic features of these regions. Additionally, we could not assign the region around cysteine 32, which is located in a loop that was shown to exhibit intermediate exchange in the context of the isolated NSH3 domain by NMR (Goudreau et al. 1994) and where electron density is missing in full-length Grb2 crystal structure (PDB 1GRI). Finally, a few isolated residues could not be assigned (0A, 14D, 90S, 103N, 114G, 132V, 135H, 199H).

Overall, 173 backbone ^{15}N - ^1H resonances were assigned (84% of the residues excluding prolines and the N-terminal residue, or 79% of all residues).

Carbon assignment HNCO and HN(CA)CO spectra allowed assignment of 86% of all ^{13}C carbons, and HNCA, HNCACB and HN(CO)CACB spectra allowed assignment of




Fig. 1 Assignment of backbone ^1H - ^{15}N pairs of full-length Grb2. **a)** Assigned ^1H - ^{15}N HSQC spectrum of [U- ^{15}N ; U- ^{13}C ; U-80% ^2H]-Grb2 (308 K, PIPES buffer pH 7.2, 600 MHz), with a zoom on the central region in **(b)**. Some of the peaks believed to correspond to the unfolded form of the NSH3 domain are marked in orange. **c** Sequence

of the construct of Grb2 that we used, corresponding to Uniprot accession number P62993 with two additional residues at the N-terminus. The missing assignments are shown in red along the sequence, while grey residues are prolines and the N-terminal residue. The limits of the three domains as defined by ProRule are given

88% of all $^{13}\text{C}_\alpha$ and 86% of all $^{13}\text{C}_\beta$ resonances, yielding good coverage of the whole sequence.

Additional peaks in the spectra The isolated NSH3 domain of Grb2 has been shown to exchange between a folded and an unfolded conformation under acidic conditions (Goudreau et al. 1994), while its ortholog in *Drosophila* (Drk) has been shown to undergo a similar equilibrium in physiological buffer (Zhang et al. 1994).

In our spectra of Grb2 at neutral pH (7.2), a few tens of peaks are present that we excluded as belonging to any unassigned residue. They are mainly located in the central part of the spectrum in the proton dimension, suggesting that they are part of an unfolded conformation. Moreover, some of these peaks could be assigned to small segments consistent with parts of the NSH3 domain sequence (assignment not reported here). Our hypothesis is that these additional peaks correspond to an unfolded form of the NSH3 domain in full-length Grb2, the other two domains being only present in their fully folded form. From peak volumes, the unfolded to folded molar ratio is of the order of 1:1.

Data deposition The ^1H , ^{13}C and ^{15}N chemical shifts have been deposited in the BioMagResBank (<http://www.bmrb.wisc.edu/>) under the BMRB accession number 50082.

Acknowledgements The authors thank François Giraud, Ewen Lescop and Nelly Morellet for spectrometer maintenance and help in operation, Annie Moretto for media and buffer preparation, and Ali Badache for fruitful discussion.

Author contributions LP performed NMR experiments and analysis. YW performed preliminary NMR experiments and analysis. LP and FG wrote the initial draft of the manuscript. AV, NA, LP and YW performed protein expression and purification. CvH conceived and supervised research, and managed funding. All authors discussed results and contributed to the final manuscript.

Funding We thank the French National Research Agency (ANR) (research Grant ANR-13-BSV8-0016 to CvH and financing YW) for providing financial support to the project. Louise Pinet was funded by a PhD grant from the Université Paris-Sud. Financial support from the TGIR-RMN-THC Fr3050 CNRS for conducting the research is gratefully acknowledged. This work was supported by the French Infrastructure for Integrated Structural Biology (FRISBI) ANR-10-INBS-05.

Data availability The presented assignments were deposited to the BMRB under the entry 50082.

Compliance with ethical standards

Conflict of interest The authors declare no conflict of interest or competing interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are

included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Buday L, Downward J (1993) Epidermal growth factor regulates p21ras through the formation of a complex of receptor, Grb2 adapter protein, and Sos nucleotide exchange factor. *Cell* 73(3):611–20
- Cheng AM, Saxton TM, Sakai R, Kulkarni S, Mbamalu G, Vogel W, Tortorice CG, Cardiff RD, Cross JC, Muller WJ, Pawson T (1998) Mammalian Grb2 regulates multiple steps in embryonic development and malignant transformation. *Cell* 95(6):793–803
- Daly RJ, Binder MD, Sutherland RL (1994) Overexpression of the Grb2 gene in human breast cancer cell lines. *Oncogene* 9(9):2723–7
- Gale NW, Kaplan S, Lowenstein EJ, Schlessinger J, Bar-Sagi D (1993) Grb2 mediates the EGF-dependent activation of guanine nucleotide exchange on Ras. *Nature* 363(6424):88–92
- Goudreau N, Cornille F, Duchesne M, Parker F, Tocque B, Garbay C, Roques BP (1994) NMR structure of the N-terminal SH3 domain of GRB2 and its complex with a proline-rich peptide from Sos. *Nat Struct Biol* 1(12):898–907
- Gu H, Neel BG (2003) The “Gab” in signal transduction. *Trends Cell Biol* 13(3):122–130
- Kohda D, Terasawa H, Ichikawa S, Ogura K, Hatanaka H, Mandiyan V, Ullrich A, Schlessinger J, Inagaki F (1994) Solution structure and ligand-binding site of the carboxy-terminal SH3 domain of GRB2. *Structure* 2(11):1029–1040
- Lowenstein EJ, Daly RJ, Batzer AG, Li W, Margolis B, Lammers R, Ullrich A, Skolnik EY, Bar-Sagi D, Schlessinger J (1992) The SH2 and SH3 domain-containing protein GRB2 links receptor tyrosine kinases to ras signaling. *Cell* 70(3):431–442
- Maignan S, Guilloteau JP, Fromage N, Arnoux B, Becquart J, Ducruix A (1995) Crystal structure of the mammalian Grb2 adaptor. *Science* 268(5208):291–293
- Ravid T, Heidinger JM, Gee P, Khan EM, Goldkorn T (2004) c-Cbl-mediated ubiquitinylation is required for epidermal growth factor receptor exit from the early endosomes. *J Biol Chem* 279(35):37153–37162
- Shen Y, Delaglio F, Cornilescu G, Bax A (2009) TALOS+: a hybrid method for predicting protein backbone torsion angles from NMR chemical shifts. *J Biomol NMR* 44(4):213–23
- Thornton KH, Mueller WT, McConnell P, Zhu G, Saltiel AR, Thanabal V (1996) Nuclear magnetic resonance solution structure of the growth factor receptor-bound protein 2 Src homology 2 domain. *Biochemistry* 35(36):11852–11864
- Vranken WF, Boucher W, Stevens TJ, Fogh RH, Pajon A, Llinas M, Ulrich EL, Markley JL, Ionides J, Laue ED (2005) The CCPN data model for NMR spectroscopy: development of a software pipeline. *Proteins* 59(4):687–696
- Wittekind M, Mapelli C, Farmer BT, Suen K-L, Goldfarb V, Tsao J, Lavoie T, Barbacid M, Meyers CA, Mueller L (1994) Orientation of peptide fragments from Sos proteins bound to the N-terminal SH3 domain of Grb2 determined by NMR spectroscopy. *Biochemistry* 33(46):13531–13539
- Yamazaki T, Zaai K, Hailey D, Presley J, Lippincott-Schwartz J, Samelson LE (2002) Role of Grb2 in EGF-stimulated EGFR internalization. *J Cell Sci* 115(Pt 9):1791–802

- Yuzawa S, Yokochi M, Hatanaka H, Ogura K, Kataoka M, Miura K, Mandiyan V, Schlessinger J, Inagaki F (2001) Solution structure of Grb2 reveals extensive flexibility necessary for target recognition. *J Mol Biol* 306(3):527–37
- Zhang O, Kay LE, Olivier JP, Forman-Kay JD (1994) Backbone ¹H and ¹⁵N resonance assignments of the N-terminal SH3 domain of drk in folded and unfolded states using enhanced-sensitivity pulsed field gradient NMR techniques. *J Biomol NMR* 4(6):845–858

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.