



HAL
open science

Exosomes: A Novel Therapeutic Paradigm for Treatment of Depression

Amanda K. A. Silva

► **To cite this version:**

Amanda K. A. Silva. Exosomes: A Novel Therapeutic Paradigm for Treatment of Depression. *Current Drug Targets*, 2020, 21, 10.2174/1389450121999201006193005 . hal-03097048

HAL Id: hal-03097048

<https://hal.science/hal-03097048>

Submitted on 5 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Title**

2 **Exosomes: A Novel Therapeutic Paradigm for Treatment of Depression**

3 **Shvetank Bhatt^{1*}, Jovita Kanoujia¹, Arghya Kusum Dhar², Surendar Arumugam³,**
4 **Amanda K. A. Silva³, Neeraj Mishra¹**

5 ¹Amity Institute of Pharmacy, Amity University Madhya Pradesh (AUMP), Gwalior-474005,
6 India

7 ²Gurunanak Institute of Pharmaceutical Science and Technology, Kolkata-700110, India

8 ³Laboratoire Matière et Systèmes Complexes (MSC), Université de Paris, UMR 7057 CNRS,
9 75205 Paris cedex 13, France

10 **Principal Corresponding Author: *Dr. Shvetank Bhatt**

11 Professor

12 Amity Institute of Pharmacy,

13 Amity University Madhya Pradesh (AUMP),

14 Gwalior-474005

15 Email: shvetankbhatt@gmail.com

16 Contact No: +919414822323

17 **Abstract**

18 Extracellular vesicles (EVs) of endocytic origin are known as exosomes. These vesicles are
19 released by cells and are accessible in biofluids, such as saliva, urine, and plasma. These
20 vesicles are made up of small RNA, DNA, proteins and play a vital role in many
21 physiological processes. In central nervous system (CNS), they participate in various
22 physiological processes such as stress of nerve cells, communication between the cells,
23 synaptic plasticity and neurogenesis. The role of exosomes in depression needs to be explored
24 further. It is known that exosomes can cross blood brain barrier (BBB), which is made up of
25 glial cells astrocytes. One of the advantages of these vesicles is that they are able to transfer
26 macromolecules like DNA, protein, mRNAs and miRNAs to recipient cells. This review
27 focuses on the potential role of exosomes in depression and their utilization as
28 atreatmentoption or diagnostic tool of depression.

29 .

30 **Key words:** Depression, Exosomes, BBB, Serotonin, Biomarker, Oxidative Stress

1 **1. Introduction**

2 **Major Depressive Disorder (MDD)**

3 Depression or MDD is the most common mental disorder that affects more than 264 million
4 people worldwide [1]. It is a leading cause of disability globally. Moreover, depression also
5 increases the risk of suicide attempts [2]. Women are more susceptible to get depression as
6 compared to men [3]. There are many psychological and pharmacological treatments
7 available for depression and most of the treatment approaches work through modulation of
8 the levels of biogenic amines in the CNS [4]. However, the available therapeutic approaches
9 are limited in their role to control resistant cases of depression as well as therapeutic efficacy
10 of medications take more time to subside the symptoms of disease [5]

11 There are various psychological, environmental and biological factors involved in the
12 pathophysiology of depression [6]. Now genetic link is also found and validated. The genes
13 related to brain derived neurotrophic factor (BDNF), 5-hydroxy tryptamine transporter (5-
14 HTT), and Norepinephrine transporter (NET) have been identified [7]. Multiple theories and
15 hypotheses are coined to explain the pathogenesis of depression in that monoamine theory is
16 widely acceptable. The monoamine hypothesis relies on the abnormally excessive
17 biotransformation of major neurotransmitters (5-HT, norepinephrine and dopamine) [8]. The
18 illustration of MDD pathogenesis involves genetic predisposition, deranged monoamine
19 synthesis/function, and altered structure/function of brain. In addition to above mentioned
20 factors, depression is also associated with increased oxidative stress due to the formation of
21 reactive oxygen species (ROS) and imbalance in oxidant and antioxidant signaling [9]. Brain
22 is more susceptible to oxidative stress due to more consumption of oxygen, higher lipid
23 contents and weaker antioxidative defence. Dysregulation of hypothalamic pituitary adrenal
24 (HPA) axis also associated with pathogenic progression of depression [10]. Moreover, in
25 depression, the levels of various inflammatory markers are increased in blood and CNS area.
26 Elevated levels of C-reactive protein and cytokines such as interleukin 6 and tumour necrosis
27 factor-alpha were observed [11]. Treatment with anti-depressants suppresses the
28 inflammatory response, whereas electroconvulsive therapy acutely increases the levels of
29 various proinflammatory cytokines [12].

30 MDD also has strong relationship with cardiovascular disorders and is linked with high death
31 rates [13]. Some previous reports have demonstrated that depression increases the risk of
32 developing cardiac disease, specifically coronary artery disease, and exacerbates the

1 prognosis after myocardial infarction [13,14,15]. Anxiety and panic disorders are another
2 disorders found in comorbidity in approximately 80% of MDD cases. MDD also increases
3 the risk of drug abuse and alcohol consumption [16,17].

4 **2. Extracellular Vesicles (EVs)**

5 Since there is no effective therapy available for the treatment of depression and associated
6 comorbid disorder, innovative approaches should be investigated. In this review we will
7 discuss the role of extracellular vesicles, including exosomes as therapeutic target and
8 biomarker for the treatment and early detection of symptoms of depression. EVs are shed by
9 most of the cells that release them into the extracellular space. These vesicles have very
10 specific and important role in signaling and communication between cells. The vesicles
11 mainly categorized in three sub types 1] apoptotic bodies (500-2000 nm); 2] microvesicles
12 (50–1000 nm); and 3] exosomes (40–200 nm) [18]. They have different biological properties.
13 Apoptotic bodies are engulfed by macrophages while microvesicles contain a range of cargo
14 that is delivered to neighbouring cells. Exosomes are developed from intraluminal vesicles
15 (ILV) via inward budding process of multivesicular body (MVB). They fuse with plasma
16 membrane that release its ILV contents as EVs called exosomes into the extracellular space
17 [19,20]. Exosomes contains proteins, lipids, and nucleic acids. They are highly enriched with
18 microRNA (miRNA). Disease states leads to altered expression of miRNA in exosomes
19 making the miRNA cargo intriguing candidates for investigation [21]. EVs are lipophilic in
20 nature and easily able to cross BBB. This specific property of exosomes makes them a target
21 of choice for the treatment or early diagnosis of CNS disorders like depression [22,23,24].

22 EVs can be isolated from serum, saliva, cerebrospinal fluid, urine, breast milk, synovial fluid,
23 effusions, semen and cell cultures [25]. The isolation of EVs is done via centrifugation at
24 100,000 g and/or mechanical filters, which are specifically designed according to diameter
25 preferred EV subtype. Limited number of research studies is conducted to evaluate the role of
26 exosomes in mental disorders.

27 **2.1 Blood Brain Barrier (BBB) Penetration and Extracellular Vesicles (EVs)**

28 Effective Delivery of drugs to the brain for the treatment of central nervous system (CNS)
29 disorders for examples neurodegenerative diseases, tumors, trauma, stroke, autoimmune
30 diseases is a major challenge [27, 28, 29, 30]. BBB is a border that limits and regulates the
31 passage of substances between the peripheral vascular circulation and the CNS, thereby

1 providing protection to the CNS from toxic substances or overactive immune responses
2 [28,31]. However the BBB permits transmembrane diffusion of lipid soluble molecules of
3 size less than 400 Da and selective transport of some compounds into and out of the brain
4 [32].

5 Two plausible transport mechanisms across the BBB were proposed namely transcellular
6 through BMECs, and paracellular through junctions between BMECs. The main components
7 of BBB are microvascular endothelial cells (BMECs), astrocytes, pericytes, the endothelial
8 basement membrane, and neighbouring neurons. A complex arrangement of tight junctions
9 (TJs) and adherens junctions (AJs) in the brain endothelial cells regulates paracellular
10 permeability [33].

11 . Around 98% of small molecule drugs and almost all of large molecule biologic drugs,
12 including recombinant proteins, monoclonal antibodies, or gene-based medicines are unable
13 to cross the BBB [34,35]. Exosomes with its contents are capable of crossing the BBB. This
14 finding has promoted biomarker research with exosomes and their use as a drug delivery
15 system. Numerous studies have demonstrated the successful brain delivery of exosomes.
16 Effective delivery of GAPDH siRNA to the brain via systemic injection of exosomes in mice
17 [36] was reported. Other experiments have been proved to be successful in delivering
18 exosomes to the mice brain via intranasal injection [37]. In another experiment using rats, a
19 fluorescently tagged protein selectively expressed in rat brain tissue was recovered in small
20 EVs having same characteristics as exosomes in their blood [38]. These studies support the
21 fact that exosomes are able to communicate from the brain to the rest of the body and they
22 can cross the BBB in a bi-directional manner. However, their exact mechanism of BBB
23 penetration remains unclear.

24 In an experiment using in vitro BBB model the transfer of EVs derived from human
25 erythrocytes was found to be dependent on the adsorptive-mediated transcytosis method of
26 transport [39]. Though the EVs crossed under healthy and inflammatory conditions, the
27 peripheral administration of lipopolysaccharides was found to considerably increase the EVs
28 movement across the BBB [39].

29 Another study revealed that in healthy and stroke-like condition exosomes crossed a BBB
30 model using transcellular BMEC endocytosis, suggesting that exosomes maintain their ability
31 to cross during stressful conditions [40]. This same study also indicated that exosomes were
32 internalized through endocytosis, and accumulate in endosomes. There was a decrease in

1 exosome transcellular migration using chlorpromazine (CPZ) an inhibitor for clathrin-
2 dependent endocytosis, which transfers clathrin from the surface of cells to intracellular
3 endosomes [40,41]. This indicates that clathrin-dependent endocytosis may be involved in
4 transportation of exosomes across the BBB [40].

5 Studies have indicated that methyl- β -cyclodextrin (M β CD), which removes cholesterol from
6 the plasma membrane [41], and filipin III, which binds to cholesterol caused a significant
7 reduction in BBB penetration of exosomes [40]. This result revealed caveolae-dependent
8 endocytosis as alternative route of exosome migration. Most likely the uptake of exosomes
9 in BMEC depends on specific ligand receptors or lipid rafts, and mechanisms of exosome
10 uptake may also depend on the cell for which they originate. It has been postulated that
11 exosomes from different cell origins have different content including protein and lipids,
12 possibly modifying their mechanism of BBB penetration [42]. Moreover, disease condition
13 may influence the mechanism of BBB crossing, as the contents of exosomes change upon
14 disease state [26, 42].

15 Studies have also proposed that exosomes play a role in increased permeability of vascular
16 barriers of the BBB. Breast cancer cells secreted exosomes uniquely expressed miR-105,
17 which directly targeted the tight junction protein ZO-1 [43]. This exosome transfer of miR-
18 105 destroyed tight junctions and the integrity of the BBB [43]. Moreover, claudin-5 (Cldn5)
19 a tight junction protein of the BBB was found to be encapsulated in exosomes [44]. A
20 loosened BBB was found in the Cldn5 knockout mice [45], suggested that exosomes carrying
21 Cldn5 might play a role in BBB integrity. It was also found that a decline in Cldn5 induced
22 depressive-like behaviors in mice, and antidepressant treatment increased Cldn5 levels and
23 promoted recovery [46]. A leaky BBB has been linked to neuroinflammation [47,48]. Thus,
24 the possible influence of exosomes on the BBB integrity also advocate a role for exosomes in
25 neuroinflammation and the pathogenesis of mental disorders [47,48] Hence, in mental
26 disorders a leaky BBB state may be caused by exosomes released from cells being influenced
27 by this disease state.

28 **2.2 Signalling mechanisms involving exosomes in the brain**

29 An important role is played by exosomes in cell to cell communication in the CNS..They are
30 involved in signalling through distal as well as neighbouring cells [49]. The exosomes work
31 as an important link for communication with similar kind of cells or also with different types

1 of cells. The release of exosomes in the CNS is controlled by glutaminergic neurons and
2 influx of Calcium ions [50,51].

3 N-methyl-D-aspartate (NMDA) is a glutaminergic receptor. The entry of calcium ion through
4 gates of this receptor triggers the neuronal release of exosomes. Other important
5 glutaminergic receptor which is involved in physiological role of exosomes is and α -amino-3-
6 hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptors [50]. After releasing,
7 exosomes can fuse with cell membrane of the recipient cell and release its constituents inside
8 the cytoplasm of the cell, transfer lipid and protein content on the cell surface. Internalisation
9 of exosomes can also take place by macropinocytosis or endocytosis process. The fusion of
10 exosomes is regulated by receptor mediated mechanisms. These diverse mechanisms for
11 transfer of constituents can initiate different signaling mechanisms in recipient cells [52].

12 Serotonin or 5-hydroxy tryptamine (5-HT) has also substantial role on the release of
13 exosomes from the non-neuronal brain cells such as microglial cells. 5-HT is a major
14 neurotransmitter which has important role in the pathophysiology of depression. According to
15 monoamine theory level of 5-HT is decreased in depression and bipolar disorders [53].
16 Serotonin can increase cytosolic levels of calcium ions, which leads to stimulation of release
17 of exosomes from primary microglia [51]. Since 5-HT is involved in release of exosomes,
18 their release is modified predominantly in case of depression. Both the factors discussed
19 above i.e. cell to cell communication and neurotransmitter mediated release of exosomes are
20 important aspects of psychopharmacology and involved in various CNS disorders including
21 depression [54]. So exosomes can be considered to play a major role in etiopathogenesis of
22 mental disorders given their prominence in the regulation of cell communication, and their
23 regulation via neurotransmitters. In addition, signaling through one nerve cell to other nerve
24 cell is important in synaptic plasticity [55].

25 **3. Role of Exosomes in Depression**

26 Depression is an important psychiatric disorder which is associated with disability and affects
27 person daily life. Diagnostic and statistical manual of mental disorders, 4th edition (DSM-IV)
28 has given 9 symptoms of depression out of which if 5 symptoms are present for more than
29 two weeks then patient can be considered as depressed. However, these have certain
30 limitations and more evaluation is required to confirm the depression cases. Multiple
31 evidence gathered from previous research suggested that estimation of various biomarkers
32 could be a useful approach for early detection of depression. Limited number of research

1 studies is conducted to evaluate the role of exosomes in mental disorders [56,57,58)
2 Particular, Banigan et al. investigated exosomes from frozen postmortem prefrontal cortex
3 to study alterations in miRNA in patients of psychoses and bipolar disorder. They found that
4 miR-497 in psychoses patients and miR-29c in bipolar patients to be upregulated compared to
5 control subjects [26]. This early piece work opens up interesting opportunities to study
6 exosomes in mental disorders, demonstrating that miRNA cargo may be interesting to be
7 evaluated in these phenotypes. The intercellular signalling of exosomes has important role in
8 normal physiological and pathological conditions in the CNS. For example glial cell
9 oligodendrocyte-derived exosomes are thought to be involved in neuroprotection against
10 cellular stress. The reason of this protective effect of exosomes is due to involvement of
11 proteins released from exosomes inside the recipient cell [56,59,60].

12 Recent data on signaling of exosomes in the CNS highlighted their role in transcriptional
13 regulation, neurogenesis, plasticity, and neuroinflammation [60]. Exosomes delivered
14 intravenously can be demonstrated to cross the BBB naturally. Exosomes from mesenchymal
15 stem cells (MSC's) demonstrate anti-inflammatory and pro-growth effects in preclinical
16 models and clinical cases reports and have been used intravenously and with intracerebral
17 and intrathecal injection..

18 Among various biomarkers microRNAs (miRNAs) emerged as an important diagnostic
19 marker that could be beneficial for patients to detect depression. These biomarkers have
20 predominant role in pathophysiology of depression [61]. These molecules affect various
21 molecular levels signaling pathways involved in progression of depression. miRNAs could be
22 used as diagnostic and therapeutic biomarkers in MDD patients. Besides miRNAs, exosomes
23 as nano- carriers could have been emerged as diagnostic biomarkers in various diseases such
24 as MDD and other CNS disorders. Intravenous delivery of exosomes can able to cross the
25 BBB [62]. Exosomes and MSC's demonstrate anti-inflammatory and pro-growth effects in
26 preclinical models and clinical cases.

27 The exosomes released from mast cells have protective effect against increased oxidative
28 stress. They have capacity to communicate a protective signal to other cells which are
29 exposed to oxidative stress and this leads to reduction in cell death [63]. Exosomes that are
30 produced by one cell under stress condition is able to induce capacity to tolerate increased
31 oxidative stress in another cell that is known as recipient cell. This effect is linked with
32 changed exosomal mRNA content that can be attenuated by reduced RNA activity through

1 exposure if UV light [64,65]. EVs and ROS are closely linked with oxidative stress. The pro-
2 oxidant conditions are involved in release of EVs in fact, NADPH oxidase and nitric oxide
3 synthase-2 (NOS-2) inhibitors inhibit the production of EVs in neutrophils [66,67].
4 According to a study conducted by Wei ZX 2020 Exosomes obtained from patients with
5 MDD caused depressive-like behaviors in mice with involvement of micro RNA regulated
6 neurogenesis.

7 Intravenous injection of Exosomes derived from blood of MDD patients into normal mice
8 exhibited the depressive phenotype behaviors as evaluated by the battery of behavioral tests
9 such as forced swim test, tail suspension test, and novelty suppressed feeding tests, and these
10 behaviours reversed by administration of exosomes derived from healthy subjects into
11 chronic unpredictable mild stress (CUMS)-treated mice. [68]. In general, secretion of
12 exosomes is increased in response to stress or pathological conditions [69].

13 .Various recent studies have observed changes in miRNAs in depressed patients, which target
14 not only important pathways associated with memory function, synaptic plasticity but also
15 formation of neurotrophic factors and modulation of immune cells[70,71]. Transfer of EVs
16 or exosomes to nerve cells occurs through supporting cells of nervous system such as
17 oligodendrocytes, microglia and astrocytes. Besides their potential as biomarkers, these
18 exosomes have a predominant role in pathophysiology and dissemination of inflammatory
19 pathways [72]. Reactive microglia were shown to release exosomes and microvesicles (MVs)
20 carrying the pro-inflammatory cytokine interleukin-1 β (IL-1 β), the IL-1 β processing enzyme
21 caspase-1, and the P2X7 receptor that may induce and propagate inflammatory reactions
22 throughout the brain [73]. Additionally, monocytes that are activated by interferon alpha
23 and/or lipopolysaccharides (LPS) release exosomes that carry altered miRNA profiles as
24 shown in Fig. 1.

25 A study data published in the *Journal of Affective Disorders* indicates a link between higher
26 levels of certain neuron-derived exosomes (NDE) and MDD [74]. In addition, a clinical study
27 has been started to evaluate the safety and efficacy of exosomes against refractory depression
28 [75]. Impaired adult hippocampal neurogenesis (AHN) is associated with depression and
29 other neurodegenerative disorders like schizophrenia [76]. Protein analysis study reveals that
30 exosomes has important role in modulation of adult neurogenesis [76]. In addition, injection
31 of known pathogen containing exosomes into the dentate gyrus is sufficient to impair AHN in
32 mice [77]. Cytokines and corticosteroids stimulate the release of glial cell astrocyte related

1 exosomes which contains several miRNAs that are critical for neurogenesis, stress response
2 and survival of neurons [76]. So the exosomes may have dual role as it may be involved in
3 maintenance and inhibition of neurogenesis in adult. Nerve cell to supporting glial cell
4 signalling is one of the important mechanisms for exosomes [78]. Injection of exosomes in
5 the tail vein of immune-challenged mice leads to increase in CNS expression of pro-
6 inflammatory cytokine mRNA and associated miRNA in mice. Protein analysis of exosomes
7 in the CNS reveals cargo involved in modulating adult neurogenesis [79].

8 **4. Exosomes as Biomarker in CNS diseases and depression**

9 Exosomes have immense capability as diagnostic tool or biomarker for early diagnosis of
10 CNS disorders such as Depression and Schizophrenia. Pathophysiological changes in
11 contents of exosomes have been observed in CNS disorders. The exosomes can be isolated
12 without much difficulty from biological fluids including urine, saliva and blood [80].
13 Moreover exosomes contents are not easily degradable by biological fluids or enzyme in our
14 body as they are protected inside the membrane [54]. In addition, *ex vivo*, these exosomes are
15 stable for longer duration can be stored long time before analysis. It is possible to trace the
16 surface marker indicating the origin for exosomes. The exosomes have great potential to
17 cross BBB and this property makes them a good biomarker to identify a CNS disorder [81].
18 Of particular interest is the ability to characterize exosomes based on their cell of origin,
19 potentially providing an extra layer of insight into the disease of interest [82]. The
20 significance of exosomes as diagnostic tool in disease like cancer is already established.
21 However their role in CNS disorders like depression is yet to be established. Many of the
22 circulating proteins and nucleic acids are diluted in the blood stream and the majority of them
23 originate from other sources besides neurons due to the tight regulation of BBB in molecule
24 transport.

25 Different types of exosomes (from multiple cells) can be segregated from biofluids using
26 different segregation techniques like ultracentrifugation, immunomagnetic beads, and
27 chromatography [83,84]. In addition, treatment with RNase, western blot analysis or mass
28 spectrometry is also used for identification/segregation of exosomes contents. Various types
29 of exosomes can be isolated from biofluids using multiple methods. Additionally, exosomes
30 have phospholipid bilayer; therefore treatment with RNase prior to use will ensure that cargo
31 used downstream was encapsulated within the vesicle [85]. Various types of exosomes may
32 be identified using western blots or mass spectrometry of proteins which are involved in

1 biogenesis of ILVs, including tetraspanins and proteins involved in the ESCRT machinery
2 needed for biogenesis [86]. It is essential to remember that these markers are not exclusively
3 associated with exosomes and further characterizations of these markers are required..

4 Exosomes derived from developing and mature neurons of hippocampus, contain L1 cell
5 adhesion molecule, and the GluR2/3 subunits of glutamate receptors, both of which are well
6 established markers of nerve cells [87,88]. Protein markers, such as glial fibrillary acidic
7 protein), glutamine aspartate transporter, and glutamine synthetase, can be used to enrich for
8 astrocytic-derived exosomes. Additionally, myelin proteo-lipid protein and 2', 3'-cyclic
9 nucleotide 3'-phosphodiesterase have been identified on oligodendrocytes derived from
10 exosomes [89]. Neural-derived exosomes from plasma have also been used in a small study
11 to evaluate protein biomarkers for patients with MDD [71]. The expression levels of
12 exosomal miRNAs vary among different biological fluids in vivo and culture medium in
13 vitro. So these changes in expression of exosomes in MDD are used to identify the early
14 stage of disorder.

15 **5. Authors insight on the topic**

16 Utilization of EVs as drug carrier has been well explored in chronic disorders such as cancer
17 and autoimmune disorders. In the future, drug loaded EVs could be explored to achieve
18 maximum therapeutic benefit sparing normal healthy cells, with less off target mediated
19 adverse effects of antidepressants with aim to target the selective affected region owing to
20 MDD induced alteration in the CNS. Disorders like depression have lack of suitable
21 diagnostic technique and treatment resistance is also more common. So, EVs maybe serve as
22 as a potential therapeutic candidate as well as biomarker, which may be effective in the field
23 of depression. For that, further studies have to be performed in order to understand the
24 homing of EVs from different cell types in MDD. In MDD, it has been documented that in
25 addition to abnormal neurotransmitter biogenesis and neurotransmitter shuttling, involvement
26 of neuronal damage is highly prevalent condition during later stage of pathophysiological
27 mechanism to disease progression. Neurogenesis capability of currently available
28 antidepressants alone or in combination are being explored and evaluated in preclinical and
29 clinical experiments. However, regenerative potential of these agents is not substantial to
30 protect the vigorous constant neurodegenerative process during later stage of MDD and
31 associated conditions. As it has been witnessed that EVs have regenerative ability and this
32 exclusive property of EVs can be exploited to prevent the ongoing neurodegenerative process

1 of MDD. Cell therapy-based treatment especially induced pluripotent stem cells exhibit
2 therapeutic activity in MDD and associated pathological conditions, however its usage for
3 these conditions have been limited by its adverse effects such as graft rejection by recipient,
4 immunogenicity reactions and low accessible and penetration to BBB. There are various
5 approaches being explored to mitigate these adverse effects caused by cell therapy. In this
6 context, there are advantages of EVs compared to cells such as reduced size, less
7 immunogenicity, easier accessibility to BBB along with carrying some essential components
8 of cell while mimicking benefits elicited by cell therapy of MDD. This lays foundation for
9 cell-free therapy for MDD associated pathological condition. Despite the expected EV
10 benefits for treating MDD, there are still unexplored uncertainties in terms of precise
11 pharmacological mechanism of action, route of administration.

12 Selectivity and accumulation of EV to intended target site of action will be hampering the
13 effort of EV as therapeutic potential in treating MDD. Though selective binding to its target
14 can be achieved by producing EV which has selective binding surface expression molecules
15 on its surface from genetically modified parental cells expressing the target selection marker.
16 There are plenty of invitro cell-based assays need to be performed to achieve the tissue and
17 target selective EV to maximize its therapeutic benefit sans adverse effect on healthy brain
18 tissues. Another obstacle to developing EV for MDD would be owing to plethora of
19 mechanism of action elicited by EV, narrow down a precise pharmacological molecular
20 mechanism action can be difficult and challenging task. Nevertheless, the MDD disease
21 pathway related molecular markers could be selected to elucidate the prime mode of
22 mechanism action of EVs are warranted using various molecule biology tools (reporter,
23 knock out or knock in based approaches. To address these issues, initiation of well-planned
24 preclinical studies including high translational value based in vitro (Target binding,
25 specificity, selectivity and internalization assays), pharmacokinetic assays (Tissue
26 distribution studies with special emphasis on central nervous system using appropriate
27 labelling agents) in vivo models of MDD, toxicology and safety pharmacology studies on
28 vital systems should be warranted. In conclusion, owing to better advantage of EV compared
29 to conventional treatment, future research EVs based therapy is expected to provide evidence
30 on their potential as standalone therapy or in combination with standard antidepressants in the
31 management of this devastating disease.

32

6. Conclusion

In the future, EVs may be useful as effective therapeutic carriers as well as biomarkers for the diagnosis of depression in early stages. However, further studies need to be done to prove the efficacy and pharmacokinetic profile of these exosomes. The exosomes have an advantage over conventional treatment to be able to cross the BBB. In depression modulation in levels of exosomes is taking place and this can be used as marker for detection of disorder in early stages. Chronic stress and inflammation also lead to modify the expression of exosomes. Exosomes are important in communication and signalling between two cells. These EVs also have role in synaptic plasticity and neurogenesis. However, some more detailed studies are required to validate the role of exosomes in MDD. In the future, exosomes may be used as effective treatment and biomarker for identification of depression and other CNS disorders.

Conflict of Interest: There is no conflict of interest among the authors.

Acknowledgement: None

Financial Assistance: Nil

References

1. <https://www.who.int/news-room/fact-sheets/detail/depression>
2. Brådvik, L. Suicide Risk and Mental Disorders. *International journal of environmental research and public health*, 2018, 15(9), 2028.
3. Albert, P. R. Why is depression more prevalent in women? *Journal of psychiatry & neuroscience*, 2015, 40(4), 219–221.
4. Heninger ,G.R.; Delgado, P.L.; Charney, D.S. The revised monoamine theory of depression: a modulatory role for monoamines, based on new findings from monoamine depletion experiments in humans. *Pharmacopsychiatry*, 1996, 29(1), 2-11.
5. Al-Harbi, K. S. Treatment-resistant depression: therapeutic trends, challenges, and future directions. *Patient preference and adherence*, 2012, 6, 369–388.
6. Hasler, G. Pathophysiology of depression: do we have any solid evidence of interest to clinicians? *World psychiatry*, 2010, 9(3), 155–161.
7. Tamatam, A.; Khanum, F.; Bawa, A. S. Genetic biomarkers of depression. *Indian journal of human genetics*, 2012, 18(1), 20–33.

- 1 8. Liu, B.; Liu, J.; Wang, M.; Zhang, Y.; Li, L. From Serotonin to Neuroplasticity:
2 Evolvement of Theories for Major Depressive Disorder. *Frontiers in cellular*
3 *neuroscience*, 2017, *11*, 305.
- 4 9. Bhatt, S.; Radhakrishnan, M.; Jindal, A.; Devadoss, T.; Dhar, A. K.
5 Neuropharmacological evaluation of a novel 5-HT₃ receptor antagonist (6g) on
6 chronic unpredictable mild stress-induced changes in behavioural and brain oxidative
7 stress parameters in mice. *Indian journal of pharmacology*, 2014, *46*(2), 191–196.
- 8 10. Keller, J.; Gomez, R.; Williams, G.; Lembke, A.; Lazzeroni, L.; Murphy, G. M. Jr.;
9 Schatzberg, A. F. HPA axis in major depression: cortisol, clinical symptomatology
10 and genetic variation predict cognition. *Molecular psychiatry*, 2017, *22*(4), 527–536.
- 11 11. Lee, C. H.; Giuliani, F. The Role of Inflammation in Depression and
12 Fatigue. *Frontiers in immunology*, 2019, *10*, 1696.
- 13 12. Dinan, T.G. Inflammatory markers in depression. *Curr Opin Psychiatry*, 2009, *22*(1),
14 32-6.
- 15 13. Dhar, A. K.; Barton, D. A. Depression and the link with cardiovascular
16 disease. *Frontiers in psychiatry*, 2016, *7*, 33.
- 17 14. Mastrogiannis, D.; Giamouzis, G.; Dardiotis, E.; Karayannis, G.; Chroub-Papavaïou,
18 A.; Kremeti, D.; Spiliopoulos, K.; Georgoulas, P.; Koutsias, S.; Bonotis, K.;
19 Mantzorou, M.; Skoularigis, J.; Hadjigeorgiou, G. M.; Butler, J.; & Triposkiadis, F.
20 (2012). Depression in patients with cardiovascular disease. *Cardiology research and*
21 *practice*, 2012, 794762.
- 22 15. Strik, J.J.; Honig, A.; Maes M. Depression and myocardial infarction: relationship
23 between heart and mind. *Prog Neuropsychopharmacol Biol Psychiatry*, 2001, *25*(4),
24 879-92.
- 25 16. Burns, L.; Teesson, M.; O'Neill, K. The impact of comorbid anxiety and depression
26 on alcohol treatment outcomes. *Addiction*. 2005, *100*(6), 787-96.
- 27 17. Hirschfeld, R. M. The Comorbidity of Major Depression and Anxiety Disorders:
28 Recognition and Management in Primary Care. *Primary care companion to the*
29 *Journal of clinical psychiatry*, 2001, *3*(6), 244–254.
- 30 18. Gheinani, A.H.; Vögeli, M.; Baumgartner, U.; Vassella, E.; Draeger, A.; Burkhard,
31 F.C.; Monastyrskaya, K. Improved isolation strategies to increase the yield and purity
32 of human urinary exosomes for biomarker discovery. *Scientific Reports*, 2018,
33 8(1),3945.
- 34 19. Lee, Y.; El Andaloussi, S.; Wood, M. J. A. Exosomes and microvesicles: extracellular
35 vesicles for genetic information transfer and gene therapy. *Human Molecular*
36 *Genetics* 2012 **21**(R1), R125–R134.
- 37 20. Johnstone, R. M.; Adam, M.; Hammond, J. R.; Orr, L.; Turbide, C. Vesicle formation
38 during reticulocyte maturation. Association of plasma membrane activities with
39 released vesicles (exosomes). *Journal of Biological Chemistry*, 1987, *262*, 9412–
40 9420.
- 41 21. Gallo, A.; Tandon, M.; Alevizos, I.; & Illei, G. G. The majority of microRNAs
42 detectable in serum and saliva is concentrated in exosomes. *PLoS ONE*, 2012, *7*,
43 e30679.

- 1 22. Chen, C.C.; Liu, L.; Ma, F.; Wong, C.W.; Guo, X.E.; Chacko, J.V.; Farhoodi, H.P.;
2 Zhang, S.X.; Zimak, J.; Ségaliny, A.; Riazifar, M.; Pham, V.; Digman, M.A.; Pone,
3 E.J.; Zhao, W. Elucidation of Exosome Migration across the Blood-Brain Barrier
4 Model In Vitro. *Cell Mol Bioeng*, 2016, 9(4), 509-529.
- 5 23. Cufaro, M. C.; Pieragostino, D.; Lanuti, P.; Rossi, C.; Cicalini, I.; Federici, L.; De
6 Laurenzi, V.; Del Boccio, P. Extracellular Vesicles and Their Potential Use in
7 Monitoring Cancer Progression and Therapy: The Contribution of
8 Proteomics. *Journal of oncology*, 2019, 2019, 1639854.
- 9 24. Yao, Z. Y., Chen, W. B., Shao, S. S., Ma, S. Z., Yang, C. B., Li, M. Z., Zhao, J. J.,
10 Gao, L.. Role of exosome-associated microRNA in diagnostic and therapeutic
11 applications to metabolic disorders. *Journal of Zhejiang University. Science*.
12 2018, 19(3), 183–198.
- 13 25. Zebrowska, A.; Skowronek, A.; Wojakowska, A.; Widlak, P.; Pietrowska, M. (2019).
14 Metabolome of Exosomes: Focus on Vesicles Released by Cancer Cells and Present
15 in Human Body Fluids. *International journal of molecular sciences*, 2019, 20(14),
16 3461.
- 17 26. Banigan, M. G.; Kao, P. F.; Kozubek, J. A.; Winslow, A. R.; Medina, J.; Costa, J.;
18 Schmitt, A.; Schneider, A.; Cabral, H.; Cagsal-Getkin, O.; Vanderburg, C. R.;
19 Delalle, I. Differential expression of exosomal microRNAs in prefrontal cortices of
20 schizophrenia and bipolar disorder patients. *PloS one*, 2013, 8(1), e48814.
- 21 27. Abbott, N.J.; Rönnbäck, L.; Hansson, E. Astrocyte-endothelial interactions at the
22 blood-brain barrier. *Nature Reviews Neuroscience*, 2006, 7(1), 41-53.
- 23 28. Obermeier, B.; Daneman, R.; Ransohoff, R.M. Development, maintenance and
24 disruption of the blood-brain barrier. *Nature Medicine*, 2013, 19(12), 1584-96.
- 25 29. Rubin, L.L.; Staddon, J.M. The cell biology of the blood-brain barrier. *Annual Review*
26 *of Neuroscience*, 1999, 22, 11–28.
- 27 30. Upadhyay, R.K. Drug delivery systems, CNS protection, and the blood brain barrier.
28 *BioMed Research International*, 2014, 2014, 869269.
- 29 31. Andreone, B.J.; Lacoste, B.; Gu, C. Neuronal and vascular interactions. *Annual*
30 *Review of Neuroscience*. 2015, 38, 25–46.
- 31 32. Sanchez-Covarrubias, L.; Slosky, L. M.; Thompson, B. J.; Davis, T. P.; Ronaldson, P.
32 T. Transporters at CNS barrier sites: obstacles or opportunities for drug delivery?
33 *Current Pharmaceutical Design*. 2014, 20(10), 1422-49.
- 34 33. Wolburg, H.; Lippoldt, A. Tight junctions of the blood-brain barrier: development,
35 composition and regulation. *Vascular Pharmacology*. 2002, 38(6), 323-37.
- 36 34. Pardridge, W.M. Drug transport across the blood-brain barrier. *Journal of Cerebral*
37 *Blood Flow & Metabolism*. 2012, 32(11), 1959-72.
- 38 35. Pardridge, W.M. The blood-brain barrier: bottleneck in brain drug development.
39 *NeuroRx*. 2005, 2(1), 3-14.
- 40 36. Alvarez-Erviti, L.; Seow, Y.; Yin, H.; Betts, C.; Lakhali, S.; Wood, M.J. Delivery of
41 siRNA to the mouse brain by systemic injection of targeted exosomes. *Nature*
42 *Biotechnology*. 2011, 29(4), 341-5.
- 43 37. Zhuang, X.; Xiang, X.; Grizzle, W.; Sun, D.; Zhang, S.; Axtell, R.C.; Ju, S.; Mu, J.;
44 Zhang, L.; Steinman, L.; Miller, D.; Zhang, H.G. Treatment of brain inflammatory

- 1 diseases by delivering exosome encapsulated anti-inflammatory drugs from the nasal
2 region to the brain. *Molecular Therapy*. 2011, 19(10), 1769-79.
- 3 38. Gómez-Molina, C.; Sandoval, M.; Henzi, R.; Ramírez, J.P.; Varas-Godoy, M.; Luarte,
4 A.; Lafourcade, C.A.; Lopez-Verrilli, A.; Smalla, K.H.; Kaehne, T.; Wyneken, U.
5 Small Extracellular Vesicles in Rat Serum Contain Astrocyte-Derived Protein
6 Biomarkers of Repetitive Stress. *International Journal of*
7 *Neuropsychopharmacology*. 2019, 22(3), 232-246.
- 8 39. Matsumoto, J.; Stewart, T.; Sheng, L.; Li, N.; Bullock, K.; Song, N.; Shi, M.; Banks,
9 W.A.; Zhang, J. Transmission of α -synuclein-containing erythrocyte-derived
10 extracellular vesicles across the blood-brain barrier via adsorptive mediated
11 transcytosis: another mechanism for initiation and progression of Parkinson's disease?
12 *Acta Neuropathologica Communications*. 2017, 5(1), 71.
- 13 40. Chen, C.C.; Liu, L.; Ma, F.; Wong, C.W.; Guo, X.E.; Chacko, J.V.; Farhoodi, H.P.;
14 Zhang, S.X.; Zimak, J.; Ségaliny, A.; Riazifar, M.; Pham, V.; Digman, M.A.; Pone,
15 E.J.; Zhao, W. Elucidation of Exosome Migration across the Blood-Brain Barrier
16 Model In Vitro. *Cellular and Molecular Bioengineering*. 2016, 9(4), 509-529.
- 17 41. Dutta, D.; Donaldson, J.G. Search for inhibitors of endocytosis: Intended specificity
18 and unintended consequences. *Cellular Logistics*. 2012, 2(4), 203-208.
- 19 42. Haraszti, R.A.; Didiot MC, Sapp E, Leszyk J, Shaffer SA, Rockwell HE, Gao F,
20 Narain NR, DiFiglia M, Kiebish MA, Aronin N, Khvorova A. High-resolution
21 proteomic and lipidomic analysis of exosomes and microvesicles from different cell
22 sources. *Journal Extracellular Vesicles*. 2016, 17(5), 32570.
- 23 43. Zhou, W.; Fong, MY.; Min, Y.; Somlo, G.; Liu, L.; Palomares, M.R.; Yu, Y.; Chow,
24 A.; O'Connor, S.T.; Chin, A.R.; Yen, Y.; Wang, Y.; Marcusson, E.G.; Chu, P.; Wu,
25 J.; Wu, X.; Li, A.X.; Li, Z.; Gao, H.; Ren, X.; Boldin, M.P.; Lin, P.C.; Wang, S.E.
26 Cancer-secreted miR-105 destroys vascular endothelial barriers to promote
27 metastasis. *Cancer Cell*. 2014, 25(4), 501-15.
- 28 44. Paul, D.; Baena, V.; Ge, S.; Jiang, X.; Jellison, E.R.; Kiprono, T.; Agalliu, D.;
29 Pachter, J.S. Appearance of claudin-5(+) leukocytes in the central nervous system
30 during neuroinflammation: a novel role for endothelial-derived extracellular vesicles.
31 *Journal of Neuroinflammation*. 2016, 13(1), 292.
- 32 45. Nitta, T.; Hata, M.; Gotoh, S.; Seo, Y.; Sasaki, H.; Hashimoto, N.; Furuse, M.;
33 Tsukita, S. Size-selective loosening of the blood-brain barrier in claudin-5-deficient
34 mice. *Journal of Cell Biology*. 2003,161(3), 653-60.
- 35 46. Menard, C.; Pfau, M.L.; Hodes, G.E.; Kana, V.; Wang, V.X.; Bouchard, S.;
36 Takahashi, A.; Flanigan, M.E.; Aleyasin, H.; LeClair, K.B.; Janssen, W.G.; Labonté,
37 B.; Parise, E.M.; Lorsch, Z.S.; Golden, S.A.; Heshmati, M.; Tamminga, C.; Turecki,
38 G.; Campbell, M.; Fayad, Z.A.; Tang, C.Y.; Merad, M.; Russo, S.J. Social stress
39 induces neurovascular pathology promoting depression. *Nature Neuroscience*. 2017,
40 20(12), 1752-1760.
- 41 47. Maes, M. The cytokine hypothesis of depression: inflammation, oxidative &
42 nitrosative stress (IO&NS) and leaky gut as new targets for adjunctive treatments in
43 depression. *Neuro Endocrinology Letters*. 2008, 29(3), 287-91.

- 1 48. Najjar, S.; Pearlman, D.M.; Alper, K.; Najjar, A.; Devinsky, O. *Neuroinflammation*
2 *and psychiatric illness. Journal of Neuroinflammation*. 2013, 10, 43
- 3 49. Zhang, G.; Yang, P. A novel cell- cell communication mechanism in the nervous
4 system: exosomes. *Journal of Neuroscience Research* 2018, 96, 45–52.
- 5 50. Lachenal, G.; Pernet-Gallay, K.; Chivet, M.; Hemming, F.J.; Belly, A.; Bodon, G.;
6 Blot, B.; Haase, G.; Goldberg, Y.; Sadoul, R. Release of exosomes from differentiated
7 neurons and its regulation by synaptic glutamatergic activity. *Molecular and Cellular*
8 *Neurosciences*, 2011, 46(2), 409-18.
- 9 51. Glebov, K.; Löchner, M.; Jabs, R.; Lau, T.; Merkel, O.; Schloss, P.; Steinhäuser, C.;
10 Walter, J. Serotonin stimulates secretion of exosomes from microglia cells. *Glia*,
11 2015; 63(4):626-34.
- 12 52. Sharma, P.; Schiapparelli, L.; Cline, H.T.; Exosomes function in cell-cell
13 communication during brain circuit development. *Curr Opin Neurobiol*,
14 2013,23(6),997–1004.
- 15 53. Cowen, P. J.; Browning, M. What has serotonin to do with depression?. *World*
16 *psychiatry: official journal of the World Psychiatric Association (WPA)*, 2015, 14(2),
17 158–160.
- 18 54. Kanninen, K.M.; Bister, N.; Koistinaho, J.; Malm, T. Exosomes as new diagnostic
19 tools in CNS diseases. *Biochimica et Biophysica Acta*, 2016, 1862(3), 403-10.
- 20 55. Chivet, M.; Hemming, F.; Fraboulet, S.; Sadoul, R. Emerging role of neuronal
21 exosomes in the central nervous system. *Frontiers in Physiology*, 2012 3, 145.
- 22 56. Saeedi, S.; Israel, S.; Nagy, C.; Turecki, G. The emerging role of exosomes in mental
23 disorders. *Translational Psychiatry*, 2019, 9, 122.
- 24 57. Brites, D.; Fernandes, A. Neuroinflammation and Depression: Microglia Activation,
25 Extracellular Microvesicles and microRNA Dysregulation. *Frontiers in cellular*
26 *neuroscience*. 2015, 9, 476.
- 27 58. Gruzdev, S.K.; Yakovlev, A.A.; Druzhkova, T.A.. The Missing Link: How Exosomes
28 and miRNAs can Help in Bridging Psychiatry and Molecular Biology in the Context
29 of Depression, Bipolar Disorder and Schizophrenia. *Cell Mol Neurobiol*, 2019, 39,
30 729–750.
- 31 59. Frühbeis, C.; Fröhlich, D.; Kuo, W. P.; Amphornrat, J.; Thilemann, S.; Saab, A. S..
32 Neurotransmitter-triggered transfer of exosomes mediates oligodendrocyte-neuron
33 communication. *PLoS Biology*, 2013, 11, e1001604.
- 34 60. Krämer-Albers, E.M.; Bretz, N.; Tenzer, S.; Winterstein, C.; Möbius, W.; Berger, H.;
35 Nave, K.A.; Schild, H.; Trotter J. Oligodendrocytes secrete exosomes containing
36 major myelin and stress-protective proteins: Trophic support for axons? *Proteomics*
37 *Clinical Applications*, 2007, 1(11), 1446-61.
- 38 61. Tavakolizadeh, J.; Roshanaei, K.; Salmaninejad, A.; Yari, R.; Nahand, J.S.; Sarkarizi,
39 H.K.; Mousavi, S.M.; Salarinia, R.; Rahmati, M.; Mousavi, S.F.; Mokhtari, R.;
40 Mirzaei, H.; MicroRNAs and Exosomes in depression: Potential diagnostic
41 biomarkers. *J Cell Biochem*, 2018, 119(5), 3783-3797.
- 42 62. Yuan, D.; Zhao, Y.; Banks, W. A.; Bullock, K. M.; Haney, M.; Batrakova, E.;
43 Kabanov, A. V. Macrophage exosomes as natural nanocarriers for protein delivery to
44 inflamed brain. *Biomaterials*, 2017, 142, 1–12.

- 1 63. Eldh, M.; Ekström, K.; Valadi, H.; Sjöstrand, M.; Olsson, B.; Jernås, M.; Lötval, J.
2 Exosomes communicate protective messages during oxidative stress; possible role of
3 exosomal shuttle RNA. *PloS one*, 2010 5(12), e15353.
- 4 64. Rule Wigginton, K.; Menin, L.; Montoya, J.P.; Kohn, T. Oxidation of Virus Proteins
5 during UV254 and Singlet Oxygen Mediated Inactivation. *Environmental Science &*
6 *Technology* 2010,44, 5437–5443.
- 7 65. Greenberg, J.R. Ultraviolet light-induced crosslinking of mRNA to proteins. *Nucleic*
8 *Acids Research* 1979;6:715–732.
- 9 66. Benedikter, B.J.; Weseler, A.R.; Wouters, E.F.M.; Savelkoul, P.H.M.; Rohde,
10 G.G.U.; Stassen, F.R.M. Redox-dependent thiol modifications: implications for the
11 release of extracellular vesicles. *Cellular and Molecular Life Sciences*, 2018, 75(13),
12 2321-2337.
- 13 67. Thom, S.R.; Bhopale, V.M.; Yang, M. Neutrophils generate microparticles during
14 exposure to inert gases due to cytoskeletal oxidative stress. *The Journal of Biological*
15 *Chemistry*, 2014,289(27),18831–18845.
- 16 68. Wei, Z.X.; Xie, G.J.; Mao, X.; Zou, X.P.; Liao, Y.J.; Liu, Q.S.; Wang, H, Cheng Y.
17 Exosomes from patients with major depression cause depressive-like behaviors in
18 mice with involvement of miR-139-5p-regulated neurogenesis.
19 *Neuropsychopharmacology*. 2020, 45(6),1050-1058.
- 20 69. Ailawadi, S.; Wang, X.; Gu, H.; Fan, G.C. Pathologic function and therapeutic
21 potential of exosomes in cardiovascular disease. *Biochim Biophys Acta*, 2015 ,
22 1852(1), 1-11.
- 23 70. Donlin-Asp, PG; Polisseni, C; Klimek, R; Heckel, A; Schuman EM Differential
24 regulation of local mRNA dynamics and translation following long-term potentiation
25 and depression. bioRxiv. doi: <https://doi.org/10.1101/2020.07.08.192369>
- 26 71. Cajigas IJ, Will T, Schuman EM. Protein homeostasis and synaptic plasticity. *EMBO*
27 *J*. 2010;29(16):2746-2752..
- 28 72. Buzas, E. I.; György, B.; Nagy, G.; Falus, A.; Gay, S. Emerging role of extracellular
29 vesicles in inflammatory diseases. *Nature Reviews Rheumatology*, 2014, 10, 356–364.
- 30 73. Frühbeis, C.; Fröhlich, D.; Krämer-Albers, E. M. (2012). Emerging roles of exosomes
31 in neuron-glia communication. *Frontiers in Physiology*, 2012, 3, 119.
- 32 74. Kuwano, N.; Kato, T.A.; Mitsuhashi, M.; Sato-Kasai, M.; Shimokawa, N.; Hayakawa,
33 K.; Ohgidani, M.; Sagata, N.; Kubo, H.; Sakurai, T.; Kanba S. Neuron-related blood
34 inflammatory markers as an objective evaluation tool for major depressive disorder:
35 An exploratory pilot case-control study. *Journal of Affective Disorders*. 2018, 240,
36 88-98.
- 37 75. <https://clinicaltrials.gov/ct2/show/NCT04202770>
- 38 76. Luarte, A.; Cisternas, P.; Caviedes, A.; Batiz, L.F.; Lafourcade, C.; Wyneken, U.;
39 Henzi, R.; Astrocytes at the Hub of the Stress Response: Potential Modulation of
40 Neurogenesis by miRNAs in Astrocyte-Derived Exosomes. *Stem Cells International*,
41 2017, 2017, 1719050.
- 42 77. Zheng, T.; Pu, J.; Chen, Y.; Guo, Z.; Pan, H.; Zhang, L.; Zhang, H.; Sun, B.; Zhang,
43 B. Exosomes Secreted from HEK293-APP Swe/Ind Cells Impair the Hippocampal
44 Neurogenesis. *Neurotoxicity Research*, 2017, 32(1), 82-93.

- 1 78. Bahrini, I.; Song, J-h.; Diez, D.; Hanayama, R. Neuronal exosomes facilitate synaptic
2 pruning by up-regulating complement factors in microglia. *Scientific Reports*,
3 2015, 5, 7989.
- 4 79. Bátiz, L. F.; Castro, M. A.; Burgos, P. V.; Velásquez, Z. D.; Muñoz, R. I.;
5 Lafourcade, C. A.; Troncoso-Escudero, P.; Wyneken, U. Exosomes as Novel
6 Regulators of Adult Neurogenic Niches. *Frontiers in cellular neuroscience*, 2016, 9,
7 501.
- 8 80. Malm, T.; Loppi, S.; Kanninen, K. M. Exosomes in Alzheimer's
9 disease. *Neurochemistry International*, 2016, 97, 193–199.
- 10 81. Grapp, M.; Wrede, A.; Schweizer, M.; Huwel, S.; Galla, HJ.; Snaidero, N.; Simons,
11 M.; Buckers, J.; Low, P.S.; Urlaub, H.; Gartner, J.; Steinfeld, R.; Choroid plexus
12 transcytosis and exosome shuttling deliver folate into brain parenchyma. *Nature*
13 *Communications*, 4 (2013), 2123.
- 14 82. Zabeo, D.; Cvjetkovic, A.; Lässer, C.; Schorb, M.; Lötvall, J.; Höög, J. L. Exosomes
15 purified from a single cell type have diverse morphology. *Journal of extracellular*
16 *vesicles*, 2017 6(1), 1329476.
- 17 83. Yu, L.L.; Zhu, J.; Liu J-X.; Jiang, F.; Ni, W.K.; Qu, L.S.; Ni, R.Z.; Lu, C.H.; Xiao M.B..
18 A Comparison of Traditional and Novel Methods for the Separation of Exosomes
19 from Human Samples. *Biomed Research International*, 2018, 1–9.
- 20 84. Li, P.; Kaslan, M.; Lee, S. H.; Yao, J.; Gao, Z. Progress in exosome isolation
21 techniques. *Theranostics*, 2017, 7, 789–804.
- 22 85. Sharples, R. A.; Scicluna, B. J.; Hill, A. F. Exosomes provide a protective and
23 enriched source of miRNA for biomarker profiling compared to intracellular and cell-
24 free blood AU - Cheng, Lesley. *J. Extracell. Vesicles 2014*, 3, 23743.
- 25 86. Lötvall, J.; Hill, A. F.; Hochberg, F.; Buzás, E. I.; Di Vizio, D.; Gardiner, C.; Gho, Y.
26 S.; Kurochkin, I. V.; Mathivanan, S.; Quesenberry, P.; Sahoo, S.; Tahara, H.;
27 Wauben, M. H.; Witwer, K. W.; & Théry, C. Minimal experimental requirements for
28 definition of extracellular vesicles and their functions: a position statement from the
29 International Society for Extracellular Vesicles. *Journal of extracellular vesicles*,
30 2014, 3, 26913.
- 31 87. Fauré, J.; Lachenal, G.; Court, M.; Hirrlinger, J.; Chatellard-Causse, C.; Blot, B.;
32 Grange, J.; Schoehn, G.; Goldberg, Y.; Boyer, V.; Kirchhoff, F.; Raposo, G.; Garin,
33 J.; Sadoul, R. Exosomes are released by cultured cortical neurones. *Molecular and*
34 *Cellular Neurosciences*, 2006, 31(4):642-8.
- 35 88. Lachenal, G.; Pernet-Gallay, K.; Chivet, M.; Hemming, F.J.; Belly, A.; Bodon, G.;
36 Blot, B.; Haase, G.; Goldberg, Y.; Sadoul R. Release of exosomes from differentiated
37 neurons and its regulation by synaptic glutamatergic activity. *Molecular and Cellular*
38 *Neurosciences*, 2011, 46(2), 409-18.
- 39 89. Krämer-Albers, E.M.; Bretz, N.; Tenzer, S.; Winterstein, C.; Möbius, W.; Berger, H.;
40 Nave K.A.; Schild, H.; Trotter J. Oligodendrocytes secrete exosomes containing
41 major myelin and stress-protective proteins: Trophic support for axons? *Proteomics*
42 *Clinical Applications*, 2007, (11), 1446-61.

43

44

45

1

2

3 **Captions for Figures**

4 **Figure 1:** Microglia activation with release of extracellular vesicles (EVs).
5 lipopolysaccharide (LPS), mitogen-activated protein kinases (MAPKs) superfamily, c-Jun N-
6 terminal kinase (JNK 1/2) and p38 proteins, nuclear factor- κ B (NF- κ B) interleukin (IL)-1 β ,
7 IL-6 and tumour necrosis factor (TNF)- α .

8 **Captions for Graphical Abstract**

9 Biosynthesis and release of extracellular vesicles:

10