



**HAL**  
open science

## **W(CO)<sub>6</sub> in cryogenic solids: A comparative study of vibrational properties**

Raphaël Thon, Wutharath Chin, Didier Chamma, Alejandro Gutiérrez Quintanilla, Michele Chevalier, Jean-Pierre Galaup, Claudine Crépin

► **To cite this version:**

Raphaël Thon, Wutharath Chin, Didier Chamma, Alejandro Gutiérrez Quintanilla, Michele Chevalier, et al.. W(CO)<sub>6</sub> in cryogenic solids: A comparative study of vibrational properties. *Journal of Luminescence*, 2017, 191, Part A, pp.78-86. 10.1016/j.jlumin.2017.01.041 . hal-03097322

**HAL Id: hal-03097322**

**<https://hal.science/hal-03097322>**

Submitted on 14 Jan 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# W(CO)<sub>6</sub> in cryogenic solids: A comparative study of vibrational properties

Raphaël Thon<sup>a,1</sup>, Wutharath Chin<sup>a,\*</sup>, Didier Chamma<sup>a</sup>, Alejandro Gutiérrez-Quintanilla<sup>a</sup>, Michèle Chevalier<sup>a</sup>, Jean-Pierre Galaup<sup>b</sup>, Claudine Crépin<sup>a</sup>

<sup>a</sup>*Institut des Sciences Moléculaires d'Orsay, CNRS, Univ. Paris-Sud, Université Paris-Saclay, F-91405 Orsay (France)*

<sup>b</sup>*Laboratoire Aimé Cotton, CNRS, ENS Cachan, Univ. Paris-Sud, Université Paris-Saclay, F-91405 Orsay (France)*

---

## Abstract

A comparative study of the vibrational properties of W(CO)<sub>6</sub> has been performed in different solids at cryogenic temperatures focusing on the IR absorption and the vibrational dynamics of the CO stretching mode of the organometallic compound. Guest-host interactions are investigated in doped solids through the linear IR spectroscopy and four-wave mixing techniques at different temperatures. We show how the host nature, the trapping site, the crystallographic ordering affect the properties of the guest molecule and in particular its vibrational dynamics.

*Keywords:* photon echo; vibrational dynamics; cryogenic solids

---

## 1. Introduction

The advent of picosecond (ps) and femtosecond (fs) laser sources in the IR domain has enabled the study of the vibrational dynamics of a wide variety of systems spanning from biological molecules [1, 2, 3, 4, 5] to material sciences [6, 7, 8] through nonlinear four-wave mixing techniques such as IR photon echo [9] or two-dimensional IR spectroscopy [10]. Thanks to the very large transition dipole

---

\*Corresponding author

*Email address:* wutharath.chin@u-psud.fr (Wutharath Chin)

<sup>1</sup>Present address: CNRS Formation Entreprises, Bât. 31, Avenue de la Terrasse, 91198 Gif-sur-Yvette Cedex (France)

moment ( $\mu = 1$  D) of the asymmetric CO stretch vibrational mode ( $T_{1u}$  mode near  $2000\text{ cm}^{-1}$ )  $\text{W}(\text{CO})_6$  organometallic complex is highly interesting since it yields a large nonlinear response. In this sense  $\text{W}(\text{CO})_6$  has been widely studied  
 10 by means of nonlinear processes in various condensed media [11, 12, 13]. In particular  $\text{W}(\text{CO})_6$  represents a very sensitive probe to characterize guest-host interactions and allows a very detailed insight into environment effects induced by the nature of the host on the molecular dynamics of  $\text{W}(\text{CO})_6$ . We have investigated the environment effects on the vibrational properties of  $\text{W}(\text{CO})_6$   
 15 in the condensed phase in different cryogenic lattices such as rare gas crystals (argon [13], krypton [14]) and molecular hosts (nitrogen [14], methane [15]).

The vibrational properties of  $\text{W}(\text{CO})_6$  have been measured by linear and nonlinear (pump-probe and photon echo techniques) IR spectroscopy at different temperatures. We thus have access to  $T_1$  population relaxation time and  $T_2$   
 20 dephasing time.  $T_2$  is related to the homogeneous broadening  $\gamma_{\text{homo}}$  through the relation:

$$\gamma_{\text{homo}} = \frac{1}{\pi T_2} = \frac{1}{2\pi T_1} + \frac{1}{\pi T_2^*} \quad (1)$$

where  $T_2^*$ , called pure dephasing time, is especially sensitive to the environment. In order to get an overview of the vibrational behavior of  $\text{W}(\text{CO})_6$  in cryogenic lattices, we performed additional experiments in neon (for probing  
 25 a third rare gas host), in parahydrogen (a specific host with quantum properties) and in carbon tetrafluoride (following the series of methane solids  $\text{CH}_4$  and  $\text{CD}_4$ ). This paper is thus dedicated to the comparison of these various results allowing for conclusive remarks on the host dependence of  $\text{W}(\text{CO})_6$  vibrational properties. After a brief description of the experimental procedures, the results in the different hosts are summarized. They are discussed in the last part before the conclusions in terms of host effects on the absorption features, vibrational population relaxation and dephasing processes. We show how the host nature, the trapping site, the crystallographic ordering affect the properties of the guest molecule and in particular its vibrational dynamics. The conclusions highlight

35 the main tendencies obtained from the comparison of all the results.

## 2. Experimental procedure and methods

The detailed experimental set-ups have been extensively described in our previous papers [14, 16] and only the main descriptions related to new results are presented.

40 In the case of  $\text{CF}_4$ , as for the experiments described in Refs. [14, 15] the matrix samples have been obtained by continuous deposition of a gaseous mixture onto a cold  $\text{CaF}_2$  window of the cryostat. Gaseous samples are prepared at room temperature by mixing  $\text{W}(\text{CO})_6$  (Sigma Aldrich) with the carrier gas with  $[\text{W}(\text{CO})_6]/[\text{Host}]$  ratio varying from  $10^{-5}$  to  $10^{-3}$ . Matrix thickness is  
45 kept below  $100 \mu\text{m}$ . Concentration and deposition times are adapted to keep an optical density of the  $T_{1u}$  mode of  $\text{W}(\text{CO})_6$  of  $\approx 1$ . Deposition temperatures can vary between 15 and 30 K depending on the host (30 K for  $\text{CF}_4$ ).

Nonlinear measurements are obtained using an IR OPA source for fs pulses (see ref. [14] for details). The fs IR OPA is combined to a 3-pulse photon echo  
50 set-up. It is pumped by a 800 nm Ti:Sa laser/amplifier system and delivers an IR pulse centered at  $1980 \text{ cm}^{-1}$ , bandwidth  $150 \text{ cm}^{-1}$ , duration 145 fs, output  $\approx 3 \mu\text{J}$ . Resulting IR beam is then split into three beams ( $\mathbf{k}_1$ ,  $\mathbf{k}_2$  and  $\mathbf{k}_3$  wave vectors) of equivalent intensities using two coated ZnSe beamsplitters. The echo signal is emitted in a well-defined direction given by phase-matching such  
55 as  $\mathbf{k}_s = \mathbf{k}_3 + \mathbf{k}_2 - \mathbf{k}_1$ . The signal beam is focused on an MCT (mercury cadmium telluride) detector. There are two independant delay stages giving access to two delay times  $\tau$  and  $T$ , called coherence time and waiting time and corresponding to the delay between the first two pulses and the last two pulses, respectively. The detected signal (quadratic detection) is:

$$S(\tau, T) = \int_0^\infty dt \left| P^{(3)}(\tau, T, t) \right|^2 \quad (2)$$

60  $P^{(3)}(\tau, T, t)$  is the third order polarization created in the sample with the wave vector  $\mathbf{k}_s$ . In a rough way, the  $S(T)$  signal at fixed  $\tau$  gives access to  $T_1$  population

lifetime and  $S(\tau)$  signal at fixed  $T$  to dephasing processes through the measure of  $T_2$  dephasing time.

Absorption spectra have been recorded using an FTIR Mattson spectrometer at  $0.25\text{ cm}^{-1}$  spectral resolution to check the quality of the sample and its evolution with temperature and time. A switching gold coated mirror brings the FTIR global beam to probe the same matrix region as in photon echo experiments. Linear and nonlinear data have been measured using the same nitrogen-cooled MCT detector.

Additional experiments in neon and parahydrogen ( $\text{pH}_2$ ) have been performed with another set-up dedicated to  $\text{pH}_2$  experiments. Hydrogen is converted to parahydrogen in a first cryostat kept at 17 K and parahydrogen vapor is directly condensed onto a cold diamond window of a second cryostat (IceOxford) at the lowest temperature of 3 K.  $\text{W}(\text{CO})_6/\text{pH}_2$  samples are obtained by flowing  $\text{pH}_2$  over a tube containing  $\text{W}(\text{CO})_6$  and carrying its vapor before the deposition. The experimental procedure to produce doped  $\text{pH}_2$  samples is the same as that described in ref. [17].  $\text{W}(\text{CO})_6/\text{Ne}$  samples are obtained in the same manner, replacing hydrogen by neon and keeping the first cryostat at 40 K to avoid any neon condensation. In this case, the first cryostat is used to trap all residual impurities. IR spectra of the samples are recorded by means of a Nicolet FTIR spectrometer at  $0.125\text{ cm}^{-1}$  spectral resolution. No time-resolved measurements have been conducted with this cryogenic set-up.

### 3. Results

#### 3.1. Rare gas matrices: Ar, Kr, Ne

$\text{W}(\text{CO})_6$  has been isolated in Ar, Kr and Ne solids. The IR spectra have been recorded in the three hosts and photon echo experiments have been performed in Ar and Kr. The IR spectrum in Kr and Ne in the region of the asymmetric  $\nu\text{CO}$  stretch of  $\text{W}(\text{CO})_6$  is displayed in Figure 1. Spectra reveal a complex absorption spreading over  $10\text{ cm}^{-1}$  in contrast to the gas phase where there is only one line at  $1998\text{ cm}^{-1}$  [18]. The complex structure reflects the complexity of

the trapping sites. With a diameter around 0.78 nm,  $\text{W}(\text{CO})_6$  is a huge impurity to trap in these crystallographic lattices. Rare gas solids have an *fcc* structure with a nearest neighbors distance  $d$  of 0.40, 0.38 and 0.32 nm for Kr, Ar and Ne respectively, i.e. much smaller than the size of  $\text{W}(\text{CO})_6$ . Consequently, the  
95 guest molecule is trapped in multi-vacancy sites. The structure of the absorption spectra can result from two effects: the presence of many different trapping sites (differing for example by the number of hosts removed), and the possible degeneracy removal of the  $T_{1u}$  mode (triply degenerated in the isolated molecule) in trapping sites of lower degree of symmetry than  $O_h$  group (the group of  
100 symmetry of the isolated molecule).

Because Kr, Ar and Ne have different lattice parameters, trapping sites should be different in the three hosts. In that sense, the Ar spectrum seems the simplest and could reflect a smaller number of different trapping sites, i.e. a better match between the size of  $\text{W}(\text{CO})_6$  and that of the vacancies [13]. Nev-  
105 ertheless spectra in the three rare gas solids show a major structure in the red side and a minor one in the blue side. The energy gap between these two bands is roughly  $5 \pm 2 \text{ cm}^{-1}$  with no obvious correlation with the lattice parameter. In Ne, the absorption spectrum spreads only over  $8 \text{ cm}^{-1}$  and this gap is reduced to  $\approx 3 \text{ cm}^{-1}$  (Figure 1). The frequency shift between matrix and gas phase  
110 ( $1998 \text{ cm}^{-1}$ ) is also the smallest. This is a well-known consequence of the lowest polarizability of Ne which induces the smallest matrix effect on the transition frequencies of guest molecules.

No significant change is observed when varying the temperature in Ar [13] and Kr [14] in terms of band position or linewidth. On the contrary, annealing at  
115 60 K in Kr leads to reorganization of the trapping sites yielding a small change in the relative intensities of the bands. We have also observed slight differences in  $\text{W}(\text{CO})_6/\text{Ar}$  spectra when varying the experimental deposition conditions. However the relative intensities of the two main features remain constant. The IR spectroscopy in Ne has been recorded at 3, 6 and 8 K and no change has  
120 been observed.

Despite complex band structures, the vibrational dynamics in Kr and Ar




Figure 1: IR spectra of  $W(CO)_6$  in (a) Kr (35 K), (b) Ne (3 K) and (c)  $pH_2$  (3 K). The inset shows a schematic of  $W(CO)_6$  (W in blue, C in grey, O in red).

occur to be rather simple. Time-resolved signals measuring  $T_1$  and  $T_2$  always exhibit long decay times in the hundreds of ps. Of course, since the IR laser pulses are spectrally broad, the time-resolved signals correspond to the nonlinear response of the entire absorption spectra. In Kr, there is no variation in the population decays when changing the temperature. The decay is not in fact mono-exponential and the  $S(T)$  signals are better fitted with two exponential contributions that take into account two different dynamics linked to the population relaxation from  $v = 1$  excited state ( $T_1$ ) and recovery to the ground state ( $T_g$ ), suggesting a long-lived intermediate level in the vibrational relaxation. We find  $T_1 = 80 \pm 20$  ps and  $T_g \approx 1.5$  ns [14]. In Ar, a single exponential is found but the signal has not been recorded until the end of the relaxation, suggesting that the long  $T_g$  time has not been actually measured. It gives a  $T_1$  population relaxation time which slightly shortens when temperature increases with  $T_1 >$  200 ps at 8 K and  $T_1 = 160$  ps at 20 K [13].

Similarly to  $S(T)$ , the  $S(\tau)$  signals in Kr show no temperature effects between 25 and 50 K. The decay of the signal yields  $100 \pm 20$  ps, equivalent to a  $\gamma_{\text{homo}}$  homogeneous broadening of  $0.106 \text{ cm}^{-1}$ . It is smaller than the width of the narrowest lines of the absorption spectrum ( $\text{FWHM} \approx 0.5 \text{ cm}^{-1}$ ): it means that the absorption structure is not composed of homogeneous substructures. In Ar,  $T_2$  values are slightly longer than in Kr, they are not constant with temperature but they are in the same order of magnitude with  $T_2$  (8 K) = 400 ps and  $T_2$  (20 K) = 200 ps. Corresponding  $\gamma_{\text{homo}}$  of 0.026 and  $0.053 \text{ cm}^{-1}$  are much less than the linewidths of the absorption structures ( $\approx 2.3 \text{ cm}^{-1}$ ). In both rare gas solids we find  $T_2$  comparable to twice  $T_1$ , meaning that dephasing processes are essentially governed by population relaxation.

### 3.2. Molecular matrices

#### 3.2.1. Nitrogen and methane

We have studied nitrogen and methane solids because the crystals also adopt an *fcc* structure similar to rare gas solids. Moreover, nearest neighbors distances  $d$  are of the same order of magnitude, and in particular close to that of Kr lattice:



$d = 0.401$  nm and  $0.417$  nm in  $N_2$  and  $CH_4$  respectively. It is thus expected that the geometry of the trapping sites are similar between these molecular solids and Kr solids. In the case of  $CH_4$ , our experimental conditions allow  
155 the exploration of different phases.  $CH_4$  solid presents two crystalline phases (both of *fcc* structure) with a temperature of phase transition at  $20.4$  K at zero pressure. From  $20.4$  K to  $90.7$  K phase I is orientationally disordered and molecules occupy  $O_h$  symmetry sites. Below  $20.4$  K in phase II, there is partial ordering of  $CH_4$  molecules due to octupole-octupole interactions with  $3/4$  of  
160 molecules undergoing hindered rotation in  $D_{2d}$  symmetry and  $1/4$  free rotation in  $O_h$  symmetry [19].

The IR absorption of  $W(CO)_6$  in  $N_2$  and  $CH_4$  is quite complex and displays structured bands as in the case of Ne, Ar and Kr. As previously, spectra show two main features centered at  $1984$  and  $1989$   $cm^{-1}$  in  $N_2$  and  $1981$  and  $1987$   
165  $cm^{-1}$  in  $CH_4$  (frequency gap  $\approx 5$   $cm^{-1}$ , as in Kr). Main differences from rare gas matrices stem from the strong temperature dependence of the IR spectra (Figure 2). Temperature effects have been of great help to assign the different substructures. In both molecular solids, four bands or substructures can be distinguished (numbered 1 to 4 in Figure 2). In  $N_2$  only band 2 exhibits a  
170 strong and reversible temperature effect while intensities of the three other bands remain almost constant. These characteristics suggest that bands 1, 3 and 4 correspond to the three components of the vibrational  $T_{1u}$  mode in sites of low symmetry. Band 2 would then correspond to sites of high symmetry where the molecule keeps its  $O_h$  symmetry [14]. In spite of a more complex absorption  
175 structure in  $CH_4$ , the comparison with  $N_2$  also leads to assign bands 1, 3 and 4 to low symmetry sites and band 2 to high symmetry sites [15]. Populations of high and low-symmetry sites are comparable in  $N_2$  and  $CH_4$  with a ratio of  $40:60$  and  $45:55$  respectively. The temperature broadening of band 2 in  $N_2$  could be due to an expected homogeneous broadening and time-resolved measurements confirm  
180 this assumption (see below). Very surprisingly, the temperature behavior seems reversed in  $CH_4$  as highlighted in Figure 2. In fact, bands 1, 2 and 3 are found to coalesce from phase II to phase I, leading to a global narrowing when




Figure 2: IR spectra of  $W(CO)_6$  in  $N_2$  at 6 and 22 K and  $CH_4$  at 8 and 22 K. Low and high temperatures are respectively plotted as black and red traces.




Figure 3:  $S(T)$  signals (normalized) of  $W(CO)_6$  in (a)  $N_2$  at 20 K and (b)  $CH_4$  at 14 K (black) and 25 K (red). The very steep rise (“spike”) at  $T = 0$  is due to additional nonlinear signals produced when the IR pulses simultaneously arrive onto the sample. It is not related to the probed dynamics [21].

temperature increases [15]. This effect is assigned to a motional narrowing due to the complete rotational disorder and the large motional fluctuations of  $CH_4$  hosts in phase I, which for this reason is also named plastic phase [20].

There is no obvious temperature effect on the vibrational population relaxation of  $W(CO)_6$  in  $N_2$  and the decay times are in the hundreds of ps, similarly to the case of Kr (Figure 3a). As in Kr two times in the population decay can also be considered, a  $T_1$  time around 180 ps and a very long recovery time  $T_g$   $> 1$  ns [14]. They represent averaged population decay times for all the sites as in Kr and Ar. On the contrary, a strong temperature effect is observed in

$\text{CH}_4$  in the  $S(T)$  signals corresponding to distinct dynamical behaviours above and below phase transition of  $\text{CH}_4$  (Figure 3b): there is one long component around 200 ps at all temperatures (same order of magnitude as  $T_1$  in  $\text{N}_2$ ) but  
 195 there is an additional shorter decay of 10 ps in phase I (red trace in Figure 3b). The weight of the short component corresponds to the weight of high symmetry sites obtained in the analysis of the absorption spectra. Our analysis of the time-resolved signals suggests that the short 10 ps component reflects  $T_1$  time of  $\text{W}(\text{CO})_6$  molecules in these high symmetry sites [15]. In these sites,  
 200  $\text{W}(\text{CO})_6$  does not significantly perturb the  $\text{CH}_4$  lattice and fast relaxation involving a methane vibron can occur in phase I because of the strong broadening of methane modes in this crystallographic phase. Note that, whatever the temperature no very long ( $> 400$  ps) decay has been measured in  $\text{CH}_4$  precluding the existence of a very long  $T_g$  time.

205 Photon echo measurements in  $\text{N}_2$  and  $\text{CH}_4$  both exhibit temperature effects. Figure 4 shows signals recorded at 9 and 20 K in  $\text{N}_2$  and at 8 and 24 K in  $\text{CH}_4$ . Experimental curves in  $\text{N}_2$  are correctly modelled when considering  $T_2$  times that depend on the family of sites. Only  $T_2$  times of high symmetry sites are strongly temperature dependent: they vary from 130 ps at 6 K to 15 ps  
 210 at 20 K. It is in perfect agreement with the temperature broadening of band 2 in the absorption, leading to a significant homogeneous contribution at 20 K. In low symmetry sites,  $T_2$  is almost constant, around 50 ps, i.e. shorter than in rare gas matrices. The temperature dependence in high symmetry sites can be described as an enhancement of the homogeneous width by coupling with a  
 215 libration mode of the nitrogen lattice [14].

In  $\text{CH}_4$ , beside the large shortening of the population relaxation time in phase I assigned to high symmetry sites,  $T_2$  is found to decrease with temperature increase in all sites. It is well depicted in Figure 4b where  $S(\tau)$  signals have been measured with long waiting times ( $T > 30$  ps) to get rid of this short 10  
 220 ps component in  $T_1$  in phase I.  $T_2$  times are found to decrease smoothly from 120-140 ps at 6 K to 25-30 ps at 30 K. The latter value is very close to twice  $T_1$  of high symmetry sites, where dephasing is governed by the loss of population




Figure 4:  $S(\tau)$  signals (normalized) of  $W(CO)_6$  in (a)  $N_2$  and (b)  $CH_4$ . Red circles and black squares represent signals at high (20 and 24 K) and low (8 and 9 K) temperatures respectively.

in phase I. The corresponding homogeneous width of  $\approx 0.4 \text{ cm}^{-1}$  at 30 K for all sites is consistent with a predominant homogeneous broadening with a band-  
width of  $0.5 \text{ cm}^{-1}$  measured on the main component of the absorption spectrum  
at 30 K. In both  $\text{N}_2$  and  $\text{CH}_4$  solids, the  $S(\tau)$  signal becomes more symmetric  
with respect to  $\tau$  (Figure 4) reflecting the increase of the homogeneous contribu-  
tion to broadening but one can notice that, whereas the rise of the signal (at  $\tau$   
 $< 0$ ) does not depend on temperature in  $\text{N}_2$ , it is longer at high temperature in  
 $\text{CH}_4$  indicating that the inhomogeneous contribution is simultaneously reduced,  
which is a signature of the motional narrowing [15].

Experiments have also been conducted in heavy methane ( $\text{CD}_4$ ) for which  
two phase transitions can be explored. In fact, the  $T_{1u}$  mode of  $\text{W}(\text{CO})_6$  is  
found in resonance with the  $2\nu_4$  overtone mode of  $\text{CD}_4$  [15]. The consequence  
is a very fast population relaxation at all temperatures ( $\approx 10 \text{ ps}$ ) which prevents  
any analysis of environment effects on pure dephasing processes in this solid.  
The absorption spectrum shows the same global structure as in  $\text{CH}_4$ , with broad  
features in all the phases, suggesting a similar analysis based on low and high  
symmetry sites with a high/low ratio in  $\text{CD}_4$  of 55:45.

### 3.2.2. Carbon tetrafluoride $\text{CF}_4$

In order to probe another crystallographic environment with molecular "spher-  
ical" hosts, we have studied  $\text{CF}_4$  solid after  $\text{CH}_4$  and  $\text{CD}_4$ .  $\text{CF}_4$  carbon tetraflu-  
oride presents a phase transition at 76.2 K (at zero pressure) [22], the high- and  
low-temperature phases are respectively designated by I and II. Phase I (or  $\beta$ ) is  
also named "plastic" phase due to orientational disorder of  $\text{CF}_4$  molecules [23],  
similarly to  $\text{CH}_4$ . It corresponds to a rhomboedric structure. Phase II (or  $\alpha$ )  
is a well-ordered crystal and the lattice structure changes to monoclinic [24].  
It is therefore an interesting medium for trapping  $\text{W}(\text{CO})_6$  in terms of lattice  
structure and orientation of molecules.

In fact, only phase II has been explored in our experiments. Attempts to  
measure spectra above 50 K failed due to low optical density of the sample  
because of evaporation of the  $\text{CF}_4$  matrix at 55 K despite a melting point at




Figure 5: IR spectra of  $W(CO)_6$  in  $CF_4$  from 10 K to 40 K (top to bottom).

89.5 K. This can be due to local heating of the matrix by the light sources. The IR spectrum is temperature dependent, as shown in Figure 5 between 10 and 40 K. At low temperature (8 K) it exhibits two main bands centered at 1987.5 and 1989  $\text{cm}^{-1}$  (bands 1 and 2) and weaker bands at 1992.9 and 1994.4  $\text{cm}^{-1}$ . The temperature dependence of the two main bands is especially pronounced. Band 1 at 1987.5  $\text{cm}^{-1}$  broadens and shows a more and more obvious structure from 8 to 20 K. It seems to contain a substructure that shifts to the blue, reaching the middle of the two bands around 22 K and joining band 2 at higher temperatures. The consequence is a red shift and an increase of band 2 at 1989  $\text{cm}^{-1}$  with temperature. At 35 K, there is only one main band at 1988.7  $\text{cm}^{-1}$ . It is found to narrow slightly from 8 to 20 K and to broaden because of the previous effect. Spectral changes observed when varying the temperature are reversible.  $\text{CF}_4$  forms a larger "sphere" (0.43 nm diameter) than methane but as in the other solids studied,  $\text{W}(\text{CO})_6$  (0.78 nm diameter) replaces more than one host in the lattice. Because of the monoclinic structure in phase II, the  $O_h$  symmetry of  $\text{W}(\text{CO})_6$  cannot be preserved and band splitting due to degeneracy removal is expected. It is thus not surprising that the spectral behavior is different in  $\text{CF}_4$ . Anyway, the comparison with the observations in  $\text{N}_2$  and  $\text{CH}_4$  suggests that the most intense band 2 at 1989  $\text{cm}^{-1}$  contains the main components of highest symmetry sites. The frequency shift of some components in the absorption spectrum toward the main one when temperature increases looks like the band coalescence observed in  $\text{CH}_4$  from phase II to phase I [15], although 35 K is rather far from the temperature of phase transition in  $\text{CF}_4$ . One must remind that a lowering of this temperature was observed in methane when the solid was doped with  $\text{W}(\text{CO})_6$  [15] and such effect can also occur in  $\text{CF}_4$ .

Unfortunately, time-resolved measurements have been very difficult to obtain due to high scattering of the samples. The best signals have been recorded in a short temperature range only (30-45 K) corresponding to the best quality samples so only global behaviors can be extracted from the signals. Nevertheless a clear shortening of  $T_2$  upon temperature increase is observed.  $T_2$  of 50 ps  $\pm$  10 ps is obtained between 10 and 20 K, and this value continuously decreases to



$\approx 20$  ps at 50 K. The  $S(\tau)$  signal at 30 K is shown in Figure 6 a. The asymmetric profile indicates a predominant inhomogeneous contribution to broadening. The deduced value of  $T_2 = 38$  ps is an average for all bands. It corresponds to a homogeneous broadening of  $0.27 \text{ cm}^{-1}$ , which is lower than the linewidth of the substructures ( $0.5 - 1 \text{ cm}^{-1}$ ) measured in the absorption spectra. The  $S(\tau)$  profile seems obviously more symmetric at 50 K (not shown), corresponding to the IR spectrum involving only one main band.  $T_1$  is measured in the hundreds of ps in the explored temperature range, without any clear temperature effect. An example is given in Figure 6b at 35 K, yielding  $T_1 = 240 \text{ ps} \pm 30 \text{ ps}$ . No obvious long component can be extracted from the  $S(T)$  curves, suggesting that there is no long  $T_g$  recovery time as in  $\text{CH}_4$ . Similarly to the case of  $\text{N}_2$ , no population relaxation enhancement takes place in  $\text{CF}_4$  while a pure dephasing time  $T_2^*$  in the tens of ps as given by eq. 1 is present, that is temperature dependent. The molecular nature of the host could be at play in pure dephasing process, as in  $\text{N}_2$  and  $\text{CH}_4$  but data are too scattered for a more precise conclusion.

### 3.3. Quantum matrix: parahydrogen $\text{pH}_2$

Only IR measurements are available in solid parahydrogen ( $\text{pH}_2$ ). Parahydrogen is a very attractive environment as it possesses unique properties. It is a quantum crystal thanks to its light  $\text{pH}_2$  mass and to the significant translational zero-point motion of the  $\text{pH}_2$  molecules about their equilibrium positions within the solid. The weak  $\text{pH}_2$ - $\text{pH}_2$  intermolecular forces, the  $J = 0$  rotational state at cryogenic temperatures for all molecules, the spherical symmetry of the ground state, the minimal cage effect contribute to the "softness" of the matrix and result in weak perturbations on the guest molecules [25]. By trapping  $\text{W}(\text{CO})_6$  molecules in  $\text{pH}_2$  we thus expect that its spectroscopy and dynamics will qualitatively differ from the other solids studied so far. Solid  $\text{pH}_2$  adopts an *hcp* lattice structure with nearest neighbors distance  $d = 0.379 \text{ nm}$ , i.e. a value very close to that of solid Ar.  $\text{W}(\text{CO})_6$  should thus replace a similar number of  $\text{pH}_2$  hosts as Ar atoms in the respective lattices. Because of its softness, site effects are usually strongly reduced in this quantum solid and absorption bands




Figure 6: Examples of time-resolved signals in  $\text{CF}_4$  doped with  $\text{W}(\text{CO})_6$  (arbitrary units), (a)  $S(\tau)$  at 30 K,  $T = 20$  ps, (b)  $S(T)$  at 35 K ("spike" at the origin as in Figure 3a).

are expected to be homogeneously broadened [26, 27, 28].

315 The IR spectrum of  $\text{W}(\text{CO})_6$  isolated in  $\text{pH}_2$  at 3 K is displayed in Figure 1c. It is simpler than in the other solids and the global feature is much narrower than in other hosts. However, it does not show a unique band, as in the isolated molecule but two doublets. The most intense doublet absorbs at 1992.2 and 1992.5  $\text{cm}^{-1}$  and the weakest one is located at 1992.8 and 1993.3  $\text{cm}^{-1}$ . The  
320 absorption spectrum exhibits a very slow time evolution, with a decrease of the bands at 1993.3 and 1992.5  $\text{cm}^{-1}$  meaning that they correspond to metastable sites. The two weaker bands have half intensities of the two stronger ones. We conclude that the two doublets correspond to two sites where the triply degenerated  $T_{1u}$  mode gives rise to two components, one being doubly degenerated  
325 (the strongest one). Such degeneracy removal is consistent with the *hcp* structure. Bands are very narrow and their actual linewidth cannot be reached with our FTIR resolution ( $\text{FWHM} < 0.125 \text{ cm}^{-1}$ ). It gives a 0.1  $\text{cm}^{-1}$  limit to the value of  $\gamma_{\text{homo}}$ . The corresponding  $T_2$  dephasing time is thus longer than 100 ps. Time-resolved experiments are necessary to obtain more accurate values.

## 330 4. Discussion

### 4.1. Spectroscopy

The complex structures in the absorption spectra in the crystalline cryogenic solids mainly reflect site effects. In amorphous solids, like xerogels or glasses, only a non structured and broad absorption band has been observed [16, 30, 31].  
335 Since this band is inhomogeneously broadened, it corresponds to a continuum of different sites. Results in different environments are summarized in Table 1. The analysis of the structures in  $\text{N}_2$  and  $\text{CH}_4$  suggests the existence of high symmetry sites where the  $O_h$  symmetry of  $\text{W}(\text{CO})_6$  is preserved [14, 15]. In *fcc* lattices, with nearest neighbors distances around 0.4 nm,  $O_h$  (or  $T_d$ ) trapping  
340 sites large enough to contain the guest molecule are obtained when removing 13 hosts (one and its 12 nearest neighbors). The size of this big hole has a diameter of around 0.77 nm, which is well adapted to  $\text{W}(\text{CO})_6$ . Therefore,

Table 1: Summary of the results on the vibrational properties for the main features of  $W(\text{CO})_6$  in different environments.

Host	T (K)	Absorption ( $\text{cm}^{-1}$ )		$T_1$ (ps)	$T_2$ (ps)	References
		Major feature	Minor feature			
Ar	7-8	1986	1991	240-270	$400 \pm 80$	Ref. [13]
	20	1986	1991	$160 \pm 10$	$200 \pm 40$	
Kr	25	1983.9	1989.3	$80 \pm 20$ &	$100 \pm 20$	Ref. [14]
				$> 1000 (T_g)$		
Ne	3 $\rightarrow$ 8	1995.9	1999	not measured		
N <sub>2</sub>	6	1984.1	1989.3	$180 \pm 30$ &	130 (high sym.) &	Ref. [14]
				$> 1000 (T_g)$	50 (low sym.)	
	20	1983.7	1989	$180 \pm 30$ &	15 (high sym.) &	
				$> 1000 (T_g)$	50 (low sym.)	
CH <sub>4</sub>	6	1980.6	1987.0	200 (low sym.)	130	Ref. [15]
	25	1980.8	1986.6	10 (high sym.) &	30	
				200 (low sym.)		
CD <sub>4</sub>	15 $\rightarrow$ 30	1980.5	1986.3	$\leq 10$	10	Ref. [15]
CF <sub>4</sub>	20 $\rightarrow$ 50	1989	1992.9	240	38	this work
pH <sub>2</sub>	3	1992.5	1993.3	not measured		
CCl <sub>4</sub>	300	1978		800	$< 1$	Ref. [29]
dibutyl- phtalate	0 $\rightarrow$ 300	1982		33	33 $\rightarrow$ 13	Ref. [11]
xerogels	4.5	1982		60-70	28	Ref. [16]
	300				4	

these " $O_h$ " trapping sites should also exist in Ar and Kr matrices and possibly  
 in Ne. Similarly to the case of  $N_2$  and  $CH_4$ , we can also conclude that in all  
 345 these *fcc* solids the weaker feature in the blue part of the spectra is assigned to a  
 non degenerated component of the  $T_{1u}$  mode of  $W(CO)_6$  in low symmetry sites.  
 A rough estimation of the integrated absorption of the weak and the strong  
 bands (separated by few wavenumbers) can give an idea of the ratio of high/low  
 symmetry sites. We find 40:60, 45:55, 45:55 in Kr, Ar and Ne respectively, it is  
 350 as expected very similar to the 40:60 and 45:55 ratios obtained in the spectral  
 analysis in  $N_2$  and  $CH_4$  respectively. The *hcp* structure of  $pH_2$  is very close  
 to the previously described *fcc* structure and with a nearest neighbors distance  
 $d$  of 0.379 nm, a hole of 13  $pH_2$  could also fit for the trapping of  $W(CO)_6$ .  
 The symmetry of such trapping site is  $D_{3h}$  in the *hcp* structure, explaining the  
 355 splitting of the  $T_{1u}$  mode in two components. The results in  $pH_2$  in that sense  
 confirm the previous assignment.

Figure 7a shows the IR spectra of  $W(CO)_6$  in the different solids studied,  
 $CH_4$ , Kr,  $N_2$ , Ar,  $CF_4$ ,  $pH_2$  and Ne from top to bottom. All these features are  
 red-shifted from the gas phase frequency ( $1998\text{ cm}^{-1}$ ) [18], meaning that the gas  
 360 to matrix shift is due to dispersive interactions. In Figure 7b is plotted the  $T_{1u}$   
 frequency measured in the different media as a function of the polarizability  $\alpha$   
 of the host. It clearly shows a correlation between  $T_{1u}$  position and  $\alpha$ , except  
 for  $CF_4$ . The weakest feature observed in all hosts is less down shifted from the  
 gas phase. It corresponds to a CO stretch mode in a specific direction in low  
 365 symmetry sites that is either weakly perturbed by the environment, or more  
 sensitive to constraints as in Ne, where it is slightly above the gas phase value  
 because of a more compact lattice ( $d = 0.32\text{ nm}$ ) inducing more repulsive forces  
 than in the other hosts. The red-shift of the  $T_{1u}$  frequency in the condensed  
 phase can reach  $20\text{ cm}^{-1}$  [30], i.e. 1% of its value. The anharmonicity of  
 370 this mode has been measured in various environments by means of oscillations  
 observed in photon echo signals and has been assigned to quantum beats between  
 the  $v = 0 \leftrightarrow v = 1$  and  $v = 1 \leftrightarrow v = 2$  transitions [11]. The anharmonicity  
 value measured in  $N_2$  is  $14.7\text{ cm}^{-1}$  [14],  $14.5\text{ cm}^{-1}$  in  $CH_4$  [15],  $14.5\text{ cm}^{-1}$  in




Figure 7: (a) IR spectra of  $W(CO)_6$  recorded in the region of  $T_{1u}$  mode in, from top to bottom,  $CH_4$ , Kr,  $N_2$ , Ar,  $CF_4$ ,  $pH_2$  and Ne. Temperature is 20 K for all solids except in  $pH_2$  and Ne where it is 3 K. The gas phase frequency ( $1998\text{ cm}^{-1}$ ) is marked with a dotted line. (b)  $\nu$  wavenumber (in  $cm^{-1}$ ) of the  $T_{1u}$  mode of  $W(CO)_6$  in the gas phase, Ne,  $pH_2$ , Ar, Kr,  $N_2$ ,  $CH_4$  and  $CF_4$  plotted as a function of the polarizability  $\alpha$  of the host ( $\text{\AA}^3$ ).

xerogels [16],  $14.7 \text{ cm}^{-1}$  in dibutylphthalate [11]. These values differ by 1.5%, to  
375 be compared with the 0.2% dispersion found for the corresponding fundamental  
frequencies.

#### 4.2. Population relaxation

Vibrational relaxation of  $\text{W}(\text{CO})_6$  in liquids was found to be strongly sol-  
vent dependent [32] because the vibrational pattern of the solvent can give  
380 intermediate levels in the vibrational relaxation depending on the vibrational  
energies of its modes compared to the probed  $T_{1u}$  mode of  $\text{W}(\text{CO})_6$ . The same  
is observed in cryogenic solids. In  $\text{CD}_4$  a vibrational mode of the host is in  
quasi-resonance with the  $T_{1u}$  mode yielding a reduced  $T_1$  value of 10 ps. We  
also suggest that the vibron of  $\text{CH}_4$  at  $1538 \text{ cm}^{-1}$  can be efficiently involved  
385 in the population relaxation of  $\text{W}(\text{CO})_6$  in high symmetry sites because this  
energy corresponds to the energy gap between the probed  $T_{1u}$  mode and low  
energy modes of  $\text{W}(\text{CO})_6$  around  $400 \text{ cm}^{-1}$ , leading also to a strongly reduced  
 $T_1$  time [15]. In all the other cases studied,  $T_1$  of 100-200 ps has been measured.  
In particular,  $T_1$  is not shortened in phase II of  $\text{CH}_4$  nor in low symmetry sites  
390 despite a rich vibrational pattern below  $1900 \text{ cm}^{-1}$ . At low temperature, the  
population relaxation pathway needs to include vibrational levels of the host of  
rather precise energies to be efficient. The same remark is verified in the case  
of  $\text{CF}_4$ . The solid possesses indeed only vibrational fundamental modes of low  
energies ( $\leq 1320 \text{ cm}^{-1}$ ) [23] which cannot fill the gap between the  $T_{1u}$  mode  
395 and low energy modes of  $\text{W}(\text{CO})_6$  [33].

The temperature dependence of the long  $T_1$  (100-200 ps) has been difficult  
to measure, with very weak changes in the short temperature range accessible  
in the experiments. Because of the experimental difficulties to obtain precise  
measurements (low laser and sample stabilities in long acquisitions of hundreds  
400 of ps), signal changes are usually in the measurement errors. A slight effect  
has been measured in Ar without the possibility to extract a temperature law.  
It means that lattice phonons are rather inefficient in the relaxation pathway,  
which is expected since the order of magnitude of their energy is once or twice

lower than the probed vibration.

405 On the other hand, 100-200 ps values are short compared with  $T_1 = 800$   
ps obtained in  $\text{CCl}_4$  at room temperature [29]. There is no obvious reason  
for  $T_1$  times to be shorter in matrices at temperatures below 20 K, especially  
in rare gas solids or  $\text{N}_2$  with no vibrational levels below  $1900 \text{ cm}^{-1}$ . In these  
solids, a longer time ( $> 1 \text{ ns}$ ) has been measured and assigned to  $T_g$  recovery  
410 time. This value compares well with the long  $T_1$  recorded in  $\text{CCl}_4$ . Assuming  
that this  $T_g$  time would correspond to the lifetime of  $v = 1$  in these cryogenic  
solids, the shorter  $T_1$  would then be assigned to the population relaxation time  
of the component of the  $T_{1u}$  mode probed in the four wave mixing experiment,  
i.e. the time to leave the level probed by the first pulse and probably to reach  
415 another component of the  $T_{1u}$  mode (through intra molecular energy transfer).  
Components of the  $T_{1u}$  mode differ by less than few wavenumbers, which is  
lower than energies of phonons. It explains that this vibrational energy transfer  
is itself inefficient. Considering that  $T_g$  times are  $T_{1u}$  lifetimes in rare gas and  
 $\text{N}_2$  solids, our experiments therefore show that the lifetime in  $\text{CH}_4$  and in  $\text{CF}_4$  is  
420 reduced (no long  $T_g$  has been detected) as expected because of the existence of  
vibrational mode of medium energies. The  $T_1$  value ( $< 100 \text{ ps}$ ) found in xerogels  
at low temperature is consistent with this latter assumption [16].

#### 4.3. Dephasing processes

Photon echo experiments give a direct access to the vibrational dephasing  
425 times. Thanks to the measurements of the population relaxation times, pure  
dephasing times are extracted by means of eq. 1. From the results reported in  
cryogenic solids, there are two kinds of environments where dephasing processes  
are governed by population relaxation. The first one is rare gas solids (Ar  
and Kr). Both  $T_1$  and  $T_2$  times are in the hundreds of ps in the explored  
430 temperature range. Because of quite large measurement errors, it is not possible  
to extract valuable values of  $T_2^*$  times, but in any case, at any temperature, we  
can conclude that  $T_2^*$  is longer than twice  $T_1$ . Population relaxation is fast in  
the second kind of environment as in  $\text{CD}_4$ , where  $T_1 \lesssim 10 \text{ ps}$  and  $T_2 \approx 10 \text{ ps}$ ,






Figure 8: Plot of  $\gamma_{homo}$  as a function of temperature in Kr rare gas solid (triangles) and in N<sub>2</sub> (squares), CH<sub>4</sub> (circles), CF<sub>4</sub> (stars) molecular hosts.

giving  $T_2^*$  longer than 20 ps, i.e. longer than twice  $T_1$ . In the high symmetry  
 435 sites of CH<sub>4</sub> in phase I,  $T_2$  cannot be measured separately from the response of  
 low symmetry sites but our signals are in full agreement with estimations of  $T_2$   
 and  $T_2^*$  equal to the above values in CD<sub>4</sub>. Otherwise,  $T_2$  and  $T_2^*$  in CH<sub>4</sub> out of  
 these specific environments are always longer than 20 ps. In the other cryogenic  
 environments,  $T_2$  is found shorter than  $T_1$  and efficient pure dephasing processes  
 440 occur ( $T_2^* < T_1$ ) due to the environment. The first important remark is that it  
 happens only with molecular hosts.

The temperature dependence of  $T_2$  and  $T_2^*$  documents the processes at play  
 in the vibrational dephasing. This temperature dependence is clearly observed  
 in N<sub>2</sub> and CH<sub>4</sub>. In N<sub>2</sub>, it concerns the temperature evolution of dephasing time  
 445 in high symmetry sites. It has not been possible to extract a clear temperature  
 dependence of  $T_2$  or  $T_2^*$  in the other sites whereas  $T_1$  and  $T_2$  values clearly  
 indicate that pure dephasing processes occur also in these sites. In contrast in  
 CH<sub>4</sub>, the temperature behavior can be followed in low symmetry sites because  
 fast population relaxation at high temperatures in high symmetry sites hides  
 450 pure dephasing processes. As in N<sub>2</sub> and CH<sub>4</sub>, pure dephasing processes are also

at play in  $\text{CF}_4$  solid. Figure 8 displays the evolution of  $\gamma_{\text{homo}}$ , linked to  $T_2$ , against temperature in high symmetry sites of  $\text{N}_2$  and low symmetry sites of  $\text{CH}_4$  compared with  $\text{CF}_4$  and Kr rare gas solid. In the latter increase of  $\gamma_{\text{homo}}$  with temperature is very weak, being more or less included in the error bars. It means that phonons related to the hosts translational motion are hardly coupled with the vibrational motion of the guest. In contrast, the strong temperature dependence in high symmetry sites of  $\text{N}_2$  where the dephasing processes at play are due to coupling with a librational mode of the solid. The behavior observed in  $\text{CH}_4$  and  $\text{CF}_4$  lies between the two cases, showing the clear influence of the molecular motions in the dephasing processes.

## 5. Conclusions

By probing the vibrational properties of a guest molecule ( $\text{W}(\text{CO})_6$ ) in diverse cryogenic lattices, we have obtained several results highlighting the specificities of the different environments explored. The IR absorption of  $\text{W}(\text{CO})_6$  always exhibits a structured feature. It is a signature of the limited number of families of trapping sites due to the crystalline environment. The simplest structure is observed in solid  $\text{pH}_2$ , a solid with specific properties minimizing site effects. Band broadening of each structure in  $\text{pH}_2$  is expected to be homogeneous but no time-resolved experiment has yet been measured to confirm this assumption. In other cryogenic matrices, some substructures possess narrow linewidths ( $\approx 0.5 \text{ cm}^{-1}$ ) but time-resolved experiments always show a significant inhomogeneous contribution to broadening. However strong homogeneous broadening is clearly observed in the absorption when temperature increases in a fraction of  $\text{N}_2$  trapping sites.

Relaxation processes are found to be host-dependent and they are moreover site dependent as proved in  $\text{N}_2$  or  $\text{CH}_4$ . Except in the case of resonances between the probed vibration and the energy levels in the host + guest system,  $T_1$  population relaxation times are long. In fact, values of 100-200 ps may reflect vibrational energy transfers between components of the  $T_{1u}$  mode of  $\text{W}(\text{CO})_6$

480 when a very long time in the ns range is detected, that would correspond to the  
v = 1 lifetime (as observed in rare gas and N<sub>2</sub> solids). Dephasing is enhanced  
in molecular hosts probably because of coupling between molecular motions.  
Acoustic phonons turn out to be very inefficient in the dephasing process, in  
contrast with phonons coming from rotational or vibrational motions of the  
485 molecular hosts. Librons and vibrons affect the vibrational dynamics and their  
effect appears to be strong in sites where the lattice is weakly perturbed, i.e.  
high symmetry sites.

### Acknowledgements

This research was supported by the RTRA Triangle de la Physique grants no.  
490 2010-004-T NOSTADYNE, 2012-053T DYMA and 2013-0436T REACMAQ.  
The authors are grateful to Justinas Ceponkus for his help in the parahydrogen  
measurements. They also thank Bernard Bourguignon and Aimeric Ouvrard for  
the use of femtosecond laser system and their technical support.

### References

- 495 [1] R. M. Hochstrasser, Special feature on Multidimensional ultrafast spec-  
troscopy, *Proceedings of the National Academy of Sciences* 104 (36) (2007)  
14190.
- [2] H. S. Chung, Z. Ganim, K. C. Jones, A. Tokmakoff, Transient 2D IR  
spectroscopy of ubiquitin unfolding dynamics., *Proceedings of the National*  
500 *Academy of Sciences of the United States of America* 104 (36) (2007) 14237–  
42.
- [3] E. H. G. Backus, R. Bloem, P. M. Donaldson, J. A. Jhalainen, R. Pfister,  
B. Paoli, A. Caffisch, P. Hamm, 2D-IR study of a photoswitchable isotope-  
505 labeled alpha-helix, *Journal of Physical Chemistry B* 114 (10) (2010) 3735–  
3740.

- [4] A. Ghosh, J.-J. Ho, A. L. Serrano, D. R. Skoff, T. Zhang, M. T. Zanni, Two-dimensional sum-frequency generation (2D SFG) spectroscopy: summary of principles and its application to amyloid fiber monolayers, *Faraday discussions* 177 (2015) 493–505.
- 510 [5] M. R. Panman, C. N. van Dijk, H. Meuzelaar, S. Woutersen, Communication: Nanosecond folding dynamics of an alpha helix: Time-dependent 2D-IR cross peaks observed using polarization-sensitive dispersed pump-probe spectroscopy, *The Journal of Chemical Physics* 142 (4) (2015) 041103.
- [6] P. M. Donaldson, P. Hamm, Gold nanoparticle capping layers: Structure, dynamics, and surface enhancement measured using 2D-IR spectroscopy, *Angewandte Chemie - International Edition* 52 (2) (2013) 634–638.
- 515 [7] R. D. Mehlenbacher, M.-Y. Wu, M. Grechko, J. E. Laaser, M. S. Arnold, M. T. Zanni, Photoexcitation dynamics of coupled semiconducting carbon nanotube thin films., *Nano Letters* 13 (4) (2013) 1495–501.
- [8] M. Grechko, Y. Ye, R. D. Mehlenbacher, T. J. McDonough, M.-Y. Wu, R. M. Jacobberger, M. S. Arnold, M. T. Zanni, Diffusion-Assisted Photoexcitation Transfer in Coupled Semiconducting Carbon Nanotube Thin Films, *Nature Communications* 8 (6) (2014) 5383–5394.
- 520 [9] W. H. Hesselink, D. A. Wiersma, Picosecond Photon Echoes Stimulated from an Accumulated Grating, *Physical Review Letters* 43 (27) (1979) 1991–1994.
- [10] M. C. Asplund, M. T. Zanni, R. M. Hochstrasser, Two-dimensional infrared spectroscopy of peptides by phase-controlled femtosecond vibrational photon echoes., *Proceedings of the National Academy of Sciences of the United States of America* 97 (15) (2000) 8219–8224.
- 530 [11] A. Tokmakoff, D. Zimdars, R. S. Urdahl, R. S. Francis, A. S. Kwok, M. D. Fayer, Infrared vibrational photon echo experiments in liquids and glasses, *The Journal of Physical Chemistry* 99 (36) (1995) 13310–13320.

- [12] K. D. Rector, M. D. Fayer, Vibrational echoes: A new approach to  
535 condensed-matter vibrational spectroscopy, *International Reviews in Physical Chemistry* 17 (3) (1998) 261–306.
- [13] M. Broquier, C. Crépin, H. Dubost, J. P. Galaup, IR spectra and vibrational dephasing of the CO stretching mode in W(CO)<sub>6</sub> doped cryogenic matrices, *Chemical Physics* 341 (1-3) (2007) 207–217.
- 540 [14] R. Thon, W. Chin, J.-P. Galaup, A. Ouvrard, B. Bourguignon, C. Crépin, Vibrational Perturbations of W(CO)<sub>6</sub> Trapped in a Molecular Lattice Probed by Linear and Nonlinear Spectroscopy., *The Journal of Physical Chemistry A* 117 (2013) 8145–8156.
- [15] R. Thon, W. Chin, D. Chamma, J. P. Galaup, A. Ouvrard, B. Bourguignon, C. Crépin, Vibrational spectroscopy and dynamics of W(CO)<sub>6</sub>  
545 in solid methane as a probe of lattice properties, *The Journal Chemical Physics* submitted.
- [16] J. P. Galaup, M. Broquier, C. Crépin, H. Dubost, J. M. Ortega, F. Chaput, J. P. Boilot, Vibrational dynamics of CO stretching in W(CO)<sub>6</sub>-doped hybrid xerogels from 5 K to room temperature with the CLIO-FEL, *Journal of Luminescence* 86 (3) (2000) 363–370.  
550
- [17] J. Ceponkus, W. Chin, M. Chevalier, M. Broquier, A. Limongi, C. Crépin, Infrared study of glycolaldehyde isolated in parahydrogen matrix., *The Journal of Chemical Physics* 133 (9) (2010) 094502.
- 555 [18] L. H. Jones, Vibrational spectra and force constants of the hexacarbonyls of chromium, molybdenum and tungsten, *Spectrochimica Acta* 19 (1) (1963) 329–338.
- [19] W. Press, A. Kollmar, CH<sub>4</sub>: tunneling states, rotations, and phase transition in a quantum molecular crystal, *Solid State Communications* 17 (1975)  
560 405–408.

- [20] D. G. Bounds, M. L. Klein, G. N. Patey, Molecular dynamics simulation of the plastic phase of solid methane, *The Journal of Chemical Physics* 72 (10) (1980) 5348.
- [21] P. Hamm, Coherent effects in femtosecond infrared spectroscopy, *Chemical Physics* 200 (3) (1995) 415–429.
- [22] Y. A. Sataty, A. Ron, F. H. Herbstein, Carbon tetrafluoride - phase II: Far infrared spectrum and crystal structure, *The Journal Chemical Physics* 62 (3) (1975) 1094–1097.
- [23] R. P. Fournier, R. Savoie, F. Bessette, A. Cabana, Vibrational Spectra of Liquid and Crystalline CF<sub>4</sub>, *The Journal of Chemical Physics* 49 (3) (1968) 1159–1164.
- [24] A. N. Fitch, J. K. Cockcroft, The structure of solid carbon tetrafluoride, *Zeitschrift fur Kristallographie* 203 (1993) 29–39.
- [25] I. F. Silvera, The solid molecular hydrogens in the condensed phase: Fundamentals and static properties, *Reviews of Modern Physics* 52 (2) (1980) 393–452.
- [26] T. Momose, H. Katsuki, H. Hoshina, N. Sogoshi, High-resolution laser spectroscopy of methane clusters trapped in solid parahydrogen, *The Journal of Chemical Physics* 107 (19) (1997) 7717–7720.
- [27] K. Yoshioka, D. T. Anderson, Infrared spectra of CH<sub>3</sub>F(ortho-H<sub>2</sub>)<sub>n</sub> clusters in solid parahydrogen, *The Journal of Chemical Physics* 119 (9) (2003) 4731–4742.
- [28] T. Momose, H. Hoshina, M. Fushitani, H. Katsuki, High-resolution spectroscopy and the analysis of ro-vibrational transitions of molecules in solid parahydrogen, *Vibrational Spectroscopy* 34 (1) (2004) 95–108.
- [29] A. Tokmakoff, B. Sauter, M. D. Fayer, Temperature-Dependent Vibrational-Relaxation in Polyatomic Liquids: Picosecond Infrared Pump-

Probe Experiments, *The Journal of Chemical Physics* 100 (12) (1994) 9035–9043.

- 590 [30] E. Heilweil, R. Cavanagh, J. Stephenson, Population relaxation of CO ( $\nu=1$ ) vibrations in solution phase metal carbonyl complexes, *Chemical Physics Letters* 134 (2) (1987) 181–188.
- [31] A. Tokmakoff, R. S. Urdahl, D. Zimdars, R. S. Francis, A. S. Kwok, M. D. Fayer, Vibrational spectral diffusion and population dynamics in a glass-forming liquid: Variable bandwidth picosecond infrared spectroscopy, *The Journal of Chemical Physics* 102 (10) (1995) 3919.
- 595 [32] A. Tokmakoff, M. D. Fayer, Homogeneous Vibrational Dynamics and Inhomogeneous Broadening in Glass-Forming Liquids - Infrared Photon-Echo Experiments from Room-Temperature to 10 K, *The Journal of Chemical Physics* 103 (8) (1995) 2810–2826.
- 600 [33] R. Amster, R. Hannan, M. Tobin, Vibrational spectra of some group VI hexacarbonyls, *Spectrochimica Acta* 19 (9) (1963) 1489–1494.

### Figure captions

Figure 1 IR spectra of  $\text{W}(\text{CO})_6$  in (a) Kr (35 K), (b) Ne (3 K) and (c)  $\text{pH}_2$  (3 K). The inset shows a schematic of  $\text{W}(\text{CO})_6$  (W in blue, C in grey, O in red).

605

Figure 2 IR spectra of  $\text{W}(\text{CO})_6$  in  $\text{N}_2$  at 6 and 22 K and  $\text{CH}_4$  at 8 and 22 K. Low and high temperatures are respectively plotted as black and red traces.

610 Figure 3  $S(T)$  signals (normalized) of  $\text{W}(\text{CO})_6$  in (a)  $\text{N}_2$  at 20 K and (b)  $\text{CH}_4$  at 14 K (black) and 25 K (red). The very steep rise (“spike”) at  $T = 0$  is due to additional nonlinear signals produced when the IR pulses simultaneously arrive onto the sample. It is not related to the probed dynamics [21].

615 Figure 4  $S(\tau)$  signals (normalized) of  $W(CO)_6$  in (a)  $N_2$  and (b)  $CH_4$ . Red circles and black squares represent signals at high (20 and 24 K) and low (8 and 9 K) temperatures respectively.

Figure 5 IR spectra of  $W(CO)_6$  in  $CF_4$  from 10 K to 40 K (top to bottom).

620

Figure 6 Examples of time-resolved signals in  $CF_4$  doped with  $W(CO)_6$  (arbitrary units), (a)  $S(\tau)$  at 30 K,  $T = 20$  ps, (b)  $S(T)$  at 35 K ("spike" at the origin as in Figure 3a).

625 Figure 7 (a) IR spectra of  $W(CO)_6$  recorded in the region of  $T_{1u}$  mode in, from top to bottom,  $CH_4$ , Kr,  $N_2$ , Ar,  $CF_4$ ,  $pH_2$  and Ne. Temperature is 20 K for all solids except in  $pH_2$  and Ne where it is 3 K. The gas phase frequency ( $1998\text{ cm}^{-1}$ ) is marked with a dotted line. (b)  $\nu$  wavenumber (in  $cm^{-1}$ ) of the  $T_{1u}$  mode of  $W(CO)_6$  in the gas phase, Ne,  $pH_2$ , Ar, Kr,  $N_2$ ,  $CH_4$  and  $CF_4$  plotted  
630 as a function of the polarizability  $\alpha$  of the host ( $\text{\AA}^3$ ).

Figure 8 Plot of  $\gamma_{homo}$  as a function of temperature in Kr rare gas solid (triangles) and in  $N_2$  (squares),  $CH_4$  (circles),  $CF_4$  (stars) molecular hosts.