



HAL
open science

Metal-free nucleophilic trifluoromethylselenolation via an iodide-mediated umpolung reactivity of trifluoromethylselenotoluenesulfonate

Kevin Grollier, Alexis Taponard, Arnaud de Zordo-Banliat, Emmanuel Magnier,
Thierry Billard

► To cite this version:

Kevin Grollier, Alexis Taponard, Arnaud de Zordo-Banliat, Emmanuel Magnier, Thierry Billard. Metal-free nucleophilic trifluoromethylselenolation via an iodide-mediated umpolung reactivity of trifluoromethylselenotoluenesulfonate. *Beilstein Journal of Organic Chemistry*, 2020, 16, pp.3032 - 3037. <10.3762/bjoc.16.252>. <hal-03097741>

HAL Id: hal-03097741

<https://hal.science/hal-03097741v1>

Submitted on 5 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



Metal-free nucleophilic trifluoromethylselenolation via an iodide-mediated umpolung reactivity of trifluoromethylselenotoluenesulfonate

Kevin Grollier^{‡1}, Alexis Taponard^{‡1}, Arnaud De Zordo-Banliat², Emmanuel Magnier² and Thierry Billard^{*1,3}

Full Research Paper

Open Access

Address:

¹Institute of Chemistry and Biochemistry (ICBMS-UMR CNRS 5246), Univ Lyon, Université Lyon 1, CNRS, CPE, INSA, 43 Bd du 11 novembre 1918, 69622 Villeurbanne, France, ²Université Paris-Saclay, UVSQ, CNRS, UMR 8180, Institut Lavoisier de Versailles, 78035 Versailles Cedex, France and ³CERMEP-In vivo imaging, Groupement Hospitalier Est, 59 Bd Pinel, 69677 Lyon, France

Email:

Thierry Billard^{*} - Thierry.billard@univ-lyon1.fr

* Corresponding author ‡ Equal contributors

Keywords:

fluorine; nucleophilic substitution; perfluoroalkylselenolation; selenium; trifluoromethylselenolation

Beilstein J. Org. Chem. **2020**, *16*, 3032–3037.
<https://doi.org/10.3762/bjoc.16.252>

Received: 23 October 2020

Accepted: 02 December 2020

Published: 10 December 2020

This article is part of the thematic issue "Organo-fluorine chemistry V".

Guest Editor: D. O'Hagan

© 2020 Grollier et al.; licensee Beilstein-Institut.
License and terms: see end of document.

Abstract

We report herein a practical method to generate CF_3Se^- (and R_FSe^-) anions from shelf-stable reagents under iodide activation. Metal-free nucleophilic trifluoromethylselenolations have been then performed with this in situ-generated anion. Perfluoroalkylselenolations have also been described.

Introduction

Because of the peculiar properties of the fluorine atom, fluorinated compounds gained a growing interest over the last decades and found applications in a large panel of fields from materials to life sciences [1-15]. Fluorinated motifs bring to molecules specific and often unique electronic and physico-chemical characteristics. In order to design new substrates with targeted properties, a modulation of the properties of the introduced substituents became fundamental. In this context, the development of innovative fluorinated groups recently emerged,

in particular by combining heteroatoms, such as chalcogens, and fluorinated moieties [16].

Despite, the negative reputation of selenium due to its toxicity at high doses, it is an essential trace element for human physiology and biochemistry [17-20]. Furthermore, selenolated compounds found valuable applications in materials [21-23], life sciences [19,20,24-29], and drug design [30-33]. Consequently, the merging of fluorinated moieties, such as CF_3 with selenium

could constitute an interesting motif in the design of new molecules, in particular in medicinal chemistry or agrochemistry. Even if, to date, there are no CF_3Se -containing pharmaceuticals registered [15], a recent work has demonstrated the promising development of trifluoromethylselenolated nonsteroidal anti-inflammatory drugs as potential anticancer drugs [34].

Over the last years, trifluoromethylselenolation reactions have gained a rising infatuation but, despite this recent interest, methods to introduce the CF_3Se group into organic substrates remain limited [35,36].

One of the simplest ways to achieve trifluoromethylselenolated compounds is the direct nucleophilic substitution of suitable leaving groups to form the $\text{CF}_3\text{Se}-\text{C}(\text{sp}^3)$ bond. This chemistry is the prerogative of the CF_3Se^- anion (Scheme 1) [37-40]. However, the formation of this selenium species requires the tedious use of red elemental selenium [41] and also suffers from stability issues. To circumvent these drawbacks, a copper com-


plex has been developed, but in this case the use of a stoichiometric amount of the metal is required [37-39].

Only a few years ago, trifluoromethylselenotoluenesulfonate (**1a**) has been developed as an efficient reagent to perform electrophilic or radical trifluoromethylselenolations [42-47]. Very recently, we demonstrated that under reductive conditions, such compounds succeeded to perform nucleophilic substitutions [48]. In this reaction, the CF_3Se^- anion was in situ generated by reduction through a double electron transfer of **1a** with TDAE (tetrakis(dimethylamino)ethylene). Even though this umpolung strategy is efficient, the use of the sensitive TDAE, a strong reducing agent, could constitute a drawback for some applications. Consequently, we decided to develop a new umpolung method in non-reductive conditions.

Results and Discussion


A few years ago, we have demonstrated that trifluoromethanesulfenamides, electrophilic trifluoromethylthiolation reagents, could also perform nucleophilic trifluoromethylthiolations through the transient formation of a CF_3SI species which presented an inverted polarity [49,50]. Based on a similar approach, we hypothesized that the CF_3SeI species could also possess the $\text{CF}_3\text{Se}^{\delta-}-\text{I}^{\delta+}$ inverted polarity. Thus, based on the previously developed conditions, reagent **1a** was reacted with benzyl bromide (**2a**) in the presence of tetrabutylammonium iodide (TBAI) in acetone at 40 °C (Table 1, entry 1).

The observed result was moderate (Table 1, entry 1). Other solvents, which led also to satisfactory yields in the “sulfur series” were then tested. Acetonitrile did not improve the yield, howev-



Scheme 1: State of the art concerning the direct nucleophilic trifluoromethylselenolation.

Table 1: Reaction between **1a** and **2a**.



entry	1a (equiv)	TBAI (equiv)	2a (equiv)	solvent	<i>T</i> (°C)	<i>t</i> (h)	3a (%) ^a
1	1	2	1	acetone	40	15	52
2	1	2	1	CH ₃ CN	40	15	54
3	1	2	1	THF	40	15	61
4	1	2	1	THF	40	4	62
5	1	2	1	THF	50	4	59
6	1	2	1	THF	60	4	54
7	1	2	1	THF	25	4	47
8	1.5	3	1	THF	40	4	55
9	1	2	2	THF	40	4	89


^aYields determined by ¹⁹F NMR spectroscopy with PhOCF₃ as an internal standard.

er, a better one was obtained in THF (Table 1, entries 2 and 3). Interestingly, a shorter reaction time (4 h instead of 15 h) provided similar results (Table 1, entries 3 and 4). At higher temperatures, the results were not improved and even a slight decrease of the yield was observed, possibly due to an increased degradation of CF_3SeI or CF_3Se^- (Table 1, entries 5 and 6). A lower temperature also decreased the yield (Table 1, entry 7). With an excess of reagents **1a** and TBAI, leading to an excess of the CF_3Se^- species, no significant improvement was observed (Table 1, entry 8). On contrary, the use of 2 equivalents of the electrophile **2a**, to ameliorate the CF_3Se^- anion trapping, had a significant effect and gave a very good yield of the product (Table 1, entry 9). With the optimal conditions in hand, the reaction was exemplified with various other electrophiles (Scheme 2).


The reaction gave generally good results with reactive electrophiles such as benzylic, allylic or propargylic ones (**3a–k**). Noteworthy, in the reaction with 2-(bromomethyl)pyridine (**2g**)

only 1 equivalent was required, maybe due to a higher reactivity. Furthermore, the reaction seems to be very sensitive to steric hindrance as illustrated by the low yield obtained for **3b**. In contrast, in the aliphatic series, only low yields were observed (**3m,n**) except for the activated α -bromo acetophenone (**3l**). This led us to suppose that CF_3Se^- might be a poor nucleophile, which is confirmed by the medium yield observed with the propargylic substrate **3k**, where the chloride starting material was used instead of the bromide as for the other compounds. Noteworthy, because of the volatility of the obtained products, the isolated yields were sometimes significantly below the NMR yields.

Higher fluorinated homologs of **1a** were also synthesized. Consequently, an extension of this method was considered with pentafluoroethylated and tridecafluorohexylated reagents **1b** and **1c**. Good yields were obtained, in particular for **5a** which constitutes, to the best of our knowledge, the first example of a direct nucleophilic tridecafluorohexylselenolation.



From a mechanistic point of view, a pathway inspired by the reaction described with trifluoromethanesulfenamides was postulated (Scheme 3) [49].



The first equivalent of iodide (from TBAI) reacts with the reagent **1a** to produce the transient species CF_3SeI with an inverted polarity on the selenium atom. This compound then undergoes the attack of the second equivalent of iodide to generate the CF_3Se^- anion with releasing of I_2 . Finally, the nucleophilic CF_3Se^- can substitute the leaving group onto the electrophilic substrate **2**. However, because of the release of I_2 , as side reaction the oxidation of the CF_3Se^- anion can be also envisaged. This was confirmed by the formation of 25–30% of $\text{CF}_3\text{SeSeCF}_3$ when 1 equivalent of **2a** was used (Table 1, entries 3 and 4). Consequently, the nucleophilic substitution is in competition with this relatively fast oxidation. By adding an excess of the electrophile **2**, the substitution is favored detrimentally to the oxidation. Nevertheless, with weaker or hindered electrophiles, the oxidation reaction is favored compared to the slower substitution. This is adequate with the observed results. Noteworthy, the supposed formation of I_2 was strengthened by the appearance of a red-brown color of the reaction media which faded after the addition of sodium thiosulfate. As demonstrated in the sulfur series [49], the in situ formation of an alkyl iodide from **2**, through a Finkelstein reaction, can be also envisaged. This would not impact the reaction pathway since the released bromide can also activate the CF_3SeI species to provide the expected CF_3Se^- anion.

Conclusion

To conclude, trifluoromethylselenotoluenesulfonate confirmed to be a versatile reagent able to perform electrophilic, radical or nucleophilic reactions depending on the conditions. The iodide-mediated, metal-free method is complementary to the previous one using TDAE. Thus, the umpolung reactivity of trifluoromethylselenotoluenesulfonate can be performed under reduc-

tive or oxidative conditions. Furthermore, this method was extended to higher fluorinated homologs allowing the first nucleophilic tridecafluorohexylselenolation.

Experimental

Typical procedure: In a 10 mL flame-dried flask tube equipped with a magnetic stirring bar was added **1a–c** (0.2 mmol, 1 equiv) followed by 0.4 mL of dry THF. Then, compound **2a–n** (0.4 mmol, 2 equiv) was added followed by TBAI (0.4 mmol, 2 equiv). The tube is then sealed and the reaction mixture stirred at 40 °C for 4 h. The conversion was checked by ^{19}F NMR spectroscopy with PhOCF_3 as internal standard. After completion, the reaction mixture was partitioned between Et_2O or pentane and water. The aqueous layer was extracted with Et_2O and pentane and the combined organic layers were dried over MgSO_4 , filtered and concentrated to dryness. The crude residue was purified by chromatography to afford the desired products **3**, **4**, or **5**.

Supporting Information

Supporting Information File 1

Additional experimental and analytical data.

[<https://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-16-252-S1.pdf>]

Funding

This work was supported by a grant from the French National Research Agency (ANR 18-CE07-0039-01). The authors are grateful to the CNRS and the French Ministry of Research for financial support. The French Fluorine Network is also acknowledged for its support.

ORCID® iDs

Kevin Grollier - <https://orcid.org/0000-0003-0850-8653>

Alexis Taponard - <https://orcid.org/0000-0001-8008-2293>

Emmanuel Magnier - <https://orcid.org/0000-0003-3392-3971>

Thierry Billard - <https://orcid.org/0000-0002-2937-9523>

References

- Smart, B. E. *J. Fluorine Chem.* **2001**, *109*, 3–11. doi:10.1016/S0022-1139(01)00375-x
- Kirsch, P. *Modern Fluoroorganic Chemistry*; Wiley-VCH: Weinheim, Germany, 2013. doi:10.1002/9783527651351
- Zhou, Y.; Wang, J.; Gu, Z.; Wang, S.; Zhu, W.; Aceña, J. L.; Soloshonok, V. A.; Izawa, K.; Liu, H. *Chem. Rev.* **2016**, *116*, 422–518. doi:10.1021/acs.chemrev.5b00392
- Meanwell, N. A. *J. Med. Chem.* **2018**, *61*, 5822–5880. doi:10.1021/acs.jmedchem.7b01788

5. Mei, H.; Han, J.; Fustero, S.; Medio-Simon, M.; Sedgwick, D. M.; Santi, C.; Ruzziconi, R.; Soloshonok, V. A. *Chem. – Eur. J.* **2019**, *25*, 11797–11819. doi:10.1002/chem.201901840
6. Han, J.; Remete, A. M.; Dobson, L. S.; Kiss, L.; Izawa, K.; Moriwaki, H.; Soloshonok, V. A.; O'Hagan, D. *J. Fluorine Chem.* **2020**, *239*, 109639. doi:10.1016/j.jfluchem.2020.109639
7. Cheng, M.; Guo, C.; Gross, M. L. *Angew. Chem., Int. Ed.* **2020**, *59*, 5880–5889. doi:10.1002/anie.201907662
8. Johnson, B. M.; Shu, Y.-Z.; Zhuo, X.; Meanwell, N. A. *J. Med. Chem.* **2020**, *63*, 6315–6386. doi:10.1021/acs.jmedchem.9b01877
9. Jeschke, P. *Pest Manage. Sci.* **2010**, *66*, 10–27. doi:10.1002/ps.1829
10. Ogawa, Y.; Tokunaga, E.; Kobayashi, O.; Hirai, K.; Shibata, N. *iScience* **2020**, *23*, 101467. doi:10.1016/j.isci.2020.101467
11. Fujiwara, T.; O'Hagan, D. *J. Fluorine Chem.* **2014**, *167*, 16–29. doi:10.1016/j.jfluchem.2014.06.014
12. Pagliaro, M.; Ciriminna, R. *J. Mater. Chem.* **2005**, *15*, 4981–4991. doi:10.1039/b507583c
13. Berger, R.; Resnati, G.; Metrangolo, P.; Weber, E.; Hulliger, J. *Chem. Soc. Rev.* **2011**, *40*, 3496–3508. doi:10.1039/c0cs00221f
14. Chopra, D.; Row, T. N. G. *CrystEngComm* **2011**, *13*, 2175–2186. doi:10.1039/c0ce00538j
15. Inoue, M.; Sumii, Y.; Shibata, N. *ACS Omega* **2020**, *5*, 10633–10640. doi:10.1021/acsomega.0c00830
16. Cahard, D.; Ma, J.-A., Eds. *Emerging Fluorinated Motifs: Synthesis, Properties and Applications*; Wiley-VCH: Weinheim, Germany, 2020. doi:10.1002/9783527824342
17. EFSA Panel on Dietetic Products, Nutrition and Allergies. *EFSA J.* **2014**, *12*, 3846. doi:10.2903/j.efsa.2014.3846
18. European Food Safety Authority (EFSA). *EFSA Supporting Publ.* **2017**, *14*, e15121E. doi:10.2903/sp.efsa.2017.e15121
19. Rayman, M. P. *Lancet* **2000**, *356*, 233–241. doi:10.1016/s0140-6736(00)02490-9
20. Mousa, R.; Notis Dardashti, R.; Metanis, N. *Angew. Chem., Int. Ed.* **2017**, *56*, 15818–15827. doi:10.1002/anie.201706876
21. Romashov, L. V.; Ananikov, V. P. *Chem. – Eur. J.* **2013**, *19*, 17640–17660. doi:10.1002/chem.201302115
22. Guo, W.; Fu, Y. *Chem. – Eur. J.* **2020**, *26*, 13322–13331. doi:10.1002/chem.202000878
23. Li, Q.; Zhang, Y.; Chen, Z.; Pan, X.; Zhang, Z.; Zhu, J.; Zhu, X. *Org. Chem. Front.* **2020**, *7*, 2815–2841. doi:10.1039/d0qo00640h
24. Mughesh, G.; du Mont, W.-W.; Sies, H. *Chem. Rev.* **2001**, *101*, 2125–2180. doi:10.1021/cr000426w
25. Rocha, J. B. T.; Piccoli, B. C.; Oliveira, C. S. *ARKIVOC* **2017**, No. ii, 457–491. doi:10.24820/ark.5550190.p009.784
26. Short, S. P.; Pilat, J. M.; Williams, C. S. *Free Radical Biol. Med.* **2018**, *127*, 26–35. doi:10.1016/j.freeradbiomed.2018.05.066
27. Solov'yev, N.; Drobyshev, E.; Bjørklund, G.; Dubrovskii, Y.; Lysiuk, R.; Rayman, M. P. *Free Radical Biol. Med.* **2018**, *127*, 124–133. doi:10.1016/j.freeradbiomed.2018.02.030
28. Antoniadou, I.; Kouskou, M.; Arsiwala, T.; Singh, N.; Vasudevan, S. R.; Fowler, T.; Cadirci, E.; Churchill, G. C.; Sharp, T. *Br. J. Pharmacol.* **2018**, *175*, 2599–2610. doi:10.1111/bph.14179
29. Jain, V. K.; Priyadarsini, K. I., Eds. *Organoselenium Compounds in Biology and Medicine: Synthesis, Biological and Therapeutic Treatments*; Royal Society of Chemistry: Cambridge, U.K., 2018. doi:10.1039/9781788011907
30. Singh, N.; Halliday, A. C.; Thomas, J. M.; Kuznetsova, O. V.; Baldwin, R.; Woon, E. C. Y.; Aley, P. K.; Antoniadou, I.; Sharp, T.; Vasudevan, S. R.; Churchill, G. C. *Nat. Commun.* **2013**, *4*, 1332. doi:10.1038/ncomms2320
31. Thangamani, S.; Younis, W.; Seleem, M. N. *Sci. Rep.* **2015**, *5*, 11596. doi:10.1038/srep11596
32. Gandin, V.; Khalkar, P.; Braude, J.; Fernandes, A. P. *Free Radical Biol. Med.* **2018**, *127*, 80–97. doi:10.1016/j.freeradbiomed.2018.05.001
33. Alcolea, V.; Pérez-Silanes, S. *Eur. J. Med. Chem.* **2020**, *206*, 112673. doi:10.1016/j.ejmech.2020.112673
34. He, X.; Zhong, M.; Li, S.; Li, X.; Li, Y.; Li, Z.; Gao, Y.; Ding, F.; Wen, D.; Lei, Y.; Zhang, Y. *Eur. J. Med. Chem.* **2020**, *208*, 112864. doi:10.1016/j.ejmech.2020.112864
35. Tlili, A.; Ismalaj, E.; Glenadel, Q.; Ghiazza, C.; Billard, T. *Chem. – Eur. J.* **2018**, *24*, 3659–3670. doi:10.1002/chem.201704637
36. Billard, T.; Toulgoat, F. When Fluorine Meets Selenium. In *Emerging Fluorinated Motifs: Synthesis, Properties and Applications*; Cahard, D.; Ma, J.-A., Eds.; Wiley-VCH: Weinheim, Germany, 2020; pp 691–721. doi:10.1002/9783527824342.ch23
37. Chen, C.; Ouyang, L.; Lin, Q.; Liu, Y.; Hou, C.; Yuan, Y.; Weng, Z. *Chem. – Eur. J.* **2014**, *20*, 657–661. doi:10.1002/chem.201303934
38. Rong, M.; Huang, R.; You, Y.; Weng, Z. *Tetrahedron* **2014**, *70*, 8872–8878. doi:10.1016/j.tet.2014.09.091
39. Yang, Y.; Lin, X.; Zheng, Z.; Lin, G.; Zhang, Y.; You, Y.; Weng, Z. *J. Fluorine Chem.* **2017**, *204*, 1–5. doi:10.1016/j.jfluchem.2017.10.001
40. Dong, T.; He, J.; Li, Z.-H.; Zhang, C.-P. *ACS Sustainable Chem. Eng.* **2018**, *6*, 1327–1335. doi:10.1021/acssuschemeng.7b03673
41. Tyrre, W.; Naumann, D.; Yagupolskii, Y. L. *J. Fluorine Chem.* **2003**, *123*, 183–187. doi:10.1016/s0022-1139(03)00118-0
42. Glenadel, Q.; Ghiazza, C.; Tlili, A.; Billard, T. *Adv. Synth. Catal.* **2017**, *359*, 3414–3420. doi:10.1002/adsc.201700904
43. Ghiazza, C.; Tlili, A.; Billard, T. *Beilstein J. Org. Chem.* **2017**, *13*, 2626–2630. doi:10.3762/bjoc.13.260
44. Ghiazza, C.; Debrauwer, V.; Billard, T.; Tlili, A. *Chem. – Eur. J.* **2018**, *24*, 97–100. doi:10.1002/chem.201705231
45. Ghiazza, C.; Khrouz, L.; Monnereau, C.; Billard, T.; Tlili, A. *Chem. Commun.* **2018**, *54*, 9909–9912. doi:10.1039/c8cc05256e
46. Ghiazza, C.; Debrauwer, V.; Monnereau, C.; Khrouz, L.; Médebielle, M.; Billard, T.; Tlili, A. *Angew. Chem., Int. Ed.* **2018**, *57*, 11781–11785. doi:10.1002/anie.201806165
47. Ghiazza, C.; Monnereau, C.; Khrouz, L.; Médebielle, M.; Billard, T.; Tlili, A. *Synlett* **2019**, *30*, 777–782. doi:10.1055/s-0037-1610347
48. Ghiazza, C.; Kataria, A.; Tlili, A.; Toulgoat, F.; Billard, T. *Asian J. Org. Chem.* **2019**, *8*, 675–678. doi:10.1002/ajoc.201900027
49. Glenadel, Q.; Bordy, M.; Alazet, S.; Tlili, A.; Billard, T. *Asian J. Org. Chem.* **2016**, *5*, 428–433. doi:10.1002/ajoc.201600003
50. Glenadel, Q.; Tlili, A.; Billard, T. *Eur. J. Org. Chem.* **2016**, 1955–1957. doi:10.1002/ejoc.201600197

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0>). Please note that the reuse, redistribution and reproduction in particular requires that the author(s) and source are credited and that individual graphics may be subject to special legal provisions.

The license is subject to the *Beilstein Journal of Organic Chemistry* terms and conditions: (<https://www.beilstein-journals.org/bjoc/terms>)

The definitive version of this article is the electronic one which can be found at:
<https://doi.org/10.3762/bjoc.16.252>