



HAL
open science

Exploring the Limits of π -Acid Catalysis Using Strongly Electrophilic Main Group Metal Complexes: The Case of Zinc and Aluminium

Jiixin Tian, Yan Chen, Marie Vayer, Alexandre Djurovic, Régis Guillot, Refka Guermazi, Samuel Dagorne, Christophe Bour, Vincent Gandon

► **To cite this version:**

Jiixin Tian, Yan Chen, Marie Vayer, Alexandre Djurovic, Régis Guillot, et al.. Exploring the Limits of π -Acid Catalysis Using Strongly Electrophilic Main Group Metal Complexes: The Case of Zinc and Aluminium. *Chemistry - A European Journal*, 2020, 26 (56), pp.12831-12838. 10.1002/chem.202001376 . hal-03097839

HAL Id: hal-03097839

<https://hal.science/hal-03097839>

Submitted on 4 Feb 2022


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Exploring the Limits of π -Acid Catalysis Using Strongly Electrophilic Main Group Metal Complexes: the case of Zinc and Aluminum

Jiaxin Tian,^[a] Yan Chen,^[a] Marie Vayer,^[a] Alexandre Djurovic,^[a] Régis Guillot,^[a] Refka Guermazi,^[b] Samuel Dagorne,^[b] Christophe Bour,^{*[a]} and Vincent Gandon^{*[a,c]}

Abstract: The catalytic activity of cationic NHC-Zn(II) and NHC-Al(III) complexes in reactions that require the electrophilic activation of soft C-C π bonds has been studied. The former proved able to act as a soft π -Lewis acid in a variety of transformations. The benefit of the bulky IPr NHC ligand was demonstrated by comparison with simple ZnX_2 salts. The tested NHC-Al(III) catalyst is not able to activate C-C π bonds but simple AlX_2^+ ions were found potent in some cases.




Scheme 1. Nonclassical carbocation intermediates in π -acid-catalyzed transformations of enynes

Introduction

π -Acid catalysis, i.e. the electrophilic activation of C-C π bonds with a Lewis acid,^[1] is a powerful way to transform alkyne, alkene or allene derivatives into valuable building blocks. Gold and platinum complexes have been the most versatile π -acids used so far.^[2] Other elements such as gallium and indium are also well-known in this field.^[3] In fact, not all π -activators behave the same way. Whereas a simple proton can promote the addition of water to alkynes through a vinyl cation intermediate, what makes gold, platinum, gallium and indium complexes rather unique is their capacity to mediate the addition of a C-C π bond to another C-C π bond through a nonclassical carbocation.^{[2], [4]} The cycloisomerization of enynes is a prototypical example of such a reactivity (Scheme 1, eq 1).^{[2a], [3c]} The classification of Lewis acids between the softest ones, which will be good for mediating enyne cycloisomerizations, and harder ones, that will be more potent for C-C π bond functionalization through classical carbocations, can be easily made by using the cycloheptatriene test that we developed.^[5] As described by Echavarren et al, 7-alkynylcycloheptatrienes are treated like enynes by gold(I) complexes (Scheme 1, eq 2)).^[6]

Likewise, we reported that the reaction of compound **1** with catalysts of the gold, platinum, mercury, gallium and indium series leads to its skeletal reorganization into the regioisomeric indenenes **2** and **3** in 1,2-dichloroethane (DCE) or toluene (Scheme 2). With harder Lewis acids of the silver, gallium, copper, and calcium series, or even a Brønsted acid such as HOTf, the bicyclic product **4** is obtained. This occurs through the rearrangement of the 7-alkynylcycloheptatriene moiety into a phenylallene,^[7] which undergoes a rapid hydroarylation by the pendant phenyl group. In DCE, a solvent that can give rise to superacids such as $HsBf_6$ in the presence of $AgSbF_6$ or SbF_5 ,^{[8], [9]} or by using $HNTf_2$, the conversion of **4** into the tricyclic compound **5** is observed with time, which corresponds to the hydroarylation of the styrene moiety through a benzylic carbocation.^[5]



Scheme 2. Isomerization pathways of the 7-alkynylcycloheptatriene compound **1** (E = CO_2Me ; LA = Lewis Acid; DCE = 1,2-dichloroethane)

[a] J. Tian,[†] Y. Chen,[†] Dr M. Vayer, Dr C. Bour, Prof. V. Gandon
Institut de Chimie Moléculaire et des Matériaux d'Orsay (ICMMO),
CNRS UMR 8182, Université Paris-Saclay, Bâtiment 420, 91405
Orsay cedex, France.

E-mail: vincent.gandon@universite-paris-saclay.fr

[†] these authors contributed equally to this work

[b] Dr. S. Dagorne, Dr. R. Guermazi
Institut de Chimie de Strasbourg, CNRS-Université de Strasbourg, 1
rue Blaise Pascal, 67000 Strasbourg, France.

[c] Prof. V. Gandon
Laboratoire de Chimie Moléculaire (LCM), CNRS UMR 9168, Ecole
Polytechnique, Institut Polytechnique de Paris, route de Saclay,
91128 Palaiseau cedex, France.

Supporting information for this article is given via a link at the end of the document.

One interest of this test is to identify new soft π -acids, as we did in the gallium(I) and indium(I) series.^{[5],[10]} Another interest is to show the efficiency of an anion exchange. Indeed, since silver salts such as AgSbF₆, which is a widely used halide abstractor, gives a different result than the softest Lewis acids (**4** vs **2/3**), this test allows to validate the formation of the desired cationic species (such as IPr-InBr₂⁺ from IPr-InBr₃ and AgSbF₆). Lastly, this test informs if the catalytic conditions release protons (**5**).


In this study, we have used the cycloheptatriene test and other reactions to analyze the catalytic behavior of other complexes of the main group metal series. We were especially interested in broadening the scope of application of π -acid catalysis to inexpensive metals such as zinc and aluminum. Regarding zinc, only simple salts of type ZnX₂ have been used so far as π -acids.^[11] We have studied the case of the NHC-stabilized zinc(II) species [IPr-Zn(C₆F₅)⁺][B(C₆F₅)₄]⁻ (Table 1, **B**) and IPr-ZnBr₂(THF) (Table 1, **C**). Synthesized by some of us,^[12] the aryl-zinc complex **B** has a higher fluoride ion affinity (FIA) than B(C₆F₅)₃ and efficiently catalyzes alkene, alkyne and CO₂ hydrosilylation. Complex **C** has thus far not been reported but the related IMes-ZnCl₂(THF) is known^[13] and the SIPr/ZnBr₂ catalytic mixture has been used for the cycloaddition of CO₂ to epoxides.^[14] Regarding aluminum complexes, we are not aware of their use as catalyst for cycloisomerization reactions. We have tested the IMes-AlBr₃ complex **D**^[15] to allow direct comparison with **C**. Both new complexes **C** and **D** were characterized by X-ray diffraction analyses.^[16]

Results and Discussion

As mentioned above, 7-alkynylcycloheptatrienes rearrange into indenenes under gold catalysis.^[6a] So did compound **1**^[5] when treated for instance by the cationic gold complex **A** (Table 1, 3 mol%), displaying a IPr NHC ligand, in DCE at 80 °C for 20 h (Table 1, entry 1). A 58/42 mixture of indenenes **2** and **3** has been isolated in 72% overall yield. The cationic IPr-stabilized aryl zinc complex **B** was then tested (entry 2). To our delight, under similar conditions, indene **2** was isolated as a sole product in 88% yield. The same result was obtained in toluene (entry 3). A better yield of 92% was reached when using 10 mol% of catalyst (entry 4). The IPr substituted zinc bromide complex **C** proved unreactive (entry 5). However, a catalytic mixture of **C** and AgSbF₆ (10 mol% each), provided a 93/7 mixture of **2** and **3**, isolated in 62% yield (entry 6). This is clearly not AgSbF₆ that is responsible for this reaction since when used alone, AgSbF₆ leads to a different product type (Scheme 2, compound **4**).^[5] Even though it could not be isolated, this clearly supports the formation of [IPr-ZnBr]⁺[SbF₆]⁻ as active species. The same transformation could not be carried out in toluene **lack of solubility of the catalyst** (entry 6). Nevertheless, the involvement of protons cannot be invoked in DCE since neither **4** nor **5** (Scheme 2) were detected. Importantly, these NHC-stabilized species **B** and **C** proved much more efficient than the simple zinc salts ZnCl₂, ZnBr₂, ZnI₂ and Zn(OTf)₂ (entries 8-13), with which low yields and regioselectivities were at best obtained

with ZnBr₂ or ZnI₂ in DCE (entries 9 and 10). It is also worthy of note that the activation of ZnBr₂ as one of these zinc salts with AgSbF₆ was not efficient, as it led to compound **4** (entry 14), i.e. the same as the one obtained in our previous study with AgSbF₆ alone.^[6a] No reaction took place with the aluminum complex **D** with or without AgSbF₆, or AlBr₃ alone (entries 15-20). As an indication of the formation of HSbF₆ in DCE, compounds **4** and **5** were isolated when using a mixture of AlBr₃ and AgSbF₆, whereas only **4** was produced in toluene (entries 21-22). **entry 22 indicates the formation of a 9/1 4/5 mixture in toluene.**


Table 1. Skeletal reorganization of **1**.




Entry	cat.	x	Solvent	Time [h]	2/3/4/5	Yield [%] ^[a]
1	A	3	DCE	20	58/42/0/0	72
2	B	3	DCE	24	100/0/0/0	88
3	B	3	toluene	24	100/0/0/0	89
4	B	10	DCE	24	100/0/0/0	92
5	C	10	DCE	24	-	.. ^[b]
6	C /AgSbF ₆	10/10	DCE	24	93/7/0/0	62
7	C /AgSbF ₆	10/10	toluene	24	-	.. ^[c]
8	ZnCl ₂	10	DCE	24	-	.. ^[b]
9	ZnBr ₂	10	DCE	24	86/14/0/0	33
10	ZnBr ₂	10	toluene	24	-	.. ^[c]
11	ZnI ₂	10	DCE	24	83/17/0/0	34
12	ZnI ₂	10	toluene	24	-	.. ^[c]
13	Zn(OTf) ₂	10	DCE	24	-	.. ^[d]
14	ZnBr ₂ /AgSbF ₆	10	DCE	24	0/0/100 ^[e] /0	49
15	D	20	DCE	24	-	.. ^[b]
16	D	20	toluene	24	-	.. ^[b]
17	D /AgSbF ₆	20	DCE	24	-	.. ^[b]
18	D /AgSbF ₆	20	toluene	24	-	.. ^[b]
19	AlBr ₃	20	DCE	24	-	.. ^[b]
20	AlBr ₃	20	toluene	24	-	.. ^[b]
21	AlBr ₃ /AgSbF ₆	20	DCE	24	0/0/68 ^[e] /32	66 ^[f]

22 AlBr₃/AgSbF₆ 20 toluene 24 0/0/90^[e]/10 40

[a] Isolated. [b] No reaction. [c] Very low conversion, trace of indene product detected. [d] Phenylallene observed, 20% yield. [e] 1:1 *EZ* mixture. [f] Determined *in situ* by ¹H NMR using CH₂Br₂ as internal standard.



The mechanism of the indene formation promoted by the model gold and zinc NHC fragments shown in Scheme 3 was computed at the BP86/def2-QZVP(Au)-6-311+G(2d,p)(other atoms)//BP86/LANL2DZ(Au,Zn)-6-31G(d,p)(other atoms). The 7-alkynylcycloheptatriene complex **a** was used as starting compound. Based on these calculations, the formation of the indene scaffold of complex **g**, corresponding to **2** in Table 1, can be explained as follows: the binding of the metal fragment to the triple bond triggers the nucleophilic attack of the middle double bond of the 1,3,5-triene fragment to give the cyclopropylcarbenoid **b**. The latter rearranges into the bicyclic allyl cation **e**. A 1,2-H shift transforms **e** into the Wheland-type complex **f**. A 1,2-proton shift finally gives rise to the indene complex **g**. For the formation of the other indene complex **d**, corresponding to product **3** in Table 1, the deciphered mechanism involves the rearrangement of **b** into the diallyl cation **c**, which undergoes a 1,2-methyl shift to give **d** (pink pathway).




Scheme 3. Calculated intermediates and transition states for the formation of indenenes from a 7-alkynylcycloheptatriene complex.

The computed free energies corresponding to Scheme 3 are collected in Table 2. While the free energies of activation all seem reasonable, there is no kinetic preference for one indene or the other in the gold series since **TS_{bc}** and **TS_{be}** lie virtually at the same value (14.0 and 14.3 kcal/mol respectively). Thus, a low selectivity is expected between the regioisomers, which is indeed the case between **2** and **3** when catalyst **A** is used (Table 1, entry 1). On the other hand, in the zinc series, the values are markedly different: 18.1 kcal/mol for **TS_{bc}** and 14.7 kcal/mol for **TS_{be}**. A high regioselectivity in favor of the indene displaying the alkyl group at the β position of the phenyl ring is therefore expected, which is corroborated by the selective formation of **2** (Table 1, entry 2).

Table 2. Computed free energies (ΔG_{298} , kcal/mol) relatively to **a**.


Entry	Computed structure	Au series	Zn series
1	TS_{ab}	11.8	10.6
2	b	3.1	7.1
3	TS_{bc}	14.0	18.1
4	c	-19.6	-25.4
5	TS_{cd}	0.8	-23.6
6	d	-55.4	-60.3
7	TS_{be}	14.3	14.7
8	e	-10.3	-4.9
9	TS_{ef}	1.2	5.9
10	f	-28.8	-34.8
11	TS_{fg}	-23.1	-29.6
12	g	-59.6	-64.1

Although the same kind of carbenoid intermediates are expected in the gold and zinc series, the zinc-derived ones seem less reactive towards double bonds. Echavarren et al reported that compound **6** transforms into tautomeric barbaralanes **7** and **8** when treated with the gold catalyst **A** (Scheme 4). They interconvert rapidly through a strain-assisted Cope rearrangement.^[6a] They likely arise from the trapping of the putative barbaralyl gold carbene. Using **B**, the cyclopropanation was not observed, although the same barbaralyl carbene explains the formation of the indene isomers. Again, the **D**/AgSbF₆ mixture did not promote any reaction in this case.



Scheme 4. Cycloisomerization of **6**.

This lack of reactivity could be due to the trigonal geometry around zinc (vs linear for gold), which brings the bulky NHC and the C₆F₅ ligands closer to the carbene center. To validate this hypothesis, DFT computations were performed at the aforementioned level of theory (Scheme 5). The formation of the polycyclic scaffold of compounds **7** and **8** shown in Scheme 4 involves two main steps: formation of the cyclopropylcarbenoid **i** and the electrophilic cyclopropanation at the carbene center to give **j**. While the first step is exergonic with gold, it remains, as found before, endergonic but attainable with zinc. There is yet a major difference between gold and zinc in the cyclopropanation step, the corresponding transition states lying at 13.4 kcal/mol and 20.6 kcal/mol respectively. We attribute this difference to the higher steric crowd around the zinc carbene center due to the trigonal geometry around the metal, vs a linear geometry with gold, which will place the bulky NHC further from the incoming double bond.



Scheme 5. Calculated key intermediates and transition states for the cycloisomerization of compound **6** (ΔG_{298} , kcal/mol).

Although the hydroarylation pathway to **4** (Scheme 2) was not observed with compound **1** under zinc catalysis (except Table 1, entry 14), such a reaction was independently tested using arenynes **11** (Table 3). This compounds was previously shown to undergo metal-catalyzed hydroarylation with PtCl₂, [RuCl₂(CO)_nL_m]₂, [RuCl₂(CO)_nL_m]₂/AgOTf and GaCl₃.^[17] The reaction was very efficient with zinc, especially with the NHC-stabilized species (entries 1 and 2). Of note, the hydroarylation catalyzed by ZnBr₂ or ZnI₂ is more efficient than with AlBr₃ (entries 3-4 vs 5), even though these simple salts could not compete with the NHC-complexes. Adding AgSbF₆ (1/1 mol% ratio) could enhance the activity of ZnBr₂ (entries 3 vs 6), but not that of ZnI₂ and AlBr₃ (entries 7 and 9 vs 4 and 5). The NHC-Al complex **D** left the starting material intact (entry 9). Since the reaction can be conducted in the presence of AgSbF₆ alone with moderate efficiency (entry 10), we can postulate the formation of IMesAlBr₂⁺ ions which are yet inactive for this transformation.

Table 3. Hydroarylation of **11**.




Entry	cat.	Yield [%] ^[a]
1	B	95
2	C /AgSbF ₆	96
3	ZnBr ₂	61
4	ZnI ₂	50
5	AlBr ₃ ^[b]	27
6	ZnBr ₂ /AgSbF ₆	79
7	ZnI ₂ /AgSbF ₆	50
8	D /AgSbF ₆	— ^[c]
9	AlBr ₃ /AgSbF ₆	22
10	AgSbF ₆	55

[a] Isolated. [b] 20 mol%. [c] no reaction.

The transfer hydrogenation of alkenes using 1,4-cyclohexadiene derivatives as H₂ surrogate is a very convenient process that can be catalyzed by gallium^[18] or boron complexes.^[19] The use of zinc in this field seems to be undocumented.^[20] The reduction of compound **13** (Table 4) was previously studied with the NHC-gallium complex [IPr-GaCl₂]⁺[SbF₆]⁻, leading to the reduced product **14** in 67% yield.^[18] Here, also we observed an increased

efficiency with the zinc catalyst **B**, which provided **14** in 80% yield (entry 1). Catalyst **C**, ZnBr₂ and ZnI₂ with AgSbF₆ worked, but the yields did not exceed 68% (entries 2, 5, and 6). It is striking to note that no reaction took place with ZnBr₂ or ZnI₂ alone (entries 3 and 4) or with the aluminum species used with AgSbF₆ (entries 7 and 8). The NHC-zinc complexes also proved much more potent than the corresponding gold complex (entry 9).

Table 4. Transfer hydrogenation of **13**.




Entry	cat.	Yield [%] ^[a]
1	B	80
2	C /AgSbF ₆	63
3	ZnBr ₂	_b,c]
4	ZnI ₂	_b,c]
5	ZnBr ₂ /AgSbF ₆	64
6	ZnI ₂ /AgSbF ₆	68
7	D /AgSbF ₆	_b,c]
8	AlBr ₃ /AgSbF ₆ ^[c]	_b,c]
9	IPr-AuCl/AgSbF ₆	37

[a] Isolated. [b] No reaction. [c] 20 mol% each.

We next turned our attention to the formation of C-O bonds. Compound **15** is an interesting substrate for the evaluation of π -acids (Table 5). As shown by Michelet, Genêt et al, with AuCl or AuCl₃ as catalysts, the two alcohol functionalities react with the alkyne moiety to furnish the bicyclic ketal **16**.^[21] A distinct behavior was reported by us when GaCl₃ was used as catalyst, as only one hydroalkoxylation took place, giving rise to a cyclic enol.^[3d] With the zinc species, similarly to gold, only product **16** was observed. It is complex **C** and AgSbF₆ that gave the best result (84% yield, entry 2). ZnBr₂ and ZnI₂ were rather deactivated this time by the presence of the silver additive (entries 3/4 vs 5/6). The latter had little influence on the activity of **D** or AlBr₃, which remained not sufficiently carbophilic to efficiently activate the alkyne functionality (entries 7 and 8). These hydroalkoxylation could well be catalyzed by an unsuspected Brønsted acid, but as mentioned above, the cycloheptatriene test has not detected the release of protons. Accordingly, it is important to note that the use of 2.2 equiv of 2,6-di-*tert*-butylpyridine had no effect on the reaction outcome. It is again, for entry 2, likely that [IPr-ZnBr]⁺[SbF₆]⁻ cette espèce peut elle être générée in situ et caractérisée par RMN ? is the

active species. We can also infer that the OH groups in **15** can be acidic enough to promote the cleavage of the Zn-C bond in **B**, yielding a less reactive zinc alkoxyde, hence the low yield in entry 1. In that respect, the zinc species used in this study do not tolerate the presence of water and therefore do not efficiently catalyze the hydration of *p*-methyl phenylacetylene, which is a major difference with gold catalysts.^[22]

Table 5. Hydroalkoxylation of **15**.




Entry	cat.	Yield [%] ^[a]
1	B	31
2	C /AgSbF ₆	84
3	ZnBr ₂	71
4	ZnI ₂	65
5	ZnBr ₂ /AgSbF ₆	17
6	ZnI ₂ /AgSbF ₆	17
7	D /AgSbF ₆ ^[b]	21
8	AlBr ₃ /AgSbF ₆ ^[b]	10

[a] Isolated. [b] 20 mol% each.

The preferred behavior of the zinc species as soft π -Lewis acids rather than hard σ -Lewis acids^[1a] was further confirmed using the β -ketoester **17a** (Table 6). It is known to undergo carbonyl-olefin metathesis by σ -Lewis acid activation of the carbonyl group.^[23] Schindler et al have shown that FeCl₃^[24] and GaCl₃^[25] are particularly efficient mediators of such transformations, including that of **17a**. However, catalysts **B**, **C**/AgSbF₆, ZnBr₂ and ZnI₂ proved unable to trigger the metathesis (entries 1-4). Only harder ZnX⁺ ions formed by halide abstraction with AgSbF₆ provided the cyclic isomers **18** and **19** in good yields (entries 5 and 6). In the other hand, the reaction was over in 4 h at 50 °C with the putative AlBr₂⁺ ion generated in situ, and the isolated yield was in the same range (entry 8). Schindler and coworkers recently reported a preprint describing an aluminum-based heterobimetallic ion pair as catalyst to promote carbonyl-olefin ring-closing metathesis.^[26]

Table 6. Carbonyl-Olefin Metathesis of **17a**.




Entry	cat.	18/19	Yield [%] ^[a]
1	B	-	_b]

2	C/AgSbF ₆	-	[b]
3	ZnBr ₂	-	[b]
4	ZnI ₂	-	[b]
5	ZnBr ₂ /AgSbF ₆	58/42	90
6	ZnI ₂ /AgSbF ₆	82/18	84
7	D/AgSbF ₆ ^[c]	-	[d]
8	AlBr ₃ /AgSbF ₆ ^[c]	67/33	79

[a] Isolated. [b] No reaction. [c] reaction performed with 20 mol% of AlBr₃/AgSbF₆ at 50 °C in 4 h. [d] no reaction.

The tandem carbonyl-olefin/transfer hydrogenation of **17** was then attempted with 1,4-cyclohexadiene (1,4-CHD) as hydrogen donor (Table 7). This tandem process was recently reported by us using IPr-GaCl₂⁺ ions as catalyst.^[27] In this reaction, **17** is first transformed into cyclopentenones (such as **18/19**), which then undergo transfer hydrogenation to cyclopentanones **20**. Thus, σ -activation of the carbonyl is required in the first step, and then a π -activation of the C=C bond is required in the second one. While zinc remained inefficient (entry 1), aluminum salts proved active. After screening the time and the equivalents of 1,4-CHD, we found that the reductive cyclization of **17a**, could be carried out in the presence of in situ generated AlBr₂⁺ ions (entries 2 and 3, 82% yield for the optimized condition). With Ar = *p*-fluoro- and *p*-chloro-benzene, the aluminum-catalyzed reaction in DCE proved to be more efficient when compared to the gallium-catalyzed one; the ethyl 2-arylcyclopentane-carboxylates **20b-c** were isolated in 44% and 60% yield respectively (entries 4-5), vs 23% and 50% respectively in our previous study.^[27] These good results contrast with those shown in Table 4, but cyclic alkenes are usually more reactive.

Table 7. Hydrogenative carbonyl-ene metathesis of **17**.



Entry	Ar	Cat.	Time (h)	Yield [%] ^{[a],[b]}
1	Ph	ZnBr ₂	24	[c]
2	Ph, 17a	AlBr ₃	4	68 (20a)
3	Ph, 17a	AlBr ₃	24	82 (20a)
4	4-ClPh, 17b	AlBr ₃	24	44 (20b)
5	4-FPh, 17c	AlBr ₃	24	60 (20c)

[a] Isolated. [b] *cis/trans dr* > 4/1. [c] complex mixture.

Conclusions

Starting from the cycloheptatrienyl test, we have revealed that cationic NHC-Zn(II) complexes can be exquisite π -acids in various reactions such as skeletal rearrangements, hydroarylations, hydroalkoxylations and transfer hydrogenations. They can even compete with NHC-Au(I) or NHC-Ga(III) catalysts sometimes, although this study has delineated some of their limitations compared to gold and gallium species. This work also shows that NHC-stabilized zinc ions are more active than simple zinc halides. Due to its proximity with zinc and gallium in the periodic table, aluminum was also tested. This time, simple AlX₂⁺ ions proved more active than a NHC-Al(III)⁺ species. They are much less versatile than the zinc complexes but have shown a promising activity in a tandem process at the border between π - and σ -Lewis acid catalysis.

Experimental Section

Experimental Details.

Acknowledgements

We are grateful to the China Scholarship Council (CSC, PhD grants for JT and YC), CNRS, MESRI (PhD grant for MV), Université Paris-Sud, ANR ANR-18-CE07-0033-01 (HICAT) and Ecole Polytechnique for their support of this work. We are thankful to the Umicore Company for a generous gift of metal complexes.

Keywords: Zinc • Aluminum • NHC ligand • Alkynes • Cycloisomerization • Carbonyl-ene metathesis

- [1] a) Y. Yamamoto, *J. Org. Chem.* **2007**, *72*, 7817; b) D. J. Gorin, F. D. Toste, *Nature* **2007**, *446*, 395-403.
- [2] a) A. Fürstner, P. W. Davies, *Angew. Chem. Int. Ed.* **2007**, *46*, 3410; b) A. S. K. Hashmi, *Chem. Rev.* **2007**, *107*, 3180-3211; c) Z. Li, C. Brouwer, C. He, *Chem. Rev.* **2008**, *108*, 3239-3265; d) A. Fürstner, *Chem. Soc. Rev.* **2009**, *38*, 3208-3221; e) N. Shapiro, F. D. Toste, *Synlett* **2010**, 675-691; f) S. P. Nolan, *Acc. Chem. Res.* **2011**, *44*, 91-100. g) C. Obradors, A. M. Echavarren, *Acc. Chem. Res.* **2014**, *47*, 902-912; h) L. Fensterbank, M. Malacria, *Acc. Chem. Res.* **2014**, *47*, 953-965; i) R. Dorel, A. M. Echavarren, *Chem. Rev.* **2015**, *115*, 9028-9072; j) D. Pflästerer, A. S. K. Hashmi, *Chem. Soc. Rev.* **2016**, *45*, 1331-1367; k) A. M. Echavarren, M. Muratore, V. López-Carrillo, A. Escríbano-Cuesta, N. Huguet, C. Obradors, *Org. React.* **2017**, *92*, 1.
- [3] a) N. Chatani, H. Inoue, T. Kotsuma, S. Murai, *J. Am. Chem. Soc.* **2002**, *124*, 10294-10295; b) Y. Miyahana, N. Chatani, *Org. Lett.* **2006**, *8*, 2155-2158; c) V. Michelet, P. Y. Toullec, J.-P. Genêt, *Angew. Chem. Int.*


- Ed.* **2008**, *47*, 4268-4315; d) H.-J. Li, R. Guillot, V. Gandon, *J. Org. Chem.* **2010**, *75*, 8435-8449; e) S. Tang, J. Monot, A. El-Hellani, B. Michelet, R. Guillot, C. Bour, V. Gandon, *Chem. Eur. J.* **2012**, *18*, 10239-10243; f) K. Surendra, E. J. Corey, *J. Am. Chem. Soc.* **2014**, *136*, 10918-10920; g) C. Bour, V. Gandon, *Coord. Chem. Rev.* **2014**, *279*, 43-57; h) C. Bour, V. Gandon, *Synlett* **2015**, *26*, 1427-1436; i) B. Michelet, G. Thiery, C. Bour, V. Gandon, *J. Org. Chem.* **2015**, *80*, 10925-10938; j) B. Michelet, S. Tang, G. Thiery, J. Monot, H. Li, R. Guillot, C. Bour, V. Gandon, *V. Org. Chem. Front.* **2016**, *3*, 1603-1613; k) J. Pérez Sestelo, L. A. Sarandeses, M. Montserrat Martínez, L. Alonso-Marañón, *Org. Biomol. Chem.* **2018**, *16*, 5733-5747.
- [4] a) R. Dorel, A. M. Echavarren, *J. Org. Chem.* **2015**, *80*, 7321-7332; b) Y. Wang, M. E. Muratore, A. M. Echavarren, *Chem. Eur. J.* **2015**, *21*, 7332-7339.
- [5] M. Vayer, R. Guillot, C. Bour, V. Gandon, *Chem. Eur. J.* **2017**, *23*, 13901-13905.
- [6] a) P. R. McGonigal, C. de León, Y. Wang, A. Homs, C. R. Solorio-Alvarado, A. M. Echavarren, *Angew. Chem. Int. Ed.* **2012**, *51*, 13093-13096; b) S. Ferrer, A. M. Echavarren, *Organometallics* **2018**, *37*, 781-786; c) M. Mato, C. García-Morales, A. M. Echavarren, *ChemCatChem* **2019**, *11*, 53-72.
- [7] S. Minegishi, J. Kamada, K. Takeuchi, K. Komatsu, T. Kitagawa, *Eur. J. Org. Chem.* **2003**, 3497-3504.
- [8] C. Bour, R. Guillot, V. Gandon, *Chem. Eur. J.* **2015**, *21*, 6066-6069.
- [9] For the formation of HOTf from AgOTf in DCE, see: T. T. Dang, F. Boeck, L. Hintermann, *J. Org. Chem.* **2011**, *76*, 9353-9361.
- [10] Z. Li, G. Thiery, M. Lichtenthaler, R. Guillot, I. Krossing, V. Gandon, C. Bour, *Adv. Synth. Catal.* **2018**, *360*, 544-549.
- [11] For selected examples, see: a) M.-Y. Lin, A. Das, R.-S. Liu, *J. Am. Chem. Soc.* **2006**, *128*, 9340-9341; b) R. Vicente, J. González, L. Riesgo, J. González, L. A. López, *Angew. Chem. Int. Ed.* **2012**, *51*, 8063-8067; c) J. González, L. A. López, R. Vicente, *Chem. Commun.* **2014**, *50*, 8536-8538; d) M. J. González, L. A. López, R. Vicente, *Tetrahedron Lett.* **2015**, *56*, 1600-1608; e) B. Wang, Y. Chen, L. Zhou, J. Wang, C.-H. Tung, Z. Xu, *J. Org. Chem.* **2015**, *80*, 12718-12724; f) L. Li, B. Zhou, Y.-H. Wang, C. Shu, Y.-F. Pan, X. Lu, L.-W. Ye, *Angew. Chem. Int. Ed.* **2015**, *54*, 8245-8249; g) B. Wang, Y. Chen, L. Zhou, J. Wang, Z. Xu, *Org. Biomol. Chem.* **2016**, *14*, 826-829; h) S. Yorimoto, A. Tsubouchi, H. Mizoguchi, H. Oikawa, Y. Tsunekawa, T. Ichino, S. Maeda, H. Oguri, *Chem. Sci.*, **2019**, *10*, 5686-5698.
- [12] D. Specklin, F. Hild, C. Fliedel, C. Gourlaouen, L. F. Veiros, S. Dagorne, *Chem. Eur. J.* **2017**, *23*, 15908-15912.
- [13] Adapted from a) D. Wang, K. Wurst, M. R. Buchmeiser, *J. Organometal. Chem.* **2004**, *689*, 2123-2130; b) A. Rit, A.-K. Wiegand, D. Mukherjee, T. P. Spaniol, J. Okuda, *Eur. J. Inorg. Chem.* **2018**, 1114-1119.
- [14] X. Liu, C. Cao, Y. Li, P. Guan, L. Yang, Y. Shi, *Synlett* **2012**, *23*, 1343-1348.
- [15] Adapted from B. Bantu, G. M. Pawar, K. Wurst, U. Decker, A. M. Schmidt, M. R. Buchmeiser, *Eur. J. Inorg. Chem.* **2009**, 1970-1976.
- [16] CCDC 1977549-1977550 contains the supplementary crystallographic data for this paper.
- [17] a) N. Chatani, H. Inoue, T. Ikeda, S. Murai, *J. Org. Chem.* **2000**, *65*, 4913-4918; b) H. Inoue, N. Chatani, S. Murai, *J. Org. Chem.* **2002**, *67*, 1414-1417; c) L. V. Graux, M. Giorgi, G. Buono, H. Clavier, *Organometallics* **2015**, *34*, 1864-1871.
- [18] B. Michelet, C. Bour, V. Gandon, *Chem. Eur. J.* **2014**, *20*, 14488-14492.
- [19] I. Chatterjee, Z.-W. Qu, S. Grimme, M. Oestreich, *Angew. Chem. Int. Ed.* **2015**, *54*, 12158-12162.
- [20] D. Wang, D. Astruc, *Chem. Rev.* **2015**, *115*, 6621-6686.
- [21] S. Antoniotti, E. Genin, V. Michelet, J.-P. Genêt, *J. Am. Chem. Soc.* **2005**, *127*, 9976-9977.
- [22] a) J. H. Teles, S. Brode, M. Chabanas, *Angew. Chem. Int. Ed.* **1998**, *37*, 1415-1418; b) G. A. Fernández, A. B. Chopra, G. F. Silbestri, *Catal. Sci. Technol.* **2016**, *6*, 1921-1929.
- [23] L. Ravindar, R. Lekkala, K. P. Rakesh, A. M. Asiri, H. M. Marwani, H.-L. Qin, *Org. Chem. Front.* **2018**, *5*, 1381-1391.
- [24] a) J. R. Ludwig, P. M. Zimmerman, J. B. Gianino, C. S. Schindler, *Nature*, **2016**, *533*, 374-379; b) L. Ma, W. Li, H. Xi, X. Bai, E. Ma, X. Yan, Z. Li, *Angew. Chem., Int. Ed.* **2016**, *55*, 10410-10413; c) C. C. McAtee, P. S. Riehl, C. S. Schindler, *J. Am. Chem. Soc.* **2017**, *139*, 2960-2963; d) J. R. Ludwig, S. Phan, C. C. McAtee, P. M. Zimmerman, J. J. Devery, C. S. Schindler, *J. Am. Chem. Soc.* **2017**, *139*, 10832-10842; h) H. Albright, P. S. Riehl, C. C. McAtee, J. P. Reid, J. R. Ludwig, L. A. Karp, P. M. Zimmerman, M. S. Sigman, C. S. Schindler, *J. Am. Chem. Soc.* **2019**, *141*, 1690-1700.
- [25] H. Albright, H. L. Vonesh, M. R. Becker, B. W. Alexander, J. R. Ludwig, R. A. Wiscons, C. S. Schindler, *Org. Lett.* **2018**, *20*, 4954-4958.
- [26] R. B. Watson, A. J. Davis, D. J. Nasrallah, J. L. Gomez-Lopez, C. S. Schindler, ChemRxiv. Preprint. <https://doi.org/10.26434/chemrxiv.9911783.v1>.
- [27] For gallium-catalyzed tandem carbonyl-olefin metathesis/transfer hydrogenation, see: A. Djurovic, M. Vayer, Z. Li, R. Guillot, J.-P. Baltaze, V. Gandon, C. Bour, *Org. Lett.* **2019**, *21*, 8132-8137.

Entry for the Table of Contents (Please choose one layout)

Layout 1:

FULL PAPER

Text for Table of Contents

*Author(s), Corresponding Author(s)****Page No. – Page No.****Title**

Layout 2:

FULL PAPER

((Insert TOC Graphic here; max. width: 11.5 cm; max. height: 2.5 cm))

*Author(s), Corresponding Author(s)****Page No. – Page No.****Title**

Text for Table of Contents