



**HAL**  
open science

# Influence of French Cued Speech on consonant production in children with cochlear implants: an ultrasound study

Laura Machart, Anne Vilain, H el ene Loevenbruck, Lucie M enard

► **To cite this version:**

Laura Machart, Anne Vilain, H el ene Loevenbruck, Lucie M enard. Influence of French Cued Speech on consonant production in children with cochlear implants: an ultrasound study. ISSP 2020 - 12th International Seminar on Speech Production, Dec 2020, Providence (virtual), United States. . hal-03098745v1

**HAL Id: hal-03098745**

**<https://hal.science/hal-03098745v1>**

Submitted on 5 Jan 2021 (v1), last revised 13 Nov 2023 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

## **Influence of French Cued Speech on consonant production in children with cochlear implants: an ultrasound study**

Laura Machart<sup>1,2</sup>, Anne Vilain<sup>2</sup>, H el ene L oevenbruck<sup>1</sup>, Lucie M enard<sup>3</sup>

<sup>1</sup>Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LPNC, 38000 Grenoble, France

<sup>2</sup>Univ. Grenoble Alpes, Grenoble INP, CNRS, GIPSA-lab, 38000 Grenoble, France

<sup>3</sup>Laboratoire de Phon etique, UQ AM, Montr eal, Canada

Although cochlear implant improves deaf children’s speech intelligibility (Turgeon et al., 2017; Grandon et al., to be published), the auditory information it provides remains degraded (Colin et al., 2017) and the perception of some acoustic features can be altered (Bouton et al., 2015). These auditory limitations may impact oral language development and lead to persistent language disorders (Geers et al., 2015). Some acoustic studies have highlighted specific impairments in the speech production of CI children (Grandon, 2016; Reidy et al., 2016). Only a few studies have investigated how CI children use their articulators to produce speech (Turgeon et al., 2017). To compensate for the degraded acoustic input some people choose to use Cued French (CF) which supplements the auditory information with a manual cue (Hage & Leybaert, 2006). CF has been shown to improve auditory sentence processing in children with hearing aids (P erier et al., 1990, Leybaert et al., 2010) and to help children build more stable phonological representations (Charlier & Leybaert, 2000). Using CF enhances speech perception even in CI children (Hage & Leybaert, 2006; Leybaert et al., 2010) and in a noisy environment (Bayard et al., 2019). It has also been suggested that CF could improve sentence production (Hage & Leybaert, 2006).

The aim of the present study is to examine the influence of CF on articulatory precision in CI children. Ultrasound imaging of the sagittal profile of the tongue provides objective information on the degree of articulatory precision. As part of an ongoing project on speech development in French, we have collected speech data on 85 normal hearing children (NH) and 17 CI children (Machart et al., 2019). Based on preliminary data analysis, in the present ultrasound study we chose to focus on the 6 consonants /t/, /k/, /s/, /ʃ/, /n/, /p/, which are frequently substituted by CI children, independently of voicing substitutions (Fig.1), and which can be distinguished by the horizontal and vertical position of the tongue. The lingual movements are recorded during the production of simple words (Picture-Naming task) each including one of the targeted consonants followed by vowel /a/. Stimuli have been chosen for their frequency and imageability: tapis /tapi/ *carpet*, carotte /kar ot/ *carrot*, sapin /sap n/ *fir tree*, chapeau /ʃapo/ *hat*, narine /narin/ *nostril*, orignal /oʁiʒal/ *moose*. Three groups of children are examined: 10 typical NH children (NH), 9 CI children with a strictly oral education (CI) and 9 CI children who benefit from a CF education (CIcf). A secondary aim of the study is to test the impact of simultaneous speech and CF production on articulatory accuracy. The CIcf group is therefore split in two subgroups: children who are not able to use CF in production (group 1: 5 children) and children who can use CF when they speak (group 2: 4 children). CIcf children group 2 are asked to utter the words and cue simultaneously (US\_CF condition). Acoustic data are transcribed using PRAAT (Boersma & Weenink) and ultrasound data are analyzed using SLURP (Laporte & M enard, 2018). The tongue position index will be the location of the highest point of the tongue on the y axis (M enard et al., 2013). The curvature index will be measured following the methods described in Dawson et al. (2015).

The hypotheses are that CI children have a better representation of speech sounds when they benefit from CF, which results in more precise articulation and lingual configurations comparable to those of children with typical development. (H1) We anticipate that CIcf children produce these contrasts more typically and more stably than CI children. More specifically, we expect accuracy scores, acoustic and articulatory parameters to be closer to the NH group in CIcf children than in CI children. (H2) Moreover, we think that using CF during production might improve articulatory control: using the hand will help in positioning the articulators resulting in higher accuracy scores, more typical acoustic parameters, and more typical lingual configurations.




Figure 1: Consonant substitutions

## REFERENCES

- Bayard, C., Machart, L., Strauß, A., Gerber, S., Aubanel, V. & Schwartz, J.L., 2019. Cued Speech enhances speech-in-noise perception in deaf or hard of hearing persons with cochlear implants or hearing aids. *Journal of Deaf Studies*.
- Boersma, P. & Weenink, D., 2019. Praat: doing phonetics by computer [Computer program]. Version 6.1.08, retrieved 5 December 2019 from <http://www.praat.org/>
- Bouton, S., Serniclaes, W., Bertoincini, J., Leuwers, C., & Cole, P. (2011). Apprentissage de la lecture et développement des habiletés associées à la réussite chez les enfants munis d'un implant cochléaire : Apport de la Langue Parlée Complétée (pp. 163-188). In J. Leybaert (Ed.), *La Langue Française Parlée Complétée : fondements et perspectives*. Solal.
- Charlier, B. L., & Leybaert, J., 2000. The rhyming skills of deaf children educated with phonetically augmented speechreading. *Quarterly Journal of Experimental Psychology Section A: Human Experimental Psychology*, 53(2), 349–375.
- Colin, S., Ecalle, J., Truy, E., Lina-Granade, G., Magnan, A., 2017. Effect of age at cochlear implantation and at exposure to Cued Speech on literacy skills in deaf children. *Res. Dev. Disabil.* 71, 61–69.
- Dawson, K. M., Tiede, M. K., & Whalen, D. H., 2016. Methods for quantifying tongue shape and complexity using ultrasound imaging. *Clin. Ling. Phon.*, 30(3–5), 328–344.
- Geers, A., Nicholas, J., Tobey, E., Davidson, L., 2015. Persistent language delay versus late language emergence in children with early cochlear implantation (pp. 155-170). *Journal of Speech, Language, and Hearing Research*, Volume 59, 2016.
- Grandon, B., 2016. *Développement typique et atypique de la production de parole : caractéristiques segmentales et intelligibilité de la parole d'enfants porteurs d'un implant cochléaire et d'enfants normo-entendants de 5 à 11 ans*. (Thèse de Linguistique). Université Grenoble Alpes.
- Grandon, B., Martinez, M. J., Samson, A. & Vilain, A., to be published. Long-term effects of cochlear implantation on the intelligibility of speech in French-speaking children. *Journal of Child Language*.
- Hage, C. & Leybaert, J., 2006. The Effect of Cued Speech on the Development of Spoken Language, in: Spencer, P.E., Marschark, M. (Eds.), *Advances in the Spoken Language Development of Deaf and Hard-of-Hearing Children*. Oxford University Press, pp. 193–211.
- Laporte, C., Ménard L., 2018. Multi-hypothesis tracking of the tongue surface in ultrasound video recordings of normal and impaired speech. *Medical Image Analysis*. 44: 98-114.
- Leybaert, J., Colin, C., Hage, C., LaSasso, C.J., 2010. Cued Speech for Enhancing Speech Perception and First Language Development of Children With Cochlear Implants. *Trends Amplif.* 14, 96–112.
- Machart, L., Meloni, G., Vilain, A., Lævenbruck, H. & Schott-Brua, V. (2019, mai). *Le développement phonologique de l'enfant implanté : l'apport de la Langue française Parlée Complétée*. VIIIème Journées de Phonétique Clinique, Mons, 75-77.
- Ménard, L., Toupin, C., Baum, S., Drouin, S., Aubin, J., & Tiede, M., 2013. Acoustic and articulatory analysis of French vowels produced by congenitally blind adults and sighted adults, *Journal of the Acoustical Society of America*, 134 (4), 2975-2987.
- Périer, O., Charlier, B., Hage, C., Alegría, J., 1990. Evaluation of the Effects of Prolonged Cued Speech Practice Upon the Reception of Spoken Language. *Cued Speech Journal*, IV, 47-59.
- Turgeon, C., Trudeau-Fisette, P., Fitzpatrick, E., & Ménard, L. (2017). Vowel intelligibility in children with cochlear implants: An acoustic and articulatory study. *Int. Journ. Ped. Otor.*, 101, 87–96.