



HAL
open science

Laboratory analysis of Martian analogues for comparison with CRISM observations for detection of polycyclic aromatic hydrocarbons

Jacqueline Campbell, B Schmitt, Olivier Brissaud, Jan-Peter Muller

► To cite this version:

Jacqueline Campbell, B Schmitt, Olivier Brissaud, Jan-Peter Muller. Laboratory analysis of Martian analogues for comparison with CRISM observations for detection of polycyclic aromatic hydrocarbons. EPSC-DPS Joint Meeting 2019, Sep 2019, Geneve, Switzerland. hal-03099240

HAL Id: hal-03099240

<https://hal.science/hal-03099240>

Submitted on 6 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Laboratory analysis of Martian analogues for comparison with CRISM observations for detection of polycyclic aromatic hydrocarbons

Jacqueline D. Campbell (1), Bernard Schmitt (2), Olivier Brissaud (2) and Jan- Peter Muller (1)

(1) Imaging Group, Mullard Space Science Laboratory, UK, (2) l'Institut de Planétologie et d'Astrophysique de Grenoble

Abstract

The results of a series of laboratory experiments are here presented which were carried out in order to generate a diagnostic infrared (IR) spectrum for Polycyclic Aromatic Hydrocarbons (PAHs) of astrobiological interest. The sensitivity of orbital observations to their possible detection will then be assessed over the Martian South Polar Residual Cap (SPRC) and Recurring Slope Lineae (RSL).

1. Introduction

PAHs are a group of chemical compounds consisting of benzene rings of hydrogen and carbon considered to be 'building blocks' for life and can also be a biomarker for extant life [1]. PAHs are abundant on Earth and in interstellar space, and have been detected on various icy moons, comets and within Martian meteorites found on Earth [2, 3]. PAHs have not been detected on the surface of Mars to date, probably due to the deleterious effects of extreme ultraviolet (UV) radiation on the surface of the planet [4]. Our hypothesis is that dynamic SPRC and RSL features may expose material that has previously been shielded from UV, and may therefore be candidate sites for PAH detection using orbital hyperspectral imaging.




Figure 1: SPRC CO₂ ice features, ESP_012784_0935 (left) and RSL surface features, ESP_034830_1670 (right). Both from HiRISE, NASA/JPL/UoA

Here we use the parameters of the (Compact Reconnaissance Imaging Spectrometer for Mars) CRISM instrument as a basis for the laboratory experiments, due to the fine spatial resolution capabilities of the instrument and the small scale of the SPRC and RSL features [5].

2. Laboratory Experiments

These experiments were performed to constrain the detectability limit of PAHs, and to establish PAH spectral features at wavelengths other than the well-known absorption feature at 3.29 μm . There is currently no existing published evidence of PAH detection within SPRC CO₂ ice features, which recede and expose dust. Ongoing research on the characterisation and nature of RSLs indicates that these dark surface streaks may be the result of upwelling of briny liquid water from beneath the Martian surface. These salty deposits could therefore be a prime site for organic detection. This work seeks to provide empirical laboratory data to improve interpretation of orbital data from CRISM, and to ascertain the effect of CO₂ ice sublimation and brine

dehydration on organic spectra, as well as provide PAH reference spectra relevant to Mars.

The site of the laboratory experiments was the “Cold Surface Spectroscopy” facility (CSS) at the Institut de Planétologie et Astrophysique de Grenoble (IPAG) Grenoble, France using the spectro-gonio radiometer and its CarboN-IR environmental cell, which have been specifically developed for studying planetary analogues.

3. Results

The detectability limit of PAHs was established within SPRC and RSL analogues, end member spectra have been established for all components of interest, and new diagnostic absorption features for PAHs have been recorded at a number of wavelengths. For RSL analogues, we found that drying brines within soil sample increased the detectability of PAHs compared with soil samples void of salt.

This work provides the data necessary to improve interpretation of orbital data, and will form the basis for continuing research of dynamic features on Mars in order to detect organic material, and contribute to the search for habitable environments on Mars.

Acknowledgements

Part of the research leading to these results has received partial funding from the European Union’s Seventh Framework Programme (FP7/2007-2013) under iMars grant agreement n° 607379; MSSL STFC Consolidated grant no. ST/K000977/1 and the first author is supported by STFC under PhD studentship no. 526933.

Europlanet 2020 RI has received funding from the European Union’s Horizon 2020 research and innovation programme under grant agreement No 654208.

References

- [1] Benner, S.A. et al. (1998) PNAS, 96,6,2425-2430
- [2] Allamandola, L.J. (2011) EAS, 46,305-317
- [3] Cruikshank. D.P. et al. (2008) Icarus, 193, 334-343
- [4] Dartnell at al. (2012) 10.1111/j.1945-5100.2012.01351.x
- [5] Murchie et al. (2007) JGR, doi:10.1029/2006JE002682