



HAL
open science

Artificial maturation of [FeFe] hydrogenase in a redox polymer film

Christina Felbek, Steffen Hardt, Cecilia Papini, Debajyoti Pramanik, Vincent Artero, Marc Fontecave, Vincent Fourmond, Nicolas Plumere, Christophe Léger

► **To cite this version:**

Christina Felbek, Steffen Hardt, Cecilia Papini, Debajyoti Pramanik, Vincent Artero, et al.. Artificial maturation of [FeFe] hydrogenase in a redox polymer film. *Chemical Communications*, 2021, 57 (14), pp.1750-1753. 10.1039/d0cc08168j . hal-03102775

HAL Id: hal-03102775

<https://hal.science/hal-03102775v1>

Submitted on 7 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Artificial maturation of [FeFe] hydrogenase in a redox polymer film

Received 00th January 20xx,
Accepted 00th January 20xx

Christina Felbek^a, Steffen Hardt^b, Cecilia Papini^c, Debajyoti Pramanik^d, Vincent Artero^d, Marc Fontecave^c, Vincent Fourmond^a, Nicolas Plumeré^{b,e}, Christophe Léger^{*a}

AUTHOR VERSION of [doi:](https://doi.org/10.1039/D0CC08168J)
[10.1039/D0CC08168J](https://doi.org/10.1039/D0CC08168J)

We demonstrate that the insertion of the dinuclear active site of [FeFe] hydrogenase into the apo-enzyme can occur when the enzyme is embedded in a film of redox polymer, under conditions of mediated electron transfer. The maturation can be monitored by electrochemistry, and is as fast as under conditions of direct electron transfer. This new approach clears further the way to the implementation of hydrogenases in large scale real life processes.

The advantages of using hydrogenases as supported catalysts of H₂ oxidation in fuel cells and H₂ photoproduction in photoelectrochemical cells have been emphasized. Indeed, these enzymes use active sites based on transition metals such as Ni and/or Fe, instead of rare and expensive metals, to reversibly catalyze the oxidation and evolution of dihydrogen. A major obstacle to be removed is the production of hydrogenases at a cost and in quantities compatible with large scale applications. Indeed, hydrogenases are complex metalloenzymes, whose biosynthesis *in vivo* is limited by the efficiency of a complex biological maturation machinery, which involves a number of other dedicated enzymes. In the particular case of the so-called [FeFe] hydrogenases, whose active site is a [Fe₂(adt)(CO)₅(CN)₂]²⁻ cluster (adt²⁻=2-aza-1,3-propanedithiolate) covalently bound to a [Fe₄S₄] cluster, a major step in the direction of large scale production has been taken by Fontecave, Happe and Lubitz and coworkers:^{1,2} they developed an artificial maturation process, which consists in 1/ producing

(potentially in large amounts) hydrogenase protein devoid of any cofactor; 2/ reconstituting the accessory FeS clusters with standard non-biological procedures; 3/ exposing the resulting solution of apo-enzyme to a synthetic dinuclear mimic of the active site, [Fe₂(adt)(CO)₄(CN)₂]²⁻. Spontaneous incorporation of the dinuclear cluster and release of one CO ligand leads to mature and active holo-hydrogenase. This very original strategy has greatly advanced the thematic field.³

The last step of the biosynthesis of the enzyme may also be performed with the apo-enzyme adsorbed onto a rotating electrode, under conditions of direct electron transfer.⁴ After the diiron complex is added in solution, the increase in catalytic current informs on the kinetics of formation of the holo enzyme (upon incorporation of the complex in the apo enzyme).⁵

Regarding the potential applications of hydrogenases, their oxygen sensitivity has also been considered a critical issue,⁶⁻¹¹ and various strategies have been proposed to get around this problem. When a redox polymer film is used to support the enzyme onto an electrode and to mediate electron transfer, hydrogen oxidation can occur in the presence of oxygen, because a self-protection mechanism prevents dioxygen from penetrating the film and damaging the enzyme.¹²⁻¹⁴ It is also possible to prepare polymer films of [FeFe] hydrogenase under aerobic conditions by protecting the anaerobically purified hydrogenase by the inhibitor sulfide,¹⁰ and then reactivating it by removing the sulfide under anaerobic conditions.¹⁵

Here we show that we can embed the apo-enzyme in a film of redox polymer and mature it by exposing the film to the dinuclear complex. The activation is fast, suggesting that the polymer does not hamper the diffusion of the complex, nor its incorporation into the apo enzyme. The current that was obtained after artificial maturation in the redox polymer was significantly larger than in equivalent experiments performed under conditions of direct electron transfer, with the apo-enzyme simply adsorbed onto an electrode. We also optimized the film formation procedure to obtain a significant current after transient exposure of the apo-hydrogenase polymer film to air.

^a CNRS, Aix-Marseille Université, Laboratoire de Bioénergétique et Ingénierie des Protéines, Marseille, France.

^b Center for Electrochemical Sciences (CES), Faculty of Chemistry and Biochemistry, Ruhr University Bochum, Universitätsstr. 150, D-44780 Bochum, Germany.

^c Laboratoire de Chimie des Processus Biologiques, Collège de France, Université Pierre et Marie Curie, CNRS UMR 8229, PSL Research University, 11 place Marcelin Berthelot, 75005 Paris, France.

^d Univ. Grenoble Alpes, CNRS, CEA, IRIG, Laboratoire de Chimie et Biologie des Métaux, 17 rue des martyrs, 38000 Grenoble, France.

^e Campus Straubing for Biotechnology and Sustainability, Technical University Munich, Schulgasse 22, 94315 Straubing, Germany.

* corresponding author: christophe.leger@imm.cnrs.fr

Electronic Supplementary Information (ESI) available: supplementary figures. See DOI: 10.1039/x0xx00000x




Fig. 1: Electrochemically-monitored time-resolved artificial maturation of anaerobically prepared films of *Me* apo-hydrogenase, under three distinct conditions: MET in a thin polymer film (green, enlarged ten times) or in a thicker polymer film (dark blue), and DET (orange, enlarged ten times). The maturation was also monitored with another thick polymer film of apo-hydrogenase exposed to air for two hours (dark blue, dashed line, see the optimization procedure in SI). In all experiments, the maturation is detected as an increase in hydrogen oxidation current after 53 nM $[\text{Fe}_2(\text{adt})(\text{CO})_4(\text{CN})_2]^{2-}$ is added to the electrochemical cell under anaerobic conditions, at $t=0$ s. $E=-160\text{mV}$ vs SHE, 40°C , pH 7, 1 atm. H_2 , $\omega = 3000$ rpm.

The experiments described here were performed with the apo-hydrogenase from *Megasphaera elsdenii* (*Me*), prepared as described previously,^{16,17} either adsorbed onto an electrode under conditions of direct electron transfer (DET),¹⁶ or embedded into a redox polymer allowing mediated electron transfer (MET) between the enzyme and the electrode. The redox polymer, whose structure is drawn in Fig. S1, consisted of a 4,4'-methylviologen redox unit attached to a polyvinyl alcohol backbone (205 kDa) by a previously published isocyanate-based coupling procedure (see methods).

We could make films of this polymer embedding the apo-hydrogenase by depositing onto a glassy carbon (GC) disk electrode a mixture of enzyme solution and polymer, and letting it dry for 24h at 4°C in a glove box filled with N_2 (but containing a significant amount of H_2). The green trace in Fig. 1 shows an example of the current response observed when such an electrode is inserted and rotated in an electrochemical cell under 1 atm. of H_2 , and poised above the H^+/H_2 Nernst potential. The current appears after a small amount of complex $[\text{Fe}_2(\text{adt})(\text{CO})_4(\text{CN})_2]^{2-}$ is added to the solution (at $t=0$). No current is observed in the absence of H_2 . Hence this current results from the catalytic oxidation of H_2 by the enzyme molecules that become active upon incorporation of the dinuclear complex. The variation in current suggests approximately first-order kinetics of complex incorporation into the apo-enzyme. Fitting an exponential function to the change in current gives the time constant of maturation, 86 s. Maturation under conditions of DET has been reported before,⁵ although not for *Me* hydrogenase. We include it here as a control and a reference. The orange trace in Fig. 1 shows the result of the same experiment as that plotted in green, but with the apo-hydrogenase from *Me* simply adsorbed onto the rotating disc electrode and undergoing direct electron transfer.

The time constant of the exponential variation of the current is very similar to that obtained with the polymer film, around 84 s. Comparing the voltammetric catalytic responses of the two films (Fig. S2, panels A & B) clearly confirms that electron transfer is mediated by the polymer in the former, and direct in the latter. Indeed, the observed catalytic wavelshapes are very different: roughly sigmoidal in the former case (Fig. S2B, obtained by subtracting the catalytic and non catalytic currents), and similar to that obtained before with the holo-enzyme adsorbed on graphite¹⁶ in the latter (Fig. S2A). The magnitude of the current was similarly low in the two cases, but we found that the MET current is significantly larger if we include a small fraction of holo-hydrogenase in the apo-hydrogenase / polymer mix (about 1/500 holo/apo) when we make the film. In that case, the resulting dry film is deep violet, as a result of catalytic reduction of the viologen in the film, in the presence of H_2 in the glove box. When this electrode is poised in the electrochemical cell under H_2 , the initial current is very small, indicating that the tiny amount of holo enzyme in the film does not give a significant catalytic current, but a large current appears when the dinuclear complex is added (dark blue trace in Fig. 1). The final current was about 20 times larger than that observed with the film produced without holo-enzyme (the green trace in Fig. 1 was enlarged ten times), in these experiments and in all others. The shape of catalytic signature (Fig. S2C) demonstrates that electron transfer is indeed mediated, not direct. Varying the drying time, we found an optimum in terms of current magnitude after two hours of drying (Fig. S3), although maximal stability is seen after 24h.

The difference in magnitude of the current is easily explained by a difference in film thickness. Consistent with this hypothesis, Fig. S4 shows the cyclic voltammograms recorded with the two matured hydrogenase/polymer films under non-catalytic conditions (under an atmosphere of argon). In both cases, the CVs consist of a pair of peaks, at an average potential of -320 ± 10 mV, which result from the oxidation and reduction of the viologen moieties in the films. When the film is made with just apo-enzyme, the peak current (i_p) is low and changes in proportion to scan rate v (as shown by the green symbols in the plot of i_p/v against v in Fig. 2); this is expected if the film is thin enough that all viologen moieties are oxidized/reduced in a single sweep. When the film of apo enzyme is made with a small amount of holo-enzyme, the non-catalytic peaks are about ten times larger (dark blue traces in Fig. S4), and the change in i_p/v against v deviates from linearity at high scan rate (dark blue symbols in Fig. 2): this indicates a transition towards a diffusion limited voltammetry, consistent with the film being thicker. We believe that the reason for this is that the viologen modified polymer is reduced when it is dissolved with holo-hydrogenase under H_2 , while the same polymer in the presence of the apo-enzyme without holo-hydrogenase is oxidized. The singly charged radical cation of the reduced viologen is more hydrophobic than the oxidized viologen dication. This high hydrophobicity enables the production of thick and adhesive films from the reduced viologen modified polymer; in contrast the oxidized form leads to thinner films

probably because of partial redissolution of the more hydrophilic oxidized polymer.

The analysis of the maturation trace (dark blue trace Fig. 1) seems to suggest that the MET maturation is faster in the thick film than in the thin film, with a time constant of 67 s, rather than 85 s, when an exponential function is fitted to the current trace (Fig. S5A). However, as explained below, this is a misinterpretation of the data. Indeed, a complication of the MET configuration compared to DET is that the interpretation of the magnitude of the current depends on film thickness. Going from very thin to very thick films, various regimes are predicted,¹⁸ with the current being either proportional to the concentration of active enzyme [E] in the film (Bartlett's "regime V"¹⁸), proportional to the square root of [E] (regime VII), or independent of [E] (regime III). This corresponds to distinct situations where the current is limited by catalysis, electron transfer, or mass transport inside the film, respectively.¹⁸ Therefore the quantitative interpretation of the change in current observed during the activation of the film requires the previous identification of the operational regime.

One strategy to determine which regime is operational consists in examining the dependence of the catalytic current on substrate (hydrogen) concentration. Again, going from very thin to very thick films, one expects the current to be proportional to either $1/(1+K_m/[H_2])$ (regime V, K_m is the Michaelis constant), $\sqrt{1/(1+K_m/[H_2])}$ (regime VII), or $[H_2]$ (regime III).¹⁸ To accurately examine the dependence of i on $[H_2]$, the effect of H_2 depletion in the solution outside the rotating film (which is all the more significant that the current is large) must be taken into account.¹⁹ We used a new method where we step the bulk concentration of H_2 using a gas mixer, record the current at different rotation rates at each H_2 concentration (a portion of the signal is shown in Fig. 3A), and calculate the H_2 concentration at the film/solution interface, $[H_2]_0$, as described in methods. Fig. 3B shows the resulting catalytic current against $[H_2]_0$; fitting a Michaelis Menten (black trace in Fig. 3B) equation returns a value of K_m around 0.13 atm of H_2 . This value is significantly lower than the value of 0.40 ± 0.05 atm. obtained under conditions of DET (Fig. 3C). In contrast, fitting the square root of a Michaelis Menten equation to the red trace in Fig. 3B returns a value of $K_m = 0.48$ atm., much closer to the expected value. This suggests that the current obtained with the thick film is proportional to the square root of the amount of active enzyme in the film, which is the case if the catalytic current is limited by electron transfer inside the film.¹⁸

The above conclusion implies that the time constant of maturation in the thick film can be obtained from the dark blue trace in Fig. 1 by fitting an exponential function to the square of the current (Fig. S5B). Doing so returns a time constant of about 87 s, which is indeed the same as in the thin film (green trace in Fig. 1) or under conditions of DET (orange trace in Fig. 1), showing that the maturation kinetics is independent of polymer thickness. The rate of activation depends on temperature, but remains the same under conditions of direct or mediated ET (see the data at 30°C in Fig. S8). The comparison with data obtained with other [FeFe] hydrogenases^{5,20} shows that the rates of maturation increase in the order *Desulfovibro*

desulfuricans < *Clostridium pasteurianum*, *M. elsdenii* < *Chlamydomonas reinhardtii*.




Fig. 2. Evaluation of the thin (green, enlarged three times) and thick (dark blue) hydrogenase/polymer films by cyclic voltammetry under non-catalytic conditions (under argon). Plot of anodic (open) and cathodic (full) peak currents divided by the square root of scan rate (\sqrt{v}), against \sqrt{v} . A straight line through the origin is expected when the film is so thin that all viologen moieties are oxidized or reduced in each sweep.

Using the above-described method, we could not prepare films of apo-hydrogenase from *M. elsdenii* under air and then activate them under anaerobic conditions, which could have been a strategy to prevent oxygen damage. This strategy is not possible with *M. elsdenii* because the apo-enzyme is O_2 sensitive (the residual activity after two hours in air is about 1%, Fig. S9) and because reductive conditions are required to stabilize thick polymer films. However, embedding the apo-enzyme in the film is advantageous in case of accidental exposure of the film to O_2 . Indeed, when a thick film of apo-hydrogenase was prepared with a small amount of holo-enzyme under anaerobic conditions, let dry under anaerobic conditions for ten minutes (during which catalytic oxidation of H_2 and reduction of the viologen stabilizes the film), exposed to air at 4°C for two hours, and finally activated under 1 atm. of H_2 , a significant current was seen (dashed line in Fig. 1, compare also the continuous and dashed blue lines in Fig. S10 for films dried for the same time); it is much more intense than that expected from the residual activity of the enzyme exposed to air for the same time: the final current is not in proportion to the remaining activity, because in these thick films, it depends on the square root of the concentration of active enzyme. If the film is prepared without a small fraction of holo-enzyme and exposed to air, no activation is detected (green dashed line in Fig. S10). These experiments show that if the thick film of apo enzyme is exposed to air for two hours before activation, a large fraction of the apo enzyme is irreversibly inactivated, but the magnitude of the H_2 current that is detected after activation remains significant, as a consequence of the favorable scaling between current and concentration of active enzyme in the thick polymer films.

Last, we note that a very small amount of dinuclear complex (53 nM) was used to activate the apo-hydrogenase films in all experiments shown here. Increasing the concentration of complex five times makes no difference (Fig. S12), which is consistent with previous results obtained under DET conditions.⁵ This shows that the affinity of the enzyme for the complex is extremely high (as observed before in DET

experiments⁵) even when the enzyme is embedded in the polymer.

From previous studies and this work, we conclude that the polymer film not only helps interfacing the holo-hydrogenase with an electrode and provides very efficient stabilization and protection from oxygen even under conditions of fuel cell operation, but can also embed the apo-enzyme and allow the dinuclear complex to reach the enzyme, without hindering this final step of the maturation process.




Fig. 3. Measurement of the K_m for H_2 , under conditions of MET with the thick hydrogenase/polymer film (A and B) and under conditions of DET (C) at 5°C , pH 7. The H_2 concentrations are expressed in units of partial pressure of H_2 , and can be converted to mM using the Henry constant for H_2 , 0.78 mM/atm. Panel A shows a portion of the experiment where we measured the H_2 oxidation current at different partial pressures of H_2 and different rotation rates. The corresponding change in current against interfacial H_2 concentration is shown in panel B. We fitted the current traces either by Michaelis Menten equation (black trace) or the square root of a Michaelis Menten equation (red trace). Panel C shows the DET current against bulk concentration of H_2 (changed by setting various partial pressures of H_2 , Fig. S6 A). Fig S6B shows a distinct DET experiment where we interpret the change in current over time as H_2 is flushed away from the cell. From the two types of DET experiments (Fig. 3C and S6B) we obtained $K_m = 0.40 \pm 0.05$ atm. H_2 .

We thank G. Caserta for performing some of the preliminary experiments and G. Kopiec for preparing the viologen modified polymer. We acknowledge support from CNRS, Aix Marseille Université, Agence Nationale de la Recherche (ANR-15-CE05-0020, Labex ARCANÉ, CBH-EUR-GS, ANR-17-EURE-0003), and the Excellence Initiative of Aix-Marseille University - A*MIDEX, a French “Investissements d’Avenir” programme (ANR-11-IDEX-0001-02, RESOLV, funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy – EXC-2033 – Projektnummer 390677874, the ERC starting grant 715900, and the ANR-DFG project SHIELDS (PL 746/2-1). The French authors are part of FrenchBIC (www.frenchbic.cnrs.fr).

Conflicts of interest

There are no conflicts to declare.

Notes and references

- G. Berggren, A. Adamska, C. Lambert, T. R. Simmons, J. Esselborn, M. Atta, S. Gambarelli, J. M. Mouesca, E. Reijerse, W. Lubitz, T. Happe, V. Artero and M. Fontecave, *Nature*, 2013, 499, 66–69.
- J. Esselborn, C. Lambert, A. Adamska-Venkatesh, T. Simmons, G. Berggren, J. Noth, J. Siebel, A. Hemschemeier, V. Artero, E. Reijerse, M. Fontecave, W. Lubitz and T. Happe, *Nat. Chem. Biol.*, 2013, 9, 607–609.
- H. Li and T. B. Rauchfuss, *J. Am. Chem. Soc.*, 2002, 124, 726–727.
- J. T. Kleinhaus, F. Wittkamp, S. Yadav, D. Siegmund and U.-P. Apfel, *Chem. Soc. Rev.*, DOI:10.1039/d0cs01089h.
- M. Sensi, M. del Barrio, C. Baffert, V. Fourmond and C. Léger, *Current Opinion in Electrochemistry*, 2017, 5, 135–145.
- C. F. Megarity, J. Esselborn, S. V. Hexter, F. Wittkamp, U.-P. Apfel, T. Happe and F. A. Armstrong, *J. Am. Chem. Soc.*, 2016, 138, 15227–15233.
- G. Goldet, C. Brandmayr, S. T. Stripp, T. Happe, C. Cavazza, J. C. Fontecilla-Camps and F. A. Armstrong, *J. Am. Chem. Soc.*, 2009, 131, 14979–14989.
- K. D. Swanson, M. W. Ratzloff, D. W. Mulder, J. H. Artz, S. Ghose, A. Hoffman, S. White, O. A. Zadvorny, J. B. Broderick, B. Bothner, P. W. King and J. W. Peters, *J. Am. Chem. Soc.*, 2015, 137, 1809–1816.
- A. Kubas, C. Orain, D. De Sancho, L. Saujet, M. Sensi, C. Gauquelin, I. Meynial-Salles, P. Soucaille, H. Bottin, C. Baffert, V. Fourmond, R. B. Best, J. Blumberger and C. Léger, *Nat. Chem.*, 2017, 9, 88–95.
- J. Esselborn, L. Kertess, U.-P. Apfel, E. Hofmann and T. Happe, *J. Am. Chem. Soc.*, 2019, 141, 17721–17728.
- P. Rodríguez-Maciá, E. J. Reijerse, M. van Gastel, S. DeBeer, W. Lubitz, O. Rüdiger and J. A. Birrell, *J. Am. Chem. Soc.*, 2018, 140, 9346–9350.
- P. Rodríguez-Maciá, L. M. Galle, R. Björnsson, C. Lorent, I. Zebger, Y. Yoda, S. P. Cramer, S. DeBeer, I. Span and J. A. Birrell, *Angew. Chem. Int. Ed Engl.*, DOI:10.1002/anie.202005208.
- N. Plumeré, O. Rüdiger, A. A. Oughli, R. Williams, J. Vivekananthan, S. Pöller, W. Schuhmann and W. Lubitz, *Nat. Chem.*, 2014, 6, 822–827.
- A. A. Oughli, F. Conzuelo, M. Winkler, T. Happe, W. Lubitz, W. Schuhmann, O. Rüdiger and N. Plumeré, *Angew. Chem. Int. Ed.*, 2015, 54, 12329–12333.
- H. Li, D. Buesen, S. Dementin, C. Léger, V. Fourmond and N. Plumeré, *J. Am. Chem. Soc.*, 2019, 141, 16734–16742.
- A. A. Oughli, S. Hardt, O. Rüdiger, J. A. Birrell and N. Plumeré, *Chem. Commun.*, 2020, 56, 9958–9961.
- G. Caserta, C. Papini, A. Adamska-Venkatesh, L. Pecqueur, C. Sommer, E. Reijerse, W. Lubitz, C. Gauquelin, I. Meynial-Salles, D. Pramanik, V. Artero, M. Atta, M. Del Barrio, B. Faivre, V. Fourmond, C. Léger and M. Fontecave, *J. Am. Chem. Soc.*, 2018, 140, 5516–5526.
- G. Caserta, A. Adamska-Venkatesh, L. Pecqueur, M. Atta, V. Artero, S. Roy, E. Reijerse, W. Lubitz and M. Fontecave, *Biochim. Biophys. Acta*, 2016, 1857, 1734–1740.
- P. N. Bartlett and K. F. E. Pratt, *J. Electroanal. Chem.*, 1995, 397, 61–78.
- M. Merrouch, J. Hadj-Said, C. Léger, S. Dementin and V. Fourmond, *Electrochim. Acta*, 2017, 245, 1059–1064.
- J. A. Birrell, K. Wrede, K. Pawlak, P. Rodríguez-Maciá, O. Rüdiger, E. J. Reijerse and W. Lubitz, *Isr. J. Chem.*, 2016, 56, 852–863.