



HAL
open science

Parametrizations of the spin density matrix

E. Di Salvo, Z.J. Ajaltouni

► **To cite this version:**

E. Di Salvo, Z.J. Ajaltouni. Parametrizations of the spin density matrix. *Annals Phys.*, 2021, 428, pp.168448. 10.1016/j.aop.2021.168448 . hal-03107960

HAL Id: hal-03107960

<https://hal.science/hal-03107960>

Submitted on 24 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Parametrizations of the Spin Density Matrix

E. Di Salvo^{a,b,1}, Z. J. Ajaltouni^{a,2}

^a Laboratoire de Physique Corpusculaire de Clermont-Ferrand,
IN2P3/CNRS Université Blaise Pascal, F-63177 Aubières Cedex, France

^b Dipartimento di Fisica dell' Università di Genova
and I.N.F.N. - Sez. Genova,
Via Dodecaneso, 33, 16146 Genova, Italy

Abstract

We propose for the spin density matrix two parametrizations which automatically fulfil the non-negativity conditions, without setting any bound on the parameters. The first one relies on a theorem, that we prove, and it is rather simple and easily adaptable to some specific reactions, where, for example, parity is conserved or angular momentum conservation entails selection rules. Moreover, in the case when the rank is less than the order of the density matrix, we show how to improve the fits to the data, either by implementing previous suggestions, or by elaborating an alternative method, for which we prove a second theorem. Our second parametrization is a variant of previous treatments, it appears suitable for some particular processes. Moreover, we discuss about the possibility of inferring the elements of the density matrix from the differential decay width. Last, we illustrate various examples of current interest, both in strong and weak decays; some of them may be helpful in the investigation of physics beyond the standard model.

PACS numbers: PACS Nos.: 23.20.En, 23.20.Js, 23.40.Bw, 24.10.-i

¹Elvio.Disalvo@ge.infn.it

²ziad@clermont.in2p3.fr

1 Introduction

The spin density matrix (SDM), which was introduced long time ago [1] to describe a mixture of pure spin states, is an essential tool for various aims, like determining the spin and the parity of the resonances [2-9], singling out some exchange mechanisms in the production reactions [8,10-19] and finding possible hints to new physics [20-22]. It concerns unstable states - to be denoted as R in the following - which are produced in some reactions, usually of the type

$$a \ b \ \rightarrow \ R \ c \ \text{or} \ R_0 \ \rightarrow \ R \ d. \quad (1)$$

The matrix elements of the SDM of R are inferred through the analysis of one of its decay modes.

This matrix is Hermitian, non-negative definite and has trace 1. It is characterized by a set of N pure, orthonormal spin states $|n\rangle$ - its eigenvectors - such that [1, 5]

$$\rho = \sum_{n=1}^N |n\rangle p_n \langle n|, \quad 0 \leq p_n \leq 1, \quad \sum_{n=1}^N p_n = 1. \quad (2)$$

Here the p_n , some of which may be zero [2], are the probabilities of the various eigenstates. If R consists of a single resonance with spin J , one has $N = 2J + 1$. However, it may happen to cope with an intermediate state that consists of two spins [23-25], of the ρ - and ω -resonances [26, 27] or of a resonance and the background [8]; in this case, it results $N = \sum_k (2J_k + 1)$, J_k being the value of each spin.

The number of non-zero eigenvalues is defined as the *rank* of the matrix, which is denoted by r , with $r \leq N$. If the mechanism, which gives rise to R , is of the type (1), an upper bound to r can be fixed [3, 28].

In a frame at rest with respect to R , the eigenstates $|n\rangle$ are not necessarily eigenstates of the operators \mathbf{J}^2 and J_z ; as an example, for a spin-3/2 resonance, the SDM may be diagonal with respect to the states

$$|3/2, 3/2\rangle, |3/2, 1/2\rangle, a|3/2, -1/2\rangle + b|3/2, -3/2\rangle, \text{ and } b|3/2, -1/2\rangle - a|3/2, -3/2\rangle, \quad (3)$$

with $|a|^2 + |b|^2 = 1$. If R involves a single spin, a given eigenstate of ρ can be reduced to an eigenstate of J_z by a rotation of the reference frame, but the unitary matrix that diagonalizes the SDM is not a rotation in the general case [29].

The SDM is generally characterized by $N^2 - 1$ real parameters. However, if parity is conserved in the production mechanism [30, 8], this number is reduced, owing to some relations between the matrix elements; moreover, if parity is conserved also in the successive decay of R , it often happens that the imaginary part of ρ cannot be measured [30, 8, 31]; last, if R is produced in a decay, the constraint of angular momentum conservation [20, 21, 32] has to be accounted for.

Obviously, all constraints that we have exposed above, including non-negativity [5, 6, 30, 8, 28, 31] and normalization condition, have to be fulfilled by a parametrization

of the SDM. A problem, which experimental physicists have to cope with, is to avoid introducing any external bounds on the parameters. Different solutions were proposed in past years [33, 30, 8]; however, a simple and sufficiently flexible parametrization seems to be still lacking [31]. Moreover, the suggestion of exploiting the rank condition for reducing the number of parameters of the SDM [5, 8] needs implementation; this problem is especially important when higher spins or more than one spin are involved. Last, the parametrizations, which were proposed in the past, concern essentially resonances which are produced in strong interactions, whereas, in the last years, one has to do with structures which arise from weak decays [20-22]: a systematic study of such cases has not yet been performed.

The aim of the present paper is to fill such gaps. In particular, we consider the SDM both for parity conserving and for parity violating processes; furthermore, we suggest two different parametrizations that automatically satisfy the above mentioned constraints. The first parametrization, which is based on a theorem, is simple and easily adaptable to different situations; moreover we implement and suggest some methods for recognizing possible null eigenstates and for reducing the number of independent parameters. The second parametrization is inspired by an unusual method [28, 15, 17-19], which takes into account the mechanism that originates the structures we want to study: indeed, in some cases, it may be convenient to parametrize the SDM as [28]

$$\rho = U\rho^{(i)}U^\dagger, \quad (4)$$

where $\rho^{(i)}$ is the initial density matrix and U a unitary operator that describes the evolution of the reaction.

Moreover, we discuss about the possibility of obtaining the elements of the SDM from data, referring in particular to the case when more spins are involved, or when parity is conserved in the production mechanism. Last, we apply our parametrizations to some reactions of interest.

Sect. 2 is dedicated to the first parametrization, for which we prove a preliminary theorem. In Sect. 3, we show how to exploit situations such that $r < N$, in part by using a second theorem. In Sect. 4, we introduce the second parametrization. Sect. 5 is devoted to a discussion about the extraction of the SDM elements from the differential decay width. Last, we illustrate a few examples in Sect. 6 and draw some conclusions in Sect. 7.

2 Parametrization of the Spin Density Matrix - I

The first parametrization is based on a theorem, which we state and prove preliminarily.

2.1 Theorem

“Consider a $N \times N$ Hermitian matrix, ρ , defined with respect to an orthonormal basis,

$$\rho = \sum_{i,j=1}^N |i\rangle \rho_{ij} \langle j|. \quad (5)$$

A necessary and sufficient condition for it to be non-negative is that

- all of its diagonal elements are non-negative and
- the Schwarz inequality [30]

$$|\rho_{ij}|^2 \leq \rho_{ii} \rho_{jj} \quad (6)$$

holds.”

Proof

a) The necessary condition is a consequence of the features of the characteristic equation of a non-negative definite Hermitian matrix [8]. However, we give a different argument, which is similar to the one proposed by Daboul [30]. If ρ is Hermitian and non-negative, we may set

$$\rho_{ij} = \sum_n U_{in}^* U_{jn} p_n, \quad \sum_n U_{in}^* U_{jn} = \delta_{ij}. \quad (7)$$

Here U is a unitary matrix³ that diagonalizes ρ and $p_n \geq 0$ is the n -th eigenvalue of ρ . Then

$$\rho_{ii} = \sum_n |U_{in}|^2 p_n \geq 0, \quad (8)$$

which proves the non-negativity of the diagonal elements of ρ . Moreover, Eq. (7) suggests to define a set of complex vectors [30]:

$$|V_i\rangle = \sum_n U_{in} \sqrt{p_n} |n\rangle. \quad (9)$$

Then, owing to the first Eq. (7), one has $\rho_{ij} = \langle V_i | V_j \rangle$ and the Schwarz inequality for the scalar product implies (6).

b) Now suppose that ρ is Hermitian, with non-negative diagonal elements and satisfies the condition (6). Therefore, we may regard each matrix element ρ_{ij} as a scalar product:

$$\rho_{ij} = \langle W_i | W_j \rangle, \quad (10)$$

where $|W_i\rangle$ ($i = 1, 2, \dots, N$) is a set of complex vectors. This set may be fixed in infinitely many ways. Indeed, we may decompose each vector $|W_i\rangle$ according to the orthonormal basis adopted for defining the elements of the SDM, Eq. (5), *i. e.*,

$$|W_i\rangle = \sum_{k=1}^N \alpha_i^k |k\rangle. \quad (11)$$

³As observed by Daboul [30], such a matrix is defined up to some given phases, $U_{jn} \rightarrow U_{jn} \exp(i\zeta_n)$

Then, Eq. (10) entails the system

$$\rho_{ij} = \sum_{k=1}^N \alpha_i^{k*} \alpha_j^k, \quad (12)$$

which is undetermined, as it consists of N^2 real equations and $2N^2$ real unknowns. We give in Appendix A one of the infinite solutions to the system.

Given any vector

$$|Y\rangle = \sum_{i=1}^N y_i |i\rangle, \quad (13)$$

Eq. (10) implies

$$\langle Y|\rho|Y\rangle = \sum_{i,j=1}^N y_i^* \langle W_i|W_j\rangle y_j = \langle Z|Z\rangle \geq 0, \quad (14)$$

where

$$|Z\rangle = \sum_{i=1}^N y_i |W_i\rangle. \quad (15)$$

This shows the non-negativity of ρ and completes the proof of our theorem.

2.2 Parametrization

The results of the previous subsection are now exploited for parametrizing the SDM of the unstable state R , which, as already explained, may consist of more spins.

Since the diagonal elements of the SDM are non-negative, we introduce N real parameters a_i , such that

$$\rho_{ii} = a_i^2 \quad \text{and} \quad \sum_{i=1}^N a_i^2 = \sum_{i=1}^N \langle W_i|W_i\rangle = 1. \quad (16)$$

Moreover, we observe, analogously to Doncel *et al.* [34], that the set of the moduli of the N vectors (9) - which characterize the SDM - can be related to the points of the surface of a hypersphere of unit radius in the N -dimensional Euclidean space R^N . Therefore, we introduce generalized spherical coordinates in that space, by means of a number of ‘angular’ parameters α_i . We propose a parametrization for the SDM of R , distinguishing between parity violation and parity conservation in the processes (1).

2.2.1 Parity Violating Processes

We define, in this case, $N - 1$ ‘angular’ parameters:

$$a_1 = \cos \alpha_1, \quad a_2 = \cos \alpha_2 \sin \alpha_1, \quad \dots \quad (17)$$

$$a_i = \cos \alpha_i \prod_{l=1}^{i-1} \sin \alpha_l \quad \dots \quad a_N = \prod_{l=1}^{N-1} \sin \alpha_l. \quad (18)$$

As regards the off-diagonal matrix elements, the Schwarz inequality (6) suggests to set

$$\rho_{ij} = |a_i||a_j| \cos \gamma_{ij} \exp(i\phi_{ij}). \quad (19)$$

2.2.2 Parity Conserving Processes

In this case, we fix a plane, say π , to be identified with the production plane [8] or, in the case of the decay, with the plane which is singled out by the momenta of R_0 and R . Two possible choices are available [35], corresponding to fixing the quantization axis respectively normal to π or lying on it. Here we focus on the latter case, which includes the helicity representation. We shall see two examples in Sect. 6. The matrix elements of the SDM are denoted as $\rho_{mm'}^{JJ'}$. Then, fixing the y -axis normally to the π -plane, and defining the reflection operator $\Pi_y = P \exp(-i\pi J_y)$, where P is the parity operator, one has

$$\Pi_y |Jm\rangle = \eta \exp[-i\pi(J-m)] |J-m\rangle; \quad (20)$$

here η is the intrinsic parity of the state that we consider. Therefore

$$(\Pi_y \rho \Pi_y^{-1})_{mm'}^{JJ'} = \eta \eta' \exp(-i\pi \Delta) \rho_{-m-m'}^{JJ'}, \quad \text{with } \Delta = J - J' - m + m', \quad (21)$$

J and J' being different spin values and m and m' their third components. Parity invariance implies

$$\rho_{mm'}^{JJ'} = \eta \eta' \exp(-i\pi \Delta) \rho_{-m-m'}^{JJ'}. \quad (22)$$

As regards the parametrization of the diagonal elements of the SDM, we distinguish two cases:

a) Odd N , corresponding to an odd number of integer spins:

$$a_1 = a_N = \frac{1}{\sqrt{2}} \cos \alpha_1, \quad a_2 = a_{N-1} = \frac{1}{\sqrt{2}} \cos \alpha_2 \sin \alpha_1, \quad \dots \quad (23)$$

$$a_i = a_{N-i+1} = \frac{1}{\sqrt{2}} \cos \alpha_i \prod_{l=1}^{i-1} \sin \alpha_l \quad \dots \quad a_{N'/2} = \prod_{l=1}^{N'/2} \sin \alpha_l, \quad (24)$$

with $N' = N - 1$.

b) Even N , in all other cases:

$$a_1 = a_N = \frac{1}{\sqrt{2}} \cos \alpha_1, \quad a_2 = a_{N-1} = \frac{1}{\sqrt{2}} \sin \alpha_1 \cos \alpha_2, \quad \dots \quad (25)$$

$$a_i = a_{N-i+1} = \frac{1}{\sqrt{2}} \cos \alpha_i \prod_{l=1}^{i-1} \sin \alpha_l \quad \dots \quad a_{N/2} = a_{N/2+1} = \frac{1}{\sqrt{2}} \prod_{l=1}^{N/2} \sin \alpha_l. \quad (26)$$

The off-diagonal elements are parametrized according to Eq. (19), taking into account the relationship (22).

If the imaginary part of ρ_{ij} is not measurable, as we shall discuss in Sect. 5, one has to set $\phi_{ij} = 0$ in the parametrization (19); incidentally, we note that the condition $(\Re \rho_{ij})^2 \leq |\rho_{ii} \rho_{jj}|$, and therefore the Schwarz inequality - which is necessary to guarantee

the non-negativity condition - is automatically fulfilled; moreover, $|\Re\rho_{ij}|$ constitutes a natural lower bound to $|\rho_{ij}|$, without introducing any external constraint [30]. One can also apply the rank condition, according to which

$$r(\Im\rho) \leq r(\Re\rho) - r(\rho), \quad (27)$$

where $\Re\rho$ and $\Im\rho$ are respectively the real and imaginary part of ρ [8].

3 Exploiting the Rank of the SDM

The complete parametrization of the SDM requires, in principle, all of the parameters that we have illustrated in the previous section. However, in some specific situations, considerable simplifications are possible, which are notably useful in the cases of high spins or of more states. Here we examine the case when the rank of the SDM is less than its order, that is, when some of the eigenvalues p_n - or equivalently the determinant and some of the principal minors - vanish.

Therefore, an important problem is to single out the kernel of the SDM [2, 3, 5, 8]. To accomplish that, one has to proceed according to more steps [8]. At first, one fits the experimental data by means of an order- N SDM, then one determines its eigenvalues, p_n , and its eigenvectors, $|n\rangle$. If some of the p_n vanish, the N -dimensional space where the SDM acts may be divided into two subspaces, the kernel, of dimension $N_k < N$, and the complementary one, whose basis is constituted by the remaining $r = N - N_k$ eigenvectors. As regards the successive steps, we indicate three different methods.

Method 1

Let

$$|k\rangle = \sum_{i=1}^N \alpha_{ki} |i\rangle, \quad k = 1, 2, \dots, N_k \quad (28)$$

be the eigenvectors of the kernel: they are expressed with respect to the basis that has been adopted in Eq. (5). The matrix

$$\rho' = U\rho U^\dagger, \quad (29)$$

with

$$U_{ki} = \delta_{ki} \quad \text{for } 1 \leq k \leq r, \quad (30)$$

$$= \alpha_{ki}^* \quad \text{for } r+1 \leq k \leq N, \quad (31)$$

results in

$$\rho'_{kl} = \rho_{kl} = \eta_{kl} \quad \text{for } 1 \leq k, l \leq r, \quad 0 \quad \text{otherwise.} \quad (32)$$

Moreover, the $r \times r$ sub-matrix η , which is defined by Eq. (32), is non-singular; it can be re-parametrized by using the procedure that we have described in the previous section.

Method 2

If the unstable state R has a fixed spin J , a single eigenstate of ρ can be reduced to an eigenstate of J_z by means of a rotation of the reference frame [29]. Indeed, any normalized vector, say,

$$|l\rangle = \sum_m \alpha_m^J |J, m\rangle, \quad \text{such that} \quad \sum_m |\alpha_m^J|^2 = 1, \quad (33)$$

may be re-written as

$$|l\rangle = U[R(\phi, \theta, 0)]|J, \bar{m}\rangle = \exp(-i\phi J_z) \exp(-i\theta J_y)|J, \bar{m}\rangle, \quad (34)$$

where $|J, \bar{m}\rangle$ is a pure spin state and ϕ and θ suitable values of the azimuthal and polar angle respectively. If $|l\rangle$ is an eigenvector that corresponds to a null eigenvalue, the transformed SDM has at least a vanishing row and column, which intersect in the main diagonal.

In this connection, we remark that, if the polarization direction of the resonance may be determined - *e. g.*, through an asymmetry in a weak decay mode -, it is convenient to rotate the reference frame so as to take the quantization axis along such a direction: the transformed SDM is diagonal.

Method 3

The third method for reducing the order of the SDM is based on a simple theorem.

Theorem: "If the Schwarz inequality is saturated for some (i, j) -pair of indices, $i \neq j$, *i. e.*,

$$|\rho_{ij}|^2 = \rho_{ii}\rho_{jj}, \quad (35)$$

the rank of the SDM is at least one unit less than its order."

Proof

We define the 2×2 matrix $\bar{\rho}$ as

$$\bar{\rho}_{11} = \rho_{ii}, \quad \bar{\rho}_{22} = \rho_{jj}, \quad \bar{\rho}_{12} = \bar{\rho}_{21}^* = \rho_{ij}. \quad (36)$$

The eigenvalues of $\bar{\rho}$ are 0 and $\rho_{ii} + \rho_{jj}$; the corresponding eigenvectors are

$$|n_1\rangle = \mathcal{N}[\sqrt{\rho_{jj}}|i\rangle - \sqrt{\rho_{ii}} \exp(-i\phi_{ij})|j\rangle] \quad (37)$$

and

$$|n_2\rangle = \mathcal{N}[\sqrt{\rho_{ii}}|i\rangle + \sqrt{\rho_{jj}} \exp(-i\phi_{ij})|j\rangle], \quad (38)$$

where ϕ_{ij} is the phase of ρ_{ij} and $\mathcal{N} = (\rho_{ii} + \rho_{jj})^{-1/2}$. The matrix $\bar{\rho}$ is diagonalized by the unitary matrix \mathcal{U} such that

$$\mathcal{U}_{11} = \mathcal{N}\sqrt{\rho_{jj}}, \quad \mathcal{U}_{12} = -\mathcal{N}\sqrt{\rho_{ii}} \exp(i\phi_{ij}), \quad \mathcal{U}_{21} = \mathcal{N}\sqrt{\rho_{ii}}, \quad \mathcal{U}_{22} = \mathcal{N}\sqrt{\rho_{jj}} \exp(i\phi_{ij}). \quad (39)$$

Therefore, defining the $N \times N$ unitary matrix U as

$$U_{ii} = \mathcal{U}_{11}, \quad U_{ij} = \mathcal{U}_{12}, \quad U_{ji} = \mathcal{U}_{21}, \quad U_{jj} = \mathcal{U}_{22}, \quad U_{lm} = \delta_{lm} \quad \text{for } l, m \neq i, j, \quad (40)$$

it follows that the new SDM $\rho' = U\rho U^\dagger$ has at least a diagonal term which vanishes. But as shown before this implies that $\rho'_{ij} = \rho'_{ji} = 0$ for a fixed i and all j and $r(\rho) \leq N - 1$.

This completes the proof and indicates how to construct, in this case, the unitary matrix for obtaining the reduction of the parameters.

This result could be generalized: any null principal minor [8] of the SDM corresponds to at least one zero eigenvalue. In particular, the most trivial principal minor, a zero in the main diagonal, implies the vanishing of the corresponding row and column, as follows automatically from Eqs. (16) and (19).

In this connection, it is worth recalling the Eberhard-Good theorem [2, 3, 5], which provides useful constraints on the spin of a resonance. In the original version [2], one considers reactions of the type (1), such that a boson resonance R is produced; the theorem states that isotropy in the decay of R implies the inequality

$$J < Q, \quad (41)$$

where J is the spin of R and Q is the product of the numbers of spin states of the other particles which are involved in the reaction. Successively, Peshkin [3] established, for any J , the more restrictive inequality

$$(2J + 1)^{-1} \leq r^{-1} \leq [\text{Tr}\rho^2]^{-1}. \quad (42)$$

Last, Minnaert [5] obtained even more stringent conditions: if $r < 2J + 1$, $2J + 1 - r$ linear combinations of the quantities $\text{Tr}\rho^k$ ($k = 1, 2, \dots, 2J + 1$) must vanish.

4 Parametrization of the Spin Density Matrix - II

The procedure, that we have just described, is especially useful in cases when it is difficult to determine *a priori* an upper bound to the rank (for example, in inclusive reactions [36, 37]), or when this bound is greater than the order of the SDM [28]. If the bound is less than the order, or if the spin structures of the initial and final state are particularly simple [13], it may be sometimes convenient to use an alternative parametrization, based on Eq. (4) [18, 28, 13]. Indeed, in a process of the type (1), this equation yields for the SDM of R

$$\rho_{kk'} = \sum_m \sum_{l'} U_{kl}^m \rho_{l'}^{(i)} U_{k'l'}^{m*}. \quad (43)$$

Here

$$U_{kl}^m = \frac{1}{\mathcal{N}_u} A_{kl}^m, \quad \mathcal{N}_u^2 = \sum_m \sum_{kl} |A_{kl}^m|^2 \quad (44)$$

and A_{kl}^m are the amplitudes of the process; m denotes the spin quantum number of the unobserved final particle (c or d) and l and l' indicate either the spin quantum number of R_0 or the pair (l_a, l_b) of the initial particles a and b .

If the number of independent amplitudes is \bar{N} , we parametrize them as $b_s \exp(i\phi_s)$, $s = 1, 2, \dots, \bar{N}$, with ϕ_s real numbers, $\phi_1 = 0$ and

$$b_1 = \cos \beta_1, \quad b_2 = \sin \beta_1 \cos \beta_2, \quad \dots \quad (45)$$

$$b_m = \cos \beta_m \prod_{l=1}^{m-1} \sin \beta_l \quad \dots \quad b_{\bar{N}} = \prod_{l=1}^{\bar{N}-1} \sin \beta_l. \quad (46)$$

Then the amplitudes are characterized by $2(\bar{N} - 1)$ real parameters, to which one has to add those of the initial SDM. Obviously, if parity is conserved, one has to take into account the corresponding symmetry relations.

In order to see whether this method is more convenient than the previous one, the overall number of such parameters has to be compared with the upper bound to the rank of the density matrix of R ; this bound is $r_i \times r_{c(d)}$, where $r_{i(c,d)}$ are respectively the ranks of the initial state and of c or d . In Sect. 6, we shall compare the two methods in some cases.

5 Discussion

The number of independent parameters, that one can extract from the normalized differential decay width, is determined by the moments, *i. e.*,

$$\frac{1}{\Gamma} \frac{d^2\Gamma}{d \cos \theta d\phi} = \sum_{L=0}^{J_m} \sum_{M=-L}^L \frac{2L+1}{4\pi} H(L, M) \mathcal{D}_{M0}^{L*}(\theta, \phi). \quad (47)$$

Here the \mathcal{D} are the Wigner rotation functions and J_m is the maximum spin of R . Moreover, one has $H(0,0) = 1$. The other moments can be extracted either by means of a best fit or by exploiting the orthogonality of the \mathcal{D}_{M0}^L functions with different L or M .

If R includes only one spin, the number of moments equals the number of parameters of the SDM that can be determined; these are related to the moments by a linear system [8, 35]. Otherwise, the best we can extract are a number of matrix elements and some linear combinations of the remaining ones [8], unless one can reduce the number of the independent parameters of the SDM, by means of some assumption.

If parity invariance holds both in the production and in the decay of R , the imaginary parts of the elements ρ_{ij} of the SDM turn out to be measurable only under very particular conditions, as shown in Appendix B:

- R includes more spins and
- there are at least two independent decay amplitudes, with a non-trivial relative phase, owing to T -odd final-state interactions [38].

Such conditions are realized, for example, in the low-energy scattering $p \bar{p} \rightarrow \Lambda \bar{\Lambda}$ [13], which involves a few partial waves; the corresponding decay amplitudes have different phases, owing to the T -odd (spin-orbit) term of the interaction between the final hadrons.

6 Examples

Here we apply our parametrizations to some concrete situations, either with parity conservation or with parity violation.

6.1 Two Spins: 1^- and 0^\mp - Parity Conservation

We consider the two cases, in order to simulate a resonance-background interference in strong decays of vector mesons; these were already considered many years ago [8], but with a different approach. One has to take into account the relation (22).

a) In the case that R includes the states 1^- and 0^- , we have $\eta' = \eta$, therefore

$$\rho_{mm'}^{JJ'} = (-)^{J-J'-m+m'} \rho_{-m-m'}^{JJ'}; \quad (48)$$

this entails the following parametrization:

$$\rho_{11}^{11} = \rho_{-1-1}^{11} = 1/2 \sin^2 \alpha_1, \quad \rho_{00}^{11} = \cos^2 \alpha_1 \cos^2 \alpha_2, \quad (49)$$

$$\rho_{10}^{11} = -\rho_{-10}^{11} = \frac{1}{\sqrt{2}} \cos \alpha_1 \sin \alpha_1 \cos \alpha_2 \cos \gamma_{10} \exp(i\varphi_{10}^{11}), \quad (50)$$

$$\rho_{1-1}^{11} = \rho_{-11}^{11} = 1/2 \sin^2 \alpha_1 \cos \gamma_{1-1}, \quad \rho_{00}^{00} = \cos^2 \alpha_1 \sin^2 \alpha_2, \quad (51)$$

$$\rho_{10}^{10} = \rho_{-10}^{10} = \frac{1}{\sqrt{2}} \cos \alpha_1 \sin \alpha_2 \cos \gamma_{10}^{10} \exp(i\varphi_{10}^{10}), \quad \rho_{00}^{10} = 0. \quad (52)$$

The remaining terms can be deduced from the previous expressions by taking into account the Hermitian character of ρ . Therefore, we have 7 parameters in all.

b) Consider now the case when R consists of the 1^- and 0^+ states. Eqs. (49) to (51) hold still true, whereas

$$\rho_{10}^{10} = -\rho_{-10}^{10} = \frac{1}{\sqrt{2}} \cos \alpha_1 \sin \alpha_2 \cos \gamma_{10}^{10} \exp(i\varphi_{10}^{10}), \quad (53)$$

$$\rho_{00}^{10} = \frac{1}{\sqrt{2}} \cos \alpha_1 \sin \alpha_2 \cos \gamma_{00}^{10} \exp(i\varphi_{00}^{10}). \quad (54)$$

Therefore, 2 more parameters are needed.

It is worth mentioning that also the SDM of an off-shell spin-1 particle is 4×4 , owing to its scalar component; however, this case concerns essentially weak decays and will be illustrated in some detail in Subsect. 6.4.

6.2 $\Lambda_b \rightarrow \Lambda J/\psi$

We consider the SDM of the J/ψ -resonance. In this case, parity is not conserved in the Λ_b decay, but the constraint of the angular momentum conservation has to be considered. Indeed, adopting the helicity formalism, one has

$$\rho_{1-1} = \rho_{-11} = 0, \quad (55)$$

since the third component of the angular momentum of Λ_b along the J/ψ momentum assumes a fixed value, either $+1/2$ or $-1/2$, but not both simultaneously. The other elements read as

$$\rho_{11} = \cos^2 \alpha_1, \quad \rho_{00} = \sin^2 \alpha_1 \cos^2 \alpha_2, \quad \rho_{-1-1} = \sin^2 \alpha_1 \sin^2 \alpha_2, \quad (56)$$

$$\rho_{10} = \rho_{01}^* = \cos \alpha_1 \sin \alpha_1 \cos \alpha_2 \sin^2 \gamma_{10} \exp(i\phi_{10}), \quad (57)$$

$$\rho_{-10} = \rho_{0-1}^* = \sin^2 \alpha_1 \cos \alpha_2 \sin^2 \gamma_{-10} \exp(i\phi_{-10}). \quad (58)$$

We have employed 6 parameters in all, one less than required in ref. 39, which is cited in the LHCb analysis of the decay [40]. It is worth noting that, according to the discussion of the previous section, all such parameters can be determined, in principle, by analyzing, *e. g.*, the decay $J/\psi \rightarrow \mu^+ \mu^-$.

Incidentally, the constraint (55) holds true for any vector boson resonance V that comes from a decay of the type

$$f_1 \rightarrow f_2 V, \quad (59)$$

where f_1 and f_2 have spin 1/2. A similar simplification occurs, *e. g.*, for the SDM of $\Delta(1232)$ in the decay

$$\Delta(1620)1/2^- \rightarrow \Delta(1232)3/2^+ \pi, \quad (60)$$

for which parity invariance must be taken into account.

Last, in the decay $\Lambda_b \rightarrow \Lambda J/\psi$, a correlation between the decay products of the two final particles may be derived, starting from the direct product $\rho_\Lambda \otimes \rho_{J/\psi}$ of the density matrices of such particles. One can proceed similarly to ref. 41, where the correlation between the two charged leptons l^+ and l^- , that result from

$$e^+ e^- \rightarrow t \bar{t} \rightarrow (b l^+ \nu_l) (\bar{b} l^- \bar{\nu}_l), \quad (61)$$

is calculated to $O(\alpha_S)$.

6.3 $t \rightarrow bW \rightarrow b\tau\nu_\tau$

In this decay, which is of the type (59), we employ the alternative method that we have described in Sect. 4. We consider, in this case, the SDM of a W -boson which is emitted in a given direction, adopting a frame at rest with respect to the top quark. It reads as [22]

$$\rho_{\mu\mu'}(\theta, \phi) = \frac{1}{4\pi} \sum_\lambda b_{\lambda\mu} \rho'_{\Lambda\Lambda'}(\theta, \phi) b_{\lambda\mu'}^*. \quad (62)$$

Here $\Lambda = \mu - \lambda$ is the spin component of t along the W momentum and μ and λ the helicities of, respectively, W and the b -quark; moreover,

$$\rho'_{\Lambda\Lambda'}(\theta, \phi) = \delta_{\Lambda\Lambda'} + \sigma_{\Lambda\Lambda'}^i P_i', \quad (63)$$

$$\vec{P}' \equiv P(\cos \theta, \sin \theta \cos \phi, \sin \theta \sin \phi); \quad (64)$$

P is the top quark polarization, $0 \leq P \leq 1$, while θ and ϕ are, respectively, the polar and the azimuthal angle of the momentum of the W -boson. Last, $b_{\lambda\mu}$ are the reduced decay amplitudes of the $t \rightarrow Wb$ decay:

$$b_{\lambda\mu} = \mathcal{N}^{-1/2} A_{\lambda\mu}, \quad \mathcal{N} = |A_{+,1}|^2 + |A_{+,0}|^2 + |A_{-,0}|^2 + |A_{-,-1}|^2. \quad (65)$$

This suggests the parametrization

$$b_{+,1} = |\cos \beta_1|, \quad (66)$$

$$b_{+,0} = |\sin \beta_1 \cos \beta_2| \exp(i\varphi_{10}) \quad (67)$$

$$b_{-,0} = |\sin \beta_1 \sin \beta_2 \cos \beta_3| \exp(i\varphi_{-10}), \quad (68)$$

$$b_{-,-1} = |\sin \beta_1 \sin \beta_2 \sin \beta_3|. \quad (69)$$

Moreover, we set

$$P = \cos^2 \beta_0. \quad (70)$$

We have assumed the phases of the amplitudes $b_{\pm, \pm 1}$ to be zero, because, in our case, they can interfere only with $b_{\pm, 0}$ respectively.

The parametrization of the W -boson SDM results in $\rho_{1-1} = 0$ and

$$\bar{\rho}_{11} = \cos^2 \beta_1 (1 + \cos^2 \beta_0 \cos \theta), \quad \bar{\rho}_{00} = \sin^2 \beta_1 (\cos^2 \beta_2 + \sin^2 \beta_2 \cos^2 \beta_3), \quad (71)$$

$$\bar{\rho}_{-1-1} = \sin^2 \beta_1 \sin^2 \beta_2 \sin^2 \beta_3 (1 - \cos^2 \beta_0 \cos \theta), \quad (72)$$

$$\bar{\rho}_{10} = |\cos \beta_1 \sin \beta_1 \cos \beta_2| [1 + \cos^2 \beta_0 \sin \theta \exp(-i\phi)] \exp(i\varphi_{10}), \quad (73)$$

$$\bar{\rho}_{-10} = |\sin \beta_1 \sin \beta_2| (|\cos \beta_3| + |\sin \beta_3|) [1 + \cos^2 \beta_0 \sin \theta \exp(i\phi)] \exp(i\varphi_{-10}), \quad (74)$$

having set $\bar{\rho} = 4\pi\rho$. As before, the remaining SDM elements are deduced from the hermiticity condition.

It is worth noting that this procedure - whose parameters, again, do not need any bound - could also be applied to the vector boson V in a decay of the type (59); it has the advantage of relating the polarization of the W to that of the parent state.

The SDM of the W -boson in the semi-leptonic decay of the top quark was parametrized also by other authors a few years ago [21, 42]. Furthermore, in ref. 42, also the more general quasi-three-body decay $t \rightarrow X_b l \nu_l$ is analyzed, in order to search for possible hints to new physics; in particular, two of the parameters, which are introduced, result to be quite sensitive to contributions beyond the standard model.

6.4 $\Lambda_b \rightarrow \Lambda_c(W^*, H^*) \rightarrow \Lambda_c \tau \nu_\tau$

The decay involves a virtual W , which includes a scalar component. For decays of this type, a complete list of the 16 matrix elements of the SDM of the virtual W is given in ref. 43. Furthermore, in the leptonic decays of the Higgs boson, the scalar component of the virtual W and Z is added in a gauge invariant manner [44]. See also ref. 22 and refs. 55 to 59 therein.

In the specific decay that we consider, the independent (reduced) decay amplitudes are 8:

$$b_{+,1}^1 = |\cos \beta_1|, \quad (75)$$

$$b_{+,0}^1 = |\sin \beta_1 \cos \beta_2| \exp(i\varphi_{+,0}^1), \quad b_{-,0}^1 = |\sin \beta_1 \sin \beta_2 \cos \beta_3| \exp(i\varphi_{-,0}^1), \quad (76)$$

$$b_{-,-1}^1 = |\sin \beta_1 \sin \beta_2 \sin \beta_3 \cos \beta_4|, \quad b_{+,0}^t = \prod_{k=1}^4 |\sin \beta_k \cos \beta_5| \exp(i\varphi_{+,0}^t), \quad (77)$$

$$b_{-,0}^t = \prod_{k=1}^5 |\sin \beta_k \cos \beta_6| \exp(i\varphi_{-,0}^t), \quad b_{+,0}^0 = \prod_{k=1}^6 |\sin \beta_k \cos \beta_7| \exp(i\varphi_{+,0}^0), \quad (78)$$

$$b_{-,0}^0 = \prod_{k=1}^7 |\sin \beta_k| \exp(i\varphi_{-,0}^0). \quad (79)$$

Here, the possible new physics is described by the amplitudes $b_{\pm,0}^0$. Moreover, analogously to the previous case, we have assumed equal to 0 the phases of $b_{\pm,\pm 1}^1$.

Then, following the same procedure and using the same notations as in the previous subsection, we find that the matrix elements of $\bar{\rho}$ read as

$$\bar{\rho}_{1-1}^1 = 0, \quad \bar{\rho}_{\pm 1 \pm 1}^1 = \xi_{\pm 1}(1 \pm P_z), \quad (80)$$

$$\bar{\rho}_{00}^{JJ'} = \eta_{00}^{JJ'} + \Delta\eta_{00}^{JJ'} P_z, \quad \bar{\rho}_{\pm 10}^{1J} = \eta_{\pm 10}^{1J} P_{\mp}. \quad (81)$$

Here J and J' run over 1, t , 0; moreover,

$$P_z = \cos^2 \beta_0 \cos \theta, \quad P_{\pm} = \cos^2 \beta_0 \sin \theta \exp(\pm i\phi); \quad (82)$$

last,

$$\xi_{\pm 1} = |b_{\pm,\pm 1}^1|^2, \quad \eta_{00}^{JJ'} = b_{+,0}^J b_{+,0}^{J'*} + b_{-,0}^J b_{-,0}^{J'*}, \quad (83)$$

$$\Delta\eta_{00}^{JJ'} = b_{-,0}^J b_{-,0}^{J'*} - b_{+,0}^J b_{+,0}^{J'*}, \quad \eta_{\pm 10}^{1J} = b_{\pm,\pm 1}^1 b_{\pm,0}^{J*}. \quad (84)$$

In this case, we have used 14 parameters in all, whereas the general treatment would require 24 parameters. But according to the considerations of Sect. 5 the number of independent parameters, that can be inferred from the differential decay width, is 8; therefore, the best fit to the data, if performed by using the above parametrization, would present some ambiguities. However, in this case, some parameters can be fixed by inserting the standard model predictions [20, 21, 42, 45]. In particular, in a previous paper [22], we showed the relationship between the tensor, that is usually employed to describe a semi-leptonic decay, and the non-covariant SDM. If at least one of the amplitudes $b_{\pm,0}^0$ is non-zero, with a non-trivial phase, it causes a T-odd component for the Λ_c polarization [46].

7 Conclusions

We have proposed two methods for parametrizing the SDM of an unstable state - consisting of one or more spins - that is produced in various reactions. Both methods satisfy automatically all of the numerous non-negativity conditions [30, 5] and are adaptable to the constraints imposed by parity [8] and angular momentum conservation. The first method is based on a theorem that we have proved and it may be applied in a simple and flexible way. Moreover, we show how to improve the fit to the data when the rank of the SDM is less than its order; we do this, either by implementing suggestions by other authors [5, 28], or with the help of a second theorem, which we have proved as well.

The second method is a variant of previous parametrizations [3, 28] and is particularly suitable under especial conditions, *e. g.*, when the structure to be analyzed derives from some decay.

We have discussed about the possibility of inferring the SDM from the differential decay width. In particular, we have examined the case where a given intermediate state

is generated and decays according to strong or electromagnetic interactions, showing that, under very particular circumstances, the imaginary part of the SDM can be measured.

Last, we have illustrated some applications; two of them, which concern the decays $t \rightarrow b W$ [20, 21] and $\Lambda_b \rightarrow \Lambda_c l^+ \nu_l$ [47-50,22], are very interesting from the viewpoint of the search for new physics beyond the standard model.

As a conclusion, we observe that our suggestions appear efficient for situations where higher spins or newly discovered structures [51] are involved.

Acknowledgments

The authors are thankful to their friend Flavio Fontanelli for useful and stimulating discussions.

Appendix A

We find an explicit solution to the N complex vectors $|W_i\rangle$ which appear in Eq. (10), *i. e.*,

$$\rho_{ij} = \langle W_i | W_j \rangle. \quad (\text{A.1})$$

This is possible owing to the Schwarz inequality, that we have assumed. The data of the system (A.1) consist of the elements of the Hermitian matrix ρ , defined with respect to an orthonormal basis $|k\rangle$ and such that

$$\rho_{ii} \geq 0 \quad \text{and} \quad |\rho_{ij}|^2 \leq \rho_{ii}\rho_{jj}. \quad (\text{A.2})$$

We assume expansions of the type

$$|W_i\rangle = \sum_{k=1}^i \alpha_i^k |k\rangle \quad (\text{A.3})$$

for any vector $|W_i\rangle$, $i = 1, 2, \dots, N$; furthermore, we establish all of the α_i^k to be real and non-negative. Then, limiting ourselves to $i \geq j$, Eq. (A.1) yields

$$\rho_{ij} = \sum_{k=1}^i \alpha_i^k \alpha_j^k. \quad (\text{A.4})$$

We show that this equation uniquely fixes all of the coefficients α_i^k of the expansions (A.3).

1) Assume, at first, that the rank of ρ is equal to N , which implies that all principal minors of the matrix are positive, in particular, $\rho_{ii} > 0$. Then we prove our statement by induction.

a) For $i=1$, Eq. (A.4) reads as

$$\alpha_1^1 = \sqrt{\rho_{11}}, \quad (\text{A.5})$$

which defines $|W_1\rangle$ through Eq. (A.3).

b) Suppose Eqs. (A.4) to be solvable with respect to α_j^k for all $k \leq j$ and all $j \leq i < N$. This amounts to asserting that all vectors

$$|W_j\rangle, \quad j \leq i, \quad (\text{A.6})$$

have been determined. Now we prove that the vector

$$|W_{i+1}\rangle = \sum_{k=1}^{i+1} \alpha_{i+1}^k |k\rangle, \quad (\text{A.7})$$

can be uniquely deduced from the system

$$\rho_{i+1j} = \sum_{k=1}^j \alpha_{i+1}^{k*} \alpha_j^k, \quad j = 1, 2, \dots, i+1, \quad (\text{A.8})$$

a complex system of $i+1$ equations, with i complex unknowns, α_{i+1}^k , $1 \leq k \leq i$, and a real one, α_{i+1}^{i+1} .

To this end, preliminarily, we show that the $i \times i$ ('triangular') matrix A , such that

$$A_{jk} = \alpha_j^k, \quad \text{with} \quad k \leq j \leq i, \quad (\text{A.9})$$

is non-singular. Consider the $i \times i$ submatrix $\tilde{\rho}$, whose matrix elements $\tilde{\rho}_{lm}$ coincide with those of ρ for $l, m \leq i$. Eq. (A.1) implies

$$\tilde{\rho}_{lm} = \langle W_l | W_m \rangle. \quad (\text{A.10})$$

The assumption of non-singularity of ρ implies the same for $\tilde{\rho}$, therefore the vectors (A.6) constitute a basis for $\tilde{\rho}$ and are linearly independent. This in turn entails the non-singularity of A^4 .

But the system (A.8) can be split into a linear subsystem with i equations and a non-linear equation:

$$\rho_{i+1j} = \sum_{k=1}^j A_{jk} \alpha_{i+1}^{k*}, \quad j = 1, 2, \dots, i, \quad \rho_{i+1i+1} = \sum_{k=1}^{i+1} |\alpha_{i+1}^k|^2. \quad (\text{A.11})$$

The non-singular character of A allows to solve the linear subsystem with respect to α_{i+1}^k , $k = 1, 2, \dots, i$. The solution can be inserted into the non-linear equation, which can be solved with respect to α_{i+1}^{i+1} . This completes the proof in the case of non-singular ρ .

2) If the rank of ρ , say r , is less than N , we perform the transformation (29), *i. e.*

$$\rho' = U \rho U^\dagger. \quad (\text{A.12})$$

The result is

$$\rho'_{lm} = \rho_{lm} = \eta_{lm} \quad \text{for} \quad 1 \leq l, m \leq r \quad \text{and} \quad 0 \quad \text{otherwise}; \quad (\text{A.13})$$

⁴As a byproduct, it is worth noting that, in this case, all α_i^i are strictly positive.

therefore η is a $r \times r$ Hermitian, non-singular matrix, to which we may apply the method described above. We find a set of linearly independent complex vectors $|w_l\rangle$, $l = 1, 2, \dots, r$. Defining

$$|W'_i\rangle = |w_i\rangle \quad \text{for } i \leq r \quad \text{and } 0 \quad \text{for } r < i \leq N, \quad (\text{A.14})$$

the transformed vectors

$$|W_i\rangle = U_{ij}|W'_j\rangle \quad (\text{A.15})$$

give the solution to the system (A.1). This completes the proof.

Appendix B

Here we discuss about the measurability of the imaginary parts of the elements of the spin density matrix (SDM) of an unstable state R that is produced and decays according to parity conserving interactions. We assume this state to include more spins and to have a two-body decay:

$$R \rightarrow a b. \quad (\text{B.1})$$

Adopting the helicity formalism, the normalized differential decay width reads as

$$\frac{1}{\Gamma} \frac{d^2\Gamma}{d \cos \theta d\phi} = \sum_{JJ'} \sum_{mm'} \sum_{\lambda_a \lambda_b} C(J, J') \rho_{mm'}^{JJ'} \mathcal{D}_{m\lambda}^{J*}(\phi, \theta, 0) \mathcal{D}_{m'\lambda}^{J'}(\phi, \theta, 0) f_{\lambda_a \lambda_b}^J f_{\lambda_a \lambda_b}^{J'*}. \quad (\text{B.2})$$

Here $C(J, J') = 1/4\pi[(2J+1)(2J'+1)]^{1/2}$, $\rho_{mm'}^{JJ'} = \langle J, m | \rho | J', m' \rangle$ is an element of the SDM, \mathcal{D} is the Wigner rotation function and

$$f_{\lambda_a \lambda_b}^J = \frac{1}{N_f} F_{\lambda_a \lambda_b}^J, \quad N_f^2 = \sum_{J, \lambda_a, \lambda_b} |F_{\lambda_a \lambda_b}^J|^2, \quad (\text{B.3})$$

the reduced decay amplitudes, with $\lambda = \lambda_a - \lambda_b$. Parity conservation implies

$$\rho_{mm'}^{JJ'} = \eta \eta' \exp(-i\pi\Delta) \rho_{-m-m'}^{J-J'}, \quad (\text{B.4})$$

$$\Delta = J - J' - m + m', \quad (\text{B.5})$$

$$f_{-\lambda_a - \lambda_b}^J f_{-\lambda_a - \lambda_b}^{J'*} = \eta \eta' \exp[-i\pi(J - J')] f_{\lambda_a \lambda_b}^J f_{\lambda_a \lambda_b}^{J'*}, \quad (\text{B.6})$$

as follows from Eqs. (20) to (22) in the text. Moreover,

$$\mathcal{D}_{m\lambda}^J(\phi, \theta, 0) = \exp(-im\phi) d_{m\lambda}^J(\theta), \quad d_{-m-\lambda}^J(\theta) = (-)^{m-\lambda} d_{m\lambda}^J(\theta). \quad (\text{B.7})$$

Therefore we may re-write Eq. (B.2) as

$$\begin{aligned} \frac{1}{\Gamma} \frac{d^2\Gamma}{d \cos \theta d\phi} = & \sum_{JJ'} \sum'_{mm'} \sum'_{\lambda_a \lambda_b} C(J, J') \{ \exp[-i(m - m')\phi] + (-)^\epsilon \exp[i(m - m')\phi] \} \\ & \times \rho_{mm'}^{JJ'} d_{m\lambda}^J(\theta) d_{m'\lambda}^{J'}(\theta) f_{\lambda_a \lambda_b}^J f_{\lambda_a \lambda_b}^{J'*}, \quad (\text{B.8}) \end{aligned}$$

where $\epsilon = \Delta + J - J' + m + m' - 2\lambda = 2(J - J' + m' - \lambda)$ and the primes in the sums indicate that they are limited to non-negative values of the indices. But ϵ is an even number, moreover only $\Im\rho_{mm'}^{JJ'}$ and $\Im(f_{\lambda_a\lambda_b}^J f_{\lambda_a\lambda_b}^{J'*})$ are odd under the simultaneous exchange $(J, m) \leftrightarrow (J', m')$, the other terms of Eq. (B.8) being even. Therefore we have

$$\frac{1}{\Gamma} \frac{d^2\Gamma}{d \cos\theta d\phi} = \sum_{JJ'} \sum_{mm'}' \sum_{\lambda_a\lambda_b}' C(J, J') \cos[(m - m')\phi] d_{m\lambda}^J(\theta) d_{m'\lambda}^{J'}(\theta) \mathcal{F}_{mm', \lambda_a\lambda_b}^{JJ'}, \quad (\text{B.9})$$

with

$$\mathcal{F}_{mm', \lambda_a\lambda_b}^{JJ'} = \Re\rho_{mm'}^{JJ'} \Re(f_{\lambda_a\lambda_b}^J f_{\lambda_a\lambda_b}^{J'*}) - \Im\rho_{mm'}^{JJ'} \Im(f_{\lambda_a\lambda_b}^J f_{\lambda_a\lambda_b}^{J'*}). \quad (\text{B.10})$$

Therefore, under the conditions that we have assumed, the imaginary part of the SDM can be measured only if more spins are involved and at least one of the relative phases between the decay amplitudes is non-trivial.

References

- [1] U. Fano: *Rev. Mod. Phys.* **29** (1957) 74
- [2] P. Eberhard and M.L. Good: *Phys. Rev.* **120** (1960) 1442
- [3] M. Peshkin: *Phys. Rev.* **123** (1961) 637
- [4] M. Ademollo *et al.*: *Phys. Rev.* **139** (1965) B1608
- [5] P. Minnaert: *Phys. Rev.* **151** (1966) 1306
- [6] R.H. Dalitz: *Nucl. Phys.* **87** (1966) 89
- [7] M.G. Doncel *et al.*: *Nucl. Phys. B* **38** (1972) 89
- [8] S.U. Chung and T.L. Trueman: *Phys. Rev. D* **11** (1975) 633
- [9] A.J. Arifi *et al.*: *Phys. Rev. D* **101** (2020) 094023
- [10] R.L. Thews: *Phys. Rev.* **188** (1969) 2264
- [11] L.E. Holloway *et al.*: *Phys. Rev. Lett.* **27** (1971) 1671
- [12] M.G. Doncel *et al.*: *Nucl. Phys. B* **119** (1977) 515
- [13] F. Tabakin and R.A. Eisenstein: *Phys. Rev. C* **31** (1985) 1858
- [14] A. Chukanov *et al.*: *Eur. Phys. Jou. C* **46** (2006) 69
- [15] H. Chen and R.G. Ping: *Phys. Rev. D* **76** (2007) 036005

- [16] C. Abelev *et al.*: *Phys. Rev. C* **77** (2008) 061902
- [17] C. Amsler *et al.*: *Eur. Phys. Jou. C* **75** (2015) 124
- [18] E. Perotti *et al.*: *Phys. Rev. D* **99** (2019) 056008
- [19] C. Zhou: *Nucl. Phys. A* **982** (2019) 559
- [20] J.A. Aguilar-Saavedra and J. Bernabeu: *Nucl. Phys. B* **840** (2010) 349
- [21] J.A. Aguilar-Saavedra and J. Bernabeu: *Phys. Rev. D* **93** (2016) 011301
- [22] Z.J. Ajaltouni and E. Di Salvo: *Eur. Phys. Jou. C* **79** (2019) 989
- [23] T. Gutsche *et al.*: *Phys. Rev. D* **87** (2013) 074031
- [24] P. Boer, T. Feldmann and D. van Dyk: *JHEP* **01** (2015) 155
- [25] T. Blake and M. Kreps: *JHEP* **11** (2017) 138
- [26] R. Enomoto and M. Tanabashi: *Phys. Lett. B* **386** (1996) 413
- [27] K. Maltman *et al.*: *Phys. Lett. B* **376** (1996) 19
- [28] C. Bourrely *et al.*: *Phys. Rep.* **59** (1982) 95
- [29] E. Leader: "Spin in Particle Physics", Cambridge University Press, 2010
- [30] J. Daboul: *Nucl. Phys. B* **4** (1967) 180
- [31] S. Dagan and Y. Dothan: *Phys. Rev. D* **26** (1982) 248
- [32] Z.J. Ajaltouni *et al.*: *Phys. Lett. B* **614** (2005) 165
- [33] D. Zwanziger: *Phys. Rev.* **136** (1964) B558
- [34] M.G. Doncel *et al.*: "Matrices densité de polarisation", summer school of Gif-sur-Yvette 1970, (Ed. R. Salmeron, Laboratoire de Physique de l'Ecole Polytechnique, Paris)
- [35] S.U. Chung: "Spin Formalisms" - Lectures given in the Academic Training Programme at CERN 1969-1970 - Geneva 1971 - CERN 71-8
- [36] J. Adam *et al.*, STAR Coll.: *Phys. Rev. D* **98** (2018) 091103
- [37] J. Adam *et al.*, STAR Coll.: *Phys. Rev. D* **98** (2018) 112009
- [38] Z.J. Ajaltouni and E. Di Salvo: *Int. Jou. Mod. Phys. E* **22** (2013) 1330006; see also refs. therein

- [39] J. Hrivnac *et al.*: *J. Phys. G* **21** 629
- [40] R. Aaij *et al.*, LHCb Coll.: *JHEP* **06** (2020) 110
- [41] L. Kaldamäe *et al.*: *Phys. Rev. D* **94** (2016) 114003
- [42] M. Fischer *et al.*: *Phys. Rev. D* **97** (2018) 093001
- [43] S. Groote *et al.*: *Eur. Phys. Jou. C* **79** (2019) 948
- [44] S. Berge *et al.*: *Phys. Rev. D* **92** (2015) 033001
- [45] M. Fischer *et al.*: *Phys. Rev. D* **65** (2002) 054036
- [46] E. Di Salvo and Z.J. Ajaltouni: *Mod. Phys. Lett. A* **32** (2017) 1750043
- [47] S. Shivashankara *et al.*: *Phys. Rev. D* **91** (2015) 115003
- [48] T. Gutsche *et al.*: *Phys. Rev. D* **91** (2015) 119907
- [49] R. Dutta: *Phys. Rev. D* **93** (2016) 054003
- [50] E. Di Salvo *et al.*: *Int. Jou. Mod. Phys. A* **33** (2018) 1850169
- [51] R. Aaij *et al.*, LHCb Coll.: *Phys. Rev. Lett.* **115**, 072001 (2015)

- Proposed two methods for parametrizing the SDM of an unstable state - consisting of one or more spins - that is produced in various reactions. Both methods satisfy automatically all of the numerous non-negativity conditions and are adaptable to the constraints imposed by parity and angular momentum conservation.
- First method → theorem; applicable in a simple and flexible way. Improvements of the fit to the data when the rank of the SDM is less than its order. Added note on Eberhard-Good theorem.
- Second method: a variant of previous parametrizations; particularly suitable under especial conditions, e. g. , when the structure to be analyzed derives from some decay.
- examined the case where a given intermediate state is generated and decays according to strong or electromagnetic interactions: under very particular circumstances, the imaginary part of the SDM can be measured.
- Illustrated some applications; two of them are very interesting from the viewpoint of the search for new physics beyond the standard model.

Ziad J. Ajaltouni: stress on the importance of the spin density matrix and of non-covariant formalism in data analysis; remarks on the feasibility of an experiment and on the measurability of some parameters.

Declaration of Interest Statement

The authors have been collaborating since 2008 on phenomenology of weak decays and methods of data analysis. The spin density matrix is an essential tool to that aim.