



HAL
open science

Cannabis use is associated with a lower risk of diabetes in chronic hepatitis C-infected patients (ANRS CO22 Hepather cohort)

Tangui Barré, Marie Libérée Nishimwe, Camelia Protopopescu, Fabienne Marcellin, Fabrice Carrat, Céline Dorival, Elisabeth Delarocque-Astagneau, Dominique G. Larrey, Marc Bourlière, Ventzislava Petrov-Sanchez, et al.

► To cite this version:

Tangui Barré, Marie Libérée Nishimwe, Camelia Protopopescu, Fabienne Marcellin, Fabrice Carrat, et al.. Cannabis use is associated with a lower risk of diabetes in chronic hepatitis C-infected patients (ANRS CO22 Hepather cohort). *Journal of Viral Hepatitis*, 2020, 27 (12), pp.1473-1483. 10.1111/jvh.13380 . hal-03109215

HAL Id: hal-03109215

<https://hal.science/hal-03109215v1>

Submitted on 19 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3
4 **Cannabis use is associated with a lower risk of diabetes in**
5
6 **chronic hepatitis C-infected patients (ANRS CO22 Hepather cohort)**
7
8

9 **Running title**
10

11 Cannabis use and diabetes in HCV patients
12
13

14
15 Tangui Barré^{1,2}, Marie Libérée Nishimwe^{1,2}, Camelia Protopopescu^{1,2}, Fabienne
16 Marcellin^{1,2}, Fabrice Carrat³, Céline Dorival⁴, Elisabeth Delarocque-Astagneau⁵, Dominique
17 Larrey⁶, Marc Bourlière⁷, Ventzislava Petrov-Sanchez⁸, Mélanie Simony⁸, Stanislas Pol^{*9},
18 Hélène Fontaine^{*10, 11} and Patrizia Carrieri^{1,2} on behalf of the ANRS/AFEF Hepather study
19 group
20
21
22
23
24

25
26
27
28
29 ¹ Aix Marseille Univ, INSERM, IRD, SESSTIM, Sciences Économiques & Sociales de
30 la Santé & Traitement de l'Information Médicale, Marseille, France;
31

32
33 ² ORS PACA, Observatoire régional de la santé Provence-Alpes-Côte d'Azur,
34 Marseille, France;
35

36
37
38 ³ Sorbonne Université, Institut National de la Santé et de la Recherche Médicale
39 (INSERM), Institut Pierre Louis d'Epidémiologie et de Santé Publique, Paris, France;
40 Assistance Publique-Hôpitaux de Paris (AP-HP), Hôpital Saint-Antoine, Unité de Santé
41 Publique, Paris, France;
42
43
44

45
46
47 ⁴ Sorbonne Université, Institut National de la Santé et de la Recherche Médicale
48 (INSERM), Institut Pierre Louis d'Epidémiologie et de Santé Publique, Paris, France;
49

50
51 ⁵ UMR1181 Biostatistique, Biomathématique, Pharmaco-épidémiologie et Maladies
52 Infectieuses (B2PHI), Institut Pasteur, Inserm, Université de Versailles-Saint-Quentin-en-
53 Yvelines (UVSQ), Paris, France;
54
55
56
57
58
59
60

1
2
3 ⁶ Service des maladies de l'appareil digestif, Hôpital Saint Eloi, IBR- Inserm
4
5 Montpellier, France ;

6
7 ⁷ Department of Hepatology and Gastroenterology, Hôpital Saint Joseph, Marseille,
8
9 France;

10
11 ⁸ ANRS (France Recherche Nord & Sud Sida-HIV Hépatites), Unit for Basic and
12
13 Clinical Research on Viral Hepatitis, Paris, France ;

14
15 ⁹ Université de Paris ; AP-HP, Hôpital Cochin, Département d'Hépatologie; INSERM
16
17 U1223, Institut Pasteur, Paris, France;

18
19 ¹⁰ AP-HP, Hôpital Cochin, Département d'Hépatologie, Paris, France;

20
21 ¹¹ INSERM U1223, Institut Pasteur; Université Paris Descartes, Paris, France.

22
23 * SP and HF made an equal contribution to the work.
24
25
26
27
28
29
30

31 Acknowledgments

32
33
34
35 The cohort received financial support from the INSERM-ANRS (France Recherche
36
37 Nord & Sud Sida-HIV Hépatites), the French ANR (Agence Nationale de la Recherche), the
38
39 French DGS (Direction Générale de la Santé), Merck Sharp and Dohme, Janssen-Cilag,
40
41 Gilead, Abbvie, Bristol-Myers Squibb, and Roche. Those funding sources had no role in the
42
43 writing of the manuscript or the decision to submit it for publication.
44
45

46
47 We thank the study participants and the participating clinicians at each site. We also
48
49 thank the INSERM-ANRS for sponsoring, funding and conducting the ANRS CO22 Hepather
50
51 cohort in collaboration with the French Association for the Study of the liver (Association
52
53 Française pour l'Etude du Foie: AFEF). Finally, our thanks to Jude Sweeney for the English
54
55 revision and editing of our manuscript.
56
57
58
59
60

ANRS-AFEF Hepather study group

Delphine Bonnet, Virginie Sicart (CHU Purpan, Toulouse, France), François Bailly, Marjolaine Beaudoin, Dominique Giboz, Kerstin Hartig-Lavie, Marianne Maynard (Hospices Civils de Lyon, Lyon, France), Morane Cavellec, Marjorie Cheraud-Carpentier, François Raffi, Florian Vivrel (Hôpital Hôtel-Dieu, Nantes, France), Jaouad Benhida, Jérôme Boursier, Paul Calès, Françoise Lunel, Frédéric Oberti (CHU Angers, Angers, France), Nathalie Boyer, Audrey Gilibert, Nathalie Giully (Hôpital Beaujon, Clichy, France), Giovanna Scoazec (Hôpital Beaujon, Clichy, France and Hôpital Henri Mondor, Créteil, France), Sandrine Fernandes, Sylvie Keser, Philippe Sultanik, Anaïs Vallet-Pichard (Hôpital Cochin, Paris, France), Juliette Foucher, Jean-Baptiste Hiriart, Aurore Mathias, Julien Vergniol (Hôpital Haut-Lévêque, Pessac, Bordeaux, France), Chrystelle Ansaldi, Laëtitia Chouquet, Emilie De Luca, Valérie Oules (Hôpital Saint Joseph, Marseille, France), Rodolphe Anty, Eve Gelsi, Régine Truchi (CHU de Nice, Nice, France), Elena Luckina, Nadia Messaoudi, Joseph Moussali, (Groupe Hospitalier Pitié-Salpêtrière, Paris, France), Barbara De Dieuleveult, Damien Labarriere, Pascal Poter, Si Nafa Si Ahmed (CHR La Source, Orléans, France), Nathalie Ganne-Carrié, Véronique Grando-Lemaire, Pierre Nahon, Alan Peltier, Judith Ung (Hôpital Jean Verdier, Bondy, France), Mariette Gougeon, Anne Guillygomarch, Caroline Jezequel (CHU Rennes, Rennes, France), Romain Moirand, Thomas F. Baumert, Michel Dofföel, Catherine Mutter, Pauline Simo-Noumbissie (Hôpitaux Universitaires de Strasbourg, Strasbourg, France), Hélène Barraud, Mouni Bensenane, Abdelbasset Nani, Sarah Hassani-Nani (CHU de Nancy, Vandoeuvre-lès-Nancy, France), Marie-Albertine Bernard (CHU de Nancy, Vandoeuvre-lès-Nancy, France and Centre Hospitalier Régional, Metz, France), Michael Bismuth, Ludovic Caillo, Stéphanie Faure, Georges-Philippe Pageaux, Marie Pierre Ripault (Hôpital Saint Eloi, Montpellier, France), Karl Barange, Christophe Bureau, Jean Marie Peron, Marie Angèle Robic, Léa Tarallo (CHU Purpan, Toulouse, France), Marine Faure, Bruno Froissart, Marie-Noëlle Hilleret, Vincent Leroy (CHU de Grenoble, Grenoble,

1
2
3 France), Odile Gorla, Victorien Grard, H el ene Montialoux (CHU Charles Nicolle, Rouen,
4
5 France), Muriel Fran ois, Christian Ouedraogo, Christelle Pauleau, Anne Varault (H opital
6
7 Henri Mondor, Cr eteil, France), Tony Andreani, B enedicte Angoulevant, Azeline Chevance,
8
9 Lawrence Serfaty (H opital Saint-Antoine, Paris, France), Teresa Antonini, Audrey Coilly,
10
11 Jean-Charles Duclos Vall e, Mariagrazia Tateo (H opital Paul Brousse, Villejuif, France),
12
13 Armand Abergel, Corinne Bonny, Chanteranne Brigitte, G eraldine Lamblin, L eon Muti
14
15 (H opital Estaing, Clermont-Ferrand, France), Abdenour Babouri, Virginie Filipe (Centre
16
17 Hospitalier R egional, Metz, France), Camille Barrault, Laurent Costes, Herv e Hag e,
18
19 Soraya Merbah (Centre Hospitalier Intercommunal, Cr eteil, France), Paul Carrier, Maryline
20
21 Debette-Gratien, J er mie Jacques (CHU Limoges, Limoges, France), Florent Artu, Val erie
22
23 Canva, S ebastien Dharancy, Alexandre Louvet (CHRU Claude Huriez, Lille, France), Marc
24
25 Bardou, Donya Da Costa Souhiel, Patrick Hillon, Marianne Latournerie (Dijon University
26
27 Hospital, Dijon, France), Yannick Bacq, Didier Barbereau, Charlotte Nicolas (CHU
28
29 Trousseau, 37044 Tours, France), Nisserine Ben Amara, Dani ele Botta-Fridlund, Isabelle
30
31 Portal (CHU Timone, Marseille, France), Moana Gelu-Simeon, Marie-Jos ee Lafrance (CHU
32
33 de Pointe- a-Pitre, Pointe- a-Pitre, Guadeloupe).
34
35

36 37 **Financial support statement (authors):**

38
39
40 FC reports grants from INSERM-ANRS, during the conduct of the study; and personal fees
41
42 from Imaxio, outside the submitted work. MB reports grants and personal fees from AbbVie
43
44 and Gilead, outside the submitted work; and personal fees from MSD, Janssen, Boehringer
45
46 Ingelheim, Intercept, and Bristol-Myers Squibb, outside the submitted work. SP has received
47
48 consulting and lecturing fees from Bristol-Myers Squibb, Janssen, Gilead, Roche, MSD, MYR-
49
50 Pharma, Shionogi, Biotest and Abbvie, as well as grants from Bristol-Myers Squibb, Gilead,
51
52 Roche and MSD. HF reports personal fees and invitations for medical meetings from Gilead,
53
54 AbbVie, Bristol-Myers Squibb, MSD, and Janssen, outside the submitted work. Other authors
55
56 declare no financial support.
57
58

59 **Trial statement**

60

1
2
3 Data were derived from a trial registered with ClinicalTrials.gov, number NCT01953458.
4
5 Written informed consent was obtained from each cohort participant before enrolment. The
6
7 protocol was designed in accordance with the Declaration of Helsinki and French law for
8
9 biomedical research, and was approved by the 'Comité de Protection des Personnes (CPP)
10
11 Ile de France 3' Ethics Committee (Paris, France) and the French Regulatory Authority
12
13 (ANSM).
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Abstract

Background & Aims: Chronic hepatitis C virus (HCV) infection is a risk factor of insulin resistance, and HCV-infected patients are at a high risk of developing diabetes. In the general population, research has shown the potential benefit of cannabis use for the prevention of diabetes and related metabolic disorders. We aimed to test whether cannabis use is associated with a lower risk of diabetes in chronic HCV-infected patients.

Methods: Chronic HCV-infected patients (n=10,445) were selected from the French national, multicenter, observational ANRS CO22 Hepather cohort. Cross-sectional data collected at cohort enrolment were used to assess the association between patients' clinical and behavioral characteristics and the risk of diabetes. Logistic regression model was performed with cannabis use as the main independent variable and a significance level set at 5%. A similar model stratified by the presence of advanced liver fibrosis (FIB-4>3.25) was also run.

Results: After multivariable adjustment, current (AOR [95%CI]: 0.49 [0.38-0.63]) and former (0.81 [0.67-0.98], p<.001) cannabis use were both associated with a reduced odds of diabetes. Conversely, male gender, tobacco use, elevated BMI, poverty, being a migrant and advanced fibrosis were associated with increased odds of diabetes. The association between cannabis use and diabetes was maintained in the stratified analysis.

Conclusions: In this large cross-sectional study of chronic HCV-infected patients, cannabis use was associated with a lower risk of diabetes independently of clinical and socio-behavioral factors. Further studies are needed to elucidate a potential

causal link and shed light on cannabis compounds and mechanisms involved in this relationship.

Keywords:

Hepatitis C, Chronic; Cannabis; Diabetes Mellitus; Marijuana Smoking; France.

Introduction

With a global prevalence of over 500 million cases in 2018, type 2 diabetes is a growing health and economic burden^{1,2}, associated with increased all-cause mortality³. In 2015, the number of hepatitis C virus (HCV) viremic infections was estimated at more than 71 million worldwide. HCV is a leading cause of death and disability, especially through cirrhosis^{4,5}. HCV and diabetes are reciprocally intertwined: HCV infection triggers diabetes (mostly type 2) and diabetes worsens HCV outcomes, for instance by increasing the risk for cirrhosis⁶. Furthermore, the treatment of diabetes in cirrhotic patients is complex⁷. Diabetes is the second most common extrahepatic manifestation of HCV infection⁸. HCV infection is assumed to lead to hepatic insulin resistance through direct impairment of the hepatocyte insulin signaling pathway and to peripheral insulin resistance through hepatic production of mediators⁹. Insulin resistance constitutes the pathological background for type 2 diabetes development, and may also trigger hepatic steatosis, the latter being also promoted directly by HCV infection¹⁰. A difference is sometimes made between type 2 diabetes and a state of impaired glucose regulation caused by loss of liver function as a consequence of cirrhosis (referred to as 'hepatogenous diabetes')⁷.

1
2
3 In high-income countries, the HCV burden is mainly felt in marginalized
4 populations¹¹ such as people who inject drugs and homeless people. In Europe,
5 injecting drug use is the main route of HCV transmission¹². Moreover, people who
6 inject drugs face many barriers to access to care^{13,14}. A higher HCV prevalence is
7 associated with lower income, lower education level and less employment¹⁵. In
8 parallel, a low socioeconomic status is associated with poorer health in general¹⁶,
9 and with higher type 2 diabetes incidence in particular¹⁷. HCV-infected patients'
10 susceptibility to diabetes may therefore be exacerbated by socioeconomic factors.
11
12

13
14 In the general population, studies examining the association between
15 cannabis use and the risk of diabetes have yielded inconclusive results, while results
16 showing the protective effect of cannabis in terms of other metabolic disorders seem
17 more conclusive¹⁸. In a cross-sectional analysis of HIV-HCV co-infected patients,
18 cannabis use, irrespective of frequency, was associated with lower odds of insulin
19 resistance after adjustment for known confounders¹⁹. In chronic HCV-infected
20 patients, cannabis use was also associated with better liver disease outcomes²⁰.
21 However, there is still a lack of data regarding the association between cannabis use
22 and the risk of type 2 diabetes in HCV mono-infected patients. Using data from the
23 French ANRS CO22 Hepather cohort, we aimed to test whether cannabis use is
24 inversely associated with type 2 diabetes in HCV-infected patients.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 Material and methods

49 Participants

50
51 ANRS CO22 Hepather is a French national, multicenter, prospective,
52 observational cohort of adult hepatitis B (HBV)- and HCV-infected patients
53 (registered with ClinicalTrials.gov, number NCT01953458). The cohort design has
54
55
56
57
58
59
60

1
2
3 been extensively described elsewhere²¹. Set up in 2012, enrolment in the study took
4 place in 32 centers throughout France. Patients were recruited during a medical
5 follow-up visit in their center (eligible patients were proposed to participate in the
6 cohort). Demographic, clinical and biological data were collected at the inclusion visit.
7
8 Patients were followed-up on a yearly basis, and supplemental data were also
9 collected during visits related to particular events (e.g., initiation of HBV or HCV
10 therapy). Written informed consent was obtained from each cohort participant before
11 enrolment. The protocol was designed in accordance with the Declaration of Helsinki
12 and French law for biomedical research, and was approved by the 'Comité de
13 Protection des Personnes (CPP) Ile de France 3' Ethics Committee (Paris, France)
14 and the French Regulatory Authority (ANSM).
15
16
17
18
19
20
21
22
23
24
25
26
27
28

29 The present study population comprised patients with chronic HCV (defined as
30 positive HCV-RNA and anti-HCV antibodies) at cohort recruitment. The main
31 exclusion criteria in Hepather were HIV co-infection, being on HCV-treatment or
32 having stopped HCV treatment for less than 3 months at recruitment. We also
33 excluded patients with no data on diabetes diagnosis (n=145) and patients with no
34 cannabis use data (n=533), leaving us with a study sample of 10,445 patients (Figure
35
36
37
38
39
40
41
42
43 1).

44 45 Data collection

46
47
48 At the enrolment visit, patients completed a face-to-face interview with their physician
49 based on a structured questionnaire. Anthropological measurements and urine and
50 blood samples were also taken.
51
52
53

54
55 Data collected in the questionnaire included: lifetime (yes/no) and current (yes/no)
56 cannabis use, current tobacco use (yes/no), current alcohol consumption (number of
57
58
59
60

1
2
3 standard drinks per day), current coffee consumption (number of cups per day),
4
5 household income (average monthly income) and own and parents' country of birth.
6
7 Body height and weight were measured. Data derived from blood samples included
8
9 platelet count (platelets/mm³), aspartate aminotransferase (AST, IU/L) and alanine
10
11 aminotransferase (ALT, IU/L).
12
13

14 15 Outcome

16
17
18 The variable 'presence of diabetes' was used as the outcome. Diabetes diagnosis
19
20 ('Yes/No') was established based on a confirmed fasting blood sugar level of at least
21
22 126 mg/dL (7.0 mmol/L) and/or uptake of antidiabetic drugs.
23
24

25 Explanatory variables

26
27
28 People who answered 'yes' to the question concerning current use of cannabis were
29
30 classified into the 'current use' category. Among people who answered 'no', those
31
32 who reported lifetime cannabis use were classified into the "lifetime use" category.
33
34 Finally, people who answered 'no' to both questions (current/lifetime use) were
35
36 classified into the 'no lifetime use' category. We therefore ended up with a three-level
37
38 cannabis variable. According to previous studies in HIV-HCV co-infected patients,
39
40 elevated coffee consumption was defined as 3 or more cups of coffee per day²². A
41
42 second threshold at 4 cups of coffee was also tested. A meta-analysis highlighted a
43
44 decrease in relative risk of diabetes by 1 cup per day increase²³. Poverty was defined
45
46 as a standard of living lower than the 2015 French poverty threshold (1,015€ per
47
48 month). Standard of living was calculated as disposable income divided by the
49
50 number of consumption units in the household. Being a migrant was defined as being
51
52 born outside France and having at least one parent also born outside France. Liver
53
54 fibrosis was assessed through FIB-4 calculation, based on age, platelet count, AST
55
56 and ALT using the following formula: (age [years] * AST [IU/L]) / (platelet count [10⁹/L])
57
58
59
60

1
2
3 * (ALT [IU/L])^{1/2}. A cut-off >3.25 was used to define advanced fibrosis²⁴. The
4
5 following variables were also tested as explanatory variables: gender, age, current
6
7 tobacco use (yes/no), current alcohol consumption (abstinent, moderate, elevated),
8
9 and body mass index (BMI) with the following WHO classes²⁵: <18.5, underweight;
10
11 ≥18.5 and <25, normal weight; ≥25 and <30, overweight; ≥30, obese. Elevated
12
13 alcohol consumption was defined as >2 standard drinks per day for women and >3
14
15 standard drinks per day for men, according to French National Authority for Health²⁶.
16
17
18
19

20 Statistical analyses

21
22 The characteristics of diabetic patients were compared with those of non-diabetic
23
24 patients using a Chi-square test (categorical variables) or Student's t-test (continuous
25
26 variables). Similarly, the characteristics of the study population were compared
27
28 between cannabis use status (using Fisher's test for continuous variables). Logistic
29
30 regression models were used to test the association between cannabis use and
31
32 diabetes at cohort entry, after adjustment for other potential predictors. Only variables
33
34 with a liberal p-value <0.20 in the univariable analyses were considered eligible for
35
36 the multivariable model²⁷. The final multivariable model was built using a backward
37
38 stepwise procedure and the likelihood ratio test (p<0.05) was used to define the
39
40 variables to be maintained in the model. An interaction between cannabis use and
41
42 BMI was tested for in the final model because cannabis use is known to be
43
44 associated with a lower BMI¹⁸. Because cirrhosis is diabetogenic⁷, similar logistic
45
46 regressions stratified by fibrosis status were also performed to test whether the
47
48 relationship between cannabis use and diabetes may differ between patients with
49
50 advanced fibrosis and other patients. Results from logistic regression models were
51
52 presented as odds ratios and adjusted odds ratios (for the multivariable model), with
53
54
55
56
57
58
59
60

1
2
3 the corresponding 95% confidence intervals. The level of significance was set at
4
5 $\alpha=0.05$ in all tests. SAS version 9.4 software was used for all analyses.
6
7

8 9 Results

10 11 12 Study population characteristics

13
14
15
16 The characteristics of the study population are presented in Tables 1 and 2.

17
18
19 The study population was mainly male (56.6%). Median age was 56 years and 13.7%
20
21 of the patients had diabetes. A majority of the study population had never used
22
23 cannabis (67.3%), 18.7% were former users and 14.0% were current users.
24
25

26 27 28 Multivariable analyses

29
30
31 Despite being significantly associated with diabetes in univariable analysis, elevated
32
33 coffee consumption (≥ 3 cups/day) was not significantly associated with the presence
34
35 of diabetes ($p=0.54$) in the multivariable analysis. Increasing the threshold to ≥ 4
36
37 cups/day did not change this result ($p=0.30$). Accordingly, this variable was removed
38
39 from the final model. Results for the final model are given in Table 3. When artificially
40
41 maintained in the model, it did not change other relationships between the explicative
42
43 and dependent variables.
44
45

46
47
48 Current and former cannabis use were both associated with a lower odds of
49
50 diabetes (compared with 'no lifetime use') (adjusted odds-ratio (OR), 95% confidence
51
52 interval (CI): 0.49 [0.38-0.63] and 0.81 [0.67-0.98], respectively). The lower odds of
53
54 diabetes in current users than in former users was statistically significant (0.60 [0.46-
55
56 0.80], $p=0.0003$). Older age and being a man were positively associated with higher
57
58 odds of diabetes. BMI classes displayed a dose-dependent relationship with the odds
59
60

1
2
3 of diabetes. Moderate alcohol consumption was associated with a 23% lower odds of
4 diabetes (compared with abstinence), while current tobacco use was associated with
5 a 17% increased odds (Table 3). Both poverty and being a migrant were positively
6 associated with the odds of diabetes, while individuals with advanced liver fibrosis
7 had a 71% increased odds of diabetes. The interaction term between cannabis use
8 and BMI was not statistically significant ($p=0.23$).

9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

When comparing individuals included in the multivariable analysis with those excluded ($n=1,230$) because of missing values, the former were older (56.8 vs. 54.8 years, $p<0.001$), more likely to be male (55.7 vs. 52.3%, $p=0.023$), less likely to be current smokers (37.2 vs. 40.5%, $p=0.028$), and more likely to have elevated coffee consumption (69.3 vs. 64.6%, $p<0.001$). When stratifying by fibrosis status, current cannabis use was associated with a lower odds of diabetes (compared with 'no lifetime use') in patients with and without advanced fibrosis (Supporting tables 1 & 2).

Discussion

Using cross-sectional data from more than 9,000 HCV-infected patients, we found that former and current cannabis use were associated with, respectively, a 19% and 51% lower risk of diabetes when compared with no lifetime cannabis use. To our knowledge, this is the first time that such an association has been highlighted for HCV-infected patients. Under the assumption that the pattern of cannabis use differs only in terms of duration between former (shorter exposure) and current users (longer exposure), our results suggest that the potential protective effect of cannabis displays a dose-dependent relationship.

Our results are consistent with a meta-analysis in the U.S general population based on the National Health and Nutrition Examination Surveys (NHANES) and the

1
2
3 National Surveys on Drug Use and Health²⁸. In that meta-analysis, recently active
4 cannabis smoking and diabetes were inversely associated (compared with no lifetime
5 cannabis use), with a summary odds-ratio of 0.7 [0.6-0.8]. This confirmed previous
6
7
8 results from NHANES data which compared lifetime use with no lifetime use²⁹.
9

10
11
12 However, the only longitudinal analyses available revealed no association between
13
14
15 cannabis use and incidence of diabetes in the general population^{30,31}, while cannabis
16
17
18 use in young adulthood was associated with prediabetes incidence in middle age³¹.

19 Beyond specificities due to HCV infection, differences in lifestyle or cannabis material
20
21
22 and practices may impair comparability between those studies coming from Sweden
23
24
25 and the U.S. Moreover, in the Swedish study the prevalence of current cannabis use
26
27
28 was very low³⁰.

29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Insulin resistance is closely related to other metabolic dysregulations and to
metabolic syndrome. The effects of cannabis use on BMI, obesity and other
metabolic dysregulations are still controversial but tend to indicate that there is a
beneficial effect^{18,32}. Several cannabis compounds exerting potential beneficial
effects on these risk factors for diabetes have been identified. The endocannabinoid
system controls energy intake via both central and peripheral mechanisms, as well as
energy storage and expenditure, with direct implications for the development of type
2 diabetes³³. Exogenous cannabinoids from cannabis are likely to interact with the
endocannabinoid system and its control of energy balance. For instance, cannabidiol,
a major non-psychoactive constituent of cannabis, decreases both body weight and
food intake in rats^{34,35}.

Given that diabetes pathogenesis in HCV-infected patients involves specific
mechanisms, the effects of cannabis use we found may be partly mediated by
mechanisms different from those observed in the general population. For instance,

1
2
3 HCV infection causes systemic inflammation³⁶, this latter being a likely contributor to
4
5 type 2 diabetes occurrence^{37,38}. As cannabidiol found in cannabis possesses anti-
6
7 inflammatory properties³⁹, this indirect action may lead to a stronger effect of
8
9 cannabis on diabetes in HCV-infected patients. Moreover, HCV-infected patients are
10
11 at greater risk of liver disease, and cirrhosis is diabetogenic⁷. Accordingly, we found
12
13 that advanced liver fibrosis is a risk factor for diabetes. Based on a large database of
14
15 USA hospital discharge records, Adejumo *et al.* found that cannabis use was
16
17 associated with a lower prevalence of liver cirrhosis among chronic HCV-infected
18
19 patients²⁰. Among users, while the difference was not statistically different,
20
21 dependent users had an even lower prevalence of liver cirrhosis than non-dependent
22
23 users. This is in line with our results showing a lower risk of diabetes among current
24
25 cannabis users than in former users. While a recent systematic review and meta-
26
27 analysis concluded that there is no association between cannabis use and
28
29 progression of liver fibrosis in HCV and HIV-HCV co-infected patients⁴⁰, the
30
31 mediating effect of fibrosis and/or cirrhosis between cannabis and diabetes deserves
32
33 further investigation. However, as the analyses in the present study were adjusted for
34
35 advanced liver fibrosis, we hypothesize that other mechanisms are involved. Patients
36
37 achieving sustained virological response (SVR) after interferon-based therapy
38
39 exhibited improvements in HOMA-IR and HOMA- β , but no impact was found on
40
41 insulin resistance⁴¹. In the era of direct-acting antivirals for HCV, SVR rates of almost
42
43 100% are being reached, inducing clear benefits on the alteration of homeostasis of
44
45 glucose metabolism associated with HCV infection, including improvement of insulin
46
47 resistance and reduced onset of *de novo* insulin resistance and type 2 diabetes in
48
49 patients without advanced cirrhosis or long-term diabetic history⁴². For HCV patients
50
51 with type 2 diabetes, whether the beneficial short-term outcomes of DAA-induced
52
53
54
55
56
57
58
59
60

1
2
3 HCV clearance are maintained and of clinical significance in the long term remains
4
5 unclear⁴³.
6
7

8 The endocannabinoid system is present in the liver and is involved in the
9 control of various hepatic functions⁴⁴. Noticeably, hepatic cannabinoid receptor CB1
10 activation contributes to obesity-related insulin resistance. Endocannabinoids may
11 also be directly involved in diabetes development, given that they may favor beta cell
12 failure through peripheral CB1 receptors⁴⁵. One can hypothesize that exogenous
13 cannabinoids from cannabis interact with peripheral and/or hepatic endocannabinoid
14 systems. For instance, chronic administration in patients with type 2 diabetes of
15 delta-9-tetrahydrocannabivarin (THCV), which is found in cannabis, leads to a
16 reduction of fasting plasma glucose and improved pancreatic β -cell function without
17 affecting body weight or appetite⁴⁶. These effects could explain the seemingly
18 protective effect of cannabis use on diabetes development, even if other cannabis
19 compounds may be involved, directly or through interaction with one another.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 The gender difference in the risk of diabetes observed in our study is well-
37 known and is mediated through endocrine and behavioral factors⁴⁷. The increased
38 risk of diabetes associated with current tobacco use has already been shown in the
39 general population⁴⁸ and seems to be mediated through an impact on body weight,
40 body composition, peripheral insulin sensitivity and pancreatic beta cell function⁴⁹.
41 Moderate alcohol consumption has previously been identified as protecting against
42 type 2 diabetes in the general population, although the exact biological mechanisms
43 involved are still unknown⁵⁰. Given the positive association previously found between
44 alcohol consumption and physical activity, the former may be a proxy of healthy
45 behavior⁵¹, while abstinent persons may include individuals who stop drinking
46 because of previous health problems.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Agardh *et al.* in their meta-analysis found an overall higher risk of diabetes in
4 groups with a low socioeconomic status, specifically in terms of educational level,
5 occupation and income¹⁷. This is in line with the associations we found between
6 poverty and the risk of diabetes. There is now mounting evidence that socioeconomic
7 factors such as income or education are risk factors for a wide range of health
8 outcomes¹⁶. For instance, it has been shown that poor people covered by
9 complementary universal health cover are frequently denied care in France⁵². With
10 regards to immigrants, in France in 2015, approximately 31% were from the
11 European Union, while 45% were from Africa⁵³. We therefore expect that a majority
12 of patients identified as migrants in our study were African. The most recent data
13 indicate that immigrant households have a higher prevalence of poverty, are
14 overrepresented in the less wealthy half of France population, and that almost a third
15 of immigrant households where the head of the household was born in Africa have a
16 standard of living lower than the first decile⁵⁴. The association we found between
17 being born outside France and the risk of diabetes is therefore certainly related to
18 socioeconomic factors, but genetic contributions may make the burden heavier for
19 some sub-populations than for others⁵⁵. We did not find a significant association
20 between elevated coffee consumption and the risk of diabetes after multivariable
21 adjustment, despite robust evidence found in the general population²³. Given that
22 elevated coffee consumption seems to protect HIV-HCV co-infected patients from
23 liver fibrosis²² and that cirrhosis is a cause of diabetes, we may hypothesize that
24 adjusting for fibrosis, as we did, lessens the protective effect of coffee on diabetes.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53
54 The main strength of the present study is the sample size and the sampling
55 procedure likely to provide data representative of HCV-infected French patients.
56
57

58 Despite its cross-sectional nature, with more than 9,000 patients and a significant
59
60

1
2
3 proportion of cannabis users, it provides the first evidence of the potential protective
4 effect of cannabis use on diabetes among HCV-infected patients. As diabetes
5 diagnosis is more likely to encourage more health-conscious behavior, it is unlikely
6 that reverse causality is involved in the association we found between cannabis use
7 and diabetes. Longitudinal studies are nonetheless required to confirm these
8 findings. Multivariable adjustment for clinical, behavioral and socio-economic
9 variables is another strength of the study. It is now well-recognized that these
10 variables impact to a great extent health outcomes. This study has limitations.
11 Collection of cannabis data during face-to-face interviews with physicians may have
12 led to underreporting of use. However, such underreporting would be unlikely to be
13 different between the two groups (i.e., with and without diabetes). Furthermore, such
14 a bias is less likely for former cannabis users. To further assess the dose-dependent
15 nature of the relationship, more precise data regarding cannabis use would have
16 been valuable. We did not have data regarding diet quality or physical activity, which
17 are known risk factor for diabetes development. However, they are related to BMI
18 which was included, and if they were available, reverse causality would have
19 impaired the interpretation (diabetic patients may be more attentive to their lifestyle).
20 Being able to take into account history of HCV treatment (information which was not
21 available in our study) would have been beneficial given that interferon-based
22 therapeutic regimens may be involved in type 1 diabetes development⁶. The lack of
23 data differentiating type 2 from type 1 diabetes also impairs the interpretation of the
24 results. However, HCV is recognized as triggering mostly type 2 diabetes and only
25 occasionally type 1 diabetes⁶. Therefore, given the high prevalence of type 2
26 diabetes among HCV infected patients and that the relationship between type 2
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 diabetes and HCV is unequivocal⁵⁶, we may expect that the vast majority of diabetes
4 cases were type 2.
5
6
7

8 9 Conclusions

10
11 In conclusion, we found that cannabis use is associated with a lower risk of
12 diabetes in HCV-infected patients. Further studies are needed to longitudinally
13 confirm these results and investigations into cannabis compounds and their biological
14 effects hold great promise for the treatment and prevention of diabetes.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

References

1. Kaiser AB, Zhang N, Pluijm WVD. Global Prevalence of Type 2 Diabetes over the Next Ten Years (2018-2028). *Diabetes*. 2018;67(Supplement 1):202-LB. doi:10.2337/db18-202-LB
2. Bommer C, Sagalova V, Heesemann E, et al. Global Economic Burden of Diabetes in Adults: Projections From 2015 to 2030. *Diabetes Care*. 2018;41(5):963-970. doi:10.2337/dc17-1962
3. Engelmann J, Manuwald U, Rubach C, et al. Determinants of mortality in patients with type 2 diabetes: a review. *Rev Endocr Metab Disord*. 2016;17(1):129-137. doi:10.1007/s11154-016-9349-0
4. Blach S, Zeuzem S, Manns M, et al. Global prevalence and genotype distribution of hepatitis C virus infection in 2015: a modelling study. *The Lancet Gastroenterology & Hepatology*. 2017;2(3):161-176. doi:10.1016/S2468-1253(16)30181-9
5. Stanaway JD, Flaxman AD, Naghavi M, et al. The global burden of viral hepatitis from 1990 to 2013: findings from the Global Burden of Disease Study 2013. *Lancet*. 2016;388(10049):1081-1088. doi:10.1016/S0140-6736(16)30579-7
6. Hammerstad SS, Grock SF, Lee HJ, Hasham A, Sundaram N, Tomer Y. Diabetes and Hepatitis C: A Two-Way Association. *Front Endocrinol (Lausanne)*. 2015;6. doi:10.3389/fendo.2015.00134
7. García-Compeán D, González-González JA, Lavallo-González FJ, González-Moreno EI, Villarreal-Pérez JZ, Maldonado-Garza HJ. Current Concepts in Diabetes Mellitus and Chronic Liver Disease: Clinical Outcomes, Hepatitis C Virus Association, and Therapy. *Dig Dis Sci*. 2016;61(2):371-380. doi:10.1007/s10620-015-3907-2
8. Younossi Z, Park H, Henry L, Adeyemi A, Stepanova M. Extrahepatic Manifestations of Hepatitis C: A Meta-analysis of Prevalence, Quality of Life, and Economic Burden. *Gastroenterology*. 2016;150(7):1599-1608. doi:10.1053/j.gastro.2016.02.039
9. Drazilova S, Gazda J, Janicko M, Jarcuska P. Chronic Hepatitis C Association with Diabetes Mellitus and Cardiovascular Risk in the Era of DAA Therapy. *Can J Gastroenterol Hepatol*. 2018;2018:6150861. doi:10.1155/2018/6150861
10. Serfaty L. Metabolic Manifestations of Hepatitis C Virus: Diabetes Mellitus, Dyslipidemia. *Clin Liver Dis*. 2017;21(3):475-486. doi:10.1016/j.cld.2017.03.004
11. Edmunds BL, Miller ER, Tsourtos G. The distribution and socioeconomic burden of Hepatitis C virus in South Australia: a cross-sectional study 2010–2016. *BMC Public Health*. 2019;19. doi:10.1186/s12889-019-6847-5
12. European Monitoring Centre for Drugs and Drug Addiction. *Hepatitis C among Drug Users in Europe: Epidemiology, Treatment and Prevention.*; 2016. http://www.emcdda.europa.eu/publications/insights/hepatitis-c-among-drug-users-in-europe_en. Accessed August 29, 2018.

13. Dillon JF, Lazarus JV, Razavi HA. Urgent action to fight hepatitis C in people who inject drugs in Europe. *Hepatology, Medicine and Policy*. 2016;1(1):2. doi:10.1186/s41124-016-0011-y
14. Neale J, Tompkins C, Sheard L. Barriers to accessing generic health and social care services: a qualitative study of injecting drug users. *Health Soc Care Community*. 2008;16(2):147-154. doi:10.1111/j.1365-2524.2007.00739.x
15. Omland LH, Osler M, Jepsen P, et al. Socioeconomic status in HCV infected patients – risk and prognosis. *Clin Epidemiol*. 2013;5:163-172. doi:10.2147/CLEP.S43926
16. Braveman P, Gottlieb L. The Social Determinants of Health: It's Time to Consider the Causes of the Causes. *Public Health Rep*. 2014;129(Suppl 2):19-31.
17. Agardh E, Allebeck P, Hallqvist J, Moradi T, Sidorchuk A. Type 2 diabetes incidence and socio-economic position: a systematic review and meta-analysis. *Int J Epidemiol*. 2011;40(3):804-818. doi:10.1093/ije/dyr029
18. Sidney S. Marijuana Use and Type 2 Diabetes Mellitus: a Review. *Curr Diab Rep*. 2016;16(11):117. doi:10.1007/s11892-016-0795-6
19. Carrieri MP, Serfaty L, Vilotitch A, et al. Cannabis Use and Reduced Risk of Insulin Resistance in HIV-HCV Infected Patients: A Longitudinal Analysis (ANRS CO13 HEPAVIH). *Clin Infect Dis*. 2015;61(1):40-48. doi:10.1093/cid/civ217
20. Adejumo AC, Adegbala OM, Adejumo KL, Bukong TN. Reduced Incidence and Better Liver Disease Outcomes among Chronic HCV Infected Patients Who Consume Cannabis. *Can J Gastroenterol Hepatol*. 2018;2018:9430953. doi:10.1155/2018/9430953
21. Pol S, Bourliere M, Lucier S, et al. Safety and efficacy of daclatasvir-sofosbuvir in HCV genotype 1-mono-infected patients. *J Hepatol*. 2017;66(1):39-47. doi:10.1016/j.jhep.2016.08.021
22. Carrieri MP, Protopopescu C, Marcellin F, et al. The impact of coffee consumption on fibrosis and steatosis in HIV-HCV co-infected patients. *J Hepatol*. November 2017. doi:10.1016/j.jhep.2017.10.025
23. Ding M, Bhupathiraju SN, Chen M, van Dam RM, Hu FB. Caffeinated and decaffeinated coffee consumption and risk of type 2 diabetes: a systematic review and a dose-response meta-analysis. *Diabetes Care*. 2014;37(2):569-586. doi:10.2337/dc13-1203
24. Sterling Richard K., Lissen Eduardo, Clumeck Nathan, et al. Development of a simple noninvasive index to predict significant fibrosis in patients with HIV/HCV coinfection. *Hepatology*. 2006;43(6):1317-1325. doi:10.1002/hep.21178
25. World Health Organization. Body mass index - BMI. <http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi>. Published July 9, 2019. Accessed July 9, 2019.
26. Haute Autorité de Santé (French National Authority for Health). *Screening tool for early detection and brief intervention (Outil d'aide au repérage précoce et à l'intervention brève)*.; 2014. https://www.has-sante.fr/jcms/c_1795221/fr/outil-d-aide-au-reperage-precoce-et-intervention-breve-alcool-cannabis-tabac-chez-l-adulte. Accessed October 24, 2019.

- 1
- 2
- 3 27. Hosmer D W, Lemeshow S. *Applied Logistic Regression*. Second edition. John Wiley & Sons, Inc.
4 <https://onlinelibrary.wiley.com/doi/book/10.1002/0471722146>. Accessed October 8, 2019.
- 5
- 6 28. Alshaarawy O, Anthony JC. Cannabis Smoking and Diabetes Mellitus: Results from Meta-
7 analysis with Eight Independent Replication Samples. *Epidemiology*. 2015;26(4):597-600.
8 doi:10.1097/EDE.0000000000000314
- 9
- 10 29. Rajavashisth TB, Shaheen M, Norris KC, et al. Decreased prevalence of diabetes in marijuana
11 users: cross-sectional data from the National Health and Nutrition Examination Survey
12 (NHANES) III. *BMJ Open*. 2012;2(1):e000494. doi:10.1136/bmjopen-2011-000494
- 13
- 14 30. Danielsson AK, Lundin A, Yaregal A, Östenson CG, Allebeck P, Agardh EE. Cannabis Use as Risk
15 or Protection for Type 2 Diabetes: A Longitudinal Study of 18 000 Swedish Men and Women. *J*
16 *Diabetes Res*. 2016;2016. doi:10.1155/2016/6278709
- 17
- 18 31. Banks MP, Pletcher MJ, Kertesz SG, Sidney S, Rana JS, Schreiner PJ. Marijuana use and risk of
19 prediabetes and diabetes by middle adulthood: the Coronary Artery Risk Development in Young
20 Adults (CARDIA) study. *Diabetologia*. 2015;58(12):2736-2744. doi:10.1007/s00125-015-3740-3
- 21
- 22 32. Ngueta G, Bélanger RE, Laouan-Sidi EA, Lucas M. Cannabis use in relation to obesity and insulin
23 resistance in the Inuit population. *Obesity (Silver Spring)*. 2015;23(2):290-295.
24 doi:10.1002/oby.20973
- 25
- 26 33. Di Marzo V, Piscitelli F, Mechoulam R. Cannabinoids and endocannabinoids in metabolic
27 disorders with focus on diabetes. *Handb Exp Pharmacol*. 2011;(203):75-104. doi:10.1007/978-
28 3-642-17214-4_4
- 29
- 30 34. Ignatowska-Jankowska B, Jankowski MM, Swiergiel AH. Cannabidiol decreases body weight gain
31 in rats: involvement of CB2 receptors. *Neurosci Lett*. 2011;490(1):82-84.
32 doi:10.1016/j.neulet.2010.12.031
- 33
- 34 35. Farrimond JA, Whalley BJ, Williams CM. Cannabinol and cannabidiol exert opposing effects on
35 rat feeding patterns. *Psychopharmacology*. 2012;223(1):117-129. doi:10.1007/s00213-012-
36 2697-x
- 37
- 38 36. Zampino R, Marrone A, Restivo L, et al. Chronic HCV infection and inflammation: Clinical impact
39 on hepatic and extra-hepatic manifestations. *World J Hepatol*. 2013;5(10):528-540.
40 doi:10.4254/wjh.v5.i10.528
- 41
- 42 37. Lontchi-Yimagou E, Sobngwi E, Matsha TE, Kengne AP. Diabetes mellitus and inflammation. *Curr*
43 *Diab Rep*. 2013;13(3):435-444. doi:10.1007/s11892-013-0375-y
- 44
- 45 38. Pedersen BK. Anti-inflammatory effects of exercise: role in diabetes and cardiovascular disease.
46 *Eur J Clin Invest*. 2017;47(8):600-611. doi:10.1111/eci.12781
- 47
- 48 39. Booz GW. Cannabidiol as an Emergent Therapeutic Strategy for Lessening the Impact of
49 Inflammation on Oxidative Stress. *Free Radic Biol Med*. 2011;51(5):1054-1061.
50 doi:10.1016/j.freeradbiomed.2011.01.007
- 51
- 52 40. Farooqui MT, Khan MA, Cholankeril G, et al. Marijuana is not associated with progression of
53 hepatic fibrosis in liver disease: a systematic review and meta-analysis. *Eur J Gastroenterol*
54 *Hepatol*. 2019;31(2):149-156. doi:10.1097/MEG.0000000000001263
- 55
- 56
- 57
- 58
- 59
- 60

- 1
- 2
- 3 41. Hu J-H, Chang M-L, Liu N-J, Yeh C-T, Huang T-J. Effect of HCV treatment response on insulin
- 4 resistance: A systematic review and meta-analysis. *Exp Ther Med*. 2019;18(5):3568-3578.
- 5 doi:10.3892/etm.2019.7995
- 6
- 7 42. Adinolfi LE, Rinaldi L, Marrone A, Giordano M. The effect of sustained virological response by
- 8 direct-acting antivirals on insulin resistance and diabetes mellitus in patients with chronic
- 9 hepatitis C. *Expert Review of Anti-infective Therapy*. 2018;16(8):595-597.
- 10 doi:10.1080/14787210.2018.1505500
- 11
- 12 43. Ribaldone DG, Sacco M, Saracco GM. The Effect of Viral Clearance Achieved by Direct-Acting
- 13 Antiviral Agents on Hepatitis C Virus Positive Patients with Type 2 Diabetes Mellitus: A Word of
- 14 Caution after the Initial Enthusiasm. *J Clin Med*. 2020;9(2). doi:10.3390/jcm9020563
- 15
- 16 44. Tam J, Liu J, Mukhopadhyay B, Cinar R, Godlewski G, Kunos G. Endocannabinoids in liver
- 17 disease. *Hepatology*. 2011;53(1):346-355. doi:10.1002/hep.24077
- 18
- 19 45. Gruden G, Barutta F, Kunos G, Pacher P. Role of the endocannabinoid system in diabetes and
- 20 diabetic complications. *Br J Pharmacol*. 2016;173(7):1116-1127. doi:10.1111/bph.13226
- 21
- 22 46. Jadoon KA, Ratcliffe SH, Barrett DA, et al. Efficacy and Safety of Cannabidiol and
- 23 Tetrahydrocannabivarin on Glycemic and Lipid Parameters in Patients With Type 2 Diabetes: A
- 24 Randomized, Double-Blind, Placebo-Controlled, Parallel Group Pilot Study. *Diabetes Care*.
- 25 2016;39(10):1777-1786. doi:10.2337/dc16-0650
- 26
- 27 47. Kautzky-Willer A, Harreiter J, Pacini G. Sex and Gender Differences in Risk, Pathophysiology and
- 28 Complications of Type 2 Diabetes Mellitus. *Endocr Rev*. 2016;37(3):278-316.
- 29 doi:10.1210/er.2015-1137
- 30
- 31 48. Pan A, Wang Y, Talaei M, Hu FB, Wu T. Relation of active, passive, and quitting smoking with
- 32 incident type 2 diabetes: a systematic review and meta-analysis. *Lancet Diabetes Endocrinol*.
- 33 2015;3(12):958-967. doi:10.1016/S2213-8587(15)00316-2
- 34
- 35 49. Maddatu J, Anderson-Baucum E, Evans-Molina C. Smoking and the Risk of Type 2 Diabetes.
- 36 *Transl Res*. 2017;184:101-107. doi:10.1016/j.trsl.2017.02.004
- 37
- 38 50. Baliunas DO, Taylor BJ, Irving H, et al. Alcohol as a risk factor for type 2 diabetes: A systematic
- 39 review and meta-analysis. *Diabetes Care*. 2009;32(11):2123-2132. doi:10.2337/dc09-0227
- 40
- 41 51. Piazza-Gardner AK, Barry AE. Examining physical activity levels and alcohol consumption: are
- 42 people who drink more active? *Am J Health Promot*. 2012;26(3):e95-104.
- 43 doi:10.4278/ajhp.100929-LIT-328
- 44
- 45 52. Le défenseur des droits. *Les Refus de Soins Opposés Aux Bénéficiaires de La CMU-C, de l'ACS et*
- 46 *de l'AME*. Paris; 2014. [https://www.cmu.fr/fichier-](https://www.cmu.fr/fichier-utilisateur/fichiers/Rapport_refus_soins_DDD.pdf)
- 47 [utilisateur/fichiers/Rapport_refus_soins_DDD.pdf](https://www.cmu.fr/fichier-utilisateur/fichiers/Rapport_refus_soins_DDD.pdf).
- 48
- 49 53. Institut National de la Statistique et des Etudes Economiques. Chap. 3.7 - Étrangers - Immigrés.
- 50 *Tableaux de l'économie française*. 2019.
- 51 <https://www.insee.fr/fr/statistiques/fichier/3696937/TEF2019.pdf>.
- 52
- 53 54. Institut National de la Statistique et des Etudes Economiques. Chap. 1.22 Niveau de vie et
- 54 pauvreté des immigrés. *Les revenus et le patrimoine des ménages*. 2018.
- 55 <https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=2&cad=rja&uact=8&ved>
- 56
- 57
- 58
- 59
- 60

1
2
3 =2ahUKEwj36GW4uHjAhXM34UKHYm0CaIQFjABegQIBBAC&url=https%3A%2F%2Fwww.insee.
4 fr%2Ffr%2Fstatistiques%2Ffichier%2F3549496%2FREVPME18_F1.22_niv-pauv-
5 immi.pdf&usg=AOvVaw1Dc_kU8Ui-TwgZ_ljrCsFK.
6

- 7
8 55. Cheng C-Y, Reich D, Haiman CA, et al. African ancestry and its correlation to type 2 diabetes in
9 African Americans: a genetic admixture analysis in three U.S. population cohorts. *PLoS ONE*.
10 2012;7(3):e32840. doi:10.1371/journal.pone.0032840
11
12 56. Antonelli A, Ferrari SM, Giuggioli D, et al. Hepatitis C virus infection and type 1 and type 2
13 diabetes mellitus. *World J Diabetes*. 2014;5(5):586-600. doi:10.4239/wjd.v5.i5.586
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Tables

Table 1: Characteristics of the study population according to diabetes status (ANRS CO22 Hepather cohort, n=10,445).

	All participants N=10,445	No diabetes N=8,988 (86.1%)	Diabetes N=1,457 (13.9%)	P-value†
	N (%)	N	%	N
Gender				
Female	4668 (44.7)	4129	88.5	539
Male	5777 (55.3)	4859	84.1	918
Cannabis use				
Never	7030 (67.3)	5883	83.7	1147
Former	1956 (18.7)	1740	89.0	216
Current	1459 (14.0)	1365	93.6	94
Tobacco use				
No or former	3928 (37.6)	3312	84.3	616
Current	6515 (62.4)	5674	87.1	841
Missing	2 (0.02)	2	100.0	.

Coffee consumption									
Low (<3 cups/day)	7185 (68.8)	6097	84.9	1088	15.1	<.001			
Elevated (≥3 cups/day)	3260 (31.2)	2891	88.7	369	11.3				
Body mass index									
Underweight (<18.5)	346 (3.3)	325	93.9	21	6.1	<.001			
Normal (≥18.5 and <25)	5259 (50.3)	4779	90.9	480	9.1				
Overweight (≥25 and <30)	3332 (31.9)	2773	83.2	559	16.8				
Obese (>30)	1414 (13.5)	1032	73.0	382	27.0				
Missing	94 (0.9)	79	84.0	15	16.0				
Alcohol consumption‡									
Abstinent	5972 (57.2)	5002	83.8	970	16.2	<.001			
Moderate	3980 (38.1)	3545	89.1	435	10.9				
Elevated	441 (4.2)	400	90.7	41	9.3				
Missing	52 (0.5)	41	78.9	11	21.2				
Poverty§									
No	6742 (64.5)	5896	87.5	846	12.6	<.001			

Yes	3360 (32.2)	2800	83.3	560	16.7	
Missing	343 (3.3)	292	85.1	51	14.9	
Being a migrant[¶]						
No	7563 (72.4)	6691	88.5	872	11.5	<.001
Yes	2869 (27.5)	2285	79.6	584	20.4	
Missing	13 (0.1)	12	92.3	1	7.7	
Advanced liver fibrosis (FIB-4>3.25)						
No	7106 (68.0)	6291	88.5	815	11.5	<.001
Yes	2511 (24.0)	1956	77.9	555	22.1	
Missing	828 (7.9)	741	89.5	87	10.5	
Age (mean years (STD))(missing=14)	56.6 (11.6)	55.9 (11.6)		60.9 (10.2)		<.001

† Chi-square test (categorical variables) or Student's t-test (continuous variable, i.e. age).

‡ Elevated alcohol consumption was defined as >2 standard drinks per day for women and >3 standard drinks per day for men. §Poverty was defined as a standard of living lower than the 2015 French poverty threshold (1,015€ per month per household consumption unit). ¶Being a migrant was defined as being born outside France and having at least one parent also born outside France. STD, standard deviation.

For Peer Review

Table 2: Characteristics of the study population according to cannabis use (ANRS CO22 Hepather cohort, n=10,445).

	Current use		Former use		No lifetime use		P-value†
	N	%	N	%	N	%	
Gender							
Female	300	6.4	545	11.7	3823	81.9	<.001
Male	1159	20.1	1411	24.4	3207	55.5	
Cannabis use							
Never	7030	100.0	
Former	.	.	1956	100.0	.	.	
Current	1459	100.0	
Tobacco use							
No or former	51	1.3	80	2.0	3797	96.7	<.001
Current	1408	21.6	1876	28.8	3231	49.6	
Missing	2	100.0	
Coffee consumption							

Low (<3 cups/day)	786	10.9	1166	16.2	5233	72.8	<.001
Elevated (≥3 cups/day)	673	20.6	790	24.2	1797	55.1	
Body mass index							
Underweight (<18.5)	103	29.8	69	19.9	174	50.3	<.001
Normal (≥18.5 and <25)	927	17.6	1053	20.0	3279	62.4	
Overweight (≥25 and <30)	335	10.1	613	18.4	2384	71.6	
Obese (>30)	89	6.2	201	14.2	1124	79.5	
Missing	5	5.3	20	21.3	69	73.4	
Alcohol consumption[‡]							
Abstinent	650	10.9	968	16.2	4354	72.9	<.001
Moderate	624	15.7	866	21.8	2490	62.6	
Elevated	171	38.8	107	24.3	163	37.0	
Missing	14	26.9	15	28.9	23	44.2	
Poverty^s							
No	792	11.8	1292	19.2	4658	69.1	<.001

Yes	634	18.9	621	18.5	2105	62.7	
Missing	33	9.6	43	12.5	267	77.8	
Being a migrant†							
No	1255	16.6	1658	21.9	4650	61.5	<.001
Yes	203	7.1	296	10.3	2370	82.6	
Missing	1	7.7	2	15.4	10	76.9	
Advanced liver fibrosis (FIB-4>3.25)							
No	1063	15.0	1351	19.0	4692	66.0	<.001
Yes	295	11.8	441	17.6	1775	70.7	
Missing	101	12.2	164	19.8	563	68.0	
Age (mean years, (STD))(missing=14)	50.6 (7.8)		52.2 (7.2)		59.0 (12.4)		<.001

† Chi-square test (categorical variables) or Fisher's test (continuous variable, i.e. age).

1
2
3 † Elevated alcohol consumption was defined as >2 standard drinks per day for women and >3 standard drinks per day for men. §Poverty was
4
5 defined as a standard of living lower than the 2015 French poverty threshold (1,015€ per month per household consumption unit). ¶Being a
6
7 migrant was defined as being born outside France and having at least one parent also born outside France. STD, standard deviation.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

For Peer Review

Table 3: Factors associated with the presence of diabetes, logistic regression models, univariable and multivariable analyses (ANRS CO22 Hepather cohort, n=10,445)

	Univariable analyses (n=10,445)			Multivariable analysis (n=9,215)		
	OR	95% CI	P-value	AOR	95% CI	P-value
Gender						
Female	1			1		
Male	1.45	1.29-1.62	<.001	1.86	1.61-2.14	<.001
Cannabis use						
No lifetime use	1			1		
Former	0.64	0.55-0.74	<.001	0.81	0.67-0.98	0.031
Current	0.35	0.28-0.44	<.001	0.49	0.38-0.63	<.001
Tobacco use						
No lifetime use or former use	1			1		

Current	0.80	0.71-0.89	<.001	1.17	1.01- 1.36	0.036
---------	------	-----------	-------	------	---------------	-------

Coffee consumption

Low (<3 cups/day)	1					
Elevated (≥ 3 cups/day)	0.72	0.63-0.81	<.001			

Body mass index

Underweight (<18.5)	0.64	0.41-1.01	0.056	0.92	0.57- 1.46	0.708
Normal (≥ 18.5 and <25)	1			1		
Overweight (≥ 25 and <30)	2.01	1.76-2.29	<.001	1.65	1.43- 1.91	<.001
Obese (>30)	3.69	3.17-4.28	<.001	3.33	2.82- 3.92	<.001

Alcohol consumption[†]

Abstinent	1			1		
Moderate	0.63	0.56-0.71	<.001	0.77	0.68- 0.89	<.001
Elevated	0.53	0.38-0.73	<.001	0.72	0.50- 1.03	0.075

Poverty[‡]

No	1			1		
Yes	1.39	1.24-1.57	<.001	1.32	1.15- 1.51	<.001
Being a migrants§						
No	1			1		
Yes	1.96	1.75-2.20	<.001	1.51	1.31- 1.74	<.001
Advanced liver fibrosis (FIB-4>3.25)						
No	1			1		
Yes	2.19	1.94-2.47	<.001	1.71	1.50- 1.95	<.001
Age (years)						
	1.04	1.03-1.05	<.001	1.04	1.03- 1.05	<.001

† Elevated alcohol consumption was defined as >2 standard drinks per day for women and >3 standard drinks per day for men. ‡Poverty was defined as a standard of living lower than the 2015 French poverty threshold (1,015€ per month per household consumption unit). §Being a migrant was defined as being born outside France and having at least one parent also born outside France. (A)OR, (adjusted) odds ratio; CI, confidence interval.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1: Flowchart of the study population (ANRS CO22 Hepather cohort).

For Peer Review




Figure 1: Flowchart of the study population (ANRS CO22 Hepather cohort).

34x30mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supporting table 1: Factors associated with the presence of diabetes in patients without advanced fibrosis, logistic regression models, univariable and multivariable analyses (ANRS CO22 Hepather cohort, n=7,106)

	Univariable analyses (n=7,106)			Multivariable analysis (n=6,813)		
	OR	95% CI	P-value	AOR	95% CI	P-value
Gender						
Female	1		<.001	1		<.001
Male	1.38	1.19-1.60		1.83	1.53-2.19	
Cannabis use						
No lifetime use	1		<.001	1		0.001
Former	0.66	0.54-0.80		0.89	0.70-1.14	
Current	0.34	0.26		0.55	0.40-0.75	
Tobacco use						
No lifetime use or former use	1		0.011	1		0.006
Current	0.83	0.71-0.96		1.30	1.08-1.57	
Coffee consumption						
Low (<3 cups/day)	1		0.010			

Elevated (≥ 3 cups/day) 0.82 0.70-0.95

Body mass index

Underweight (< 18.5) 0.63 0.34-1.16 $< .001$ 0.79 0.42-1.49 $< .001$

Normal (≥ 18.5 and < 25) 1 1

Overweight (≥ 25 and < 30) 2.37 1.20-2.82 1.89 1.57-2.28

Obese (> 30) 4.55 3.75-5.54 3.94 3.20-4.87

Alcohol consumption†

Abstinent 1 $< .001$ 1 $< .001$

Moderate 0.56 0.48-0.66 0.66 0.55-0.78

Elevated 0.41 0.25-0.67 0.52 0.30-0.89

Poverty‡

No 1 $< .001$ 1 0.008

Yes 1.33 1.15-1.55 1.27 1.06-1.52

Being a migrant§

No 1 $< .001$ 1 $< .001$

1
2
3 Yes 2.00 1.72-2.32 1.56 1.3-1.87
4
5

6
7 **Age (years)** 1.05 1.04-1.05 <.001 1.05 1.04-
8 1.06 <.001
9
10

11 † Elevated alcohol consumption was defined as >2 standard drinks per day for women and
12 >3 standard drinks per day for men. ‡Poverty was defined as a standard of living lower than
13 the 2015 French poverty threshold (1,015€ per month per household consumption unit).
14 §Being a migrant was defined as being born outside France and having at least one parent
15 also born outside France. (A)OR, (adjusted) odds ratio; CI, confidence interval.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supporting table 2: Factors associated with the presence of diabetes in patients with advanced fibrosis, logistic regression models, univariable and multivariable analyses (ANRS CO22 Hepather cohort, n=2,511)

	Univariable analyses (n=2,511)			Multivariable analysis (n=2,410)		
	OR	95% CI	P-value	AOR	95% CI	P-value
Gender						
Female	1		<.001	1		<.001
Male	1.39	1.15-1.69		1.87	1.50-2.33	
Cannabis use						
No lifetime use	1		<.001	1		<.001
Former	0.62	0.47-0.81		0.66	0.49-0.89	
Current	0.330	0.23-0.50		0.39	0.25-0.60	
Tobacco use						
No lifetime use or former use	1		<.001			
Current	0.72	0.60-0.87				
Coffee consumption						
Low (<3 cups/day)	1		0.214			

Elevated (≥ 3 cups/day)	0.85	0.66-1.20				
Body mass index						
Underweight (< 18.5)	0.88	0.45-1.69	$< .001$	1.14	0.56-2.31	$< .001$
Normal (≥ 18.5 and < 25)	1			1		
Overweight (≥ 25 and < 30)	1.50	1.21-1.85		1.31	1.04-1.64	
Obese (> 30)	2.57	1.99-3.31		2.43	1.85-3.19	
Alcohol consumption†						
Abstinent	1		0.101			
Moderate	0.89	0.73-1.10				
Elevated	0.64	0.41-1.00				
Poverty‡						
No	1		0.008	1		0.011
Yes	1.30	1.07-1.58		1.34	1.07-1.67	
Being a migrant§						
No	1		$< .001$	1		0.001

1						
2						
3					1.16-	
4	Yes	1.89	1.55-2.29		1.45	
5						1.81
6						
7						
8						1.04-
9	Age (years)	1.02	1.01-1.03	<.001	1.05	0.001
10						1.06
11						
12						
13						

† Elevated alcohol consumption was defined as >2 standard drinks per day for women and
 >3 standard drinks per day for men. ‡Poverty was defined as a standard of living lower than
 the 2015 French poverty threshold (1,015€ per month per household consumption unit).
 §Being a migrant was defined as being born outside France and having at least one parent
 also born outside France. (A)OR, (adjusted) odds ratio; CI, confidence interval.

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60