



HAL
open science

Exact observability properties of subelliptic wave and Schrödinger equations

Cyril Letrouit

► **To cite this version:**

Cyril Letrouit. Exact observability properties of subelliptic wave and Schrödinger equations. 2021. hal-03109787v1

HAL Id: hal-03109787

<https://hal.science/hal-03109787v1>

Preprint submitted on 14 Jan 2021 (v1), last revised 11 Oct 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXACT OBSERVABILITY PROPERTIES OF SUBELLIPTIC WAVE AND SCHRÖDINGER EQUATIONS

CYRIL LETROUIT

ABSTRACT. In this survey paper, we report on recent works concerning exact observability (and, by duality, exact controllability) properties of subelliptic wave and Schrödinger-type equations. These results illustrate the slowdown of propagation in directions transverse to the horizontal distribution. The proofs combine sub-Riemannian geometry, semi-classical analysis, spectral theory and non-commutative harmonic analysis.

CONTENTS

1. Introduction	1
2. Subelliptic wave equations are never observable	17
3. Subellipticity and strength of propagation: Baouendi-Grushin Schrödinger equations	25
4. Subelliptic Schrödinger equation via non-commutative harmonic analysis	31
5. Perspectives and open problems	39
References	41

1. INTRODUCTION

1.1. Controllability and observability. The problem of (exact) controllability of PDEs, which has been intensively studied in the past decades, is the following: given a manifold M , a subset $\omega \subset M$, a time $T > 0$ and an operator A acting on functions on M , the study of exact controllability consists in determining whether, for any initial state u_0 and any final state u_1 , there exists f such that the solution of

$$(1.1) \quad \partial_t u = Au + \mathbf{1}_\omega f, \quad u|_{t=0} = u_0$$

in M is equal to u_1 at time T . Here, $\mathbf{1}_\omega$ is the characteristic function of ω . In other words, exact controllability holds if it is possible, starting from any initial state, to reach any final state just acting on ω during a time T . The general answer depends on the time T , the control set ω , the operator A , and the functional spaces in which u_0 , u_1 and f live. This problem is relevant in many physical situations: typical examples are the control of the temperature of a room by a heater, or the acoustic insulation of a room just by acting on a small part of it.

By duality (Hilbert Uniqueness Method, see [Lio88]), the exact controllability property is equivalent to some inequality of the form

$$(1.2) \quad \exists C_{T,\omega} > 0, \quad \forall u_0, \quad \|u_0\|^2 \leq C_{T,\omega} \int_0^T \|\mathbf{1}_\omega u(t)\|^2 dt,$$

where u is the solution of the adjoint equation $(\partial_t + A^*)u = 0$ with initial datum u_0 (here again, one should specify functional spaces). This is called an observability inequality. In other words, controllability holds if and only if any solution of $(\partial_t + A^*)u = 0$ can be detected from ω , in a “quantitative way” which is measured by the constant $C_{T,\omega}$. This paper is devoted to the study of observability for equations of wave-type, Schrödinger-type or heat-type, i.e. we consider the equations

$$(1.3) \quad (\partial_{tt}^2 - \mathcal{L})u = 0 \quad (\text{Wave-type}),$$

$$(1.4) \quad (i\partial_t - \mathcal{L})u = 0 \quad (\text{Schrödinger-type}),$$

$$(1.5) \quad (\partial_t - \mathcal{L})u = 0 \quad (\text{Heat-type})$$

for various time-independent operators \mathcal{L} on M .¹ By duality, all the observability results presented here imply exact controllability results as explained above, but we won't state them for the sake of simplicity.

1.2. Observability of classical PDEs. Let us present a first series of results, dating back to the 1990's, which concern the observability problem in case M is a compact Riemannian manifold with a metric g and with boundary $\partial M \neq \emptyset$, $\mathcal{L} = \Delta_g$ is the Laplace-Beltrami operator on (M, g) and the equation is one of the three equations (1.3), (1.4) or (1.5), with Dirichlet boundary conditions $u|_{\partial M} = 0$. We deal with these three problems in this order, following the chronology of the results.

Throughout this section, (M, g) is a fixed manifold with boundary $\partial M \neq \emptyset$ and $\mathcal{L} = \Delta_g$. In this section, the notation dx stands for the associated Riemannian volume $dx = d\text{vol}_g(x)$.

1.2.1. Observability of the Riemannian wave equation. Let us start with the wave equation (1.3) with initial data $(u_{t=0}, \partial_t u|_{t=0}) = (u_0, u_1) \in H^1(M) \times L^2(M)$ and Dirichlet boundary conditions. The energy of a solution, which is conserved along the flow, is

$$E(u(t)) = \int_M (|\nabla_g u(t, x)|^2 + |\partial_t u(t, x)|^2) dx$$

which is in particular equal to the initial energy $\|\nabla u_0\|_{L^2(M)}^2 + \|u_1\|_{L^2(M)}^2$. Let $T > 0$ and ω be a measurable subset. The observability inequality reads as follows:

$$(1.6) \quad E(u(0)) \leq C \int_0^T \int_\omega |\partial_t u(t, x)|^2 dx dt.$$

¹The wave equation involves a ∂_{tt}^2 term, and thus does not enter, strictly speaking, the framework given by equation 1.1. However, it is possible to give a common framework for all three equations, at the cost of being a bit more abstract. See [Cor07, Section 2.3] for a general introduction.

Note that thanks to the conservation of energy, the left hand side of (1.2) has been replaced by the energy of the initial datum.

We set $P = \partial_{tt}^2 - \Delta_g$ (which is a second-order pseudo-differential operator), whose principal symbol is

$$p_2(t, \tau, x, \xi) = -\tau^2 + g^*(x, \xi)$$

with τ the dual variable of t and g^* the principal symbol of $-\Delta_g$. In $T^*(\mathbb{R} \times M)$, the Hamiltonian vector field \vec{H}_{p_2} associated with p_2 is given by $\vec{H}_{p_2}f = \{p_2, f\}$. Since $\vec{H}_{p_2}p_2 = 0$, we get that p_2 is constant along the integral curves of \vec{H}_{p_2} . Thus, the characteristic set $\mathcal{C}(p_2) = \{p_2 = 0\}$ is preserved under the flow of \vec{H}_{p_2} . Null-bicharacteristics are then defined as the maximal integral curves of \vec{H}_{p_2} which live in $\mathcal{C}(p_2)$. In other words, the null-bicharacteristics are the maximal solutions of

$$\begin{cases} \dot{t}(s) = -2\tau(s), \\ \dot{x}(s) = \nabla_{\xi} g^*(x(s), \xi(s)), \\ \dot{\tau}(s) = 0, \\ \dot{\xi}(s) = -\nabla_x g^*(x(s), \xi(s)), \\ \tau^2(0) = g^*(x(0), \xi(0)). \end{cases}$$

It is well-known that the projection $x(s)$ of a bicharacteristic ray $(x(s), \xi(s))$ traveled at speed 1 is a geodesic in M , i.e., a curve which realizes the minimal distance between any two of its points which are close enough.

Let us also mention the fact that at the boundary of M , the above definition of null-bicharacteristics has to be completed (yielding the so-called Melrose-Sjöstrand flow): due to trajectories which “graze” along the boundary, one cannot always define the null-bicharacteristics which touch the boundary by reflexion, and we refer the reader to [MS78] and [LRLTT17] for more on this subject. In these papers, a notion of “generalized bicharacteristics” is defined, which explains how to define bicharacteristics at the boundary. For us, this will only be useful to give a precise statement for Theorem 1.

Definition 1.1. *Let $T > 0$ and $\omega \subset M$ be a measurable subset. We say that the Geometric Control Condition holds in time T in ω , and we write $(GCC)_{\omega, T}$, if for any projection γ of a bicharacteristic ray traveled at speed 1, there exists $t \in (0, T)$ such that $\gamma(t) \in \omega$.*

The following result states that the observability of (1.3) is (more or less) equivalent to the geometric condition $(GCC)_{\omega, T}$. It illustrates the finite speed of propagation for waves.

Theorem 1 ([BLR92], [BG97], [HPT19]). *Assume that $\omega \neq \emptyset$ is open and that $(GCC)_{\omega, T}$ holds. Assume also that no generalized bicharacteristic has a contact of infinite order with $(0, T) \times \partial M$. Then (1.6) holds, i.e., the wave equation (1.3) is observable in time T on ω . Conversely, if the wave equation (1.3) is observable in time T , then $(GCC)_{T, \bar{\omega}}$ holds, where $\bar{\omega}$ denotes the closure of ω .*

Note that the second statement in the last theorem is not the exact converse of the first one, since it involves the closure $\bar{\omega}$ and not simply ω . This is due to the phenomenon of grazing rays: if there exists a ray γ which does not enter ω but which touches the boundary $\partial\omega$, so that the geometric control condition is not satisfied, it can however happen (notably if the flow is “stable” close from the ray) that observability holds, see [Leb92a, Section VI.B] for an example.

Considering solutions of (1.3) of the form $e^{it\sqrt{\lambda}}\varphi$ where φ is an eigenfunction of $-\Delta_g$ corresponding to the eigenvalue λ , the following result follows from Theorem 1:

Corollary 1.2. *Assume that $\omega \neq \emptyset$ is open and that there exists $T > 0$ such that $(GCC)_{\omega,T}$ holds. Then, for any eigenfunction φ of $-\Delta_g$, there holds*

$$\int_{\omega} |\varphi(x)|^2 dx \geq C \int_M |\varphi(x)|^2 dx.$$

In particular, $\text{supp}(\varphi) = M$.

All the observability inequalities stated in this survey paper yield similar lower bounds, but we will not state them thereafter.

Remark 1.3 (Gaussian beams). *The fact that $(GCC)_{\bar{\omega},T}$ is a necessary condition for observability can be understood as follows. If $(GCC)_{\bar{\omega},T}$ does not hold, then let $\gamma : [0, T] \rightarrow M$ be a geodesic which does not enter $\bar{\omega}$. By compactness, there exists $\varepsilon > 0$ such that $\gamma|_{[0, T]}$ does not meet an ε -neighborhood of $\bar{\omega}$. Then, one can construct a sequence of solutions $(u_n)_{n \in \mathbb{N}}$ of the wave equation whose initial energy $E(u_n(0))$ is normalized to 1, and with energy $E(u(t))$ localized around $\gamma(t)$ at any time $t \in [0, T]$: quantitatively, the energy of u_n outside a tubular neighborhood of γ of size ε tends to 0 as $n \rightarrow +\infty$. This disproves the observability inequality (1.6). The sequence $(u_n)_{n \in \mathbb{N}}$, if taken as a Gaussian profile centered at a point describing γ , is called a Gaussian beam.*

1.2.2. *Observability of the Riemannian Schrödinger equation.* In case of the Schrödinger equation (1.4), the observability inequality reads as follows:

$$(1.7) \quad \|u_0\|_{L^2(M)}^2 \leq C \int_0^T \int_{\omega} |u(t, x)|^2 dx dt.$$

Indeed, as for the wave equation (1.3), the L^2 -norm of the solution is preserved along the flow, so that $\|u(T)\|_{L^2} = \|u_0\|_{L^2}$. A sufficient condition for observability is the following:

Theorem 2 ([Leb92b] and Appendix of [DGL06]). *Assume that $\omega \neq \emptyset$ is open and that $(GCC)_{\omega, T'}$ holds for some $T' > 0$. Then (1.7) holds, i.e., the Schrödinger equation (1.4) is observable in any time $T > 0$ on ω .*

The interplay between T' and T in the above result is due to the fact that the Schrödinger equation “propagates at infinite speed” so that no matter how large T' is, observability holds

in any time $T > 0$ if $(GCC)_{\omega, T'}$ holds. This contrasts with the finite speed of propagation of the wave equation.

The converse of the above theorem, namely to find necessary conditions on (ω, T) for (1.7) to hold, is notoriously a difficult problem. The main results in this direction are for the torus (see [Jaf90], [BZ12], [AM14]), and in Riemannian manifolds with negative curvature (see [DJN19]), where (1.7) holds for any non-empty open subset ω and any time $T > 0$. Indeed, it is expected that if the geodesic flow of the background geometry is unstable, solutions of (1.4) are more “delocalized” than those of (1.3) for example. See also the case of the disk [ALM16].

1.2.3. Observability of the Riemannian heat equation. Let us end with the heat equation. The observability inequality reads as follows:

$$(1.8) \quad \|u(T)\|_{L^2(M)}^2 \leq C \int_0^T \int_{\omega} |u(t, x)|^2 dx dt.$$

Theorem 3 ([LR95]). *Let $\omega \neq \emptyset$ be open and $T > 0$. Then (1.8) holds, i.e., the heat equation (1.5) is observable in time T on ω .*

Note that no geometric condition on ω is required in this case. This result illustrates the infinite speed of propagation of the heat equation.

The works presented hereafter address that same problem of observability of linear PDEs, but with focus on subelliptic PDEs, meaning that the Laplace-Beltrami operator is replaced in these PDEs by a sub-Laplacian, which is a subelliptic operator. The next subsection is thus devoted to introduce the main objects of study, namely sub-Laplacians.

1.3. Sub-Riemannian geometry and sub-Laplacians. After the founding work of Lars Hörmander, and with the development of sub-Riemannian geometry since the 1980’s, subelliptic operators have been considered as a natural generalization of elliptic operators. In particular, sub-Laplacians, which we will define soon, are natural generalizations of the Laplace-Beltrami operator. Therefore, the question of observability/controllability of evolution PDEs driven by sub-Laplacians has been investigated since a decade, with a particular focus on parabolic (or heat-type) equations. In this survey, we mainly focus on wave-type and Schrödinger-type subelliptic equations, for which the first results appeared in 2019.

1.3.1. Sub-Laplacians. Instead of defining subelliptic operators in full generality, we prefer here to focus only on sub-Laplacians. Simple examples of sub-Laplacians exist, but we make a small detour through sub-Riemannian geometry in order to define them with generality. Readers only interested in the results of Sections 3 and 4 could skip this part, and focus on Examples 1.7 to 1.10 which are sufficient for these sections. The books [Mon02] and [ABB19] are clear and detailed introductions to sub-Riemannian geometry.

Definition 1.4. *A sub-Riemannian structure is a triple (M, \mathcal{D}, g) where*

- M is a smooth connected compact manifold of dimension $n \in \mathbb{N}^*$ with or without boundary;
- \mathcal{D} is a smooth horizontal distribution on M , i.e., a smooth assignment $M \ni x \mapsto \mathcal{D}_x \subset T_x M$ (possibly with non-constant rank) which satisfies the Hörmander condition

$$(1.9) \quad \text{Lie}(\mathcal{D}) = TM,$$

i.e., any element of TM can be obtained as the iterated bracket of elements of \mathcal{D} .

- g is a Riemannian metric on \mathcal{D} .

In case $\mathcal{D} = TM$, we recover traditional Riemannian manifolds. In the general case where $\mathcal{D} \subset TM$, the set $TM \setminus \mathcal{D}$ can be understood as the directions where the metric g takes the value $+\infty$. A well-known theorem, due to Chow and Rashevskii, asserts that any two points can be joined by a path, i.e., a continuous function $\gamma : [0, 1] \rightarrow M$ with derivative $\dot{\gamma}(t)$ contained in $\mathcal{D}_{\gamma(t)}$ for almost any $t \in [0, 1]$. In other words, the sub-Riemannian distance

$$d_g(x_0, x_1) = \inf \left\{ \int_0^1 \sqrt{g_{\gamma(t)}(\dot{\gamma}(t), \dot{\gamma}(t))} dt \mid \gamma(0) = x_0, \gamma(1) = x_1, \dot{\gamma}(t) \in \mathcal{D}_{\gamma(t)} \text{ a.s. for } t \in [0, 1] \right\}$$

is finite for any $x_0, x_1 \in M$.

When moving in a sub-Riemannian structure, \mathcal{D} should be understood as the “set of allowed directions for the motion”, and, although it is not possible to move directly in directions of $TM \setminus \mathcal{D}$, Chow-Rashevskii’s theorem asserts that any two points can be joined by a path. This is due to “indirect motions”, that is, paths which describe spirals turning around a fixed forbidden direction of $TM \setminus \mathcal{D}$ and thus advancing in this direction (although indirectly).

Definition 1.5. *The step k of a sub-Riemannian structure (M, \mathcal{D}, g) is the least integer $k \in \mathbb{N}$ such that $\mathcal{D}^k = TM$, where, for $j \in \mathbb{N}^*$, \mathcal{D}^j is defined through the recursive relation $\mathcal{D}^j = \mathcal{D}^{j-1} + [\mathcal{D}, \mathcal{D}^{j-1}]$ and $\mathcal{D}^1 = \mathcal{D}$.*

Remark 1.6. *More generally, the step k_x can be defined at any point $x \in M$, just by considering \mathcal{D}_x^j instead of \mathcal{D}^j in the above definition.*

To a sub-Riemannian structure (M, \mathcal{D}, g) and an arbitrary smooth volume μ on M , we can associate a sub-Laplacian $\Delta_{g,\mu}$ as follows:

$$(1.10) \quad \Delta_{g,\mu} = - \sum_{i=1}^m X_i^* X_i = \sum_{i=1}^m X_i^2 + \text{div}_\mu(X_i) X_i$$

where $(X_i)_{1 \leq i \leq m}$ denotes a local g -orthonormal frame such that $\mathcal{D} = \text{Span}(X_1, \dots, X_m)$ and the star designates the transpose in $L^2(M, \mu)$. The divergence with respect to μ is defined by $L_X \mu = (\text{div}_\mu X) \mu$, where L_X stands for the Lie derivative. Up to a lower order term, a sub-Laplacian is thus a “sum of squares”.

1.3.2. *Examples.* We now give a few examples of sub-Laplacians which we shall study in the sequel. In these examples, we sometimes invert the point of view, giving first the sub-Laplacian, from which the distribution \mathcal{D} and the metric g can be easily deduced.

Example 1.7. On $M = (-1, 1)_x \times \mathbb{T}_y$ where $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$, we set $\Delta_G = \partial_x^2 + x^2\partial_y^2$. This sub-Laplacian is the so-called Baouendi-Grushin operator, sometimes improperly called simply Grushin operator (see [Gar17, Section 11]). In this case, $\mathcal{D} = \text{Span}(\partial_x, x\partial_y)$ and $\mathcal{D}^2 = \text{Span}(\partial_x, \partial_y) = TM$. In particular, $\mathcal{D} = TM$ outside the line $\{x = 0\}$. Also, μ is the Lebesgue measure and g is defined by saying that $(\partial_x, x\partial_y)$ is an orthonormal frame for g . The structure (M, \mathcal{D}, g) has step 2 on the line $\{x = 0\}$ and step 1 outside this line. Since this sub-Riemannian structure is ‘‘Riemannian’’ outside this line, the Baouendi-Grushin operator is sometimes called ‘‘almost-Riemannian’’.

Example 1.8. More generally, for $\gamma \geq 0$ (not necessarily an integer), one can consider $\Delta_\gamma = \partial_x^2 + |x|^{2\gamma}\partial_y^2$ on the same manifold $M = (-1, 1)_x \times \mathbb{T}_y$. For $\gamma \in \mathbb{N}$, the associated sub-Riemannian structure has step $k = \gamma + 1$. Note that for $\gamma \notin \mathbb{N}$, the relation (1.9) is not necessarily satisfied, but we include this class of examples in our study since our computations allow to handle them.

Example 1.9. Given $d \in \mathbb{N}^*$, one can also define a sub-Laplacian arising from the Heisenberg group \mathbf{H}^d of dimension $2d + 1$. Recall that the Heisenberg group \mathbf{H}^d is \mathbb{R}^{2d+1} endowed with the group law $(x, y, z) \cdot (x', y', z') := (x + x', y + y', z + z' + \frac{1}{2} \sum_{j=1}^d (x_j y'_j - x'_j y_j))$, where $x, y, x', y' \in \mathbb{R}^d$ and $z, z' \in \mathbb{R}$. Taking the left-quotient of \mathbf{H}^d by the discrete subgroup $\tilde{\Gamma} = (\sqrt{2\pi}\mathbb{Z})^{2d} \times \pi\mathbb{Z}$, we obtain a compact manifold $M = \tilde{\Gamma} \backslash \mathbf{H}^d$. Let

$$X_j = \partial_{x_j} - \frac{y_j}{2}\partial_z, \quad Y_j = \partial_{y_j} + \frac{x_j}{2}\partial_z, \quad \text{for } j = 1, \dots, m,$$

which are left-invariant and can be thus considered as vector fields on the quotient manifold M . Finally, we define the sub-Laplacian

$$\Delta_{\tilde{\Gamma} \backslash \mathbf{H}^d} = \sum_{j=1}^d X_j^2 + Y_j^2.$$

Since $[X_j, Y_j] = \partial_z$ for any j , this sub-Laplacian is naturally associated with a (compact) step 2 sub-Riemannian structure.

Example 1.10. Heisenberg-type groups are generalizations of Heisenberg groups. They were first introduced in [Kap80], where the fundamental solution of the associated sub-Laplacians, which is particularly simple, was computed. These groups give rise to step 2 sub-Riemannian structures, which center can be of dimension $p > 1$ (whereas the center of the Heisenberg group \mathbf{H}^d is of dimension 1). For a precise definition, see [Kap80].

Example 1.11. Contact sub-Laplacians arise from a particular type of sub-Riemannian structures of step 2. Let (M, \mathcal{D}, g) be a (metric) contact structure, i.e., a sub-Riemannian structure of odd dimension $n = 2m + 1$ such that there exists a 1-form α on M , with $\mathcal{D} = \text{Ker}(\alpha)$

and $\alpha \wedge (d\alpha)^m \neq 0$ at any point of M . Then, for any smooth volume μ , the associated sub-Laplacian $\Delta_{g,\mu}$ is called a contact sub-Laplacian. A typical example is given by the Heisenberg sub-Laplacian $\Delta_{\mathbf{H}^d}$ defined above. The study of eigenfunctions of general 3D contact sub-Laplacians has been undertaken in [CdVHT18].

Example 1.12. Magnetic Laplacians are also sub-Laplacians, and we focus here on a simple family of examples. In \mathbb{R}^3 with coordinates x, y, z , we consider the two vector fields $X_1 = \partial_x - A_x(x, y)\partial_z$ and $X_2 = \partial_y - A_y(x, y)\partial_z$ where A_x, A_y are functions which do not depend on z . The magnetic Laplacian is then $\Delta = X_1^2 + X_2^2$. The 1-form $A = A_x dx + A_y dy$ is called the connection form, and the 2-form $B = dA$ is called the magnetic field. The modulus $|b|$ of the function b defined by the relation $B = b dx \wedge dy$ is called the intensity of the magnetic field. Taking a quotient or assuming that $|b|$ is bounded away from 0 at infinity, it is possible to assume that the sub-Laplacian Δ has a compact resolvent.

Magnetic Laplacians were used by Montgomery to prove the existence of abnormal minimizers in some sub-Riemannian geometries (see [Mon94]): abnormal minimizers are local minimizers of the sub-Riemannian distance which are not projections of bicharacteristics, and they show up for example as zero curves of the intensity b . Subsequently, Montgomery showed that the spectral asymptotics of Δ are very different depending on the fact that b vanishes or not (see [Mon95]).

Remark 1.13. *In all the previous examples, as well as in the general definition of sub-Laplacians given above, the volume μ is assumed to be smooth. However, one can also define sub-Laplacians for non-smooth volumes μ . This is natural when the sub-Riemannian distribution is singular, for example $\mathcal{D} = \text{Span}(\partial_x, x\partial_y)$ in \mathbb{R}^2 (see Example 1.7), since the Popp measure, which is an intrinsic measure defined on sub-Riemannian manifolds (see [Mon02, Section 10.6]), “blows up”. The associated sub-Laplacian is then unitarily equivalent to a sub-Laplacian with smooth volume plus a singular potential. The essential self-adjointness of some of these sub-Laplacians has been studied for example in [BL13]. When they are essentially self-adjoint, the controllability/observability of the associated evolution equations can also be studied: this is an open question which we do not address here.*

1.3.3. *Hypoellipticity and subellipticity.* Two notions are often used to qualify the smoothing properties of sub-Laplacians: hypoellipticity and subellipticity. Here, we briefly recall their definitions and explain why they are not exactly equivalent.

Definition 1.14. *A (pseudo-)differential operator A with C^∞ coefficients in M is hypoelliptic in M if for all $u \in \mathcal{D}'(M)$ and $x \in M$, if $Au \in C^\infty$ near x , then $u \in C^\infty$ near x .*

Hypoellipticity appeared naturally in the work of Kolmogorov [Kol34] on the motion of colliding particles when he wrote down the equation

$$\partial_t u - \mathcal{L}u = f \quad \text{where} \quad \mathcal{L} = x\partial_y + \partial_x^2.$$

Indeed, the operator \mathcal{L} is hypoelliptic.

Definition 1.15. A formally selfadjoint (pseudo-)differential operator $A : C^\infty(M) \rightarrow C^\infty(M)$ of order 2 is said to be subelliptic if there exist $s, C > 0$ such that

$$(1.11) \quad \|u\|_{H^s(M)}^2 \leq C((Au, u)_{L^2(M)} + \|u\|_{L^2(M)}^2)$$

for any $u \in C^\infty(M)$.

Using (1.9), Hörmander was able to prove that any sub-Laplacian $\Delta_{g,\mu}$ is hypoelliptic (see [Hör67] and [HN05, Chapter 2]). His proof relies on the fact that $\Delta_{g,\mu}$ is subelliptic; indeed, the optimal s in (1.11) is $1/k$, where k is the step of the associated sub-Riemannian structure, as proved by Rothschild and Stein [RS76, Theorem 17 and estimate (17.20)].

Conversely, note that an hypoelliptic “sum of squares” (i.e., an operator of the form (1.10) which is hypoelliptic) does not necessarily satisfy the Lie bracket assumption (1.9): given a smooth function $a : \mathbb{R} \rightarrow \mathbb{R}$ vanishing at infinite order at 0 but with $a(s) > 0$ for $s \neq 0$, the sub-Laplacian $\Delta = \partial_{x_1}^2 + a(x_1)^2 \partial_{x_2}^2$ on $\mathbb{R}_{x_1 x_2}^2$ is hypoelliptic although (1.9) fails (see [Fed71] and [Mor78]).

Let us finally mention that some operators A satisfy the property that if Au is real-analytic, then u is real-analytic: they are called *analytic hypoelliptic*. The so-called Trèves conjecture describes a possible link between analytic hypoellipticity of an operator and the absence of abnormal geodesics (see [Tre99] for the conjecture and [ABM18] for more recent results).

We end this section with a remark concerning the compactness of the manifolds M considered in this survey.

Remark 1.16. Because of the physical nature of the problems studied in control/observability theory, most equations are set in compact manifolds, and this survey is no exception to the rule. Even in Example 1.9, the sub-Laplacian is defined on a compact quotient of the Heisenberg group. Together with the hypoellipticity, the compactness of the underlying manifold implies that all sub-Laplacians have a compact resolvent, and thus a discrete spectrum, which is of importance for deriving properties of eigenfunctions from observability results.

1.4. Observability of subelliptic PDEs: known results. This section is devoted to stating results which were previously known in the literature about controllability/observability of subelliptic PDEs. All PDEs we consider are well-posed in natural energy spaces which we do not systematically recall.

1.4.1. Subelliptic heat equations. Let us start with the result proved in [BCG14], which concerns the heat equation (1.5) where $\mathcal{L} = \Delta_\gamma$ is the Baouendi-Grushin-type operator introduced in Example 1.8 for some $\gamma > 0$. The open subset of observation $\omega \subset (-1, 1) \times \mathbb{T}$ they consider is a vertical strip of the form $(a, b) \times \mathbb{T}$ where $0 < a < b < 1$. The observability inequality is (1.8), with the modification that u runs over the set of solutions of (1.5) with $\mathcal{L} = \Delta_\gamma$. The authors prove the following result, to be compared with Theorem 3:

Theorem 4 ([BCG14]). *Let $\gamma > 0$ and ω be as above. Then*

- *If $\gamma \in (0, 1)$, then for any $T > 0$, (1.8) holds;*
- *If $\gamma = 1$, i.e., $\Delta_\gamma = \Delta_G$, then there exists $T_0 > 0$ such that (1.8) holds if $T > T_0$ and does not hold if $T < T_0$;*
- *If $\gamma > 1$, then, for any $T > 0$, (1.8) fails.*

The proof is done by establishing an infinite number of Carleman inequalities for operators $-\partial_x^2 + n^2 x^2$ for $n \in \mathbb{Z}$, with bounds uniform in n . It was proved in [BMM15] that in case $\gamma = 1$, the minimal time T_0 is equal to $a^2/2$. The fact that $T_0 \geq a^2/2$ can be seen by using explicit eigenfunctions of Δ_γ .

Koenig studied the observability of (1.5) with $\mathcal{L} = \Delta_G$, but for another geometry of the observation set ω : this time, it is a horizontal band of the form $(-1, 1) \times I$ where I is a proper open subset of \mathbb{T} .

Theorem 5 ([Koe17]). *Let $\omega = (-1, 1) \times I$ where I is a proper open subset of \mathbb{T} . Then (1.8) fails for any $T > 0$.*

The proof of this result relies on the non-observability of the 1D half-heat equation $\partial_t u + |\partial_x|u = 0$ and on techniques coming from complex analysis where the complex variable is $z = e^{-t+iy}$.

Although the observability properties of the heat equation driven by general hypoelliptic operators are still mysterious, we list here a few works addressing this question. The recent works [Lis20], [BDE20] and [DK20] continue and generalize the analysis of [BCG14] and [Koe17] on the control of the Baouendi-Grushin heat equation. Besides, [BC17] establishes the existence of a minimal time of observability, as in the second point of Theorem 4, for the heat equation driven by the Heisenberg sub-Laplacian of Example 1.9. Let us finally mention the papers [DR20] and [BPS18] which also deal with controllability issues for hypoelliptic parabolic equations.

The above theorems show that some subelliptic heat equations driven by simple sub-Laplacians require a larger time to be observable than the usual Riemannian heat equation, and observability may even fail in any time $T > 0$. As we will see, this is a very general phenomenon for subelliptic evolution PDEs, at least for subelliptic wave equations and (some) Schrödinger-type equations. Our results, however, do not shed any new light on subelliptic heat equations, which remain mysterious due to the lack of “general arguments” which would not rely on geometric and analytic features specific to very particular sub-Laplacians.

1.4.2. Approximate observability of subelliptic PDEs. Recently, Laurent and Léautaud have studied the observability of subelliptic PDEs but with focus on a different notion of observability, called approximate observability. As before, all their results can also be stated in terms of a dual notion, called approximate controllability, however we will not even mention these dual

statements in order to keep the presentation as simple as possible. Their results are quantitative, in the sense that they give explicit bounds on the observability constants involved in their results. The next paragraphs are devoted to a brief description of their results (see [LL20]).

Let us consider a sub-Riemannian structure (M, \mathcal{D}, g) and an associated sub-Laplacian $\Delta_{g,\mu}$ as in (1.10). We assume that the manifold M (assumed to have no boundary, $\partial M = \emptyset$), the smooth volume μ and the vector fields X_i are all *real-analytic*. For $s \in \mathbb{R}$, the operator $(1 - \Delta_{g,\mu})^{\frac{s}{2}}$ is defined thanks to functional calculus, and we consider the (adapted) Sobolev spaces

$$\mathcal{H}^\ell(M) = \{u \in \mathcal{D}'(M), (1 - \Delta_{g,\mu})^{\frac{\ell}{2}}u \in L^2(M)\}$$

with the associated norm $\|u\|_{\mathcal{H}^\ell(M)} = \|(1 - \Delta_{g,\mu})^{\frac{\ell}{2}}u\|_{L^2(M)}$.

Theorem 6 ([LL20]). *Let ω be a non-empty open subset of M and let $T > \sup_{x \in M} d_g(x, \omega)$. We denote by k the step of the sub-Riemannian structure (M, \mathcal{D}, g) . Then there exist $c, C > 0$ such that*

$$(1.12) \quad \|(u_0, u_1)\|_{\mathcal{H}^1 \times L^2} \leq C e^{c\Lambda^k} \|u\|_{L^2((-T, T) \times \omega)}, \quad \text{with } \Lambda = \frac{\|(u_0, u_1)\|_{\mathcal{H}^1 \times L^2}}{\|(u_0, u_1)\|_{L^2 \times \mathcal{H}^{-1}}}$$

for any solution u of (1.3) on $(-T, T)$ such that $(u, \partial_t u)|_{t=0} = (u_0, u_1) \in \mathcal{H}^1(M) \times L^2(M)$.

The above result in particular implies unique continuation (and quantifies it): if $u = 0$ in $(-T, T) \times \omega$, then $u \equiv 0$. However, the exact observability inequality which we shall study (see (1.6)) is a stronger requirement than (1.12), in particular because of the presence of the ‘‘typical frequency of the datum’’ Λ in the right-hand side of (1.12). The techniques used for proving Theorem 6 are totally different from those we present in the sequel.

1.4.3. Observability of Baouendi-Grushin Schrödinger equation. The recent work [BS19] is the first one dealing with exact observability of a subelliptic Schrödinger equation, namely in the context of Example 1.7 with observation set given by a horizontal band as in Theorem 5. The observability inequality is given by (1.7), except that u runs over the solutions of the Schrödinger equation driven by the sub-Laplacian Δ_G .

Theorem 7 ([BS19]). *Let $M = (-1, 1) \times \mathbb{T}$ and $\Delta_G = \partial_x^2 + x^2 \partial_y^2$. Let $\omega = (-1, 1) \times I$ where $I \subsetneq \mathbb{T}$ is open. Let $T_0 = \mathcal{L}(\omega)$ be the length of the maximal sub-interval contained in $\mathbb{T} \setminus I$. Then, the observability property (1.7) holds if and only if $T > T_0$.*

Again, this result shows the existence of a minimal time of control which contrasts with the ‘‘infinite speed of propagation’’ illustrated by Theorem 2. Its proof relies on fine semi-classical analysis, somehow linked to that explained in Section 3.

1.4.4. Non-linear subelliptic PDEs. Although this survey is devoted only to *linear* subelliptic PDEs, let us say a word about *non-linear* subelliptic PDEs. To study the cubic Grushin-Schrödinger equation $i\partial_t u - (\partial_x^2 + x^2 \partial_y^2)u = |u|^2 u$, Patrick Gérard and Sandrine Grellier introduced a toy model, the cubic Szegő equation, which models the interactions between the

nonlinearity and the lack of dispersivity of the linear equation (already visible in the above Theorem 7). In [GG10], they put this equation into a Hamiltonian framework and classify the traveling waves for this equation, which are related to the traveling waves resulting from (3.3).

1.5. Main results. Let us now present the main results contained in the papers [Let20b], [LS20] and [FKL20]. All of them illustrate the slowdown of propagation of evolution PDEs in directions transverse to the distribution: in a nutshell, observability will require a much longer time to hold for subelliptic PDEs than for elliptic ones, and this time will be even larger when the step k of the underlying sub-Riemannian structure is larger. All our results are summarized in Figure 1 at the end of this section.

1.5.1. First main result. We start with a general result on subelliptic wave equations. Let (M, \mathcal{D}, g) be a sub-Riemannian structure such that $\partial M \neq \emptyset$ and μ be a smooth volume on M . We consider $\Delta = -\sum_{i=1}^m X_i^* X_i$ the associated sub-Laplacian, where $(X_i)_{1 \leq i \leq m}$ is a local g -orthonormal frame such that $\mathcal{D} = \text{Span}(X_1, \dots, X_m)$ (locally). The sub-Riemannian gradient is defined by the formula

$$\nabla^{\text{sR}} \phi = \sum_{i=1}^m (X_i \phi) X_i.$$

Consider the wave equation

$$(1.13) \quad \begin{cases} \partial_{tt}^2 u - \Delta u = 0 & \text{in } (0, T) \times M \\ u = 0 & \text{on } (0, T) \times \partial M, \\ (u|_{t=0}, \partial_t u|_{t=0}) = (u_0, u_1) \end{cases}$$

where $T > 0$, and the initial data (u_0, u_1) are in an appropriate energy space. The natural energy of a solution u of the sub-Riemannian wave equation (1.13) is

$$E(u(t, \cdot)) = \frac{1}{2} \int_M (|\partial_t u(t, x)|^2 + |\nabla^{\text{sR}} u(t, x)|^2) d\mu(x).$$

Observability holds in time T_0 on ω if there exists $C > 0$ such that for any solution u of (1.13),

$$(1.14) \quad E(u(0)) \leq C \int_0^{T_0} \int_{\omega} |\partial_t u(t, x)|^2 d\mu(x) dt.$$

Theorem 8 ([Let20b]). *We assume that the set of all $x \in M$ such that $\mathcal{D}_x \neq T_x M$ is dense in M . Let $T_0 > 0$ and let $\omega \subset M$ be a measurable subset such that $M \setminus \omega$ has nonempty interior. Then the subelliptic wave equation (1.13) is not exactly observable on ω in time T_0 .*

Theorem 8 can be reformulated as follows: subelliptic wave equations are never observable. The condition that the set of all $x \in M$ such that $\mathcal{D}_x \neq T_x M$ is dense in M means that (M, \mathcal{D}, g) is nowhere Riemannian; this assumption is absolutely necessary since otherwise, locally, (1.13) would be the usual elliptic wave equation, and its observability properties would depend on the GCC, as stated in Theorem 1.

The key ingredient in the proof of Theorem 8 is that the GCC fails for any time $T_0 > 0$: in other words, there exist geodesics which spend a time greater than T_0 outside ω . Then, the Gaussian beam construction described in Remark 1.3 allows to contradict the observability inequality (1.14).

1.5.2. *Second main result.* Our second main result, obtained in collaboration with Chenmin Sun, sheds a different light on Theorem 8. For this second statement, we consider the generalized Baouendi-Grushin operator of example 1.8, i.e. $\Delta_\gamma = \partial_x^2 + |x|^{2\gamma} \partial_y^2$ on $M = (-1, 1)_x \times \mathbb{T}_y$, and we assume that $\gamma \geq 1$ (not necessarily an integer). We also consider the Schrödinger-type equation with Dirichlet boundary conditions

$$(1.15) \quad \begin{cases} i\partial_t u - (-\Delta_\gamma)^s u = 0 \\ u|_{t=0} = u_0 \in L^2(M) \\ u|_{x=\pm 1} = 0 \end{cases}$$

where $s \in \mathbb{N}$ is a fixed integer. Given an open subset $\omega \subset M$, we say that (1.15) is observable in time $T_0 > 0$ in ω if there exists $C > 0$ such that for any $u_0 \in L^2(M)$,

$$(1.16) \quad \|u_0\|_{L^2(M)}^2 \leq C \int_0^{T_0} \|e^{-it(-\Delta_\gamma)^s} u_0\|_{L^2(\omega)}^2 dt.$$

Our second main result, obtained in collaboration with Chenmin Sun, roughly says that observability holds if and only if the subellipticity (measured by the step $\gamma + 1$ in case $\gamma \in \mathbb{N}$), is not too strong compared to the strength of propagation s :

Theorem 9 ([LS20]). *Assume that $\gamma \geq 1$. Let $I \subsetneq \mathbb{T}_y$ be a strict open subset, and let $\omega = (-1, 1)_x \times I$. Then, for $s \in \mathbb{N}$, we have:*

- (1) *If $\frac{1}{2}(\gamma + 1) < s$, (1.15) is observable in ω for any $T_0 > 0$;*
- (2) *If $\frac{1}{2}(\gamma + 1) = s$, there exists $T_{\text{inf}} > 0$ such that (1.15) is observable in ω for T_0 if and only if $T_0 > T_{\text{inf}}$;*
- (3) *If $\frac{1}{2}(\gamma + 1) > s$, for any $T_0 > 0$, (1.15) is not observable in ω .*

The case $s = 1/2$ corresponds to wave equations. Strictly speaking, it is not covered by Theorem 9 since s is assumed to belong to \mathbb{N} in this theorem, but we see that for any positive γ it is roughly related to Point (3), and we thus recover the intuition given by Theorem 8 that subelliptic wave equations should not be observable. The case $\gamma = s = 1$ allows to recover Theorem 7, except that we do not find with our method the critical time T_{inf} . Let us also notice that if $\gamma \in \mathbb{N}$, since $\gamma + 1$ is the step of the sub-Laplacian Δ_γ , the number $\frac{1}{2}(\gamma + 1)$ appearing in Theorem 9 coincides with the exponent known as the gain of Sobolev derivatives in subelliptic estimates (see Section 1.3.3).

1.5.3. *Third main result.* Finally, our third main result, obtained in collaboration with Clotilde Fermanian Kammerer, illustrates how tools coming from noncommutative harmonic analysis

can be used to analyze sub-Laplacians and the associated evolution equations. Our main message is that *a pseudodifferential calculus “adapted to the sub-Laplacian” can be used to prove controllability and observability results for subelliptic PDEs* (instead of the usual pseudodifferential calculus used for example to prove Theorem 9). As we will see, in the present context, once defined this natural pseudodifferential calculus and the associated semi-classical measures (which relies essentially on functional analysis arguments), observability results follow quite directly.

To relate this last result to the previous ones, let us say that it is roughly linked to the critical case $s = \gamma = 1$ of Point 2 of Theorem 9, i.e., to the case where subelliptic effects are exactly balanced by the strength of propagation of the equation. Indeed, we consider the usual Schrödinger equation ($s = 1$) in some particular non-commutative Lie groups, called H-type groups (see Example 1.10), which have step 2 (corresponding to $\gamma = 1$ for Baouendi-Grushin operators). As in Point 2 of Theorem 9, we establish that under some geometric conditions on the set of observation ω , observability holds if and only if time is sufficiently large. The main difference with Theorem 9 relies in the tools used for the proof, which could lead to different generalizations. For example, the tools employed in this section allow to handle the case with analytic potential, see (1.17) below. Also, with these tools, we could imagine to prove observability results for higher-step nilpotent Lie groups, but it requires to know explicit formulas for their representations, since they determine the propagation properties of the semi-classical measures we construct (see Proposition 4.3).

To keep the presentation as simple as possible, we will present our last result only for the Heisenberg groups \mathbf{H}^d of Example 1.9, and not for general H-type groups (which are handled in [FKL20]). By doing so, we avoid defining general H-type groups, while keeping the main message of this work, namely the use of noncommutative harmonic analysis for proving observability inequalities.

Using the notations of Example 1.9, we consider $M = \tilde{\Gamma} \backslash \mathbf{H}^d$ together with the equation

$$(1.17) \quad i\partial_t u + \frac{1}{2}\Delta_M u + \mathbb{V}u = 0$$

on M , where \mathbb{V} is an analytic function defined on M . The factor $\frac{1}{2}$ in front of Δ_M plays no role, we put it here just to keep the same conventions as in [FKL20].

The Schrödinger equation (1.17) is observable in time T on the measurable set U if there exists a constant $C_{T,U} > 0$ such that

$$(1.18) \quad \forall u_0 \in L^2(M), \quad \|u_0\|_{L^2(M)}^2 \leq C_{T,U} \int_0^T \left\| e^{it(\frac{1}{2}\Delta_M + \mathbb{V})} u_0 \right\|_{L^2(U)}^2 dt.$$

Recall that Theorem 2 asserts that, in the Riemannian setting and without potential, the observability of the Schrödinger equation is implied by the Geometric Control Condition (GCC), which says that any trajectory of the geodesic flow enters U within time T . Using normal geodesics, one can also define a sub-Riemannian geodesic flow (see Section 2.1) but in some

directions of the phase space (called degenerate directions in the sequel), its trajectories are stationary. For them, we thus need to replace GCC by another condition. In the case of the Heisenberg group \mathbf{H}^d , there is only one such direction, thought as “vertical” since it is related to the ∂_z vector field.

The Heisenberg group \mathbf{H}^d comes with a Lie algebra \mathfrak{g} . Via the exponential map

$$\text{Exp} : \mathfrak{g} \rightarrow \mathbf{H}^d$$

which is a diffeomorphism from \mathfrak{g} to \mathbf{H}^d , one identifies \mathbf{H}^d and \mathfrak{g} as a set and a manifold. Moreover, \mathfrak{g} is equipped with a vector space decomposition

$$\mathfrak{g} = \mathfrak{v} \oplus \mathfrak{z},$$

such that $[\mathfrak{v}, \mathfrak{v}] = \mathfrak{z} \neq \{0\}$ and \mathfrak{z} (of dimension 1) is the center of \mathfrak{g} . We define a scalar product on \mathfrak{z} by saying that ∂_z has norm 1, which allows to identify \mathfrak{z} to its dual \mathfrak{z}^* . We also fix an orthonormal basis $V = (V_1, \dots, V_{2d})$ of \mathfrak{v} .

We consider the “vertical” flow map (also called “Reeb”, in honor of Georges Reeb) on $M \times \mathfrak{z}^*$:

$$\Phi_0^s : (x, \lambda) \mapsto (\text{Exp}(sd\mathcal{Z}^{(\lambda)}/2)x, \lambda), \quad s \in \mathbb{R}$$

where, for $\lambda \in \mathfrak{z}$, $\mathcal{Z}^{(\lambda)}$ is the element of \mathfrak{z} defined by $\lambda(\mathcal{Z}^{(\lambda)}) = |\lambda|$ (or equivalently, $\mathcal{Z}^{(\lambda)} = \lambda/|\lambda|$ after identification of \mathfrak{z} and \mathfrak{z}^*). We introduce the following H-type geometric control condition. **(H-GCC)** The measurable set U satisfies **H-type GCC** in time T if

$$\forall (x, \lambda) \in M \times (\mathfrak{z}^* \setminus \{0\}), \quad \exists s \in (0, T), \quad \Phi_0^s((x, \lambda)) \in U \times \mathfrak{z}^*.$$

The flow Φ_0^s thus replaces the geodesic flow in the degenerate direction.

Definition 1.17. We denote by $T_{\text{GCC}}(U)$ the infimum of all $T > 0$ such that H-type GCC holds in time T (and we set $T_{\text{GCC}}(U) = +\infty$ if H-type GCC does not hold in any time).

We also consider the additional assumption:

(A) All (normal) geodesics reach U in finite time (“sub-Riemannian GCC”).²

Normal geodesics are projections of bicharacteristics, see Section 2.1 for a complete definition. It is worth mentioning that, in Heisenberg groups, all geodesics are normal.

Theorem 10 ([FKL20]). Let $U \subset M$ be open and denote by \overline{U} its closure.

- (1) Assume that U satisfies **(A)** and that $T > T_{\text{GCC}}(U)$, then the observability inequality (1.18) holds, i.e. the Schrödinger equation (1.17) is observable in time T on U .
- (2) Assume $T \leq T_{\text{GCC}}(\overline{U})$, then the observability inequality (1.18) fails.

²We can reformulate Assumption (A) as follows: for any $(x, \omega) \in M \times \mathfrak{v}^*$ such that $|\omega| = 1$, there exists $s \in \mathbb{R}$ such that $\text{Exp}(s\omega \cdot V)x \in U$. Here, $\omega \cdot V = \sum_{j=1}^{2d} \omega_j V_j$ where ω_j denote the coordinates of ω in the dual basis of V and it is assumed that $\sum_{j=1}^{2d} \omega_j^2 = 1$.

This statement looks like Theorem 1 which holds for elliptic waves. In some sense, “the Schrödinger equation in Heisenberg groups looks like an elliptic wave equation”, a phenomenon which was already pointed out by authors studying Strichartz estimates, see [BGX00] and [BFKG16] for example.

Let us also say that, as already mentioned, Theorem 10 holds more generally in quotients of H-type groups.

To conclude, let us draw a table summing up most of the results presented in this introduction:

	Elliptic	Step 2	Step 4	Step > 4
Waves and half-waves ($s = 1/2$)	T_{inf} (under GCC)	∞	∞	∞
Schrödinger ($s = 1$)	0 (under GCC)	T_{inf}	∞	∞
bi-Schrödinger ($s = 2$)	0 (under GCC)	0	T_{inf}	∞
Heat	0	T_{inf} or ∞	?	?

FIGURE 1. Observability of subelliptic PDEs depending on the step.

If the results are established only in particular cases, they are in blue.

The first line is covered by Theorems 1 and 8, the second line by Theorems 2, 9 and 10, the third line by Theorem 9 and the fourth line by Theorems 3, 4 and 5 (see also Corollary 3.2). For the last two interrogation marks, see Section 5.1. Note that we illustrated Theorem 9 with the bi-Schrödinger equation, but we could have done it for a general s .

1.6. Organization of the survey. The goal of this survey is to give an overview of the ideas behind the three main results (and their proofs), namely Theorems 8, 9 and 10, to point out their common features, to develop ideas which were not necessarily written in the papers, and to explain how the tools developed along the proofs could be generalized.

In Sections 2, 3 and 4, we explain respectively the main lines of the proofs of Theorems 8, 9 and 10. Therefore, Section 2 is quite geometric and presents for example the notion of nilpotentization of a sub-Riemannian structure; Section 3 is more “semi-classical” and illustrates how resolvent estimates can be used to prove observability inequalities; and Section 4 is also “semi-classical”, but the pseudo-differential operators used in this section are adapted to the group structure (that is, they come from non-commutative Fourier analysis). Sections 3 and 4, although proving quite similar results, call for very different generalizations, which are discussed notably in Remark 4.4.

In Section 5, we finally list a few natural open questions and directions of research which follow from our work.

Acknowledgments. I first thank Clotilde Fermanian Kammerer and Chenmin Sun for these very interesting collaborations, and my PhD advisors Emmanuel Trélat and Yves Colin de

Verdière who taught me so much about the subject. I also thank Clotilde Fermanian, Chenmin Sun and Emmanuel Trélat for their suggestions and comments on a preliminary version of this survey. While working on the series of works reported here, I benefited from the interactions with many other mathematicians, among whom I am particularly grateful to Richard Lascar and Véronique Fischer for several interesting discussions. I was partially supported by the grant ANR-15-CE40-0018 of the ANR (project SRGI).

2. SUBELLIPTIC WAVE EQUATIONS ARE NEVER OBSERVABLE

In this section, we explain ideas of proof for Theorem 8. Therefore, we assume that the set of all $x \in M$ such that $\mathcal{D}_x \neq T_x M$ is dense in M . Let also $T_0 > 0$ and $\omega \subset M$ be a measurable subset such that $M \setminus \omega$ has nonempty interior. Under these assumptions, Theorem 8 will follow from Propositions 2.1 and 2.2. The first of these propositions says that given any open set of M , there exists a normal geodesic of (M, \mathcal{D}, g) traveled at speed 1 which does not leave this open set on the time-interval $[0, T_0]$: this phenomenon is not true in Riemannian manifolds but is true in sub-Riemannian manifolds under the above assumptions. We will apply this proposition to an open set lying at positive distance from ω , and thus isolate a geodesic which remains far from ω . The second proposition tells us that, as in the elliptic setting, it is possible to construct a sequence of solutions of the wave equation whose energy concentrates along this geodesic. This last fact contradicts the observability inequality.

Before stating these propositions, let us mention that, as in the elliptic setting, a normal geodesic is the projection of a null-bicharacteristic associated to the principal symbol p_2 of Δ (see Definition 2.3).

Proposition 2.1. *For any $x \in M$ and any open neighborhood V of x in M (with the initial topology on M), there exists a non-stationary normal geodesic $t \mapsto x(t)$ (traveled at speed 1) of (M, \mathcal{D}, g) such that $x(t) \in V$ for any $t \in [0, T_0]$.*

Proposition 2.2. *Let $[0, T_0] \ni t \mapsto x(t)$ be a non-stationary normal geodesic (traveled at speed 1) which does not meet ∂M . Then there exists a sequence $(u_k)_{k \in \mathbb{N}^*}$ a sequence of solutions of the wave equation (1.13) such that*

- *The energy of u_k is bounded below with respect to k and $t \in [0, T_0]$:*

$$(2.1) \quad \exists A > 0, \forall t \in [0, T_0], \quad \liminf_{k \rightarrow +\infty} E(u_k(t, \cdot)) \geq A.$$

- *The energy of u_k is small off $x(t)$: for any $t \in [0, T_0]$, we fix V_t an open subset of M for the initial topology of M , containing $x(t)$, so that the mapping $t \mapsto V_t$ is continuous (V_t is chosen sufficiently small so that this makes sense in a chart). Then*

$$(2.2) \quad \sup_{t \in [0, T_0]} \int_{M \setminus V_t} (|\partial_t u_k(t, x)|^2 + |\nabla^{sR} u_k(t, x)|^2) d\mu(x) \xrightarrow{k \rightarrow +\infty} 0.$$

The “easy part” is Proposition 2.2, while the proof of Proposition 2.1 is more complicated. Therefore, we start with a sketch of proof for Proposition 2.2 in Section 2.1, and we give ideas for the proof of Proposition 2.1 in Section 2.2.

2.1. Gaussian beams in sub-Riemannian geometry. We set $P = \partial_{tt}^2 - \Delta$ and we consider the associated Hamiltonian

$$p_2(t, \tau, x, \xi) = -\tau^2 + g^*(x, \xi)$$

with τ the dual variable of t and g^* the Hamiltonian (or principal symbol) associated to $-\Delta$. For $\xi \in T^*M$, we have

$$g^* = \sum_{i=1}^m h_{X_i}^2.$$

Here, given any smooth vector field X on M , we denoted by h_X the Hamiltonian function (momentum map) on T^*M associated with X defined in local (x, ξ) -coordinates by $h_X(x, \xi) = \xi(X(x))$. Then g^* is both the principal symbol of $-\Delta$, and also the cometric associated with g .

In $T^*(\mathbb{R} \times M)$, the Hamiltonian vector field \vec{H}_{p_2} associated with p_2 is given by $\vec{H}_{p_2}f = \{p_2, f\}$. Since $\vec{H}_{p_2}p_2 = 0$, we get that p_2 is constant along the integral curves of \vec{H}_{p_2} . Thus, the characteristic set $\mathcal{C}(p_2) = \{p_2 = 0\}$ is preserved under the flow of \vec{H}_{p_2} . Null-bicharacteristics are then defined as the maximal integral curves of \vec{H}_{p_2} which live in $\mathcal{C}(p_2)$. In other words, the null-bicharacteristics are the maximal solutions of

$$(2.3) \quad \begin{cases} \dot{t}(s) = -2\tau(s), \\ \dot{x}(s) = \nabla_{\xi} g^*(x(s), \xi(s)), \\ \dot{\tau}(s) = 0, \\ \dot{\xi}(s) = -\nabla_x g^*(x(s), \xi(s)), \\ \tau^2(0) = g^*(x(0), \xi(0)). \end{cases}$$

This definition needs to be adapted when the null-bicharacteristic meets the boundary ∂M , but in the sequel, we only consider solutions of (2.3) on time intervals where $x(t)$ does not reach ∂M .

In the sequel, we take $\tau = -1/2$, which gives $g^*(x(s), \xi(s)) = 1/4$. This also implies that $t(s) = s + t_0$ and, taking t as a time parameter, we are led to solve

$$(2.4) \quad \begin{cases} \dot{x}(t) = \nabla_{\xi} g^*(x(t), \xi(t)), \\ \dot{\xi}(t) = -\nabla_x g^*(x(t), \xi(t)), \\ g^*(x(0), \xi(0)) = \frac{1}{4}. \end{cases}$$

In other words, the t -variable parametrizes null-bicharacteristics in a way that they are traveled at speed 1.

Definition 2.3. *A normal geodesic is the projection $t \mapsto x(t)$ of a null-bicharacteristic, (i.e., a solution of (2.4)).*

Normal geodesics live in the “elliptic part” of g^* , i.e., where $g^* \neq 0$; this is the key point in the proof of Proposition 2.2. Indeed, this result is well-known in the elliptic context, it is due to Ralston [Ral82], and already Hörmander noted that his argument extended to non-elliptic operators, as long as we were working in the elliptic part of the symbol (see [Hör07, Chapter 24.2]).

Let us start the proof of Proposition 2.2. Taking charts of M , we can assume $M \subset \mathbb{R}^n$. In the sequel, we change a bit the notations: we use $x = (x_0, x_1, \dots, x_n)$ where $x_0 = t$ in the earlier notations, and we set $x' = (x_1, \dots, x_n)$. Similarly, we take the notation $\xi = (\xi_0, \xi_1, \dots, \xi_n)$ where $\xi_0 = \tau$ previously, and $\xi' = (\xi_1, \dots, \xi_n)$. The bicharacteristics are parametrized by s as in (2.3), and without loss of generality, we only consider bicharacteristics with $x(0) = 0$ at $s = 0$, which implies in particular $x_0(s) = s$ because of our choice $\tau^2(s) = g^*(x(s), \xi(s)) = 1/4$. In the sequel, a null-bicharacteristic $s \mapsto (x(s), \xi(s))$ is fixed, with $x(0) = 0$.

We take

$$(2.5) \quad v_k(x) = k^{\frac{n}{4}-1} a_0(x) e^{ik\psi(x)}.$$

where the phase $\psi(x)$ is quadratic,

$$\psi(x) = \xi'(s) \cdot (x' - x'(s)) + \frac{1}{2}(x' - x'(s)) \cdot M(s)(x' - x'(s)),$$

for $x = (t, x') \in \mathbb{R} \times \mathbb{R}^n$ and s such that $t = t(s)$. This choice of v_k is the so-called WKB ansatz, and it only yields *approximate solutions* of the wave equation (1.3). Indeed, there holds

$$(2.6) \quad \partial_{tt}^2 v_k - \Delta v_k = (k^{\frac{n}{4}+1} A_1 + k^{\frac{n}{4}} A_2 + k^{\frac{n}{4}-1} A_3) e^{ik\psi}$$

with

$$\begin{aligned} A_1(x) &= p_2(x, \nabla \psi(x)) a_0(x) \\ A_2(x) &= L a_0(x) \quad (L \text{ is a linear transport operator}) \\ A_3(x) &= \partial_{tt}^2 a_0(x) - \Delta a_0(x). \end{aligned}$$

If we take ψ to be complex-valued (by choosing a complex-valued matrix $M(s)$), then $e^{ik\psi}$ looks like a Gaussian centered at $x'(s)$, for any s . Hence, for large k , we see on (2.6) is naturally small outside the geodesic $s \mapsto x'(s)$, and the only place where it may be large is precisely on the geodesic. In order for v_k to be an approximate solution of the wave equation, it is thus sufficient to have A_1, A_2, A_3 vanish at sufficiently high order on the geodesic curve, and this is achieved by choosing adequately $M(s)$ (which is a complex-valued matrix varying continuously with s) and $a_0(x)$.

More precisely, in order to achieve the quantitative bound $\|\partial_{tt}^2 v_k - \Delta v_k\|_{L^1((0,T);L^2(M))} \leq Ck^{-\frac{1}{2}}$, one has to

- take M as a solution of the Riccati equation

$$\frac{dM}{ds} + MCM + B^T M + MB + A = 0$$

where A, B, C are explicit functions of s which can be expressed in terms of the second derivatives of p_2 along $s \mapsto (x(s), \xi(s))$;

- take a_0 so that $a_0(x(0)) \neq 0$ and La_0 vanishes along $s \mapsto x(s)$, i.e., $La_0(x(s)) = 0$. This is possible since L is a linear transport operator.

The first point yields that A_1 together with its first and second derivatives vanish along the geodesic curve $s \mapsto x(s)$, and the second point implies that A_2 vanishes along this same curve. Beside implying that v_k is an approximate solution of the wave equation, these choices guarantee that the energy of v_k concentrates (uniformly in s) along the geodesic:

$$\sup_{t \in [0, T]} \int_{M \setminus V_t} (|\partial_t v_k(t, x)|^2 + |\nabla^{sR} v_k(t, x)|^2) d\mu(x) \xrightarrow{k \rightarrow +\infty} 0$$

where V_t is defined in Proposition 2.2. Note also that the bound

$$\exists A > 0, \forall t \in [0, T_0], \quad \liminf_{k \rightarrow +\infty} E(v_k(t, \cdot)) \geq A$$

is satisfied thanks to the normalizing constant $k^{\frac{n}{4}-1}$ in the definition of v_k (2.5).

In order to pass from approximate to exact solutions of the wave equation, one chooses u_k to have the same initial data as v_k and to be an exact solution of the wave equation (1.3). Then, the Gronwall lemma ensures that (2.1) and (2.2) are satisfied, and Proposition 2.2 is proved.

Let us end this section with two other possible points of view on Proposition 2.2 and its above proof. First, it can be reformulated in terms of propagation of Lagrangian spaces, as written in [Let20b]. This point of view was developed for example by Hörmander in [Hör07, Chapter 24]. Another reformulation is that microlocal defect measures propagate in the elliptic part of the symbol as for the usual elliptic wave equation: in other words, one could take a sequence of initial data concentrating microlocally on the starting point $(x(0), \xi(0))$ of the geodesic and prove that the microlocal defect measure associated to the solutions propagates following the geodesic flow (see [Gér91]). But the Gaussian beam construction is interesting in its own and particularly simple, this is why we presented this point of view here.

2.2. Spiraling geodesics. Proposition 2.1 can be easily seen to hold in the Heisenberg group, as shown in Section 2.2.1; the proof then consists in extending its validity to the so-called “nilpotent approximation”, and then to any sub-Riemannian structure satisfying the assumptions of Theorem 8.

2.2.1. Spiraling geodesics in the 3D flat Heisenberg case. We consider the three-dimensional manifold with boundary $M_1 = (-1, 1)_{x_1} \times \mathbb{T}_{x_2} \times \mathbb{T}_{x_3}$, where $\mathbb{T} = \mathbb{R}/\mathbb{Z} \approx (-1, 1)$ is the 1D torus. We endow M_1 with the vector fields $X_1 = \partial_{x_1}$ and $X_2 = \partial_{x_2} - x_1 \partial_{x_3}$ and we set $\mathcal{D}_1 = \text{Span}(X_1, X_2)$, with the metric g_1 being defined by the fact that (X_1, X_2) is a g_1 -orthonormal frame of \mathcal{D}_1 . Then, $(M_1, \mathcal{D}_1, g_1)$ is a sub-Riemannian structure, which we will call in the sequel

the ‘‘Heisenberg manifold with boundary’’. This is a variant of Example 1.9 which has no boundary. We endow M with an arbitrary smooth volume μ and we denote by

$$(2.7) \quad g_{\text{Heis}}^* = \xi_1^2 + (\xi_2 - x_1 \xi_3)^2$$

the 3D flat Heisenberg Hamiltonian.

The geodesics we consider are given by

$$\begin{aligned} x_1(t) &= \varepsilon \sin(t/\varepsilon) \\ x_2(t) &= \varepsilon \cos(t/\varepsilon) - \varepsilon \\ x_3(t) &= \varepsilon(t/2 - \varepsilon \sin(2t/\varepsilon)/4). \end{aligned}$$

They spiral around the x_3 axis $x_1 = x_2 = 0$.

Here, one should think of ε as a small parameter. In the sequel, we denote by x_ε the geodesic with parameter ε . The associated momenta are

$$(2.8) \quad \xi_1 = \frac{1}{2} \cos(t/\varepsilon), \quad \xi_2 = 0 \quad \text{and} \quad \xi_3 = \frac{1}{2\varepsilon},$$

and we can check that that $g_{\text{Heis}}^* \equiv 1/4$. The constant ξ_3 is a kind of rounding number reflecting the fact that the geodesic spirals at a certain speed around the x_3 axis. To obtain a geodesic which makes smaller spirals, we choose a larger covector belonging to \mathcal{D}^\perp .

To prove Proposition 2.1 in the case of the Heisenberg manifold, we can assume without loss of generality that V contains 0. Then, given any $T_0 > 0$, for ε sufficiently small, we have $x_\varepsilon(t) \in V$ for every $t \in (0, T_0)$. This proves Proposition 2.1 in this case.

We proceed by approximation in order to find similar geodesics in arbitrary sub-Riemannian manifolds: for that, we need to use the fact that any sub-Riemannian structure can be locally approximated by an other sub-Riemannian structure which is nilpotent. This is the so-called nilpotentization procedure, which dates back to [RS76]. The reader can refer to [ABB19, Chapter 10] and [Jea14, Chapter 2] for recent introductions to the subject.

Essentially, the nilpotentization procedure consists in a truncation in the Taylor series of the vector fields X_1, \dots, X_m which define the sub-Laplacian. However, to obtain good properties of the ‘‘truncated’’ (or nilpotentized) structure, one needs to do the Taylor expansion in a good system of coordinates. Those systems are called ‘‘privileged coordinates’’. To introduce them properly, we need a few definitions; in particular we need to define the sub-Riemannian dilations which also play a key role in Section 4. Some readers might prefer to start with an illustrating example: it is provided at the end of the next section, as Example 2.4.

2.2.2. Nilpotent approximation. We follow the notations of Definition 1.5. This yields, for any point $q \in M$, a flag

$$\{0\} = \mathcal{D}_q^0 \subset \mathcal{D}_q^1 \subset \dots \subset \mathcal{D}_q^{k_q-1} \subsetneq \mathcal{D}_q^{k_q} = T_q M.$$

For $0 \leq i \leq k_q$, we set $n_i(q) = \dim \mathcal{D}_q^i$, and the sequence $(n_i(q))_{0 \leq i \leq k_q}$ is called the growth vector at point q . A point q' is called regular if the growth vector $(n_i(q'))_{0 \leq i \leq r(q')}$ at q' is constant for q' in a neighborhood of q .

A family (Z_1, \dots, Z_n) of n vector fields is said to be adapted to the sub-Riemannian flag of (M, \mathcal{D}, g) at q if it is a frame of $T_q M$ at q and if, for any $i \in \{1, \dots, k_q\}$, the vectors $Z_1, \dots, Z_{n_i(q)}$ at q span \mathcal{D}_q^i .

Given an adapted frame (Z_1, \dots, Z_n) at q , we consider the exponential coordinates of the second kind defined as the inverse of the local diffeomorphism

$$(x_1, \dots, x_n) \mapsto \exp(x_1 Z_1) \circ \dots \circ \exp(x_n Z_n)(q).$$

This is an example of “privileged coordinates” at q , and we shall use them in the sequel.

We define the non-decreasing sequence of weights $w_i(q)$ for $1 \leq i \leq n$ as follows. Given any $1 \leq i \leq n$, there exists a unique $1 \leq j \leq n$ such that $n_{j-1}(q) + 1 \leq i \leq n_j(q)$. We set $w_i(q) = j$. For example, for any q in the Heisenberg manifold, $w_1(q) = w_2(q) = 1$ and $w_3(q) = 2$: indeed, the coordinates x_1 and x_2 have “weight 1”, while the coordinate x_3 has “weight 2” since ∂_{x_3} requires a bracket to be generated.

The above privileged coordinates at q yield a smooth mapping $\psi_q : U \rightarrow \mathbb{R}^n$, where U is a neighborhood of q in M , with $\psi_q(q) = 0$. For every $\varepsilon \in \mathbb{R} \setminus \{0\}$, we consider the dilation $\delta_\varepsilon : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined by

$$\delta_\varepsilon(x) = (\varepsilon^{w_1(q)} x_1, \dots, \varepsilon^{w_n(q)} x_n)$$

for every $x = (x_1, \dots, x_n)$. A dilation δ_ε acts also on functions and vector fields on \mathbb{R}^n by pull-back: $\delta_\varepsilon^* f = f \circ \delta_\varepsilon$ and $\delta_\varepsilon^* X$ is the vector field such that $(\delta_\varepsilon^* X)(\delta_\varepsilon^* f) = \delta_\varepsilon^*(Xf)$ for any $f \in C^1(\mathbb{R}^n)$.

Coming back to the vector fields X_1, \dots, X_m which span \mathcal{D} , we write the Taylor expansion close to q

$$(\psi_q)_* X_i(x) \sim \sum_{\alpha, j} a_{\alpha, j} x^\alpha \partial_{x_j}.$$

We can write X_i as a formal series

$$(2.9) \quad X_i = X_i^{(-1)} + X_i^{(0)} + X_i^{(1)} + \dots$$

where $X_i^{(s)}$ is a homogeneous vector field of degree s , meaning that

$$\delta_\varepsilon^*(\psi_q)_* X_i^{(s)} = \varepsilon^s (\psi_q)_* X_i^{(s)}.$$

For $1 \leq i \leq m$, we set $\widehat{X}_i^q = (\psi_q)_* X_i^{(-1)}$, which is a vector field on \mathbb{R}^n . Then \widehat{X}_i^q is homogeneous of degree -1 with respect to dilations, i.e., $\delta_\varepsilon^* \widehat{X}_i^q = \varepsilon^{-1} \widehat{X}_i^q$ for any $\varepsilon \neq 0$. Also, $(\widehat{X}_1^q, \dots, \widehat{X}_m^q)$ generates a nilpotent Lie algebra of step k_q (see [Jea14, Proposition 2.3]).

The nilpotent approximation $(\widehat{M}^q, \widehat{\mathcal{D}}^q, \widehat{g}^q)$ of (M, \mathcal{D}, g) at q is then defined as $\widehat{M} \simeq \mathbb{R}^n$ endowed with the sub-Riemannian distribution $\widehat{\mathcal{D}}^q = \text{Span}(\widehat{X}_1^q, \dots, \widehat{X}_m^q)$ and the sub-Riemannian metric \widehat{g}^q on $\widehat{\mathcal{D}}^q$ verifying $\widehat{g}^q(\widehat{X}_i^q, \widehat{X}_j^q) = g_q(X_i, X_j)$.

Example 2.4. To illustrate the previous definitions, we reproduce here the example [Jea14, Example 2.8]. We consider the vector fields on $\mathbb{R}^2 \times \mathbb{T}$ given by $X_1 = \cos(\theta)\partial_x + \sin(\theta)\partial_y$ and $X_2 = \partial_\theta$. Since $[X_1, X_2] = \sin(\theta)\partial_x - \cos(\theta)\partial_y$, the weights are $(1, 1, 2)$ at any point. At $q = 0$, the vector fields ∂_x , ∂_θ and ∂_y define an adapted frame, with associated system of privileged coordinates (x, θ, y) . Writing down the Taylor expansions of X_1 and X_2 in these coordinates, we get the homogeneous components

$$X_1^{(-1)} = \partial_x + \theta\partial_y, \quad X_1^{(0)} = 0, \quad X_1^{(1)} = -\frac{\theta^2}{2}\partial_x - \frac{\theta^3}{6}\partial_y, \quad \dots$$

and $X_2^{(-1)} = X_2 = \partial_\theta$. Hence, the nilpotent approximation of (X_1, X_2) at $q = 0$ in coordinates (x, θ, y) is

$$\widehat{X}_1 = \partial_x + \theta\partial_y, \quad \widehat{X}_2 = \partial_\theta.$$

It generates a nilpotent Lie algebra of step 2 since all brackets of length ≥ 3 between \widehat{X}_1 and \widehat{X}_2 vanish.

2.2.3. Conclusion of the proof. Coming back to the proof of Proposition 2.1, we note that there exist $q \in V$ and $X_1, X_2 \in \mathcal{D}$ such that $[X_1, X_2](q) \notin \mathcal{D}_q$. This is because otherwise $[\mathcal{D}, \mathcal{D}] \subset \mathcal{D}$ in V , and therefore the rank condition (1.9) cannot hold in V . We are now reduced to prove Proposition 2.1 for $x = q$.

The proof of Proposition 2.1 now decomposes into two steps:

- Reduce to the nilpotentized case;
- Prove Proposition 2.1 for the nilpotentized structure $(\widehat{M}^q, \widehat{D}^q, \widehat{g}^q)$.

We will focus on the second point since the first point follows from similar considerations. Hence, the rest of this section is devoted to explaining the ideas required for proving Proposition 2.1 for the nilpotentized structure $(\widehat{M}^q, \widehat{D}^q, \widehat{g}^q)$.

The nilpotent Hamiltonian is denoted by

$$(2.10) \quad \widehat{g}^* = \sum_{i=1}^m h_{\widehat{X}_i}^2$$

where we used the notation h_Z for vector fields Z which was introduced at the beginning of Section 2.1. As explained at the end of Section 2.2.1, the proof consists in identifying a ‘‘Heisenberg sub-structure’’ in $(\widehat{M}^q, \widehat{D}^q, \widehat{g}^q)$. This terminology, which we will not make very explicit in the sequel, roughly means that the Hamiltonian (2.10) contains a hidden Heisenberg Hamiltonian (2.7).

The Baouendi-Grushin case of Example 1.7 does not contain a ‘‘Heisenberg sub-structure’’, but it contains one if we add a supplementary variable: to pass from Baouendi-Grushin to Heisenberg, one ‘‘adds a coordinate’’. For more general sub-Riemannian structures, this process of adding variables to obtain a free nilpotent Lie algebra is called desingularization, and it was introduced in [RS76].

To simplify the notations, and since for the rest of this section we are working in the nilpotentized structure, we simply denote by (M, \mathcal{D}, g) the nilpotentized structure $(\widehat{M}^q, \widehat{\mathcal{D}}^q, \widehat{g}^q)$.

Following Remark 1.6, we denote by k the step (or non-holonomic order) of \mathcal{D} (i.e., $\widehat{\mathcal{D}}^q$) at 0 (which is equal to the step at q before the nilpotentization). Let $\mathcal{L} = \mathcal{L}(1, \dots, m)$ be the free Lie algebra generated by $\{1, \dots, m\}$. We denote by \mathcal{L}^s the subspace generated by elements of \mathcal{L} of length $\leq s$, and by \tilde{n}_s the dimension of \mathcal{L}^s . Recall that in our notations M stands for \widehat{M}^q and we desingularize around 0.

Lemma 2.5 ([RS76], [Bel96](Section 7.3), [Jea14](Lemma 2.5, Theorem 2.9 and Remark 2.9)). *Let $\widetilde{M} = M \times \mathbb{R}^{\tilde{n}_r - n}$. Then there exist a neighborhood $\widetilde{U} \subset \widetilde{M}$ of $(0, 0)$, a neighborhood $U \subset M$ of 0 with $U \times \{0\} \subset \widetilde{U}$, local coordinates (u, v) on \widetilde{U} , and smooth vector fields on \widetilde{U} ,*

$$\widetilde{X}_i(u, v) = X_i(u) + \sum_{j=n+1}^{\tilde{n}_k} b_{ij}(u, v) \partial_{v_j}, \quad i = 1, \dots, m,$$

such that

- Every \tilde{q} in \widetilde{U} is regular (see beginning of Section 2.2.2);
- For any $1 \leq i \leq m$, we have $d\tilde{\pi}(\widetilde{X}_i) = X_i$, where $\tilde{\pi} : \widetilde{M} \rightarrow M$ is the canonical projection;
- $\text{Lie}(\widetilde{X}_1, \dots, \widetilde{X}_m)$ is a free nilpotent Lie algebra of step k ;

For examples illustrating this desingularization procedure, see [Jea14, Examples 2.15 and 2.16].

We gather in the notations Y_1, \dots, Y_N the vector fields $\widetilde{X}_3, \dots, \widetilde{X}_m$ and all brackets involving the vector fields $\widetilde{X}_1, \dots, \widetilde{X}_m$ except $[\widetilde{X}_1, \widetilde{X}_2]$. In other words, if we set $\mathcal{Y} = \text{Span}(Y_1, \dots, Y_N)$, then \mathcal{Y} together with $\widetilde{X}_1, \widetilde{X}_2$ and $[\widetilde{X}_1, \widetilde{X}_2]$ form a basis of the free Lie algebra $\text{Lie}(\widetilde{X}_1, \dots, \widetilde{X}_m)$.

We consider the coordinates given by

$$(x_1, x_2, x_3, y_1, \dots, y_N) \mapsto \exp(x_1 \widetilde{X}_1) \exp(x_2 \widetilde{X}_2) \exp(-x_3 [\widetilde{X}_1, \widetilde{X}_2]) \exp(y_1 Y_1) \dots \exp(y_N Y_N).$$

These are exponential coordinates of the second kind, hence privileged coordinates (see Section 2.2.2). Due to the particular triangular structure of these coordinates, the following relations hold

$$\begin{aligned} \widetilde{X}_1 &= \partial_{x_1} \\ \widetilde{X}_2 &= \partial_{x_2} && \text{on } x_1 = 0 \\ [\widetilde{X}_1, \widetilde{X}_2] &= -\partial_{x_3} && \text{on } x_1 = x_2 = 0 \\ Y_\ell &= \partial_{y_\ell} && \text{on } x_1 = x_2 = x_3 = y_1 = \dots = y_{\ell-1} = 0. \end{aligned}$$

Then, the key observation is that, due to the development (2.9) of each vector field into homogeneous components, there holds

$$\begin{aligned}\tilde{X}_1 &= \partial_{x_1} \\ \tilde{X}_2 &= \partial_{x_2} - x_1 \partial_{x_3} \\ \tilde{X}_j &= 0 \pmod{\mathcal{Y}} \text{ for any } j \geq 3.\end{aligned}$$

(see the full proof in Section 3.2 of [Let20b]).

We denote by \tilde{g}^* the Hamiltonian associated to $(\tilde{M}, \tilde{D}, \tilde{g})$. Due to the above considerations,

$$(2.11) \quad \tilde{g}^*(x, y, \xi, \eta) = \xi_1^2 + (\xi_2 - x_1 \xi_3)^2 + \sum_{i=1}^{m-2} \left(\sum_{\ell=3}^N \alpha_\ell^i \eta_\ell \right)^2$$

where $\xi_i = h_{\tilde{X}_i}$ and $\eta_\ell = h_{Y_\ell}$, and α_ℓ^i are constants. The first two terms in the right-hand side are what we called the ‘‘Heisenberg sub-structure’’.

Exactly as in the Heisenberg case where we chose the momenta (2.8) to obtain a spiraling geodesic, we consider here $\tilde{\lambda}_0 \in T_0^* \tilde{M}$ with $\langle \tilde{\lambda}_0, \tilde{X}_1 \rangle = 1/2$, $\langle \tilde{\lambda}_0, \tilde{X}_2 \rangle = 0$, $\langle \tilde{\lambda}_0, [\tilde{X}_1, \tilde{X}_2] \rangle = (2\varepsilon)^{-1}$ and $\langle \tilde{\lambda}_0, Y \rangle = 0$ for any $Y \in \mathcal{Y}$. For this choice, the last sum in the right-hand side of (2.11) gives no contribution to the geodesics, and thus we have found with this choice of $\tilde{\lambda}_0$ a spiraling geodesic in \tilde{M} . It now suffices to project it on M to conclude the proof of Proposition 2.1.

3. SUBELLIPTICITY AND STRENGTH OF PROPAGATION: BAOUENDI-GRUSHIN SCHRÖDINGER EQUATIONS

3.1. Motivation. It is well-known that the Riemannian wave equation propagates at finite speed and that the Riemannian Schrödinger equation propagates at infinite speed. For subelliptic equations, the propagation ‘‘at null speed’’ of the wave equation (shown to be a general phenomenon in Section 2), and the existence of travelling waves solutions of the Heisenberg Schrödinger equation (see [BGX00]) motivated us to undertake a general study of the propagation speed of subelliptic equations depending on two parameters: the step k and the strength of propagation s . This last parameter, which will be defined below, is equal to 1/2 for the wave equation and to 1 for the Schrödinger equation.

As a first step in this study, we took the model family of operators given by Example 1.8, i.e., we consider for $\gamma \geq 0$ (not necessarily an integer) the sub-Laplacian $\Delta_\gamma = \partial_x^2 + |x|^{2\gamma} \partial_y^2$ on the manifold $M = (-1, 1)_x \times \mathbb{T}_y$. In case $\gamma \in \mathbb{N}$, the step is given by $k = \gamma + 1$. Then, for $s \in \mathbb{N}$, we consider the equation (1.15). The strength of propagation is the parameter s appearing in this equation; this terminology seemed natural to us since equations with large s tend to propagate more quickly. The domain of observation is a strip $\omega = (-1, 1) \times I$ where $I \subset \mathbb{T}$ as mentioned in the introduction. Depending on γ and s , Theorem 9 says if the observability inequality (1.16) holds or not: in particular, it makes appear a threshold $s = \frac{\gamma+1}{2}$ at which subellipticity effects and propagation effects balance each other.

The proof of Theorem 9 splits into two parts:

- (1) To disprove observability when $s < \frac{1}{2}(\gamma + 1)$ or when $s = \frac{1}{2}(\gamma + 1)$ and time is small (i.e., for Point (3) and part of Point (2)), we construct solutions of (1.15) which propagate along the vertical line $\{x = 0\}$. This is different from what was done in Section 2: in Section 2, we were working in the elliptic part of the symbol of the sub-Laplacian, whereas here the constructed solutions propagate in the characteristic manifold (where the sub-Laplacian is not elliptic). Thus, the present construction is more involved and less general than that of Section 2, but these difficulties are unavoidable since we are dealing with Schrödinger equations whose speed of propagation in the elliptic part of the symbol is infinite. This Gaussian beam construction is described in Section 3.2
- (2) To prove observability when $s > \frac{1}{2}(\gamma + 1)$ or when $s = \frac{1}{2}(\gamma + 1)$ and time is large (i.e., for Point (1) and part of Point (2)), we establish a sharp resolvent estimate, i.e., a time-independent inequality which describes the maximal concentration outside ω of an approximate eigenfunction of Δ_γ . By a classical argument due to Nicolas Burq and Maciej Zworski (see [BZ04]), this implies observability. Indeed, this method of proof gives as corollaries an observability result for the heat equation and a decay estimate for the damped wave equation, both driven by the sub-Laplacian Δ_γ . The resolvent estimate is presented and commented in Section 3.3, and (sketchly) proved in Section 3.4.

3.2. Vertical Gaussian beams for contradicting observability. For proving Point (3) and Point (2) for small times, it is indeed sufficient to deal with the “critical” case $s = \frac{1}{2}(\gamma + 1)$. Point (3) then follows immediately from the abstract result [Mil12, Corollary 3.9]: if (1.15) was observable for some $T > 0$ and some $s < \frac{\gamma+1}{2}$, then it would be observable in any time for $s = \frac{1}{2}(\gamma + 1)$, which would be a contradiction. Hence, in the sequel, we assume $s = \frac{1}{2}(\gamma + 1)$. Moreover, we also assume that $\gamma > 1$; for the case $s = \gamma = 1$, the argument is slightly different, see [BS19].

The non-observability part of Point (2) immediately follows from the following proposition:

Proposition 3.1. *There exist $T_0 > 0$ and a sequence of solutions $(v_n)_{n \in \mathbb{N}}$ of (1.15) with initial data $(v_n^0)_{n \in \mathbb{N}}$ such that $\|v_n^0\|_{L^2(M)} = 1$ and*

$$(3.1) \quad \int_0^{T_0} \int_\omega |v_n(t, x, y)|^2 dx dy dt \xrightarrow{n \rightarrow +\infty} 0.$$

Let us explain the intuition behind this result on the simpler example of the Grushin-Schrödinger equation

$$(3.2) \quad i\partial_t u + \Delta_G u = 0$$

where $\Delta_G = \partial_x^2 + x^2 \partial_y^2$ has been introduced in Example 1.7. We consider this equation in $\mathbb{R}_t \times \mathbb{R}_x \times \mathbb{R}_y$ instead of $\mathbb{R}_t \times (-1, 1)_x \times \mathbb{T}_y$ since computations become simpler in this (non-compact) setting. Making a Fourier expansion in the y variable, we see that we obtain from

$-\Delta_G$ a family of harmonic oscillators $-\partial_x^2 + x^2\eta^2$ whose associated eigenvalues are $(2m+1)|\eta|$, $m \in \mathbb{N}$. In other words,

$$(3.3) \quad L^2(\mathbb{R}^2) = \bigoplus_{\pm} \bigoplus_{m \in \mathbb{N}} V_m^{\pm}, \quad \Delta_G|_{V_m^{\pm}} = \pm i(2m+1)\partial_y,$$

and thus the solution of (3.2) is obtained by solving an infinite number of transport equations along the y -axis, traveling at speed $2m+1$ for $m \in \mathbb{N}$. More explicitly, the solution of (3.2) in $\mathbb{R}_t \times \mathbb{R}_x \times \mathbb{R}_y$ is

$$u(t, x, y) = \frac{1}{\sqrt{2\pi}} \sum_{m=0}^{\infty} \int_{\mathbb{R}} e^{-it(2m+1)|\eta| + iy\eta} \widehat{u}_{0,m}(\eta) h_m(\sqrt{|\eta|x}) d\eta$$

where h_m is the m -th eigenfunction of the harmonic oscillator $-\partial_x^2 + x^2$ on \mathbb{R} (Hermite function). Our goal is to find solutions whose energy is concentrated in $M \setminus \omega$ during a long time, therefore these solutions should travel slowly along the y -axis. It will be the case if we manage to construct solutions corresponding to low values of m ; indeed Burq and Sun [BS19, Section 9] showed how to construct such solutions whose only non-vanishing component is the “mode” $m = 0$ (but η takes different values).

The case of the equation $i\partial_t u - (-\Delta_\gamma)^s u = 0$ (or, more precisely, (1.15)) is more involved but it is based on the same idea. The computations are less explicit since the eigenfunctions of Δ_γ are less explicit than those of Δ_G , but the knowledge of the behaviour at infinity of the first eigenfunction of the harmonic oscillator is indeed sufficient to do the computations. This (positive) normalized ground state of the operator $Q_\gamma = -\partial_z^2 + |z|^{2\gamma}$ on \mathbb{R}_z is denoted by ϕ_γ and it satisfies

$$Q_\gamma \phi_\gamma = \mu_0 \phi_\gamma.$$

The normalized ground state of the operator $P_{\gamma,w} = -\partial_x^2 + |x|^{2\gamma} w^2$ on \mathbb{R}_x is then

$$p_\gamma(w, x) = |w|^{\frac{1}{2(\gamma+1)}} \phi_\gamma(|w|^{\frac{1}{\gamma+1}} x).$$

and the associated eigenvalue is $\lambda_\gamma(w) = \mu_0 |w|^{\frac{2}{\gamma+1}}$. The relation between the different variables is $z = |w|^{\frac{1}{\gamma+1}} x$, and w will be taken to equal to η , the Fourier dual variable of y .

We assume without loss of generality that $\omega = \mathbb{R}_x \times I_y$, where $I = (a, b)$, with $0 < a < b \leq +\infty$. Let us fix $T_0 < a/\mu_0^s$.

We take $\psi \in C_c^\infty(\frac{1}{2} \leq |\eta| \leq 1)$, a sequence $h_n \rightarrow 0$, and we consider

$$v_n(t, x, y) = h_n^{\frac{1}{2}} \int_{\mathbb{R}} \psi(h_n \eta) e^{iy\eta - it\mu_0^s |\eta|} p_\gamma(\eta, x) d\eta,$$

which is a solution of (1.15) (recall that $s = (\gamma+1)/2$). It has only high η -modes thanks to the cutoff $\psi(h_n \eta)$, and each of these modes travels at speed μ_0^s along the y -axis since $2s/(\gamma+1) = 1$. By the Plancherel theorem, we can see that $\|v_{n,0}\|_{L^2} \gtrsim 1$, so that we just have to prove (3.1).

The key intuition is that, for any $t \in \mathbb{R}_+$, the mass of $v_n(t, \cdot, \cdot)$ is concentrated near $(x, y) = (0, \mu_0^s t)$, and thus outside ω for $t \leq T_0$. To prove it, one uses the Poisson formula, which yields

$$v_n(t, x, y) = \sum_{m \in \mathbb{Z}} \widehat{K_{t,x,y}^{(n)}}(2\pi m)$$

where $\Phi_m(t, y, w) = yw - 2\pi mw - t\mu_0^s w$ and

$$\widehat{K_{t,x,y}^{(n)}}(2\pi m) = h_n^{\frac{1}{2}} \int_{\mathbb{R}} \psi(h_n w) p_\gamma(w, x) e^{i\Phi_m(t,y,w)} dw.$$

Then, (3.1) follows from the fact that for $|y| \geq a$ and $t \leq T_0$, each $\widehat{K_{t,x,y}^{(n)}}(2\pi m)$ is small.

3.3. Resolvent estimate. This subsection is devoted to explaining how time-independent estimates can be relevant for proving observability inequalities (which inherently involve a time variable), and to apply it to our particular goal of proving Point (1) and part of Point (2) of Theorem 9. The starting point of the analysis is the following theorem due to Nicolas Burq and Maciej Zworski, which will be commented after its statement:

Theorem 11 ([BZ04]). *Let $P(h)$ be self-adjoint on some Hilbert space \mathcal{H} with densely defined domain \mathcal{D} and $A(h) : \mathcal{D} \rightarrow \mathcal{H}$ be bounded. Assume that uniformly for $\tau \in I = [-b, -a] \subset \mathbb{R}$, we have the following resolvent inequality*

$$(3.4) \quad \|u\|_{\mathcal{H}} \leq \frac{G(h)}{h} \|(P(h) + \tau)u\|_{\mathcal{H}} + g(h) \|A(h)u\|_{\mathcal{H}}$$

for some $1 \leq G(h) \leq O(h^{-N_0})$. Then there exist constants $C_0, c_0, h_0 > 0$, such that for every $T(h)$ satisfying

$$\frac{G(h)}{T(h)} < c_0,$$

we have, for all $0 < h < h_0$

$$\|\psi(P(h))u\|_{\mathcal{H}}^2 \leq C_0 \frac{g(h)^2}{T(h)} \int_0^{T(h)} \|A(h) e^{-\frac{itP(h)}{h}} \psi(P(h))u\|_{\mathcal{H}}^2 dt,$$

where $\psi \in C_c^\infty((a, b))$.

This statement, proved in [BZ04, Section 3], calls for several comments:

- Typically, $P(h) = -h^2 \Delta$, or, in our case $P(h) = -h^2 \Delta_\gamma$ with domain

$$D(\Delta_\gamma) = \{u \in \mathcal{D}'(M) : \partial_x^2 u, |x|^{2\gamma} \partial_y^2 u \in L^2(M) \text{ and } u|_{\partial M} = 0\}.$$

Therefore, the term $(P(h) + \tau)u$ gives an estimate on how far from a true eigenfunction u is. The small parameter h is called semi-classical parameter, and it naturally shows up when studying Schrödinger equations.

- The operator $A(h)$ is the observation operator, and in our case it does not depend on h : it is equal to the characteristic function $\mathbf{1}_\omega$ of ω .

- The expression $\psi(P(h))$ is defined thanks to functional calculus, and $T(h)$ has to be understood as the time-scale at which the observability inequality is valid. The parameter τ shows up naturally when taking the Fourier transform in time of the equation: it is the dual variable of t .

In our case, the resolvent estimate (or quasimode estimate) (3.4) takes the following form:

Theorem 12. *Let $\gamma \geq 1$ and let ω contain a horizontal strip $(-1, 1) \times I$. There exist $C, h_0 > 0$ such that for any $v \in D(\Delta_\gamma)$ and any $0 < h \leq h_0$, there holds*

$$\|v\|_{L^2(M)} \leq C(h^{-(\gamma+1)})\|(h^2\Delta_\gamma + 1)v\|_{L^2(M)} + \|v\|_{L^2(\omega)}.$$

The exponent $h^{-(\gamma+1)}$ is optimal. It may seem strange that the parameter s does not appear in the above resolvent inequality. However, we can deduce directly from Theorem 12 another resolvent inequality where s appears:

$$\|u\|_{L^2(M)} \leq C(h^{-(\gamma+1)})\|((-h^2\Delta_\gamma)^s - 1)u\|_{L^2(M)} + \|u\|_{L^2(\omega)}.$$

From this and Theorem 11, one obtains the following *spectrally localized observability inequality*:

$$(3.5) \quad \|u_h\|_{L^2(M)}^2 \leq C \int_0^T \|e^{-it'(-\Delta_\gamma)^s} u_h\|_{L^2(\omega)}^2 dt'$$

where $u_h = \psi((-h^2\Delta_\gamma)^s)u$ with $\psi \in C_c^\infty((1/2, 2))$. This is not totally immediate, the precise argument is written in [LS20, Section 3.1]. From (3.5), we can conclude thanks to a standard unique continuation argument the proof of Theorem 9, that is, we transform this inequality for u_h (which is spectrally localized) to the inequality (1.16) which holds for any initial datum u_0 . This unique continuation argument is based on the fact that if u_* is an eigenfunction of $(-\Delta_\gamma)^s$ which vanishes on ω , then it vanishes on the whole manifold M . We refer to [LS20, Section 3.2] for full details.

3.4. Proof of Theorem 12. The proof of Theorem 12 is quite long, and we will only describe here its main lines. It consists in decomposing solutions according to the joint eigenspaces of $|D_y|$ and $-\Delta_\gamma$ (it has already been done in Section 3.2); then we distinguish between several “regimes” depending on the value of $|D_y|$ compared to the value of $-\Delta_\gamma$, and this corresponds to different geometric propagations of solutions. This strategy is inspired by that of [BS19], but here we work in a time-independent setting since we seek to prove a resolvent estimate. This fine analysis of the different microlocal regimes should be compared with the non-commutative approach of Section 4, see Remark 4.4.

The proof is by contradiction. Assume that there exists a sequence $(v_h)_{h>0}$ such that

$$(3.6) \quad \|v_h\|_{L^2(M)} = 1, \quad \|v_h\|_{L^2(\omega)} = o(1), \quad \|f_h\|_{L^2(M)} = o(h^{\gamma+1})$$

where $f_h = (h^2\Delta_\gamma + 1)v_h$, and we seek for a contradiction, which would prove Theorem 12. A short argument shows that we can furthermore assume that $v_h = \psi(h^2\Delta_\gamma)v_h$ where $\psi \in C_c^\infty((-\infty, 0))$ verifies: $\psi \equiv 1$ near -1 and $\psi = 0$ outside $(-2, -\frac{1}{2})$. Here and in all the sequel,

we use functional calculus to define expressions such as $\psi(h^2\Delta_\gamma)$. The equality $v_h = \psi(h^2\Delta_\gamma)v_h$ means that all frequencies (computed with respect to Δ_γ) of v_h are comparable to $-h^{-2}$.

We use a decomposition of v_h as $v_h = v_h^1 + v_h^2 + v_h^3 + v_h^4$ where

$$\begin{aligned} v_h^1 &= (1 - \chi_0(b_0hD_y))v_h, & v_h^2 &= (\chi_0(b_0hD_y) - \chi_0(b_0^{-1}hD_y))v_h \\ v_h^3 &= (\chi_0(b_0^{-1}hD_y) - \chi_0(h^\epsilon D_y))v_h, & v_h^4 &= \chi_0(h^\epsilon D_y)v_h, \end{aligned}$$

that is, a decomposition according to the dual Fourier variable of y . Here, $b_0 \ll 1$ and ϵ will be fixed later. Choosing a good cut-off $\chi_0 \in C_c^\infty(\mathbb{R})$, the proof consists in showing that $\|v_h^j\|_{L^2(M)} = o(1)$ for $j = 1, 2, 3, 4$, which contradicts (3.6). The methods used for each j are quite different, and roughly correspond to the different behaviours of geodesics according to their momentum $\eta \sim D_y$. More precisely:

- v_h^1 corresponds to large $|D_y|$ (i.e., large η -momenta). For example, the initial data of the vertical Gaussian beams constructed in Section 3.2 satisfy $v_h = v_h^1$. To prove that $\|v_h^1\|_{L^2(M)} = o(1)$, we use the positive commutator method. This method dates back at least to [Hör71, Section 3.5] and has been widely used, for example for proving propagation of singularities for the wave equation. Here, it is based on the relation

$$[\Delta_\gamma, x\partial_x + (\gamma + 1)y\partial_y] = 2\Delta_\gamma.$$

Using this relation and computed in two different ways the expression

$$([h^2\Delta_\gamma + 1, \chi_\epsilon(x)\phi(y)(x\partial_x + (\gamma + 1)y\partial_y)]v_h^1, v_h^1)_{L^2(M)}$$

for some well-chosen cut-offs χ_ϵ and ϕ , it is possible to deduce that $\|v_h^1\|_{L^2(M)} = o(1)$.

- v_h^2 , v_h^3 and v_h^4 are “microlocalized” in the elliptic part of the symbol of $-\Delta_\gamma$. In some sense, this implies that if we restrict to functions v_h such that $v_h^1 = 0$, a better resolvent estimate should hold, showing observability of the Schrödinger equation in any time, as in the Riemannian case.
- To prove that $\|v_h^2\|_{L^2(M)} = o(1)$, we consider a defect measure associated to $(v_h^2)_{h>0}$. The key point is that it is invariant along geodesics of the sub-Riemannian metric which reach ω in finite time (it gives no mass to other geodesics), thus it is null since $\|v_h^2\|_{L^2(\omega)} = o(1)$. This argument is similar in spirit to the construction done in [Leb92b] for the Riemannian Schrödinger equation: v_h^2 corresponds to the truly elliptic (or semi-classical) regime.
- The defect measures associated to v_h^3 and v_h^4 are more complicated to handle since they are invariant along the horizontal geodesics $y = \text{const}$ (which do not enter ω). In some sense, being an approximate solution of $(h^2\Delta_\gamma + 1)v = 0$ excludes the possibility of “scarring” along a single horizontal geodesic; equivalently, for such quasimodes, the mass of v_h^3 and v_h^4 in M is controlled by their mass in ω . To establish such properties, several tools are available: for example, the papers [BZ12] and [AM14] deal with similar issues for the elliptic Schrödinger equation in tori, using either normal form arguments

or 2-microlocal techniques. The argument we present in [LS20] relies on the positive commutator method and a similar normal form argument as in [BZ12].

3.5. Further comments and open questions. Together with Corollary 2 in [DM12], the resolvent estimate proved in Theorem 12 implies an observability result for heat-type equations:

Corollary 3.2. *Assume that $\gamma \geq 1$ and that ω contains a horizontal strip $(-1, 1)_x \times I$. For any $s > \frac{1}{2}(\gamma + 1)$ and any $T_0 > 0$, observability for the heat equation with Dirichlet boundary conditions*

$$(3.7) \quad \begin{cases} \partial_t u + (-\Delta_\gamma)^s u = 0 \\ u|_{t=0} = u_0 \in L^2(M) \\ u|_{x=\pm 1} = 0 \end{cases}$$

holds in time T_0 . In other words, there exists $C > 0$ such that for any $u_0 \in L^2(M)$, there holds

$$\|e^{-T_0(-\Delta_\gamma)^s} u_0\|_{L^2(M)}^2 \leq \int_0^{T_0} \|e^{-t(-\Delta_\gamma)^s} u_0\|_{L^2(\omega)}^2 dt.$$

Let us finally mention a few open questions raised by our study:

- What happens if $0 < \gamma < 1$, a case which is not covered by Theorem 9? Indeed, the sub-Laplacian Δ_γ on $\mathbb{R} \times \mathbb{T}$ is not essentially self-adjoint (see [BP16]), which means that the Schrödinger evolution is not well defined if we do not impose any additional boundary condition on $\{x = 0\}$.
- Is it possible to generalize Theorem 9 to other sub-Laplacians? It seems so that Point (3) might be the easiest one to generalize: very roughly, as seen in Sections 2 and 3.2, some kind of “normal form” or “approximation” argument for the sub-Laplacian could be relevant since $s < \frac{1}{2}(\gamma + 1)$ leaves some space for perturbative arguments.
- Is it possible to construct in a more robust way solutions of subelliptic Schrödinger-type equations which are microlocally concentrated in the cone where the principal symbol of the sub-Laplacian vanishes?

4. SUBELLIPTIC SCHRÖDINGER EQUATION VIA NON-COMMUTATIVE HARMONIC ANALYSIS

The last result we present in this survey, namely Theorem 10, may seem difficult to approach because of the massive use of noncommutative Fourier analysis all along the statement and proofs. Our goal here is to give some keys which could facilitate reading, and to explain why this theory is adapted to analyze subelliptic PDEs.

The plan of this section is as follows: according to the theory of non-commutative harmonic analysis, we define in Section 4.1 the (operator-valued) Fourier transform (4.1), based on the unitary irreducible representations of the group, recalled in (4.4), which form an analog to the usual frequency space. Then, in Section 4.2, we follow the same path as for the usual Fourier transform: we use the Fourier inversion formula (4.5) to define in (4.6) a class of symbols and

the associated semi-classical pseudo-differential operators in (4.7). From this, we deduce the existence of semi-classical measures, which have additional invariance properties when they are associated to solutions of the Schrödinger equation. This allows to prove the first part of Theorem 10. The second part of this theorem is proved using solutions of the Schrödinger equation which propagate along the vertical direction: although this is very close in spirit to the vertical Gaussian beam construction of Section 3.2, we have developed in [FKL20] a more robust framework for these “wave-packet solutions”, based on the same non-commutative harmonic analysis tools as before.

The material of Sections 4.1 and 4.2 borrows many ideas from [FKF20] (and of course from [FKL20]). In the sequel we use the notations of Section 1.5.3.

Additionally, note that the element $(p, q, z) = (p_1, \dots, p_d, q_1, \dots, q_d, z)$ of \mathbf{H}^d can be written

$$(p, q, z) = \exp_{\mathbf{H}^d}(p_1 X_1 + \dots + p_d X_d + q_1 Y_1 + \dots + q_d Y_d + z Z)$$

and with the Baker-Campbell-Hausdorff formula we recover the group law given in Example 1.9.

4.1. Noncommutative Fourier analysis. We aim at defining a Fourier transform adapted to M , and at proving the associated Fourier inversion formula. This is standard, and the main references are [CG04] and [Tay86] (see also Appendix A of [FKL20]).

The usual Fourier transform $\widehat{f}(\lambda) = \int_{\mathbb{R}^d} f(x) e^{-ix\lambda} dx$ is replaced in this non-commutative setting by the formula

$$(4.1) \quad \mathcal{F}f(\lambda) := \int_{\mathbf{H}^d} f(x) (\pi_x^\lambda)^* dx.$$

To give it a sense, we need to explain how to define dx and to give a sense to λ and $(\pi_x^\lambda)^*$.

To define dx , we recall that in Section 1.5.3, we defined a scalar product on \mathfrak{z} , for which ∂_z has norm 1. We also need a scalar product on \mathfrak{v} , which is obtained by saying that the $2d$ vector fields

$$X_j = \partial_{x_j} + \frac{y_j}{2} \partial_z, \quad Y_j = \partial_{y_j} - \frac{x_j}{2} \partial_z, \quad j = 1, \dots, d$$

form an orthonormal basis. This allows to consider the Lebesgue measure $dv dz$ on $\mathfrak{g} = \mathfrak{v} \oplus \mathfrak{z}$. Via the identification of \mathbf{H}^d with \mathfrak{g} by the exponential map, this induces a Haar measure dx on \mathbf{H}^d and on its quotient M . Thus, we can integrate functions on \mathbf{H}^d , define Lebesgue spaces, etc as usual.

Finally, in formula (4.1), $(\pi_x^\lambda)^*$ is an operator. In particular, $\mathcal{F}f(\lambda)$ is operator-valued. The operator $(\pi_x^\lambda)^*$ is the adjoint of the irreducible representations π_x^λ of \mathbf{H}^d .

Definition 4.1. *The (strongly continuous) unitary representations of a locally compact topological group G are the homomorphisms $\pi : G \rightarrow U(\mathcal{H}_\pi)$ where $U(\mathcal{H}_\pi)$ is the group of unitary operators on a Hilbert space \mathcal{H}_π , which satisfy that $g \mapsto \pi(g)\xi$ is a norm continuous function for every $\xi \in \mathcal{H}_\pi$.*

A unitary representation is called irreducible if the only closed linear subspaces of \mathcal{H}_π invariant under $\pi(g)$ for all $g \in G$ are 0 and \mathcal{H}_π . The set of all unitary irreducible representations (modulo unitary equivalence) is denoted by \widehat{G} .

A specificity of Heisenberg groups, and more generally of H-type groups, is that their irreducible representations can be explicitly computed, thanks to Kirillov's theory. We will neither enter the details of Kirillov's theory nor show the computations specific to H-type groups (see Appendix A of [FKL20]), but only give the explicit expression of irreducible representations:

- For $\lambda \in \mathfrak{z}^* \setminus \{0\} \sim \mathbb{R} \setminus \{0\}$ and $x = (p, q, z) \in \mathbf{H}^d$, we consider the operator π_x^λ defined by

$$(4.2) \quad \pi_x^\lambda \Phi(\xi) = e^{i(\lambda(z) + \frac{1}{2}|\lambda|p \cdot q + \text{sgn}(\lambda)\sqrt{|\lambda|\xi \cdot q})} \Phi\left(\xi + \sqrt{|\lambda|}p\right),$$

which acts on functions $\Phi \in L^2(\mathbb{R}^d)$ (in (4.2), sgn is the sign function). Then, $\pi^\lambda(\cdot) = \pi_x^\lambda$ defines a unitary representation of \mathbf{H}^d which is infinite dimensional (since the Hilbert space $L^2(\mathbb{R}^d)$ has infinite dimension) and which can be proved to be irreducible.

- For $\omega \in \mathfrak{v}^*$ and $x = \text{Exp}(V + Z) \in \mathbf{H}^d$ with $V \in \mathfrak{v}$ and $Z \in \mathfrak{z}$, we set

$$(4.3) \quad \pi_x^{0,\omega} = e^{i\omega(V)}$$

and $\pi^{0,\omega}(\cdot) = \pi_x^{0,\omega}$ can thus be seen as a 1-dimensional representation over the Hilbert space $\mathcal{H}_{(0,\omega)} = \mathbb{C}$.

Then, the set $\widehat{\mathbf{H}}^d$ of all unitary irreducible representations modulo unitary equivalence is parametrized by $(\mathfrak{z}^* \setminus \{0\}) \sqcup \mathfrak{v}^*$ (see Proposition 2.4 and Theorem 2.5 in Chapter 1 of [Tay86]):

$$(4.4) \quad \widehat{\mathbf{H}}^d = \{\text{class of } \pi^\lambda : \lambda \in \mathfrak{z}^* \setminus \{0\}\} \sqcup \{\text{class of } \pi^{0,\omega} : \omega \in \mathfrak{v}^*\}.$$

The formula (4.1) defines the Fourier transform for $\lambda \in \mathfrak{z}^* \setminus \{0\}$, but we need to complete it with a formula for $\omega \in \mathfrak{v}^*$:

$$\widehat{f}(0, \omega) = \mathcal{F}f(0, \omega) := \int_{\mathbf{H}^d} f(x) (\pi_x^{(0,\omega)})^* dx.$$

With these definitions at hand, one can prove an inversion formula for $f \in \mathcal{S}(\mathbf{H}^d)$ and $x \in \mathbf{H}^d$:

$$(4.5) \quad f(x) = c_0 \int_{\mathfrak{z}^* \setminus \{0\}} \text{Tr} \left(\pi_x^\lambda \mathcal{F}f(\lambda) \right) |\lambda|^d d\lambda,$$

where Tr denotes the trace of operators of $\mathcal{L}(L^2(\mathbb{R}^d))$ (see [Tay86, Theorem 2.7 in Chapter 1]). Note that in this inversion formula, the finite dimensional irreducible representations given by (4.3) are absent.

4.2. Symbols and semi-classical measures. Starting from the Fourier inversion formula (4.5), we define symbols (of pseudo-differential operators) on M as a class of functions on $M \times \widehat{\mathbf{H}^d}$. The set $M \times \widehat{\mathbf{H}^d}$ is interpreted as the phase space of M , in analogy with the fact that $\mathbb{T}^d \times \mathbb{R}^d$ is the phase space of the torus \mathbb{T}^d .

To start, we note that the set of functions on the quotient $M = \widetilde{\Gamma} \backslash \mathbf{H}^d$ is in one-to-one relation with the set of $\widetilde{\Gamma}$ -left periodic functions on \mathbf{H}^d , i.e., functions on \mathbf{H}^d such that

$$\forall x \in \mathbf{H}^d, \quad \forall \gamma \in \widetilde{\Gamma}, \quad f(\gamma x) = f(x).$$

We consider the class of symbols \mathcal{A}_0 of fields of operators defined on $M \times \widehat{\mathbf{H}^d}$ by

$$\sigma(x, \lambda) \in \mathcal{L}(L^2(\mathbb{R}^d)), \quad (x, \lambda) \in M \times \widehat{\mathbf{H}^d},$$

that are smooth in the variable x and Fourier transforms of functions of the set $\mathcal{S}(\mathbf{H}^d)$ of Schwartz functions on \mathbf{H}^d in the variable λ : for all $(x, \lambda) \in M \times \widehat{\mathbf{H}^d}$,

$$(4.6) \quad \sigma(x, \lambda) = \mathcal{F}\kappa_x(\lambda)$$

where $\kappa(\cdot) \in \mathcal{C}^\infty(M, \mathcal{S}(\mathbf{H}^d))$. One can easily check that these symbols form an algebra (which is a motivation for introducing them as Fourier transforms of functions κ_x).

As already mentioned in Section 2.2.2), there is a natural family of dilations on \mathbf{H}^d defined as

$$\delta_\varepsilon(x, y, z) = (\varepsilon x, \varepsilon y, \varepsilon^2 z), \quad (x, y, z) \in \mathbf{H}^d, \quad \varepsilon > 0.$$

If $\varepsilon > 0$, we associate with κ_x (and thus with $\sigma(x, \lambda)$) the function κ_x^ε defined on \mathbf{H}^d by

$$\kappa_x^\varepsilon(\cdot) = \varepsilon^{-Q} \kappa_x(\delta_{\varepsilon^{-1}}(\cdot)),$$

We then define the semi-classical pseudo-differential operator $\text{Op}_\varepsilon(\sigma)$ via the identification of functions f on M with $\widetilde{\Gamma}$ -left periodic functions on \mathbf{H}^d recalled above:

$$(4.7) \quad \text{Op}_\varepsilon(\sigma)f(x) = \int_{\mathbf{H}^d} \kappa_x^\varepsilon(y^{-1}x)f(y)dy.$$

The operator $\text{Op}_\varepsilon(\sigma)$ is well-defined as an operator on M . Indeed,

$$\text{Op}_\varepsilon(\sigma)f(\gamma x) = \text{Op}_\varepsilon(\sigma)f(x).$$

It is not difficult to check that these definitions yield a “good symbolic calculus”: for example, the family of operators $(\text{Op}_\varepsilon(\sigma))_{\varepsilon>0}$ is uniformly bounded in $\mathcal{L}(L^2(M))$. This allows to define semi-classical measures (see [Gér91] and [Bur97] for similar propositions in the Euclidean framework):

Proposition 4.2. *Let $(u^\varepsilon)_{\varepsilon>0}$ be a bounded family in $L^\infty(\mathbb{R}, L^2(M))$. There exist a sequence $(\varepsilon_k) \in (\mathbb{R}_+^*)^\mathbb{N}$ with $\varepsilon_k \xrightarrow[k \rightarrow +\infty]{} 0$ and a map $t \mapsto \Gamma_t d\gamma_t$ in $L^\infty(\mathbb{R}, \widetilde{\mathcal{M}}_{ov}^+(M \times \widehat{\mathbf{H}^d}))$ such that we have*

for all $\theta \in L^1(\mathbb{R})$ and $\sigma \in \mathcal{A}$,

$$(4.8) \quad \int_{\mathbb{R}} \theta(t) (\text{Op}_{\varepsilon_k}(\sigma) u^{\varepsilon_k}(t), u^{\varepsilon_k}(t))_{L^2(M)} dt \xrightarrow{k \rightarrow +\infty} \int_{\mathbb{R} \times M \times \widehat{\mathbf{H}}^d} \theta(t) \text{Tr}(\sigma(x, \lambda) \Gamma_t(x, \lambda)) d\gamma_t(x, \lambda) dt.$$

Here \mathcal{A} is the “closure” of \mathcal{A}_0 in some sense made precise in [FKF20].

Here are a few comments on this statement:

- We denote by $\widetilde{\mathcal{M}}_{ov}(M \times \widehat{\mathbf{H}}^d)$ the set of pairs (γ, Γ) where γ is a positive Radon measure on $M \times \widehat{\mathbf{H}}^d$ and $\Gamma = \{\Gamma(x, \lambda) \in \mathcal{L}(L^2(\mathbb{R}^d)) : \lambda \in \widehat{\mathbf{H}}^d\}$ is a measurable field of trace-class operators such that

$$\|\Gamma d\gamma\|_{\mathcal{M}} := \int_{M \times \widehat{\mathbf{H}}^d} \text{Tr}(|\Gamma(x, \lambda)|) d\gamma(x, \lambda) < \infty.$$

Here, as usual, $|\Gamma| := \sqrt{\Gamma \Gamma^*}$. Finally, we say that a pair (γ, Γ) in $\widetilde{\mathcal{M}}_{ov}(M \times \widehat{\mathbf{H}}^d)$ is positive when $\Gamma(x, \lambda) \geq 0$ for γ -almost all $(x, \lambda) \in M \times \widehat{\mathbf{H}}^d$. In this case, we write $(\gamma, \Gamma) \in \widetilde{\mathcal{M}}_{ov}^+(M \times \widehat{\mathbf{H}}^d)$.

- This set $\widetilde{\mathcal{M}}_{ov}(M \times \widehat{\mathbf{H}}^d)$ can be identified to the topological dual of the algebra of symbols \mathcal{A} (to be rigorous, it requires to take the quotient by a relation of equivalence on the pairs (γ, Γ)). This is why it naturally appears as a limit of the left-hand side of (4.8).
- Hence, the semi-classical measures that we consider here are operator-valued, whereas semi-classical measures are mostly scalar in the literature, see for example [Leb92b]. This operator-valued feature is fundamental since it is due to non-commutativity of nilmanifolds, and it is a consequence of the original features of Fourier analysis on nilpotent groups seen in Section 4.1.
- The integral in the time variable in (4.8) may seem weird at first sight, and indeed it is possible to define semi-classical measures of functions on M (thus, time-independent). However, our goal here is to study semi-classical measures associated to solutions of the non-semi-classical Schrödinger equation (1.17) (a semi-classical Schrödinger equation would have the form $i\varepsilon \partial_t u + \varepsilon^2 \Delta u = 0$). For such equations, it is difficult to derive results for the semi-classical measures at each time t (see also [AM14]). However, one can prove results for the time-averaged semi-classical measures, and this is why we define these time-averaged measures in Proposition 4.2.
- Sections 4.1 and 4.2 up to Proposition 4.2 could be generalized to all graded Lie groups through the generalization of the tools we use (see Remarks 3.3 and 4.4 in [FKF19]). However, the next proposition, namely Proposition 4.3, is specific to H-type groups (in particular, Heisenberg groups) since its proof uses the explicit expressions of irreducible representations seen in Section 4.1.

The semi-classical measures associated (by Proposition 4.2) to families of solutions to the Schrödinger equation (1.17) have special features, which are the subject of Proposition 4.3 below. To state it properly, we need some definitions.

In the (non compact) group \mathbf{H}^d , the operator

$$H(\lambda) = |\lambda| \sum_{j=1}^d \left(-\partial_{\xi_j}^2 + \xi_j^2 \right)$$

is the Fourier resolution of the sub-Laplacian $-\Delta_{\mathbf{H}^d}$ above $\lambda \in \mathfrak{z}^* \setminus \{0\}$, meaning that

$$\forall f \in \mathcal{S}(\mathbf{H}^d), \quad \mathcal{F}(-\Delta_{\mathbf{H}^d} f)(\lambda) = H(\lambda) \mathcal{F}(f)(\lambda).$$

Up to a constant, this is a quantum harmonic oscillator with discrete spectrum $\{|\lambda|(2n+d), n \in \mathbb{N}\}$ and finite dimensional eigenspaces. For each eigenvalue $|\lambda|(2n+d)$, we denote by $\Pi_n^{(\lambda)}$ and $\mathcal{V}_n^{(\lambda)}$ the corresponding spectral orthogonal projection and eigenspace.

Proposition 4.3. *Assume $\Gamma_t d\gamma_t$ is associated with a family of solutions to (1.17).*

(1) *For $(x, \lambda) \in M \times \mathfrak{z}^*$*

$$(4.9) \quad \Gamma_t(x, \lambda) = \sum_{n \in \mathbb{N}} \Gamma_{n,t}(x, \lambda) \quad \text{with} \quad \Gamma_{n,t}(x, \lambda) := \Pi_n^{(\lambda)} \Gamma_t(x, \lambda) \Pi_n^{(\lambda)}.$$

Moreover, the map $(t, x, \lambda) \mapsto \Gamma_{n,t}(x, \lambda) d\gamma_t(x, \lambda)$ defines a continuous function from \mathbb{R} into the set of distributions on $M \times (\mathfrak{z}^ \setminus \{0\})$ valued in the finite dimensional space $\mathcal{L}(\mathcal{V}_n^{(\lambda)})$ which satisfies*

$$(4.10) \quad \left(\partial_t - \left(n + \frac{d}{2} \right) \mathcal{Z}^{(\lambda)} \right) (\Gamma_{n,t}(x, \lambda) d\gamma_t(x, \lambda)) = 0$$

(2) *For $(x, (0, \omega)) \in M \times \mathfrak{v}^*$, the scalar measure $\Gamma_t d\gamma_t$ is invariant under the flow*

$$\Xi^s : (x, \omega) \mapsto (x \text{Exp}(s\omega \cdot V), \omega).$$

Here, $\omega \cdot V = \sum_{j=1}^{2d} \omega_j V_j$ where ω_j denote the coordinates of ω in the dual basis of V .

The relations (4.9) and (4.10) describe the same phenomenon as (3.3): the Schrödinger equation behaves as a superposition of waves traveling at different speeds along the vertical axis.

4.3. Proof of Theorem 10. We finally explain how Theorem 10 follows from Proposition 4.3.

4.3.1. Proof of Point (1) of Theorem 10. As in Section 3.3, to prove Point (1) of Theorem 10, it is sufficient to prove a *spectrally localized* observability inequality. Let $h > 0$ and $\psi \in C_c^\infty((1/2, 2), [0, 1])$. Using functional calculus, we set

$$(4.11) \quad \mathcal{P}_h f = \psi \left(-h^2 \left(\frac{1}{2} \Delta_M + \mathbb{V} \right) \right) f, \quad f \in L^2(M).$$

We seek to prove

$$(4.12) \quad \|\mathcal{P}_h u_0\|_{L^2(M)}^2 \leq C_0 \int_0^T \left\| e^{it(\frac{1}{2}\Delta_M + \mathbb{V})} \mathcal{P}_h u_0 \right\|_{L^2(U)}^2 dt.$$

We argue by contradiction. If (4.12) is false, then there exist $(u_0^k)_{k \in \mathbb{N}}$ and $(h_k)_{k \in \mathbb{N}}$ such that $u_0^k = \mathcal{P}_{h_k} u_0^k$,

$$(4.13) \quad \|u_0^k\|_{L^2(M)} = 1 \quad \text{and} \quad \int_0^T \|u_k(t)\|_{L^2(U)}^2 dt \xrightarrow{k \rightarrow +\infty} 0.$$

where

$$u_k(t) = e^{it(\frac{1}{2}\Delta_M + \mathbb{V})} \mathcal{P}_{h_k} u_0^k.$$

We consider (after extraction of a subsequence if necessary), the semi-classical measure $\Gamma_t d\gamma_t$ of $u_k(t)$ given by Proposition 4.2 and satisfying the properties listed in Proposition 4.3. The goal is to prove that $\gamma_t \equiv 0$.

Using the second part of (4.13), one obtains

$$\int_0^T \int_{U \times \widehat{\mathbf{H}}^d} \text{Tr}(\Gamma_t(x, \lambda)) d\gamma_t(x, \lambda) dt = 0$$

i.e., $\gamma_t \equiv 0$ above U . Setting $\gamma_{n,t}(x, \lambda) = \text{Tr}(\Gamma_{n,t}(x, \lambda)) \gamma_t(x, \lambda)$, and using the positivity of Γ_t , one can deduce that

$$(4.14) \quad \int_{U \times \mathfrak{J}^*} d\gamma_{n,t}(x, \lambda) = 0, \quad \text{for almost every } t \in [0, T], \quad \forall n \in \mathbb{N}.$$

The transport equation (4.10) tells us that $\gamma_{n,t}$ travels at speed $n + \frac{d}{2}$ along the z -axis, hence not slower than $\gamma_{0,t}$. Using (H-GCC) together with (4.14), we get that $\gamma_{n,t} \equiv 0$ for any $n \in \mathbb{N}$, hence $\gamma_t \equiv 0$. This contradicts the conservation of energy (i.e., the first part of (4.13)). Thus, (4.12) is proved.

Remark 4.4. *It is tempting to compare the approaches developed in Sections 3 and 4, which share the common goal of proving observability inequalities for subelliptic Schrödinger equations. On one side, the semi-classical measures of Section 4 seem particularly adapted: once defined the operator-valued Fourier transform, the definitions of symbols and semi-classical measures are natural since they are modeled on the Euclidean case. But this approach has the drawback to require the knowledge of global objects on the manifold (the representations), and for the moment their local and geometric aspects are not sufficiently well understood to handle more general geometric situations. On the contrary, the proof of Section 3, which uses usual pseudo-differential tools (i.e., Euclidean or Riemannian ones), is directly linked with the underlying geometry (see Section 3.4) but we see the limits of these tools in the fact that already for the simple Baouendi-Grushin models, the computations are sophisticated and sometimes heavy.*

4.3.2. *Proof of Point (2) of Theorem 10.* As for Point (3) of Theorem 9, to disprove the observability inequality (1.18), we construct a particular family of solutions of the Schrödinger equation (1.17), called wave packets (and which are somehow related to the vertical Gaussian beams of Section 3.2).

In the Euclidean context, given $(x_0, \xi_0) \in \mathbb{R}^d \times \mathbb{R}^d$ and $a \in \mathcal{S}(\mathbb{R}^d)$, the associated wave packet is the family (indexed by ε) of functions

$$(4.15) \quad u_{\text{eucl}}^\varepsilon(x) = \varepsilon^{-d/4} a\left(\frac{x-x_0}{\sqrt{\varepsilon}}\right) e^{\frac{i}{\varepsilon} \xi_0 \cdot (x-x_0)}, \quad x \in \mathbb{R}^d.$$

The oscillation along ξ_0 is forced by the term $e^{\frac{i}{\varepsilon} \xi_0 \cdot (x-x_0)}$ and the concentration on x_0 is performed via $a(\cdot/\sqrt{\varepsilon})$ at the scale $\sqrt{\varepsilon}$ for symmetry reasons: the ε -Fourier transform of $u_{\text{eucl}}^\varepsilon$, $\varepsilon^{-d/2} \widehat{u}_{\text{eucl}}^\varepsilon(\xi/\varepsilon)$ presents a concentration on ξ_0 at the scale $\sqrt{\varepsilon}$. Taking a compactly supported in the interior of a unit cell for the torus, one can generalize their definition to the case of the torus by extending them by periodicity.

To perform a similar construction in the non-commutative setting, more precisely in \mathbf{H}^d , we replace $a(\cdot/\sqrt{\varepsilon})$ by

$$a_\varepsilon(x) = a(\delta_{\varepsilon^{-1/2}}(x))$$

for some $a \in \mathcal{C}_c^\infty(G)$, and the oscillations $e^{\frac{i}{\varepsilon} \xi_0 \cdot (x-x_0)}$ by

$$e_\varepsilon(x) = (\pi_x^{\lambda_\varepsilon} \Phi_1, \Phi_2), \quad \lambda_\varepsilon = \frac{\lambda_0}{\varepsilon^2}$$

where $\lambda_0 \in \mathfrak{z}^*$ and $\Phi_1, \Phi_2 \in \mathcal{S}(\mathbb{R}^d)$.

Using the multiplication on the left by elements of $\widetilde{\Gamma}$, one can define a periodization operator \mathbb{P} which associates to functions on \mathbf{H}^d whose support is contained in a unit cell of M the $\widetilde{\Gamma}$ -left periodic function obtained by periodization. We restrict to $\varepsilon \in (0, 1)$ We consider a unit cell of M , i.e., a subset \mathcal{B} of \mathbf{H}^d which is a neighborhood of $\mathbf{1}_{\mathbf{H}^d}$ and such that $\cup_{\gamma \in \widetilde{\Gamma}} (\gamma \mathcal{B}) = \mathbf{H}^d$ and we choose functions a that are in $\mathcal{C}_c^\infty(\mathcal{B})$.

Proposition 4.5. *Let $\Phi_1, \Phi_2 \in \mathcal{S}(\mathbb{R}^d)$, $a \in \mathcal{C}_c^\infty(\mathcal{B})$, $x_0 \in M$, $\lambda_0 \in \mathfrak{z}^* \setminus \{0\}$. Then, there exists $\varepsilon_0 > 0$ such that the family $(v^\varepsilon)_{\varepsilon \in (0, \varepsilon_0)}$ defined by*

$$v^\varepsilon(x) = |\lambda_\varepsilon|^{d/2} \varepsilon^{-p/2} \mathbb{P}(e_\varepsilon a_\varepsilon)(x_0^{-1}x),$$

has only one semi-classical measure $\Gamma d\gamma$ where

$$(4.16) \quad \gamma = c_a \delta(x - x_0) \otimes \delta(\lambda - \lambda_0), \quad c_a = \|\Phi_2\|^2 \int_{G_3} |a(x_3)|^2 dx_3,$$

and Γ is the operator defined by

$$\Gamma \Phi = \frac{(\Phi, \Phi_1)}{\|\Phi_1\|^2} \Phi_1, \quad \forall \Phi \in L^2(\mathbb{R}^d).$$

What is really important in the above proposition is that γ is concentrated on a single point of the phase space.

In the rest of the proof, we say that the family v^ε is a wave packet on M with cores (x_0, λ_0) , profile a and harmonics (Φ_1, Φ_2) , and we write

$$v^\varepsilon = WP_{x_0, \lambda_0}^\varepsilon(a, \Phi_1, \Phi_2) = |\lambda_\varepsilon|^{d/2} \varepsilon^{-p/2} \mathbb{P}(e_\varepsilon a_\varepsilon)(x_0^{-1}x).$$

Remark 4.6. In [FKL20] (see Section 4 and Appendix C), we develop a more general theory of wave packets, notably showing that the structure of wave packets is preserved by the evolution under the Schrödinger flow.

We take as initial data in (1.17) a wave packet u_0^ε in M with harmonics given by the first Hermite function h_0 :

$$u_0^\varepsilon = WP_{x_0, \lambda_0}^\varepsilon(a, h_0, h_0).$$

We denote by $u^\varepsilon(t)$ the associated solution, $u^\varepsilon(t) = e^{it(\frac{1}{2}\Delta_M + \mathbb{V})}u_0^\varepsilon$. Our choice of harmonics for u_0^ε guarantees that the semi-classical measure $\Gamma_t d\gamma_t$ associated to these solutions, when decomposed according to Proposition 4.3, has only one non-vanishing component, which corresponds to $n = 0$. In other words,

$$(4.17) \quad \gamma_t(x, \lambda) = c \delta \left(x - \text{Exp} \left(t \frac{d}{2} \mathcal{Z}^{(\lambda)} \right) x_0 \right) \otimes \delta(\lambda - \lambda_0)$$

and $\Gamma_{0,t}$ is the orthogonal projector on h_0 .

Now, using the assumptions made in Point (2) of Theorem 10, there exists a continuous function $\phi : M \rightarrow [0, 1]$ such that $\phi(\Phi_0^s(x_0, \lambda_0)) = 0$ for any $s \in [0, T]$ and $\phi = 1$ on $\bar{U} \times \mathfrak{z}^*$. From this, we deduce

$$0 \leq \int_0^T \int_U |u^\varepsilon(t, x)|^2 dx dt \leq \int_0^T \int_M \phi(x) |u^\varepsilon(t, x)|^2 dx dt \xrightarrow{\varepsilon \rightarrow 0} \int_0^T \int_{M \times \mathfrak{z}^*} \phi(x) d\gamma_t(x, \lambda) dt = 0.$$

Therefore, the observability inequality (1.18) cannot hold.

5. PERSPECTIVES AND OPEN PROBLEMS

In the field of subelliptic PDEs, and notably concerning controllability/observability, several interesting questions remain unanswered, and this concluding section lists a few of them.

5.1. Observability of the heat equation for other sub-Laplacians. As explained in Section 1.4.1, the observability properties of subelliptic heat equations are known only in particular geometries. More general results would require a deeper understanding of the geometric meaning of the solutions constructed in [BCG14] or [Koe17]. Let us formulate two conjectures:

- (1) For any sub-Laplacian of step 2, if $M \setminus \omega$ has non-empty interior, observability of the associated heat equation fails for sufficiently small times $T > 0$;
- (2) For any sub-Laplacian of step ≥ 3 , if $M \setminus \omega$ has non-empty interior, observability of the associated heat equation fails for any time $T > 0$.

These conjectures are inspired by the results mentioned in Section 1.4.1 and by the paper [LL20] (see notably Section 1.4).

5.2. Observability of Schrödinger for other sub-Laplacians. Even in the Riemannian case, the observability properties of the Schrödinger equation remain mysterious: although (GCC) is known to be a sufficient condition for observability, it is not a necessary condition (see Section 1.2.2). In the sub-Riemannian case, the problem is even “more open”, since no general sufficient condition is known for the moment, except trivial ones: only very particular geometries have been explored (see Theorems 7, 9 and 10), and they rely on tools which are not robust enough to cover general (in particular non-flat) sub-Riemannian geometries.

5.3. Propagation of singularities for subelliptic PDEs. Observability properties of the wave and Schrödinger equations are related to propagation of singularities of their solutions (see for example [BLR92]). The propagation of singularities for sub-Riemannian wave equations has been addressed in a series of papers by Melrose, R. Lascar and B. Lascar, culminating with the general result of [Mel84]; but, at least to the author, the proof of this last result is not limpid, and it would be useful to rewrite it in a more comprehensive way.

Besides, the understanding of propagation of singularities for sub-Riemannian Schrödinger equations surely requires the introduction of a notion of singularity “adapted to the sub-Riemannian geometry”, i.e., taking into account the number of brackets needed to generate each direction.

5.4. Trace formulas. For most (sub-)Laplacians Δ , we do not have access to the knowledge of the full spectrum, i.e., to all eigenvalues. But it is sometimes possible to compute quantities of the form

$$\sum_{n \in \mathbb{N}} f(\lambda_n)$$

where f is a (possibly complex-valued) function and λ_n describes the spectrum (with multiplicities) of $-\Delta$, i.e., $-\Delta\varphi_n = \lambda_n\varphi_n$ for smooth functions φ_n . Classical choices for f are the following: $f(x) = e^{-tx}$ (heat equation), $f(x) = |x|^{-s}$ (zeta functions), $f(x) = \cos(t\sqrt{x})$ (wave equation), $f(x) = e^{-itx/h}$ (semi-classical Schrödinger equation).

The literature on trace formulas in Riemannian manifolds is vast. But in the sub-Riemannian case, only few trace formulas have been established, and most of them are formulated with the heat kernel. It would be of interest to prove trace formulas for other kernels.

5.5. Eigenfunctions and quasimodes of sub-Laplacians. The properties of eigenfunctions and quasimodes of sub-Laplacians remain widely unknown. Beside the concentration results given by observability properties (see for example Corollary 1.2), one could expect to characterize the weak limits of high-frequency eigenfunctions (or of the square of their modulus) in the limit where the eigenvalue tends to $+\infty$: these weak limits are known as Quantum Limits and they were widely studied in the Riemannian case. In [CdVHT18], the authors undertook their study in the sub-Riemannian case, proving that they concentrate (except for a null-density subsequence) on the characteristic cone Σ , and also showing a Quantum Ergodicity result valid

for 3D contact sub-Laplacians with ergodic Reeb flow. Their study was pursued in [Let20a], in the case where a commutativity assumption on the vector fields involved in the definition of the sub-Laplacian is satisfied: then, techniques coming from joint spectral calculus can be applied. In particular, we characterized all Quantum Limits of a family of sub-Laplacians obtained as a product of flat 3D Heisenberg sub-Laplacians. But, for example, the higher-dimensional (non-flat) contact case remains open.

5.6. Lines of attack. For the understanding of the problems mentioned in Sections 5.2, 5.3, 5.4, 5.5, and also for the derivation of WKB solutions of subelliptic equations, we imagine two possible lines of attack, which are somehow parallel:

- The derivation of microlocal normal forms for families of sub-Laplacians. By microlocal normal forms, we mean for example that some sub-Laplacians can be written, possibly after conjugation by a Fourier Integral Operator, as products or sums of simple lower-order operators: see [CdVHT18] for the microlocal normal form of 3D contact sub-Laplacians, and [Let20a] for higher-dimensional examples. These microlocal normal forms encapsulate many spectral properties of sub-Laplacians; they allow to reinterpret some of the results presented in this survey and we hope they could help to attack the above issues.
- The use of noncommutative harmonic analysis. If we are working directly in a nilpotent Lie group as in Section 4, it seems that the representations of the group, particularly if they are known explicitly, could be a starting point for tackling the above issues. But more generally, since tangent spaces of sub-Riemannian manifolds come naturally with the structure of (quotients of) nilpotent Lie groups with dilations, this approach might serve for the same issues in general sub-Riemannian manifolds.

REFERENCES

- [ABB19] Andrei Agrachev, Davide Barilari, and Ugo Boscain. *A comprehensive introduction to sub-Riemannian geometry*, volume 181. Cambridge University Press, 2019.
- [ABM18] Paolo Albano, Antonio Bove, and Marco Mughetti. Analytic hypoellipticity for sums of squares and the Treves conjecture. *Journal of Functional Analysis*, 274(10):2725–2753, 2018.
- [ALM16] Nalini Anantharaman, Matthieu Léautaud, and Fabricio Macià. Wigner measures and observability for the Schrödinger equation on the disk. *Inventiones mathematicae*, 206(2):485–599, 2016.
- [AM14] Nalini Anantharaman and Fabricio Macià. Semiclassical measures for the Schrödinger equation on the torus. *Journal of the European Mathematical Society*, 16(6):1253–1288, 2014.
- [BC17] Karine Beauchard and Piermarco Cannarsa. Heat equation on the Heisenberg group: Observability and applications. *Journal of Differential Equations*, 262(8):4475–4521, 2017.
- [BCG14] Karine Beauchard, Piermarco Cannarsa, and Roberto Guglielmi. Null controllability of Grushin-type operators in dimension two. *Journal of the European Mathematical Society*, 16(1):67–101, 2014.
- [BDE20] Karine Beauchard, Jérémie Dardé, and Sylvain Ervedoza. Minimal time issues for the observability of Grushin-type equations. volume 70, pages 247–312, 2020.

- [Bel96] André Bellaïche. The tangent space in sub-Riemannian geometry. In *Sub-Riemannian geometry*, pages 1–78. Springer, 1996.
- [BFKG16] Hajer Bahouri, Clotilde Fermanian Kammerer, and Isabelle Gallagher. Dispersive estimates for the Schrödinger operator on step-2 stratified Lie groups. *Analysis & PDE*, 9(3):545–574, 2016.
- [BG97] Nicolas Burq and Patrick Gérard. Condition nécessaire et suffisante pour la contrôlabilité exacte des ondes. *Comptes Rendus de l'Académie des Sciences-Series I-Mathematics*, 325(7):749–752, 1997.
- [BGX00] Hajer Bahouri, Patrick Gérard, and Chao-Jiang Xu. Espaces de Besov et estimations de Strichartz généralisées sur le groupe de Heisenberg. *Journal d'Analyse Mathématique*, 82(1):93–118, 2000.
- [BL13] Ugo Boscain and Camille Laurent. The Laplace-Beltrami operator in almost-Riemannian geometry. *Annales de l'institut Fourier*, 63(5):1739–1770, 2013.
- [BLR92] Claude Bardos, Gilles Lebeau, and Jeffrey Rauch. Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary. *SIAM Journal on Control and Optimization*, 30(5):1024–1065, 1992.
- [BMM15] Karine Beauchard, Luc Miller, and Morgan Morancey. 2D Grushin-type equations: minimal time and null controllable data. *Journal of Differential Equations*, 259(11):5813–5845, 2015.
- [BP16] Ugo Boscain and Dario Prandi. Self-adjoint extensions and stochastic completeness of the Laplace–Beltrami operator on conic and anticonic surfaces. *Journal of Differential Equations*, 260(4):3234–3269, 2016.
- [BPS18] Karine Beauchard and Karel Pravda-Starov. Null-controllability of hypoelliptic quadratic differential equations. *Journal de l'École polytechnique – Mathématiques*, 5:1–43, 2018.
- [BS19] Nicolas Burq and Chenmin Sun. Time optimal observability for Grushin Schrödinger equation. *arXiv preprint arXiv:1910.03691*, 2019.
- [Bur97] Nicolas Burq. Mesures semi-classiques et mesures de défaut. *Astérisque*, 245:167–195, 1997.
- [BZ04] Nicolas Burq and Maciej Zworski. Geometric control in the presence of a black box. *Journal of the American Mathematical Society*, 17(2):443–471, 2004.
- [BZ12] Nicolas Burq and Maciej Zworski. Control for Schrödinger operators on tori. *Mathematical Research Letters*, 19(2):309–324, 2012.
- [CdVHT18] Yves Colin de Verdière, Luc Hillairet, and Emmanuel Trélat. Spectral asymptotics for sub-Riemannian Laplacians, I: Quantum ergodicity and quantum limits in the 3-dimensional contact case. *Duke Mathematical Journal*, 167(1):109–174, 2018.
- [CG04] Laurence Corwin and Frederick P. Greenleaf. *Representations of nilpotent Lie groups and their applications: Volume 1, Part 1, Basic theory and examples*, volume 18. Cambridge University Press, 2004.
- [Cor07] Jean-Michel Coron. *Control and nonlinearity*. Number 136. American Mathematical Soc., 2007.
- [DGL06] Belhassen Dehman, Patrick Gérard, and Gilles Lebeau. Stabilization and control for the nonlinear Schrödinger equation on a compact surface. *Mathematische Zeitschrift*, 254(4):729–749, 2006.
- [DJN19] Semyon Dyatlov, Long Jin, and Stéphane Nonnenmacher. Control of eigenfunctions on surfaces of variable curvature. *arXiv preprint arXiv:1906.08923*, 2019.
- [DK20] Michel Duprez and Armand Koenig. Control of the Grushin equation: non-rectangular control region and minimal time. *ESAIM: Control, Optimisation and Calculus of Variations*, 26:3, 2020.
- [DM12] Thomas Duyckaerts and Luc Miller. Resolvent conditions for the control of parabolic equations. *Journal of Functional Analysis*, 263(11):3641–3673, 2012.
- [DR20] Jérémie Dardé and Julien Royer. Critical time for the observability of Kolmogorov-type equations. *arXiv preprint arXiv:2006.01753*, 2020.
- [Fed71] VS Fedii. On a criterion for hypoellipticity. *Mathematics of the USSR-Sbornik*, 14(1):15, 1971.

- [FKF19] Clotilde Fermanian Kammerer and Véronique Fischer. Semi-classical analysis on H-type groups. *Science China Mathematics*, 62(6):1057–1086, 2019.
- [FKF20] Clotilde Fermanian Kammerer and Véronique Fischer. Quantum evolution and sub-Laplacian operators on groups of Heisenberg type. *To appear in Journal of spectral theory (arXiv:1910.14486)*, 2020.
- [FKL20] Clotilde Fermanian Kammerer and Cyril Letrouit. Observability and controllability for the Schrödinger equation on quotients of groups of Heisenberg type. *arXiv preprint arXiv:2009.13877*, 2020.
- [Gar17] Nicola Garofalo. Fractional thoughts. *arXiv preprint arXiv:1712.03347*, 2017.
- [Gér91] Patrick Gérard. Microlocal defect measures. *Communications in Partial Differential Equations*, 16(11):1761–1794, 1991.
- [GG10] Patrick Gérard and Sandrine Grellier. The cubic Szegő equation. volume 43, pages 761–810, 2010.
- [HN05] Bernard Helffer and Francis Nier. Hypoelliptic estimates and spectral theory for Fokker-Planck operators and Witten Laplacians. *Lecture Notes in Mathematics*, 1862, 2005.
- [Hör67] Lars Hörmander. Hypoelliptic second order differential equations. *Acta Mathematica*, 119(1):147–171, 1967.
- [Hör71] Lars Hörmander. On the existence and the regularity of solutions of linear pseudodifferential equations. *L'Enseignement Mathématique*, 17(2), 1971.
- [Hör07] Lars Hörmander. *The analysis of linear partial differential operators III: Pseudo-differential operators*. Springer Science & Business Media, 2007.
- [HPT19] Emmanuel Humbert, Yannick Privat, and Emmanuel Trélat. Observability properties of the homogeneous wave equation on a closed manifold. *Communications in Partial Differential Equations*, 44(9):749–772, 2019.
- [Jaf90] Stéphane Jaffard. Contrôle interne exact des vibrations d’une plaque rectangulaire. *Portugaliae Mathematica*, 47(4):423–429, 1990.
- [Jea14] Frédéric Jean. *Control of nonholonomic systems: from sub-Riemannian geometry to motion planning*. Springer, 2014.
- [Kap80] Aroldo Kaplan. Fundamental solutions for a class of hypoelliptic PDE generated by composition of quadratic forms. *Transactions of the American Mathematical Society*, 258(1):147–153, 1980.
- [Koe17] Armand Koenig. Non-null-controllability of the Grushin operator in 2D. *Comptes Rendus Mathématique*, 355(12):1215–1235, 2017.
- [Kol34] Andrey Kolmogoroff. Zufällige Bewegungen (zur Theorie der brownischen Bewegung). *Annals of Mathematics*, pages 116–117, 1934.
- [Leb92a] Gilles Lebeau. Control for hyperbolic equations. *Journées équations aux dérivées partielles*, pages 1–24, 1992.
- [Leb92b] Gilles Lebeau. Contrôle de l’équation de Schrödinger. *Journal de Mathématiques Pures et Appliquées*, 71(3):267–291, 1992.
- [Let20a] Cyril Letrouit. Quantum limits of sub-Laplacians via joint spectral calculus. *arXiv preprint arXiv:2007.00910v2*, 2020.
- [Let20b] Cyril Letrouit. Subelliptic wave equations are never observable. *arXiv preprint arXiv:2002.01259v2*, 2020.
- [Lio88] Jacques-Louis Lions. Contrôlabilité exacte, perturbations et stabilisation de systèmes distribués. Tome 1. *RMA*, 8, 1988.
- [Lis20] Pierre Lissy. A non-controllability result for the half-heat equation on the whole line based on the prolate spheroidal wave functions and its application to the Grushin equation. *Hal preprint <https://hal.archives-ouvertes.fr/hal-02420212/file/PSWF1D.pdf>*, 2020.

- [LL20] Camille Laurent and Matthieu Léautaud. Tunneling estimates and approximate controllability for hypoelliptic equations. *To appear in Mem. Amer. Math. Soc.*, 2020.
- [LR95] Gilles Lebeau and Luc Robbiano. Contrôle exact de l'équation de la chaleur. *Communications in Partial Differential Equations*, 20(1-2):335–356, 1995.
- [LRLTT17] Jérôme Le Rousseau, Gilles Lebeau, Peppino Terpolilli, and Emmanuel Trélat. Geometric control condition for the wave equation with a time-dependent observation domain. *Analysis & PDE*, 10(4):983–1015, 2017.
- [LS20] Cyril Letrouit and Chenmin Sun. Observability of Baouendi-Grushin-type equations through resolvent estimates. *arXiv preprint arXiv:2010.05540*, 2020.
- [Mel84] Richard Melrose. Propagation for the wave group of a positive subelliptic second-order differential operator. *Taniguchi Symp. HERT Katata*, pages 181–192, 1984.
- [Mil12] Luc Miller. Resolvent conditions for the control of unitary groups and their approximations. *Journal of Spectral Theory*, 2(1):1–55, 2012.
- [Mon94] Richard Montgomery. Abnormal minimizers. *SIAM Journal on Control and Optimization*, 32(6):1605–1620, 1994.
- [Mon95] Richard Montgomery. Hearing the zero locus of a magnetic field. *Communications in Mathematical Physics*, 168(3):651–675, 1995.
- [Mon02] Richard Montgomery. *A tour of subriemannian geometries, their geodesics and applications*. Number 91. American Mathematical Soc., 2002.
- [Mor78] Yoshinori Morimoto. On the hypoellipticity for infinitely degenerate semi-elliptic operators. *Journal of the Mathematical Society of Japan*, 30(2):327–358, 1978.
- [MS78] Richard B. Melrose and Johannes Sjöstrand. Singularities of boundary value problems. I. *Communications on Pure and Applied Mathematics*, 31(5):593–617, 1978.
- [Ral82] James Ralston. Gaussian beams and the propagation of singularities. *Studies in partial differential equations*, 23(206):C248, 1982.
- [RS76] Linda Preiss Rothschild and Elias M. Stein. Hypoelliptic differential operators and nilpotent groups. *Acta Mathematica*, 137(1):247–320, 1976.
- [Tay86] Michael Eugene Taylor. *Noncommutative harmonic analysis*. Number 22. American Mathematical Soc., 1986.
- [Tre99] François Trèves. Symplectic geometry and analytic hypo-ellipticity. In *Proceedings of Symposia in Pure Mathematics*, volume 65, pages 201–219. American Mathematical Society, 1999.

(C. Letrouit) SORBONNE UNIVERSITÉ, UNIVERSITÉ PARIS-DIDEROT SPC, CNRS, INRIA, LABORATOIRE JACQUES-LOUIS LIONS, ÉQUIPE CAGE, F-75005 PARIS.

DMA, ÉCOLE NORMALE SUPÉRIEURE, CNRS, PSL RESEARCH UNIVERSITY, 75005 PARIS.

Email address: `letrouit@ljl11.math.upmc.fr`