



HAL
open science

Combining human and snail indicators for an integrative risk assessment of metal(loid)-contaminated soils

M. Louzon, B. Pauget, A. Pelfrêne, F. Gimbert, A. de Vaufleury

► To cite this version:

M. Louzon, B. Pauget, A. Pelfrêne, F. Gimbert, A. de Vaufleury. Combining human and snail indicators for an integrative risk assessment of metal(loid)-contaminated soils. *Journal of Hazardous Materials*, 2020, pp.124182. 10.1016/j.jhazmat.2020.124182 . hal-03109798

HAL Id: hal-03109798

<https://hal.science/hal-03109798v1>

Submitted on 10 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 Combining human and snail indicators for an integrative risk assessment of metal(loid)- 2 contaminated soils

3 Authors: M. Louzon¹, B. Pauget², A. Pelfrène³, F. Gimbert¹, A. de Vaufleury¹

4 ¹ UMR CNRS 6249 Chrono-Environment, University of Franche-Comté, 16 route de Gray,
5 25030 Besançon Cedex, France

6 ² TESORA, Le Visium, 22 Avenue Aristide Briand, 94110 Arcueil, France.

7 ³ Yncrea Hauts-de-France, ULR 4515, Laboratoire Génie Civil et géo-Environnement
8 (LGCgE), F-59000 Lille, France.

9 Corresponding author: annette.devaufleury@univ-fcomte.fr

10 **Keywords:** bioavailability, bioaccessibility, metal(loid)s, hazard quotient, carcinogenic risk
11 *Cantareus aspersus*

12 Abstract

13 With the new soil uses such as land restoration and to protect wilderness, the human
14 health risk assessment (HHRA) and environmental risk assessment (ERA) should be
15 combined. Based on the relationships demonstrated between an indicator of soil quality, the
16 land snail, and human exposure, the aim of this study is to examine the snail and human risk
17 indicators twenty-nine soils contaminated by metal(loid)s. HHRA was evaluated by both
18 hazard quotient and carcinogenic risk. When the human health indicators were ranked as
19 uncertain, they were weighted by bioaccessibility to refine the risk assessment. The ERA was
20 performed with risk coefficient after *ex situ* snail exposure. The results showed strong and
21 novel relationships between human health and environmental risk indicators that had never
22 been found before. For 62 % of the soils, both indicators revealed either a confirmed risk or
23 an uncertain level of risk. **Overall pollutants present greater risk for human than for
24 environment, with 55 vs 28 % of the studied soils classified in the proven risk, respectively.**
25 An original integrative risk assessment of polluted soils has been proposed, that shall help
26 setting up relevant strategies to manage contaminated soils considering not only human but
27 also environmental indicators of risk.

28

29 1. Introduction

30 The One Health concept demonstrates that human, animal and ecosystem health are
31 interrelated, notably concerning infectious diseases and chemical contamination ([Aguirre et](#)

32 [al., 2016](#); [Lebov et al., 2017](#); [Destoumieux-Garzon et al., 2018](#); [Harrison et al., 2019](#)). This
33 concept is not currently applied to the management of contaminated soils in France, in
34 verifying the compatibility of soil uses (*e.g.*, community gardens, recreational areas), or in
35 assessing the environmental risk ([MEEM, 2017](#)). Moreover, the remediation of industrial and
36 urban wastelands requires the assessment of both the human and ecological health risks in the
37 context of increasing alternative uses to both limit urban sprawl and protect the wilderness
38 (such as through land restoration approaches). In most cases, only the total concentration of
39 contaminants in soil is considered ([Jia et al., 2018](#)), although according to the bioavailability
40 concept, only a fraction of these contaminants is available and may therefore be assimilated
41 by organisms, leading to adverse effects ([Allan, 2002](#); [ISO 17402, 2008](#); [ISO 19204, 2017](#)).
42 To determine an accurate assessment of risk, there is a real need to refine human health and
43 environmental risk assessments (HHRA and ERA, respectively) to account for the
44 bioavailable concentrations of contaminants and to combine these assessments to propose an
45 integrative risk assessment (IRA) of contaminated soils ([Alvarenga et al., 2018](#); [Louzon et al.,](#)
46 [2020a](#)). This IRA aims associating HHRA and ERA in the frame of the One health concept
47 applied to the management of terrestrial pollutions. Nowadays, HHRA is very often the
48 driving endpoint in the risk assessment of contaminated soils, while the ERA is not
49 systematically carried out in the current methodology for the management of these soils. Yet,
50 usually HHRA and ERA are in most of the cases studied separately. In France, the method to
51 assess the compatibility of the toxicological risks with the current uses of the site, that is
52 called “the interpreting the state of the environment (ISE)” may be used to perform the HHRA
53 ([MEEM, 2017](#)). These toxicological risks are assessed by the comparison of hazard quotients
54 (HQ) and carcinogenic risk (CR) to toxicological reference values (TRV). To refine the
55 method of exposure and risk assessment (especially when the values of HQ and CR showed
56 uncertain levels of risk), the bioaccessible concentrations of pollutants measured in soils

57 according to validated and standardized protocols may be used (Pelfrêne et al., 2013; MEEM,
58 2017; ISO 17924, 2018). In this case, total soil concentrations are weighted by the
59 bioaccessible fraction determined using the UBM (unified bioaccessibility method) test
60 (MEEM, 2017). Some terrestrial bioassays and bioindicators can be utilized to perform an
61 ERA using for example the risk coefficient (RC) based on the identification of excess
62 transfers and toxicity of contaminants to the land snail (Pauget and de Vaufleury, 2015;
63 Louzon et al., 2020b). As previously determined in soils contaminated by As, Cd and Pb,
64 strong relationships exist between oral bioaccessibility to humans and bioavailability to snails
65 (Louzon et al., 2020a), making these methodologies relevant candidates to perform an IRA
66 for these metal(loid)s. The IRA methodology requires established amplitudes of the responses
67 of indicators used and the convergences and/or divergences between them. The goals of this
68 study were to examine indicators of human health and environmental risks of contaminated
69 soils with using the land snail, *Cantareus aspersus*, as a bioindicator of soil quality (de
70 Vaufleury, 2015; Itziou et al., 2018; Carbone and Faggio, 2019; Baroudi et al., 2020; Radwan
71 et al., 2020), to evaluate their complementarity and propose a way to combine them to
72 perform IRA of contaminated soils. This novel evaluation was conducted in accordance with
73 the French ISE methodology (MEEM, 2017), which is based on comparisons of the state of
74 the environment (*i.e.*, the risk of soil contamination) and the current anthropic uses (HHRA)
75 and the transfers to organisms of the soil fauna (ERA).

76 **2. Experimental**

77 **2.1. Soils**

78 Twenty-nine soils contaminated with different mixtures and concentrations of As, Cd
79 and Pb were sampled (0–25 cm deep) in France after humus removal (**Table 1**). Next, these
80 soils were dried (< 40°C) and sieved through 250 µm. The physicochemical parameters of
81 these soils (clay, silt, sand, organic matter (OM), carbonate (CaCO₃) contents, pH_{water},

82 cationic exchange capacity (CEC), and the oxidizable cations of aluminium (Alox) and iron
83 (Feox)) were previously characterized by Louzon et al. (2020a) and were summarized in **table**
84 **S1**. The minimal and maximal values of the physicochemical parameters were 4.3 and 8.3 for
85 pH water, 4.3 cmol kg⁻¹ and 38.8 cmol kg⁻¹ for CEC, 0.020 cmol kg⁻¹ and 0.054 cmol kg⁻¹
86 for Alox, 0.005 cmol kg⁻¹ and 0.008 cmol kg⁻¹ for Feox, < 1.00 g kg⁻¹ and 169 g kg⁻¹ for
87 CaCO₃ content, 15.4 g kg⁻¹ and 360 g kg⁻¹ for OM content, 68.0 g kg⁻¹ and 416 g kg⁻¹ for
88 clays, 75.0 g kg⁻¹ and 734 g kg⁻¹ for silts and 66.0 g kg⁻¹ and 837 g kg⁻¹ for sands. The degree
89 of soil contamination (**Table 1**) were previously characterized in Louzon et al. (2020a), and
90 the min and max concentrations were 3.28 mg kg⁻¹ dry weight (dw) and 228 mg kg⁻¹ dw for
91 As, 0.10 mg kg⁻¹ dw and 257 mg kg⁻¹ dw for Cd and 17.8 mg kg⁻¹ dw and 8971 mg kg⁻¹ dw
92 for Pb.

93 **2.2. Human health risk assessment**

94 HHRA was performed in accordance with the French methodology (MEEM, 2017).
95 On the basis of the concentration of each of the metal(loid)s in the soils, the average daily
96 dose ((ADD) in mg kg⁻¹ of body weight day⁻¹) was calculated according to the ingestion of
97 soil particles (Liang, 2017; Jia et al., 2018) with generic input data from the ISE calculation
98 grid (MEDD, 2007a). More precisely, the data considered that 91 mg of soil (Stanek et al.,
99 2001; MEEM, 2017) could be consumed daily by a child (15 kg) over 340 days for 6 years of
100 theoretical exposure (**Eq. 1**).

$$101 \quad ADD = \frac{CS \times IR \times EF \times ED}{BW \times AT} \times 10^{-6} \quad (1)$$

102 where:

103 CS: concentration of contaminant in soil (mg kg⁻¹)

104 IR: ingestion rate of soil (mg day⁻¹)

105 EF: exposure frequency (day year⁻¹)

106 ED: exposure duration (years)

107 BW: body weight (kg)

108 AT: averaging time (days) for non-carcinogenic contaminants: ED (years) x 365 days year⁻¹
109 (Bevan and Harrison, 2017) or for carcinogenic contaminants: 70 years (*i.e.*, lifetime) x 365
110 days year⁻¹.

111 With ADD, both the threshold effects by calculating the hazard quotient (HQ) and the non-
112 threshold effects by calculating the carcinogenic risk (CR) were considered by using the
113 toxicological reference values (TRV) of each contaminant (Table 2) recommended by
114 ANSES (2020) and INERIS (2018) (Eqs. 2 and 3) (MEDD, 2007b; Liang et al., 2017; Jia et
115 al., 2018), where TRV is expressed in mg kg⁻¹ day⁻¹ for threshold chemicals and in (mg kg⁻¹
116 day⁻¹)⁻¹ for carcinogens.

$$117 \quad HQ = \frac{ADD}{TRV} \quad (2)$$

$$118 \quad CR = ADD \times TRV \quad (3)$$

119 The interpretation of HQ and CR for As, Cd and Pb was ranked into three levels: no risk,
120 uncertain risk and potential risk (Table 3). When values of HQ and/or CR are uncertain
121 (Table 3), ADD can be weighted by the percentage of bioaccessible fraction measured with
122 UBM (ISO 17924, 2018; Louzon et al., 2020a). The bioaccessible fractions (expressed in %)
123 of the metal(loid)s corresponded to the ratio between the measured concentrations and soil
124 concentrations (expressed in mg kg⁻¹ dw). To refine the exposure assessment, bioaccessibility
125 can be introduced into HHRA by using a relative bioavailability factor (RB) to quantify the
126 difference in bioavailability of a metal(loid) between the soil matrix and the reference matrix
127 (on which the TRV was built). This factor can be then used in the calculation of ADD (Eq. 4).

$$128 \quad ADD_{adjusted} = ADD \times RB = \frac{ADD \times RB_{soil}}{RB_{TRV}} = ADD \times \frac{BA_{soil} \times fa_{soil}}{BA_{TRV} \times fa_{TRV}} \quad (4)$$

129 where RB_{soil} and RB_{TRV} are the absolute bioavailability of the metal(loid) in the soil matrix
130 and the reference matrix, respectively (unitless); BA_{soil} and BA_{TRV} is the absolute
131 bioaccessibility of the metal(loid) in the soil matrix and the reference matrix, respectively
132 (unitless); fa_{soil} and fa_{TRV} are the absorbed fraction of the metal(loid) extracted from the soil
133 matrix and the reference matrix, respectively (unitless). Based on [INVS \(2012\)](#), it was found
134 that for As and Cd: (i) fa_{soil} and fa_{TRV} were similar, and (ii) the absolute bioaccessibility **in the**
135 **gastric (G) phase (i.e., stomachal phase)** of these elements in drinking water (i.e., fa_{TRV}) was
136 close to 100 %. Thus, for As and Cd, the bioaccessible fraction was directly used to weight
137 the ADD (**Eq. 5**).

$$138 \quad ADD_{adjusted} (As \text{ or } Cd) = ADD \times BA_{soil} \quad (5)$$

139 A similar approach was used for Pb. The TRV established by WHO was calculated
140 considering all sources of Pb (**water, air, and food**) with the aim of not exceeding a blood
141 level of $50 \mu\text{g L}^{-1}$ in a child. The calculation was performed on the basis of: (i) a Pb uptake of
142 20 to 80 % in children's diets ([Ziegler et al., 1978](#); [De Michele, 1984](#)) and (ii) a Pb retention
143 of 30 % in the body. According to [Oomen et al. \(2006\)](#), the absolute bioavailability of Pb in
144 the reference matrix can be estimated to average 40 %. Thus, the adjusted ADD for Pb can be
145 determined by considering the maximum value of the absorption (80 %) for all soils (**Eq. 6**)
146 ([INVS, 2012](#)).

$$147 \quad ADD_{adjusted} (Pb) = ADD \times \frac{BA_{soil} \times 0.8}{0.4} = ADD \times 2 \times BA_{soil} \quad (6)$$

148 For a conservative estimate, the phase (**gastric or gastro-intestinal, i.e., G or GI**) in which oral
149 bioaccessibility **is the highest** was used to weight the ADD ([INVS, 2012](#)). Finally, the ISE
150 was performed for each soil to obtain the HHRA by considering the worst interpretation of
151 risk indicators for all contaminants (e.g., if HQ and CR for As and Pb rank a soil as no risk,

152 but its HQ for Cd rank the soil as proven risk, the soil is then considered to have risk due to
153 the Cd) using the criteria summarized in **table 3**.

154 **2.3. Environmental risk assessment**

155 The RCs for environmental risk assessments were obtained using the snail *ex situ* test
156 over 28 days (Louzon et al., 2020b). Sub-adult snails (*Cantareus aspersus*) were exposed as
157 described in Louzon et al. (2020a). To assess environmental risk by RC calculation based on
158 bioaccumulated concentrations, the following formula was used with the toxicity point (TP)
159 from ATSDR (2015) and *ex situ* threshold guide values (TGV) according to Louzon et al.
160 (2020b) and **table 2 (Eq. 7)**.

$$161 \quad RC = \left(\left(\frac{\text{Median concentration of contaminant in snail viscera at 28 days}}{TGV} \right) - 1 \right) \times TP \quad (7)$$

162 The ERA was performed for each soil by the evaluation of the risk of transferred metallic
163 elements (ERITME) index based on the sum of the RCs (Pauget and de Vaufleury, 2015) (**Eq.**
164 **8**). The interpretation of the level of risk as a function of RC and/or ERITME values is
165 possible on the basis of ecotoxic effects on the kinetics of sexual maturity (Louzon et al.,
166 *submitted*) (**Table 3**).

$$167 \quad ERITME = RC_{As} + RC_{Cd} + RC_{Pb} \quad (8)$$

168 **2.4. Statistical analysis**

169 The statistical analysis was performed with R (version 3.4.2) (R Core Team, 2018). The
170 assessment of the relationships between the indicators of human health and environmental
171 risks were performed using Spearman's correlation. The relationships between the total
172 concentrations of metal(loid)s in the soils and the excess metal(loid) transfers to snails (C_{28}
173 $_{days} > TGV$) were evaluated by comparisons of the identity line ($y = x$) of the ratio between the
174 total concentration in the soils and the SGV (soil guide values) (Pauget et al., 2013) as a

175 function of the ratio between the internal concentration in the snails after 28 days of exposure
176 and the **TGV**.

177 **3. Results and discussion**

178 **3.1. Human health risk assessment**

179 HQ and CR were calculated for each soil to estimate human health risk (**Table 4**). By
180 considering the total concentrations of metal(loid)s in the twenty-nine soil samples, the HQ
181 ranged from 0.04 to 2.86 for As, from 0 to 4.04 for Cd and from 0.16 to 80.5 for Pb,
182 **respectively. The CR ranged from 2.38×10^{-6} to 1.65×10^{-4} for As and 7.33×10^{-8} to $3.69 \times$**
183 **10^{-5} for Pb.** Overall, for the HHRA, the HQ and CR indicators demonstrated divergence in
184 several cases for the same metal(loid)s. Values of these indicators were uncertain for 16, 9
185 and 16 soils for the HQ of As, Cd and Pb, respectively, and 27 and 20 soils for the CR of As
186 and Pb, respectively (in **table 3**, the uncertainty area concerns values between 0.2 to 5 for HQ
187 and 10^{-6} to 10^{-4} for CR).

188 According to **INVS (2012)**, to refine the risk assessment, the HQ adjusted and the CR
189 adjusted were calculated on the basis of the highest bioaccessible fraction obtained between
190 the G and the GI phases (**Table 4**). The adjustment of ADD for Pb can be made by
191 considering the maximum value of the absorption (80 %) and the mean bioavailability (40 %).
192 For As, the bioaccessible fractions were higher in the GI phase than in the G phase for 9 soils
193 (O1, O12, O13, O14, O16, O18, O19, O20 and O22), whereas for Cd and Pb, the
194 bioaccessible fraction was systematically higher in the G phase than in the GI phase (**Table**
195 **1**). For these metals, the higher bioaccessibility in the G phase is explained by the acidic pH
196 of the stomach-simulated fluids, which contributes to higher solubilization of metals (**Ruby et**
197 **al., 1996; Wragg et al., 2011**). For As, the relatively similar level of bioaccessibility between
198 the G and GI phases could be explained by the reduced adsorption and precipitation reactions

199 caused by the particularly geochemical behaviour of this metalloid in the simulated fluid
200 conditions at a neutral pH (Ruby et al., 1996; Wragg et al., 2011; Rahman et al., 2017).

201 For As, after adjustments of the values in the uncertainty area, the HQ and CR adjusted
202 ranged from 0.02 to 2.86 and 4.28×10^{-7} to 1.65×10^{-4} , respectively, leading to decreased
203 numbers of soils (7 and 23, respectively) with uncertain risk (Table 4). In the current
204 management of As polluted soils CR is shown to be present in very low concentrations in soil,
205 notably at the SGV of As in soils (25 mg kg^{-1}). Indeed, the soil O7 for example, presented a
206 measured concentration of As of 12.8 mg kg^{-1} (Table 1), and the unadjusted and adjusted CRs
207 were 9.27×10^{-6} and 3.24×10^{-6} , respectively (Table 4), despite the low bioaccessible
208 fractions (35 % and 25 %, respectively for the G and the GI phases). In this example, the
209 adjustment of the CR by bioaccessibility is not sufficient to reduce the uncertainty of the risk
210 to the level of “no risk” (Table 4). The adjustment of the HQ in the area of uncertain risk for
211 Cd by bioaccessibility provides a HQ ranging from 0 to 4.04 with a decrease of 3 % in the
212 number of soils with uncertain risk (*i.e.*, 31 % and 28 % of the soils with uncertain risk for
213 HQ and HQ adj., respectively) (Table 4). The decrease in the number of soils with uncertain
214 risk for the HQ of As was higher than for Cd in relation to the lower gastric bioaccessibility of
215 As versus Cd in the soils of this study (G bioaccessible median fraction for As and Cd of 32
216 % and 84 %, respectively) (Table 1). For As, and to a lesser extent for Cd, these results
217 validated the inclusion of bioaccessibility to improve the interpretation of human health risks,
218 both in terms of toxicological threshold effects and carcinogenic effects. For Pb, the adjusted
219 HQ and CR ranged from 0.16 to 80.5 and from 7.33×10^{-8} to 7.38×10^{-5} , respectively (Table
220 4). However, due to the correction applied to obtain a better ADD consideration for Pb, the
221 use of an orally bioaccessible fraction to refine the HQ or CR of Pb seems inefficient, unless
222 the bioaccessible fraction was smaller than 50 % (see eq. 6 in 2.2. Materials and methods
223 section).

224 The results of the ISE showed that none of the studied soils presented no risk to human health,
225 notably due to the contamination of these soils by Pb. However, after the soils with uncertain
226 risk were ranked based on adjustments of HQ and CR, two soils became no risk (O16 and
227 O22). For an ISE based on results that were risk adjusted or non-risk adjusted with
228 bioaccessibility, the results showed that 11 and 18 soils, respectively, had uncertain risk,
229 leading to a soil status modification of 24 % (**Table 4**). The relevance of adjusting HQ and
230 CR with the bioaccessible fraction to decrease the uncertainty regarding the risks of
231 contaminated soils (carcinogenic or not) to human health was observed, particularly for soil
232 contaminated by As (**Table 4**). However, because Pb and its adjustment with **eq. 6** mainly
233 drive the ISE, the bioaccessibility fraction in this study was not sufficient to decrease the
234 global risk for HHRA for most of the soils. The use of the bioaccessible G fraction is
235 questionable for physiological reasons. Indeed, metal(loid) absorption mainly occurs in the
236 intestine ([Mushak, 1991](#); [Diamond et al., 1997](#)). Currently, in the regulations ([INVS, 2012](#)),
237 the most penalizing phase (often the G phase) is considered. Consequently, by considering the
238 G phase, which is in the most of the cases, the most penalizing to be more protective and
239 decrease the uncertainty regarding risk, the risk can be overestimated. Due to the
240 consideration of the absorption abilities of Pb in the human digestive tract (**eq. 5**), the oral
241 bioaccessibility fraction improved the HHRA and demonstrated human health risks in 5
242 additional soils (**Table 4**). However, the G phase is the main phase considered to decrease the
243 uncertainty caused by Pb in relation to the stronger bioaccessibility of Pb in the stomach
244 (**Table 1**). For physiological reasons, if the gastro-intestinal phase is considered (*i.e.*, the main
245 site of contaminant assimilation), the results could be different. Indeed, the median (min,
246 max) of the bioaccessible fractions of Pb in the gastric and gastrointestinal phases was 84 %
247 (17 %, 100 %) and 5 % (0 %, 45 %), respectively (**Table 1**).

248 **3.2. Environmental risk assessment**

249 The snail *ex situ* test was used to assess the environmental risk of these contaminated
250 soils by measuring the bioaccumulated concentrations of metal(loid)s in the visceral mass
251 (**Table 1**). The results showed that the concentrations of metal(loid)s ranged from 0.098 to
252 21.9 mg kg⁻¹ of dw viscera for As, from 2.53 to 333 mg kg⁻¹ for Cd and from 1.42 to 856 mg
253 kg⁻¹ for Pb. An inter-soil variability of bioaccumulated concentrations in *C. aspersus* was
254 found and this variability reflected the influence of the soil contamination (concentration of
255 metal(loid)s in the soils and mixtures) and the influence of the physicochemical parameters of
256 the soils on metal(loid) mobility and transfer, such as the silt content, the cationic exchange
257 capacity and the organic matter content (Pauget et al., 2012; Louzon et al., 2020a). On the
258 basis of the *ex situ* threshold guide values (TGV), RCs were calculated for each soil and
259 metal(loid) and summed for the assessment of risk by the ERITME index calculation (**Table**
260 **5**). RCs ranged from 0 to 35500 for As, 0 to 23400 for Cd and 0 to 37400 for Pb. ERITME for
261 the three metal(loid)s studied varied from 0 to 96300 (**Table 3**). Finally, according to the
262 bioavailability of As, Cd and Pb for the land snail, 11, 10 and 8 soils were classified as
263 without risks, with uncertain risk and with proven environmental risk, respectively (**Table 5**).
264 The relationships between the ratio of the concentrations of metal(loids) in the soils with the
265 soil guide values (SGV) and the ratio of the concentration of metal(loid)s in the viscera of
266 snails after 28 days of exposure to the threshold guide value (when the ratios between the
267 internal concentrations and TGV are higher than 1, a bioavailability of contaminant is
268 characterized) are shown in **figure 1**. Soils below the identity line ($y = x$) showed a high
269 bioavailability of metal(loids)s in snails based on internal concentrations of the metal(loid)s
270 measured in the snails, while soils above the line showed low bioavailability of excess
271 metal(loids)s in the soils to snails (**Figure 1**). This inter-soil variability according to the total
272 concentrations in soils and in the snail is related to the concept of the bioavailability. More
273 precisely, for As transfer to snails, this variability results from the positive influence of silt

274 and the negative influence of organic matter content in soils (Louzon et al., 2020a). The
275 values of RCs and the ERITME index constituted a basis to prioritize the management of
276 polluted soils (*e.g.*, remediation, adjustment of use) according to the environmental risk on the
277 basis of the toxicity of metal(loid)s transferred to snails (Pauget and de Vaufleury, 2015;
278 Mariet et al., 2017; Louzon et al., 2020b). For example, by considering the ERITME values
279 for soils O28 and O7 (724 and 12328, respectively), the O7 soil presented a higher
280 environmental risk than O28, and its management should be prioritized to limit environmental
281 risk, although the soil O7 was less contaminated (Table 5). Considering the variability of the
282 values of RC that reflect the soil contamination and their physico-chemical properties (Tables
283 1 and 5), the bioavailability assessment is a way to refine environmental impact assessments
284 according to the specificity of each site in terms of physicochemical parameters (Bradham et
285 al., 2006; Pauguet et al., 2013).

286 3.3. Correlations between human health and environmental risk indicators

287 Spearman's correlation coefficients (ρ) were calculated between human health risk
288 indicators (HQ or CR) either adjusted or not adjusted for bioaccessibility if the values were in
289 the area of uncertain risk and the environmental risk indicator assessed with land snails (RC),
290 and are summarized in table 6. Strong correlations were demonstrated between RC and
291 values of human health risk indicators with significant ρ between 0.57 to 0.93. For HQ, the
292 adjustment with bioaccessibility increased the correlations with RC for Pb (+2 %) but not for
293 Cd, and it decreased the correlations with RC for As (-10 %). For CR, the adjustment with
294 bioaccessibility increased the correlations with RC for As (+13 %) and for Pb (+3 %) (Table
295 6). Overall, the correlations between human health and environmental risk indicators were
296 improved by considering oral bioaccessibility for humans. On the basis of the correlations
297 obtained, the relationships discovered between oral bioaccessibility and the bioavailability of
298 metal(loid)s in soils as bioindicators of soil quality by Rahman et al. (2017) and Louzon et al.

299 (2020a) can be extended between human health (non-carcinogenic and carcinogenic) and
300 environmental health risk indicators.

301 **3.4. Integrative risk assessment**

302 According to the strong correlations observed, the interpretation of the assessment of
303 the human health, whether adjusted or not adjusted with oral bioaccessible fraction and
304 environmental risk was compared for each soil to evaluate similarities and discrepancies
305 (Figure 2). In figure 2, on the basis of the values used to interpret the indicators of risk
306 (Table 3), the worst level of human health risk with HQ and/or CR was used to rank the soils.
307 The comparison between HHRA or HHRA adj. and ERA showed that 52 % and 45 % of the
308 soils, respectively, had the same level of risks (Figure 2). These differences can be related to
309 the parameters used to consider the absorption of Pb (see eq. 6 in subsection 2.2. of the
310 Materials and methods) that increases the values of HQ and CR adjusted. For example,
311 without adjustment, the HQ for Pb of soil O1 was 3.86 (uncertain risk), while with
312 adjustment, the HQ was 6.25 (proven risk) (Table 4).

313 These results demonstrated convergences and some differences in terms of interpretation
314 between the conclusions of the ERA and the HHRA of the soils. For example, existing human
315 activities at the site of soil O28 appear inappropriate due to the state and the fate of the soil
316 contamination, while this does not seem to be the case for the soil fauna organisms. Indeed,
317 the RC of Pb in the O28 and O14 soils differed greatly (724 and 11560, respectively) with
318 various environmental risk interpretations (no risk and uncertain risk) (Table 4), while their
319 total soil concentrations were higher (467 mg kg⁻¹ and 719 mg kg⁻¹) (Table 1). Their human
320 health risk was also higher (HQ for Pb of 4.19 and 6.45 for O28 and O14 soils, respectively).
321 The HQ of the Pb in these soils was close to 5, and classifying the soils as having high
322 uncertain and proven risks, respectively (Table 4). This example illustrates the different
323 conclusions that can be drawn between HHRA and ERA. For HHRA, these two soils are

324 classified as **high uncertain and proven risk, respectively**, which is in agreement with the alert
325 threshold of saturnism risk (300 mg kg⁻¹) and its vigilance threshold (100 mg kg⁻¹) (HCSP,
326 2014), which implies management of soil pollution (notably by considering the G phase). For
327 soil fauna, soils O14 and O28 were classified as either having uncertain risk and no risk,
328 respectively.

329 The overall assessment of the environmental risk cannot be performed only on the basis of the
330 response of one bioindicator. However, the use of an accumulation bioindicator that shows the
331 transfer and the bioavailability of contaminants provides essential data regarding risk. This
332 information constitutes the necessary elements of decisions-making to help manage the
333 pollution of the contaminated soils. The IRA was performed for each soil by considering the
334 worst interpretation of the risk to be the most protective for the health of the ecosystem
335 including humans (One Health). Finally, of the twenty-nine investigated soils, often highly
336 contaminated with As, Cd and Pb (in terms of mixture and concentration), fifteen soils
337 presented risks and fourteen were of uncertain risk (**Figure 2**). For these last fourteen soils,
338 supplementary investigations are required to refine the level of risk, potentially by using
339 others bioindicators such as earthworms (Fründ et al., 2011) and specific assays, to assess for
340 example the carcinogenic (Vasseur and Bonnard, 2014; de Souza et al., 2015; Liu et al., 2015)
341 and endocrine disruptor (Höss and Weltje, 2007; Mertl et al., 2014; Druart et al., 2017)
342 potentials of these soils (ISO 19204, 2017). **These approaches that would possibly involve**
343 **integrated biomarker responses (IBR) can be coupled with the ERITME index as**
344 **demonstrated by Mleiki et al. (2020) for another snail species, *Cantareus apertus*.** According
345 to the cost of the decontamination (Lombi and Hamon, 2005), the prioritization of the
346 remediation of the soils could be ranked according to the values of the risk index (HQ adj.,
347 CR adj. and ERITME) to better assess the adequacy between the uses of the soils and the risks
348 (MEEM, 2017) and to protect terrestrial ecosystems (Pauget et al., 2013). Currently, the

349 management of contaminated soils is predominantly based on HHRA. When HHRA
350 concludes that a risk to humans exists, decontamination or other management methodologies
351 should be used. When remediation solutions are **too much expensive** or when the economic
352 balance of the site redevelopment project does not cover the management overhead, the
353 project can be abandoned and the site remains unclaimed. The assessment of the
354 environmental risk and the pollutant transfers in the environment developed in this study
355 reveal that, although a risk to humans was identified, the contaminants were not necessarily
356 bioavailable (environmental and/or toxicological bioavailability) to an organism of the soil
357 fauna (see yellow stars in **figure 2**). Consequently, the anthropic uses of soil and the risks
358 can be assessed by considering the bioaccessibility to humans and the bioavailability to snails
359 (**Figure 3**). In certain cases, if the IRA shows a higher risk for human health with the current
360 use ($ISE > \text{thresholds}$) but not for the environment ($ERITME < \text{threshold}$), in the future,
361 through developing this methodology to establish a management plan (by using quantitative
362 human health risk assessment), certain sites can be managed by avoiding human activities
363 (*e.g.*, restricting access). In cases such as urban wastelands, the pollutants remain in place to
364 promote land restoration that can increase the biodiversity and/or other alternative uses, such
365 as the production of renewable energy from ecological compensation.

366 **4. Conclusion**

367 This study provides essential knowledge to begin building bridges between human
368 health and environmental risk indicators in the context of the One Health approach and the
369 increasing global environmental crisis. **For the first time, methodologies to determine the
370 level of risk for both risk indicators were applied simultaneously to polluted soils. Strong
371 correlations are found between the values of risk indicators for human and environmental
372 health for the three metal(loid)s studied, but in most cases, their interpretation does not
373 indicate the same level of risk. Unexpectedly, for about two-third of the studied soils both**

374 indicators revealed either a confirmed risk or an uncertain level of risk. Another development
375 is that we found that overall polluted soils present greater risk for human than for the
376 environmental bioindicator, with 55 % versus 28 % of the studied soils classified in the
377 proven risk, respectively. The methodology we presented is one way to assess the integrative
378 risk of polluted soils by considering bioaccessibility for humans and bioavailability for
379 bioindicators to better consider the specificity of each site and to protect the health of the
380 ecosystem that encompasses humans and all other life forms.

381 **5. Declaration of competing interest**

382 The authors declare no conflicts of interest.

383 **6. Acknowledgements**

384 The authors would like to thank ADEME for funding the COMBINE study
385 programme (Coupler des indicateurs chimiques et Biologiques pour une évaluation
386 Intégrative des risques sanitaires et Environnementaux [grant number 1572C0310]) and the
387 ODESSA programme (Optimisation de la gestion Des Sites et Sols pollués par une mesure
388 simple de bioAccessibilité [grant number 1472C0046]). The authors thank Hubert Leprond,
389 Nadia Crini and Francis Douay for their kind advice about our work and Dominique Rieffel
390 and Caroline Amiot for their technical assistance.

391 **7. References**

- 392 Aguirre, A.A., Beasley, V.A., Auspurger, T., Benson, W.H., Whaley, J., Basu, N., 2016.
393 One health – Transdisciplinary opportunities for SETAC leadership in integrating and
394 improving the health of people, animals and the environment. *Environ. Toxicol.*
395 *Chem.* **35**, 2383-2391. <https://doi.org/10.1002/etc.3557>
- 396 Alvarenga, P., Clemente, R., Garbisu, C., Becerril, J.M., 2018. **Chapter 3 – Indicators for**
397 **monitoring mine site rehabilitation, in: Prasad, M.N.V., de Campos Favas, P.J., Maiti**

398 S.K. (Eds.), *Bio-Geotechnologies for Mine Site Rehabilitation*. Elsevier, Amsterdam,
399 pp. 49-66. <https://doi.org/10.1016/B978-0-12-812986-9.00003-8>

400 Allan, H.E., 2002. *Bioavailability of metals in terrestrial ecosystems: importance of*
401 *partitioning for bioavailability to invertebrates, microbes and plants*. Society of
402 *Environmental Toxicology and Chemistry, Pensacola*.

403 ANSES, 2020. Liste des valeurs toxicologiques de référence (VTR) construites par l'Anses.
404 Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du
405 travail, Paris. [https://www.anses.fr/fr/content/liste-des-valeurs-toxicologiques-de-](https://www.anses.fr/fr/content/liste-des-valeurs-toxicologiques-de-r%C3%A9f%C3%A9rence-vtr-construites-par-l%E2%80%99anses)
406 [r%C3%A9f%C3%A9rence-vtr-construites-par-l%E2%80%99anses](https://www.anses.fr/fr/content/liste-des-valeurs-toxicologiques-de-r%C3%A9f%C3%A9rence-vtr-construites-par-l%E2%80%99anses)

407 ATSDR, 2015. Detailed Data Table for the 2015 Priority List of Hazardous Substances that
408 will be the subject of toxicological profile. *Agency for Toxic Substances and Disease*
409 *Registry, Atlanta, Georgia, pp. 22*.

410 Baroudi, F., Al Alam, J., Faljoun, Z., Millet, M., 2020. Snail as sentinel organism for
411 monitoring the environmental pollution; a review. *Ecol. Indic.* 113, 106240.
412 <https://doi.org/10.1016/j.ecolind.2020.106240>

413 Bevan, R.J., Harrison, P.T.C., 2017. Threshold and non-threshold chemical carcinogens: A
414 survey of the present regulatory landscape. *Regul. Toxicol. Pharmacol.* 88, 291-302.
415 <https://doi.org/10.1016/j.yrtph.2017.01.003>

416 Bradham, K.D., Dayton, E.A., Basta, N.T., Schroder, J., Payton, M., Lanno, R.P., 2006.
417 Effect of soil properties on lead bioavailability and toxicity to earthworms. *Environ.*
418 *Toxicol. Chem.* 25, 769-775. <https://doi.org/10.1897/04-552R.1>

419 Carbone, D., Faggio, C., 2019. *Helix aspersa* as sentinel of development damage for
420 biomonitoring purpose: A validation study. *Molec. Reprod. Dev.* 86, 1283-1291.
421 <https://doi.org/10.1002/mrd.23117>

422 Charrier, M., Brune, A., 2003. The gut microenvironment of helioid snails (Gastropods:
423 Pulmonata): *in situ* profiles of pH, oxygen and hydrogen determined by
424 microsensors. *Can. J. Zool.* 81, 928-935. <https://doi.org/10.1139/z03-071>

425 De Michele, S.J., 1984. Nutrition of lead. *Comp. Biochem. Physiol. Part A* 78, 401-408.
426 [https://doi.org/10.1016/0300-9629\(84\)90567-x](https://doi.org/10.1016/0300-9629(84)90567-x)

427 de Vaufléury, A., 2015. Landsnail for ecotoxicological assessment of chemicals and soil
428 contamination, in: Armon, R.H., Hänninen, O. (Eds.), *Environmental Indicators*.
429 Springer, Dordrecht, pp. 345-391. https://doi.org/10.1007/978-94-017-9499-2_23

430 Denys, S., Caboche, J., Tack, K., Rychen, G., Wragg, J., Cave, M., Jondreville, C., Feidt, C.,
431 2012. *In vivo* validation of the unified BARGE method to assess the bioaccessibility
432 of arsenic, antimony, cadmium and lead in soils. *Environ. Sci. Technol.* 46, 6252-
433 6260. <https://doi.org/10.1021/es3006942>

434 Destoumieux-Garzon, D., Mavingui, P., Boetsch, G., Boissier, J., Darriet, F., Duboz, P.,
435 Fritsch, C., Giraudoux, P., Le Roux, F., Morand, S., Paillard, C., Pontier, D., Sueur,
436 C., Voituron, Y., 2018. The One Health Concept: 10 years old and a long road ahead.
437 *Front. Vet. Sci.* 5, 14. <https://doi.org/10.3389/fvets.2018.00014>

438 Diamond, G.L., Goodrum, P.E., Felter, S.P., Ruoff, W.L., 1998. Gastrointestinal absorption
439 of metals. *Drug Chem. Toxicol.* 21, 223-251.
440 <https://doi.org/10.3109/01480549809011649>

441 Druart, C., Gimbert, F., Scheifler, R., de Vaufléury, A., 2017. A full life-cycle bioassay with
442 *Cantareus aspersus* shows reproductive effects of a glyphosate-based herbicide
443 suggesting potential endocrine disruption. *Environ. Pollut.* 226, 240-249.
444 <https://doi.org/10.1016/j.envpol.2017.03.061>

445 Fründ, H.C., Graefe, U., Tisher, S., 2011. Earthworms as bioindicators of soil quality, in:
446 Karaca, A. (Ed.), *Biology of Earthworms, Soil Biology 24*. Springer, Dordrecht, pp.
447 261-278. https://doi.org/10.1007/978-3-642-14636-7_16

448 Gomot-de Vaufleury, A., 2000. Standardized growth toxicity testing (Cu, Zn, Pb and
449 Pentachlorophenol) with *Helix aspersa*. *Ecotoxicol. Environ. Safe.* 46, 41-50.
450 <https://doi.org/10.1006/eesa.1999.1872>

451 Harrison, S., Kivuti-Bitok, L., Macmillan, A., Priest, P., 2019. EcoHealth and One Health: A
452 theory-focused review in response to calls for convergence. *Environ. Int.* 132,
453 105058. <https://doi.org/10.1016/j.envint.2019.105058>

454 HCSP, 2014. Expositions au plomb: Détermination de nouveaux objectifs de gestion.
455 Rapport. *Haut Conseil de la santé publique, Paris, pp. 101.*

456 Höss, S., Weltje, L., 2007. Endocrine disruption in nematodes: effects and mechanisms.
457 *Ecotoxicology* 16, 15-28. <https://doi.org/10.1007/s10646-006-0108-y>

458 INERIS, 2018. Rapport d'étude. Bilan des choix de VTR disponibles sur le portail des
459 substances chimiques de l'INERIS. *DRC-17-163632-11568A. Institut national de*
460 *l'environnement industriel et des risques, Verneuil-en-Halatte, pp. 62.*

461 INVS, 2012. Guide pratique. Quantités de terre et poussières ingérées par un enfant de
462 moins de 6 ans et bioaccessibilité des polluants : état des connaissances et
463 propositions. *Institut de Veille Sanitaire, Paris, pp. 85.*

464 ISO 17402, 2008. Soil Quality. Requirements and guidance for the selection and application
465 of methods for the assessment of bioavailability of contaminants in soil and soil
466 materials. International Standardisation Organisation, Geneva.
467 <https://www.iso.org/standard/38349.html>

468 ISO 17924, 2018. Soil Quality. Assessment of human exposure from ingestion of soil and
469 soil material – Procedure for the estimation of the human

470 bioaccessibility/bioavailability of metals in soil. International Standardisation
471 Organisation, Geneva. <https://www.iso.org/standard/64938.html>

472 ISO 19204, 2017. Soil Quality. Procedure for site-specific ecological risk assessment of soil
473 contamination (soil quality TRIAD approach). International Standardisation
474 Organisation, Geneva. <https://www.iso.org/standard/63989.html>

475 Itziou, A., Patsis, P.A., Dimitriadis, V.K., 2019. Introduction of the land snail *Cornu*
476 *aspersum* as a bioindicator organism of terrestrial pollution with the use of a suite of
477 biomarkers. *Toxicol. Environ. Chem.* 100, 717-736.
478 <https://doi.org/10.1080/02772248.2019.1588274>

479 Jia, Z., Li, S., Wang, L., 2018. Assessment of soil heavy metals for eco-environment and
480 human health in a rapidly urbanization area of the upper Yangtze Basin. *Sci. Rep.* 8,
481 3256. <https://doi.org/10.1038/s41598-018-21569-6>

482 Lebov, J., Grieger, K., Womack, D., Zaccaro, D., Whitehead, N., Kowalczyk, B.,
483 MacDonalad, P.D.M., 2017. A framework for One Health research. *One Health* 3,
484 44-50. <https://doi.org/10.1016/j.onehlt.2017.03.004>

485 Liang, Y., Yi, X., Dang, Z., Wang, Q., Luo, H., Tang, J., 2017. Heavy metal contamination
486 and health risk assessment in the vicinity of a tailing pond in Guangdong, China. *Int.*
487 *J. Environ. Res. Public Health* 14, 1557. <https://doi.org/10.3390/ijerph14121557>

488 Liu, L., Chen, L., Floehr, T., Xiao, H., Bluhm, K., Hollert, H., Wu, L. 2015. Assessment of
489 the mutagenicity of sediments from Yangtze river estuary using *Salmonella*
490 *typhimurium*/microcosme assay. *PLoS One* 10, e0143522.
491 <https://doi.org/10.1371/journal.pone.0143522>

492 Lombi, E., Hamon, R.E., 2005. Remediation of polluted soils. In Hillel D (eds)
493 Encyclopedia of Soils in the Environment. Academic Press, Cambridge, pp 379-385.
494 <https://doi.org/10.1016/B0-12-348530-4/00087-4>

495 Louzon, M., Pelfrêne, A., Pauget, B., Gimbert, F., Morin-Crini, N., Douay F., de Vaufleury,
496 A., 2020a. Bioaccessibility of metal(loid)s in soils to humans and their
497 bioavailability to snails: a way to associate human health and ecotoxicological risk
498 assessment? *J. Hazard. Mater.* 384, 121432.
499 <https://doi.org/10.1016/j.hazmat.2019.121432>

500 Louzon, M., Pauget, B., Gimbert, F., Morin-Crini, N., de Vaufleury, A., 2020b. *Ex situ*
501 environmental risk assessment of polluted soils using threshold guide values for the
502 land snail *Cantareus aspersus*. *Sci. Total Environ.* 721, 137789
503 <https://doi.org/10.1016/j.scitotenv.2020.137789>

504 Louzon, M., Devalloir, Q., Gimbert, F., Pauget, B., Rieffel, D., de Vaufleury, A. From
505 bioavailability to risk assessment of polluted soil using snails: Link between excess
506 of transfer and inhibition of sexual maturation. *Environ. Sci. Pollut. Res.* Submitted

507 Mariet, A.L., Pauget, B., de Vaufleury, A., Bégeot, C., Walter-Simonnet, A.V., Gimbert, F.,
508 2017. Using bioindicators to assess the environmental risk of past mining activities in
509 the Vosges Mountains (France). *Ecol. Indic.* 75, 17-26.
510 <https://doi.org/10.1016/j.ecolind.2016.11.042>

511 MEDD, 2007a. L'interprétation de l'état des milieux – Description – Grille de calcul.
512 Ministère de l'Ecologie et du Développement Durable, Paris, calculs grid xls.
513 [http://ssp-infoterre.brgm.fr/sites/default/files/upload/documents/iemgc_v0-](http://ssp-infoterre.brgm.fr/sites/default/files/upload/documents/iemgc_v0-022007.xls)
514 [022007.xls](http://ssp-infoterre.brgm.fr/sites/default/files/upload/documents/iemgc_v0-022007.xls)

515 MEDD, 2007b. La démarche d'interprétation de l'état des milieux. Ministère de l'Ecologie
516 et du Développement Durable, Paris, pp. 42. [http://ssp-](http://ssp-infoterre.brgm.fr/sites/default/files/upload/documents/iem_v0-022007.pdf)
517 [infoterre.brgm.fr/sites/default/files/upload/documents/iem_v0-022007.pdf](http://ssp-infoterre.brgm.fr/sites/default/files/upload/documents/iem_v0-022007.pdf)

518 MEEM, 2017. Méthodologie Nationale de Gestion des Sites et Sols Pollués. Direction
519 générale de la prévention des risques. Ministère de l'Environnement, de l'Energie et

520 de la Mer, Paris, pp. 128. <http://ssp->
521 infoterre.brgm.fr/sites/default/files/upload/documents/methodo_ssp_2017.pdf

522 Mertl, J., Kirchnawy, C., Osorio, V., Grininger, A., Richter, A., Bergmair, J., Pyerin, M.,
523 Washüttl, M., Tacker, M., 2014. Characterization of estrogen and androgen activity
524 of food contact materials by different *in vitro* bioassays (YES, YAS, ER α and AR
525 CALUX) and chromatographic analysis (GC-MS, HPLC-MS). *PLoS One* 9,
526 [e100952](https://doi.org/10.1371/journal.pone.0100952). <https://doi.org/10.1371/journal.pone.0100952>

527 Mleiki, A., El Menif, N.J., Marigomez, I., 2020. Integrative assessment of the biological
528 responses elicited by metal pollution in the green garden snail, *Cantareus apertus*:
529 Laboratory and field studies. *Ecol. Indic.* 117, 106589.
530 <https://doi.org/10.1016/j.ecolind.2020.106589>

531 Mushak, P., 1991. Gastro-intestinal absorption of lead in children and adults: Overview of
532 biological and biophysico-chemical aspects. *Chem. Spec. Bioavailab.* 3, 87-104.
533 <https://doi.org/10.1080/09542299.1991.11083160>

534 Oomen, A., Brandon, E., Swartjes, F.A., Sips, A., 2006. How can information on oral
535 bioavailability improve human health risk assessment for lead-contaminated soils?
536 Implementation and scientific basis. RIVM report, pp. 108.
537 <https://www.rivm.nl/bibliotheek/rapporten/711701042.pdf>

538 Ottosen, L.M., Hansen, H.K., Jensen, P.E., 2009. Relation between pH and desorption of
539 Cu, Cr, Zn and Pb from industrially polluted soils. *Water Air Soil Pollut.* 201, 295-
540 304. <https://doi.org/10.1007/s11270-008-9945-z>

541 Pauget, B., Gimbert, F., Coeurdassier, M., Scheifler, R., de Vaufleury, A., 2011. Use of
542 chemical methods to assess Cd and Pb bioavailability to the snail *Cantareus*
543 *aspersus*: A first attempt taking into account soil characteristics. *J. Hazard. Mater.*
544 192, 1804-1811. <https://doi.org/10.1016/j.jhazmat.2011.07.016>

545 Pauget, B., Gimbert, F., Scheifler, R., Coeurdassier, M., de Vaufleury, A., 2012. Soil
546 parameters are key factors to predict metal bioavailability to snails based on
547 chemical extractant data. *Sci. Total Environ.* 431, 413-425.
548 <https://doi.org/10.1016/j.scitotenv.2012.05.048>

549 Pauget, B., Gimbert, F., Coeurdassier, M., Crini, N., Pérès, G., Faure, O., Douay, F., Hitmi,
550 A., Beguiristain, T., Alaphilippe, A., Guernion, M., Houot, S., Legras, M., Vian, J.F.,
551 Hedde, M., Bispo, A., Grand, C., de Vaufleury, A., 2013. Ranking field site
552 management priorities according to their metal transfer to snails. *Ecol. Indic.* 29,
553 445–454. <https://doi.org/10.1016/j.ecolind.2013.01.012>

554 Pauget, B., de Vaufleury, A., 2015. The SET and ERITME indices: Integrative tools for the
555 management of polluted sites. *Ecol. Indic.* 53, 206–210.
556 <https://doi.org/10.1016/j.ecolind.2015.01.037>

557 Pelfrène, A., Douay, F., Richard, A., Roussel, H., Girondelot, B., 2013. Assessment of
558 potential health risk for inhabitants living near a former lead smelter. Part 2: site-
559 specific human health risk assessment of Cd and Pb contamination in kitchen
560 gardens. *Environ. Monit. Assess.* 185, 2999-3012. [https://doi.org/10.1007/s10661-](https://doi.org/10.1007/s10661-012-2767-x)
561 [012-2767-x](https://doi.org/10.1007/s10661-012-2767-x)

562 Pelfrène, A., Douay, F., 2018. Assessment of oral and lung bioaccessibility of Cd and Pb
563 from smelter-impacted dust. *Environ. Sci. Pollut. Res.* 25, 3718-3730.
564 <https://doi.org/10.1007/s11356-017-0760-1>

565 R Core Team, 2018. R: A language and environment for statistical computing. R Foundation
566 for statistical computing, Vienna.

567 Radwan, M.A., El-Gendy, K.S., Gad, A.F., 2020. Biomarker responses in terrestrial
568 gastropods exposed to pollutants: A comprehensive review. *Chemosphere* 257,
569 127218. <https://doi.org/10.1016/j.chemosphere.2020.127218>

570 Rahman, M.S., Reichelt-Brushet, A.J., Clark, M.W., Far zana, T., Yee, L.H., 2017. Arsenic
571 bio-accessibility and bioaccumulation in aged pesticide contaminated soils: A
572 multiline investigation to understand environmental risk. *Sci. Total Environ.* 581-
573 582, 782-793. <https://doi.org/10.1007/s10661-012-2767-x>

574 Ruby, M.V., Davis, A., Schoof, R., Eberle, S., Sellstone, C.M., 1996. Estimation of lead and
575 arsenic bioavailability using a physiologically based extraction test. *Environ. Sci.*
576 *Technol.* 30, 422-430. <https://doi.org/10.1021/es950057z>

577 Stanek, E.J., Calabrese, E.J., Zorn, M., 2001. Soil ingestion distributions for Monte Carlo
578 risk assessment in children. *Hum. Ecol. Risk Assess. Int. J.* 7, 357-368.
579 <https://doi.org/10.1080/20018091094402>

580 Vasseur, P., Bonnard, M., 2014. Ecogenotoxicology in earthworms: A review. *Curr. Zool.*
581 60, 255-272. <https://doi.org/10.1093/czoolo/60.2.255>

582 Wragg, J., Cave, M., Basta, N., Brandon, E., Casteel, S., Denys, S., Gron, C., Oomen, A.,
583 Reimer, K., Tack, K., Van de Wiele, T., 2011. An inter-laboratory trial of the unified
584 BARGE bioaccessibility method for arsenic, cadmium and lead in soil. *Sci. Total*
585 *Environ.* 409, 4016-4030. <https://doi.org/10.1016/j.scitotenv.2011.05.019>

586 Ziegler, E.E., Edwards, B.B., Jensen, R.L., Mahaffey, K.R., Fomon, S.J., 1978. Absorption
587 and retention of lead by infants. *Pediatric Res.* 12, 29-34.
588 <https://doi.org/10.1203/00006450-197801000-00008>

Figure 1: Relationships between the ratio of [metal(loid)s]soils/SGV(metal(loid)s) ($\text{mg kg}^{-1} \text{ dw}$) and the ratio of [metal(loid)s]snails/TGV(metal(loid)s) ($\text{mg kg}^{-1} \text{ dw}$) (SGV and TGV were summarized in Louzon et al. (2020b); dotted line represents the identity line $y=x$).




Figure 2: Summary of the HHRA with ISE and ISE adj. and the ERA with ERITME for each of the soils in the study (ISE were adjusted with the oral bioaccessible fraction when HQ or CR were in the area of uncertain risk using the classification in the **table 3**. Yellow stars indicate soils where human health risk is evidenced or is uncertain and the level of environmental risk is no risked. The result of the risk ranking for ISE in this figure is based on the most negative interpretation of the risk for HQ and CR for all of the metal(loid)s studied. ISE, ISE adj. and ERA with ERITME are presented by colored bar plots in blue, orange and green, respectively).




Figure 3: Development of a management plan combining HHRA and ERA (according to results of this study and adapted from the graphical abstract in Louzon et al. (2020a)).




Table 1: Concentrations of As, Cd and Pb in soils and their bioaccessible fractions (gastric and gastro-intestinal) measured using the UBM test and the concentrations bioaccumulated over 28 days in the viscera of snails (mg kg⁻¹ dw). Total concentrations of metal(loid)s in soils and their measured bioaccessible fractions are noted as follows for each soil: C_{soil}(G phase|GI phase).

Soil	[As]soil	[As]snail	[Cd]soil	[Cd]snail	[Pb]soil	[Pb]snail
O1	13.4 (38 % 38 %)	0.298	1.32 (95 % 31 %)	5.02	430 (81 % 26 %)	84.6
O2	11.5 (32 % 29 %)	0.298	1.91 (73 % 21 %)	5.63	320 (68 % 2 %)	43.3
O3	228 (100 % 49 %)	21.9	257 (100 % 55 %)	333	6256 (100 % 39 %)	856
O4	74.4 (57 % 38 %)	1.29	48.0 (86 % 25 %)	60.0	5224 (76 % 7 %)	652
O5	108 (60 % 49 %)	2.25	85.7 (89 % 34 %)	73.6	8971 (100 % 20 %)	731
O6	139 (80 % 64 %)	19.2	109 (96 % 45 %)	307	2184 (93 % 5 %)	783
O7	12.8 (35 % 25 %)	0.298	4.20 (92 % 25 %)	18.6	385 (88 % 2 %)	267
O8	5.64 (21 % 21 %)	0.298	0.64 (85 % 39 %)	5.33	30.7 (66 % 10 %)	5.21
O9	23.3 (45 % 42 %)	3.93	7.34 (84 % 37 %)	32.0	534 (96 % 2 %)	208
O10	48.6 (73 % 52 %)	2.31	27.8 (100 % 40 %)	71.9	2233 (100 % 44 %)	503
O11	27.9 (50 % 49 %)	3.78	18.2 (85 % 36 %)	52.5	1260 (88 % 10 %)	431
O12	7.74 (14 % 16 %)	0.298	1.42 (73 % 34 %)	12.5	222 (89 % 45 %)	171
O13	12.6 (27 % 29 %)	0.262	6.50 (74 % 17 %)	28.3	484 (94 % 9 %)	162
O14	23.4 (38 % 65 %)	2.02	15.4 (87 % 57 %)	46.7	719 (85 % 2 %)	271
O15	13.5 (38 % 27 %)	0.482	1.30 (78 % 63 %)	6.04	177 (64 % 0 %)	21.0
O16	5.52 (15 % 16 %)	0.098	0.64 (85 % 32 %)	4.80	17.8 (60 % 0 %)	4.12
O17	17.8 (11 % 8 %)	0.098	0.200 (77 % 60 %)	2.56	33.7 (41 % 2 %)	1.42
O18	74.4 (8 % 8 %)	1.39	0.520 (66 % 26 %)	3.20	46.6 (33 % 1 %)	2.56
O19	10.1 (13 % 14 %)	0.098	4.49 (71 % 28 %)	12.1	117 (72 % 2 %)	37.4
O20	26.0 (23 % 23 %)	1.98	0.370 (71 % 34 %)	4.59	150 (70 % 2 %)	38.9
O21	11.3 (49 % 44 %)	2.23	2.32 (83 % 31 %)	11.5	288 (84 % 1 %)	83.7
O22	3.28 (16 % 18 %)	0.098	0.235 (87 % 57 %)	4.24	19.7 (72 % 18 %)	12.1
O23	57.0 (30 % 25 %)	1.48	0.102 (82 % 35 %)	2.53	660 (17 % 3 %)	39.1
O24	12.0 (34 % 24 %)	0.829	3.57 (96 % 29 %)	12.4	303 (86 % 2 %)	35.6
O26	13.3 (43 % 28 %)	0.098	0.450 (100 % 61 %)	3.20	312 (88 % 3 %)	58.0
O27	28.5 (24 % 16 %)	0.098	7.43 (73 % 38 %)	8.81	1882 (68 % 7 %)	137
O28	25.8 (29 % 23 %)	0.098	2.60 (84 % 28 %)	5.52	467 (63 % 9 %)	25.5
O29	36.1 (24 % 16 %)	0.379	27.2 (49 % 34 %)	28.3	5258 (99 % 32 %)	820
O30	28.3 (27 % 22 %)	1.16	12.8 (81 % 41 %)	20.4	3241 (97 % 24 %)	730
Min.	3.28 (8 % 8 %)	0.098	0.102 (49 % 17 %)	2.53	17.8 (17 % 0 %)	1.42
Med.	23.4 (32 % 25 %)	0.656	3.89 (84 % 34 %)	12.3	426 (84 % 5 %)	110
Max.	228 (100 % 65 %)	21.9	257 (100 % 63 %)	333	8971 (100 % 45 %)	856

Table 2: Toxicological reference values (TRV) for As, Cd and Pb (ingestion) in mg kg⁻¹ day⁻¹ (ANSES, 2020; INERIS, 2018) for no carcinogenic risks (with threshold) and carcinogenic risk (CR), toxicity point (TP) (ATSDR, 2015) and *ex situ* threshold guide values (TGV) for snails in mg kg⁻¹ (Louzon et al., 2020b)

Soil	TRV with threshold	Toxicological target	TRV for CR	<i>Ex situ</i> TGV	TP
As	0.00045	Skin	1.5	0.364	600
Cd	0.00036	Kidney	ND	5.60	400
Pb	0.00063	Kidney	0.0085	9.06	400

ND: not determined

Table 3: Interpretation of values of hazard quotient (HQ), cancer risk (CR) (MEEM, 2017) and risk coefficient (RC) (Louzon et al., submitted)

Risk	HQ	CR	RC
No	< 0.2	< 10^{-6}	< 2574
Uncertain	0.2-5	10^{-6} - 10^{-4}	2574-22720
Proven	> 5	> 10^{-4}	> 22720

Table 4: Assessment of human health risk with hazard quotient (HQ), cancer risk (CR) and interpretation the state of the environment (ISE) (when HQ or CR were in the area of uncertain risk, they were weighted (HQ adj. or CR adj.) by bioaccessible fraction. The result of the ISE is the most conservative level of risk for all the metal(loid)s studied according to the ISE grid of interpretation summarized in table 3).

Soil	As				Cd				Pb				ISE	
	HQ	CR	HQ adj.	CR adj.	HQ	CR	HQ adj.	CR adj.	HQ	CR	HQ adj.	CR adj.	risk	risk adj.
O1	0.17	9.76 x 10 ⁻⁶	no. adj.	3.71 x 10 ⁻⁶	0.02	no TRV	no adj.	no TRV	3.86	1.77 x 10 ⁻⁶	6.25	2.87 x 10 ⁻⁶	uncertainty	proven
O2	0.15	8.36 x 10 ⁻⁶	no. adj.	2.68 x 10 ⁻⁶	0.03	no TRV	no adj.	no TRV	2.87	1.32 x 10 ⁻⁶	3.90	1.80 x 10 ⁻⁶	uncertainty	uncertainty
O3	2.86	1.65 x 10 ⁻⁴	2.86	no adj.	4.04	no TRV	4.04	no TRV	56.09	2.57 x 10 ⁻⁵	no adj.	5.14 x 10 ⁻⁵	proven	proven
O4	0.93	5.40 x 10 ⁻⁵	0.53	3.08 x 10 ⁻⁵	0.75	no TRV	0.65	no TRV	46.82	2.15 x 10 ⁻⁵	no adj.	3.27 x 10 ⁻⁵	proven	proven
O5	1.36	7.85 x 10 ⁻⁵	0.82	4.71 x 10 ⁻⁵	1.35	no TRV	1.20	no TRV	80.49	3.69 x 10 ⁻⁵	no adj.	7.38 x 10 ⁻⁵	proven	proven
O6	1.75	1.01 x 10 ⁻⁴	1.40	no adj.	1.71	no TRV	1.64	no TRV	19.61	9.02 x 10 ⁻⁶	no adj.	1.68 x 10 ⁻⁵	proven	proven
O7	0.16	9.27 x 10 ⁻⁶	no. adj.	3.24 x 10 ⁻⁶	0.07	no TRV	no adj.	no TRV	3.45	1.59 x 10 ⁻⁶	6.07	2.80 x 10 ⁻⁶	uncertainty	proven
O8	0.07	4.10 x 10 ⁻⁶	no. adj.	8.61 x 10 ⁻⁷	0.01	no TRV	no adj.	no TRV	0.27	1.27 x 10 ⁻⁷	0.36	no adj.	uncertainty	uncertainty
O9	0.29	1.70 x 10 ⁻⁵	0.13	7.65 x 10 ⁻⁶	0.12	no TRV	no adj.	no TRV	4.79	2.20 x 10 ⁻⁶	9.20	4.22 x 10 ⁻⁶	uncertainty	proven
O10	0.61	3.53 x 10 ⁻⁵	0.45	2.58 x 10 ⁻⁵	0.44	no TRV	0.44	no TRV	20.02	9.18 x 10 ⁻⁶	no adj.	1.84 x 10 ⁻⁵	proven	proven
O11	0.35	2.03 x 10 ⁻⁵	0.18	1.02 x 10 ⁻⁵	0.29	no TRV	0.25	no TRV	11.33	5.19 x 10 ⁻⁶	no adj.	9.13 x 10 ⁻⁶	proven	proven
O12	0.10	5.63 x 10 ⁻⁶	no. adj.	9.01 x 10 ⁻⁷	0.02	no TRV	no adj.	no TRV	1.99	9.10 x 10 ⁻⁷	3.54	no adj.	uncertainty	uncertainty
O13	0.16	9.18 x 10 ⁻⁶	no. adj.	2.66 x 10 ⁻⁶	0.10	no TRV	no adj.	no TRV	4.34	1.99 x 10 ⁻⁶	8.16	3.74 x 10 ⁻⁶	uncertainty	proven
O14	0.30	1.70 x 10 ⁻⁵	0.20	1.11 x 10 ⁻⁵	0.24	no TRV	0.21	no TRV	6.45	2.96 x 10 ⁻⁶	no adj.	5.03 x 10 ⁻⁶	proven	proven
O15	0.17	9.84 x 10 ⁻⁶	no. adj.	3.74 x 10 ⁻⁶	0.02	no TRV	no adj.	no TRV	1.59	7.29 x 10 ⁻⁷	2.04	no adj.	uncertainty	uncertainty
O16	0.07	4.01 x 10 ⁻⁶	no. adj.	6.42 x 10 ⁻⁷	0.01	no TRV	no adj.	no TRV	0.16	7.33 x 10 ⁻⁸	no adj.	no adj.	uncertainty	no risk
O17	0.22	1.29 x 10 ⁻⁵	0.02	1.42 x 10 ⁻⁶	0.00	no TRV	no adj.	no TRV	0.31	1.39 x 10 ⁻⁷	0.25	no adj.	uncertainty	uncertainty
O18	0.93	5.40 x 10 ⁻⁵	0.07	4.32 x 10 ⁻⁶	0.01	no TRV	no adj.	no TRV	0.42	1.92 x 10 ⁻⁷	0.28	no adj.	uncertainty	uncertainty
O19	0.12	7.34 x 10 ⁻⁶	no. adj.	1.03 x 10 ⁻⁶	0.07	no TRV	no adj.	no TRV	1.05	4.81 x 10 ⁻⁷	1.51	no adj.	uncertainty	uncertainty
O20	0.33	1.89 x 10 ⁻⁵	0.08	4.35 x 10 ⁻⁶	0.01	no TRV	no adj.	no TRV	1.35	6.18 x 10 ⁻⁷	1.89	no adj.	uncertainty	uncertainty
O21	0.14	8.21 x 10 ⁻⁶	no. adj.	4.02 x 10 ⁻⁶	0.03	no TRV	no adj.	no TRV	2.58	1.18 x 10 ⁻⁶	4.33	1.98 x 10 ⁻⁶	uncertainty	uncertainty
O22	0.04	2.38 x 10 ⁻⁶	no. adj.	4.28 x 10 ⁻⁷	0.00	no TRV	no adj.	no TRV	0.17	8.11 x 10 ⁻⁸	no adj.	no adj.	uncertainty	no risk
O23	0.72	4.14 x 10 ⁻⁵	0.22	1.24 x 10 ⁻⁵	0.00	no TRV	no adj.	no TRV	5.92	2.71 x 10 ⁻⁶	no adj.	9.21 x 10 ⁻⁷	proven	proven
O24	0.15	8.69 x 10 ⁻⁶	no. adj.	2.95 x 10 ⁻⁶	0.06	no TRV	no adj.	no TRV	2.72	1.25 x 10 ⁻⁶	4.68	2.15 x 10 ⁻⁶	uncertainty	uncertainty
O26	0.17	9.68 x 10 ⁻⁶	no. adj.	4.16 x 10 ⁻⁶	0.01	no TRV	no adj.	no TRV	2.80	1.28 x 10 ⁻⁶	4.93	2.25 x 10 ⁻⁶	uncertainty	uncertainty
O27	0.36	2.07 x 10 ⁻⁵	0.09	4.97 x 10 ⁻⁶	0.12	no TRV	no adj.	no TRV	16.88	7.75 x 10 ⁻⁶	no adj.	1.05 x 10 ⁻⁵	proven	proven
O28	0.32	1.88 x 10 ⁻⁵	0.09	5.45 x 10 ⁻⁶	0.04	no TRV	no adj.	no TRV	4.19	1.92 x 10 ⁻⁶	5.28	2.42 x 10 ⁻⁶	uncertainty	proven
O29	0.46	2.62 x 10 ⁻⁵	0.11	6.29 x 10 ⁻⁶	0.43	no TRV	0.21	no TRV	47.15	2.17 x 10 ⁻⁵	no adj.	4.30 x 10 ⁻⁵	proven	proven
O30	0.36	2.06 x 10 ⁻⁵	0.10	5.56 x 10 ⁻⁶	0.20	no TRV	0.16	no TRV	29.04	1.33 x 10 ⁻⁵	no adj.	2.58 x 10 ⁻⁵	proven	proven

The level of risk is interpreted according to three categories: no risk, uncertain risk and proven risk, labelled as green, orange and red, respectively.

Table 5: Assessment of the environmental risk with risk coefficient (RC) and evaluation of the risk of transferred metallic elements (ERITME) index.

Soil	RC As	RC Cd	RC Pb	ERITME
O1	0	0	3336	3336
O2	0	4	1512	1516
O3	35520	23400	37400	96320
O4	1524	3880	28400	33804
O5	3108	4840	31880	39828
O6	31080	21520	34160	86760
O7	0	928	11400	12328
O8	0	0	0	0
O9	5880	1884	8800	16564
O10	3210	4720	21800	29730
O11	5640	3352	18640	27632
O12	0	492	7160	7652
O13	0	1620	6760	8380
O14	2730	2936	11560	17226
O15	192	32	528	752
O16	0	0	0	0
O17	0	0	0	0
O18	1692	0	0	1692
O19	0	464	1252	1716
O20	2664	0	1316	3980
O21	3078	420	3296	6794
O22	0	0	136	136
O23	1842	0	1328	3170
O24	768	484	1172	2424
O26	0	0	2160	2160
O27	0	228	5640	5868
O28	0	0	724	724
O29	24	1620	35800	37444
O30	1314	1056	31840	34210

The level of risk is interpreted according to three categories: no risk, uncertain risk and proven risk, labelled as green, orange and red, respectively.

Table 6: Spearman correlation coefficients (ρ) for each metal(loids) and each risk index (adjusted or not adjusted with bioaccessible fraction if indicators were in the area of uncertainty risk. Significance of ρ was assessed with p-values and signified with stars: “*”, “**” and “***” for < 0.05, < 0.01 and < 0.001, respectively).

	RC As	RC Cd	RC Pb
HQ	0.67***	0.93***	0.90***
HQ adj.	0.57**	0.93***	0.92***
CR	0.67***	ND	0.90***
CR adj.	0.80***	ND	0.93***

HQ: hazard quotient, HQ adj.: hazard quotient adjusted, CR: carcinogenic risk, CR adj.: carcinogenic risk adjusted, RC: risk coefficient, ND = not determined because no TRV for CR of Cd exist.

Graphical abstract

