



Axion search with BabyIAXO in view of IAXO

A. Abeln, K. Altenmüller, S. Arguedas Cuendis, E. Armengaud, D. Attié, S. Aune, S. Basso, L. Bergé, B. Biasuzzi, P.T.C. Borges de Sousa, et al.

► To cite this version:

A. Abeln, K. Altenmüller, S. Arguedas Cuendis, E. Armengaud, D. Attié, et al.. Axion search with BabyIAXO in view of IAXO. 40th International Conference on High Energy Physics (ICHEP 2020), Jul 2020, Prague (virtual), Czech Republic. pp.631. hal-03113465

HAL Id: hal-03113465

<https://hal.science/hal-03113465>

Submitted on 18 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Axion search with BabyIAXO in view of IAXO

A. Abeln,¹ K. Altenmüller,² S. Arguedas Cuendis,³ E. Armengaud,⁴ D. Attié,⁴
 S. Aune,⁴ S. Basso,⁵ L. Bergé,⁶ B. Biasuzzi,⁴ P. T. C. Borges De Sousa,³ P. Brun,⁴
 N. Bykovskiy,³ D. Calvet,⁴ J. M. Carmona,² J. F. Castel,² S. Cebrián,² V. Chernov,^{7,8}
 F. E. Christensen,⁹ M.M. Civitani,⁵ C. Cogollos,^{10,11} T. Dafni,² A. Derbin,¹²
 K. Desch,¹³ D. Díez,² M. Dinter,²¹ B. Döbrich,³ I. Drachnev,¹² A. Dudarev,³
 L. Dumoulin,⁶ D. D. M. Ferreira,⁹ E. Ferrer-Ribas,⁴ I. Fleck,¹⁴ J. Galan,^{2,*}
 D. Gascón,^{10,11} L. Gastaldo,¹ M. Giannotti,¹⁵ Y. Giomataris,⁴ A. Giuliani,⁶
 S. Gninenko,⁸ J. Golm,^{3,16} N. Golubev,⁸ L. Hagge,²¹ J. Hahn,¹⁴ C. J. Hailey,¹⁷
 D. Hengstler,¹ P. L. Henriksen,⁹ R. Iglesias-Marzoa,¹⁸ F. J. Iguaz-Gutierrez,¹⁹
 I. G. Irastorza,² C. Iñiguez,¹⁸ K. Jakovčić,²⁰ J. Kaminski,¹³ B. Kanoute,¹⁹
 S. Karstensen,²¹ L. Kravchuk,⁸ B. Lakić,²⁰ T. Lasserre,⁴ P. Laurent,⁴ O. Limousin,⁴
 A. Lindner,²¹ M. Loidl,²² I. Loms kaya,¹² G. López-Alegre,¹⁸ B. Lubsandorzhiev,⁸
 K. Ludwig,²¹ G. Luzón,² C. Malbrunot,³ C. Margalejo,² A. Marin-Franch,¹⁸
 S. Marnieros,⁶ F. Marutsky,²¹ J. Mauricio,^{10,11} Y. Menesguen,²² M. Mentink,³
 F. Mescia,^{10,11} J. Miralda-Escudé,^{10,23} H. Mirallas,² J. P. Mols,⁴ V. Muratova,¹²
 X. F. Navick,⁴ C. Nones,⁴ A. Notari,^{10,11} A. Nozik,^{7,8} L. Obis,² C. Oriol,⁶ F. Orsini,¹⁹
 A. Ortiz de Solórzano,² S. Oster,²¹ H. P. Pais Da Silva,³ V. Pantuev,⁸
 T. Papaevangelou,⁴ G. Pareschi,⁵ K. Perez,²⁴ O. Pérez,² E. Picatoste,^{10,11}
 M. J. Pivovarov,^{26,25} D. V. Poda,⁶ J. Redondo,² A. Ringwald,²¹ M. Rodrigues,²²
 F. Rueda-Teruel,¹⁸ S. Rueda-Teruel,¹⁸ E. Ruiz-Choliz,²⁷ J. Ruz,²⁶ E. O. Saemann,²¹
 J. Salvado,^{10,11} T. Schiffer,¹³ S. Schmidt,¹³ U. Schneekloth,²¹ M. Schott,²⁷ L. Segui,⁴
 F. Tavecchio,⁵ H. H. J. ten Kate,³ I. Tkachev,⁸ S. Troitsky,⁸ D. Unger,¹ E. Unzhakov,¹²
 N. Ushakov,⁸ J. K. Vogel,²⁶ D. Voronin,⁸ A. Weltman,²⁸ U. Werthenbach,¹⁴
 W. Wuensch³ and A. Yanes-Díaz¹⁸

¹Heidelberg University, Kirchhoff Institute for Physics

²Center for Astroparticles and High Energy Physics (CAPA), Universidad de Zaragoza, 50009 Zaragoza, Spain

³CERN - European Organization for Nuclear Research, Geneva, Switzerland

⁴IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France

⁵INAF, Italian National Institute for Astrophysics, Osservatorio Astronomico di Brera, Milano/Merate, Italy

⁶Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France

*Speaker

- ⁷*Moscow Institute of Physics and Technology, Moscow, Russia*
- ⁸*Institute for Nuclear Research of the Russian Academy of Sciences, 60th October Anniversary Prospect 7A, 117312, Moscow, Russia*
- ⁹*Technical University of Denmark, DTU Space Kgs. Lyngby, Denmark*
- ¹⁰*Institut de Ciències del Cosmos, Universitat de Barcelona, Barcelona, Spain*
- ¹¹*Departament de Física Quàntica i Astrofísica, Universitat de Barcelona, Barcelona, Spain*
- ¹²*Petersburg Nuclear Physics Institute - NRC Kurchatov Institute, Gatchina, 188300, Russia*
- ¹³*Physikalisches Institut der Universität Bonn, Nussallee 12, 53115 Bonn, Germany*
- ¹⁴*Center for Particle Physics Siegen, Siegen University, Siegen, Germany*
- ¹⁵*Physical Sciences, Barry University, 11300 NE 2nd Ave., Miami Shores, FL 33161, USA*
- ¹⁶*Institute for Optics and Quantum Electronics, Friedrich Schiller University Jena, Jena, Germany*
- ¹⁷*Columbia Astrophysics Laboratory, New York, U.S.A.*
- ¹⁸*Centro de Estudios de Física del Cosmos de Aragón, Plaza San Juan, Teruel, Spain*
- ¹⁹*Synchrotron SOLEIL, 91192 Gif-sur-Yvette, France*
- ²⁰*Rudjer Bošković Institute, Bijenička cesta 54, 10000 Zagreb, Croatia*
- ²¹*Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany*
- ²²*CEA, LIST, Laboratoire National Henri Becquerel (LNE-LNHB), F-91191 Gif-sur-Yvette, France*
- ²³*ICREA, Barcelona, Spain*
- ²⁴*Massachusetts Institute of Technology, 77 Massachusetts Ave., Cambridge, MA 02139, USA*
- ²⁵*SLAC National Accelerator Laboratory, Menlo Park, CA, U.S.A.*
- ²⁶*Lawrence Livermore National Laboratory, Livermore, CA, U.S.A.*
- ²⁷*Johannes Gutenberg University, Mainz, Germany*
- ²⁸*High Energy Physics, Cosmology & Astrophysics Theory (HEPCAT) group, University of Cape Town, Private Bag, 7700 Rondebosch, South Africa*
- E-mail: javier.galan@unizar.es*

Axions are a natural consequence of the Peccei-Quinn mechanism, the most compelling solution to the strong-CP problem. Similar axion-like particles (ALPs) also appear in a number of possible extensions of the Standard Model, notably in string theories. Both axions and ALPs are very well motivated candidates for Dark Matter, and in addition, they would be copiously produced at the sun's core. A relevant effort during the last decade has been the CAST experiment at CERN, the most sensitive axion helioscope to-date. The International Axion Observatory (IAXO) is a large-scale 4th generation helioscope. As its primary physics goal, IAXO will look for solar axions or ALPs with a signal to background ratio of about 5 orders of magnitude higher than CAST. Recently the IAXO collaboration has proposed an intermediate experimental stage, BabyIAXO, conceived to test all IAXO subsystems (magnet, optics, detectors and sun-tracking systems) at a relevant scale for the final system and thus serve as pathfinder for IAXO but at the same time as a fully-fledged helioscope with record and relevant physics reach in itself with potential for discovery. BabyIAXO was endorsed by the Physics Review committee of DESY last May 2019. Here we will review the status and prospects of BabyIAXO and its potential to probe the most physics motivated regions of the axion & ALPs parameter space.

*40th International Conference on High Energy physics - ICHEP2020
July 28 - August 6, 2020
Prague, Czech Republic (virtual meeting)*

1. Introduction

The axion is a hypothetical particle that emerged as a natural solution to the *strong CP problem*. The problem arises from the fact that the description of strong interactions given by the quantum chromodynamics (QCD) theory does not forbid such violation. Furthermore, since electroweak interactions violate CP it is difficult to understand why strong interactions do not. The Peccei-Quinn mechanism to restore CP conservation at the strong sector [1, 2] developed into an interesting outcome, the axion, a new boson arising naturally in the theory [3, 4]. The phenomenological properties of the axion are described through a unique parameter, the scale factor f_a , which must be much greater than the electroweak scale in order to circumvent the constraints imposed by accelerator based searches. Both, the couplings of axions to ordinary matter, and the axion mass, m_a , are inversely proportional to f_a . The parameter space of axions is further constrained by cosmological and astrophysical arguments, such as the evolution of stars or the dark matter content of the universe. As a consequence of its very weak couplings the axion results to be a long-lived particle, and therefore, under certain conditions it is considered to be a potential *dark matter* candidate.

Two main theoretical models are considered as a reference for axion searches, *hadronic axions* or *KSVZ axions* [5, 6], and *DSFZ* or *GUT axions* [7, 8], which do not couple to hadrons at tree level. Furthermore, a plethora of axion-like particles (ALPs) emerge in different extensions to the SM. Their similarity with standard QCD axions makes that IAXO will be able to exploit its physics program to explore ALPs hinted regions other than the theoretical QCD models [9, 10]

2. BabyIAXO and IAXO Physics potential

IAXO is a 4th generation helioscope that will push the sensitivity to solar axions to unprecedented levels, surpassing those of the most sensitive axion helioscope up to date, the CERN Axion Solar Telescope (CAST) [11]. The final IAXO configuration will allow to explore a vast region of the axion parameter space, covering theoretically QCD favored regions, stellar cooling hints as the horizontal branch stars [12] or white dwarf (WD) cooling anomalies [13], and the universe gamma-rays transparency anomaly reported by HESS [14] and MAGIC [15], between other hints. Figure 1 shows those hints and the sensitivities reached by BabyIAXO and IAXO. BabyIAXO, being a prototype of the full IAXO, will cover itself an interesting new unexplored region in the parameter space, reaching $g_{a\gamma} \sim 1.5 \times 10^{-11} \text{ GeV}^{-1}$ for masses up to 0.25 eV, including the favored QCD axion region. See reference [16] for a full review.

3. Design and construction of BabyIAXO

BabyIAXO will be hosted by the DESY research facility after the recent endorsement by the DESY Physics Review committee in May 2019. Most of the BabyIAXO systems (cryogenics, magnet, optics, detector systems) are in an advanced mature state and moving towards the construction phase. BabyIAXO will benefit from the existence of a Medium-Sized Telescope (MST) [17] instrument, commissioned by the Cherenkov Telescope Array (CTA) group at DESY, to serve as a dedicated driving system for BabyIAXO. The most critical element of the system is the magnet system which is being designed and commissioned at CERN [18].




Figure 1: The prospects of the different IAXO setups and its sensitivity to $g_{a\gamma}$ as a function of the axion mass, m_a , including different hints and models described in the text [9].

The figure of merit (FOM) of the experiment can be factorized on different experimental systems [16], such as the magnet, $f_M = B^2 L^2 A$, the combination of optic and detectors systems, $f_{DO} = \frac{\epsilon_d \epsilon_o}{\sqrt{(ba)}}$, and tracking exposure, $f_T = \sqrt{(\epsilon_t t)}$. The total FOM, $f_{FOM} = f_M f_{DO} f_T$, will be roughly proportional to the expected number of photons converted from axions that can be detected, and that will be 5 orders of magnitude higher than present measurements for the IAXO+ prospects, being IAXO+ the ultimate reach of the IAXO R&D program where technology is pushed to the limits. Table 1 shows a summary of the experimental setup parameters that apply to each sensitivity plot shown previously in Figure 1.

Parameter [units]	BabyIAXO	IAXO	IAXO+
B [T]	~ 2	~ 2.5	~ 3.5
L [m]	10	22	22
A [m ²]	0.77	2.3	3.9
f_M [T ² m ⁴]	~ 230	~ 6000	~ 64000
b [cm ⁻² keV ⁻¹ s ⁻¹]	10^{-7}	10^{-8}	10^{-9}
ϵ_d	0.7	0.8	0.8
ϵ_o	0.35	0.7	0.7
a [cm ²]	2×0.3	8×0.15	8×0.15
f_{DO} [keV ^{1/2} s ^{1/2}]	1000	5112	16166
ϵ_t	0.5	0.5	0.5
t [year]	1.5	3	5
f_T [year ^{1/2}]	0.87	1.22	1.58

Table 1: A comparison of the relevant experimental parameters from BabyIAXO to IAXO+.

An important upgrade for IAXO is the magnet contribution to the f_{FOM} , that provides to the magnetic field configuration with a much bigger aperture, and additional bores, with respect to its predecessor, CAST. After the magnetic field, and driving system (which provides also a longer tracking exposure, ~ 18 hours-per-day) have been fixed, the next system to enhance axion searches is the x-ray detector system which is under active design, as shown in Figure 2.




Figure 2: The undergoing CAD design for the shielding of detectors including an active VETO system .

In that sense, extensive studies are taking place in order to better understand the nature of the background to achieve the low levels required. The optimization of the active VETO tagging system to different sources, as cosmic neutrons, gammas and muons (see Figure 3) is crucial for the success of IAXO, in order to achieve the background level of $10^{-8} \text{keV}^{-1} \text{cm}^{-2} \text{s}^{-1}$, and ultimately a level of 10^{-9} leading to the detection of just ~ 0.5 background counts if the prospects for IAXO+ scenario are achieved. BabyIAXO will be no doubt an excellent center of operations to demonstrate and optimize all the final systems that will be finally used to construct IAXO.




Figure 3: *Geant4* Monte Carlo simulations of the detector, shielding VETO systems using RESTSoft [19]. From left to right, a simulated muon, neutron and high-energy gamma. Green lines describe the trajectories of gammas and black lines of neutrons.

References

- [1] R. Peccei and H.R. Quinn, *CP conservation in the presence of instantons*, *Phys. Rev. Lett.* **38** (1977) 1440.

- [2] R. Peccei and H.R. Quinn, *Constraints imposed by CP conservation in the presence of instantons*, *Phys. Rev.* **D16** (1977) 1791.
- [3] S. Weinberg, *A New Light Boson?*, *Phys. Rev. Lett.* **40** (1978) 223.
- [4] F. Wilczek, *Problem of Strong P and T Invariance in the Presence of Instantons*, *Phys. Rev. Lett.* **40** (1978) 279.
- [5] J.E. Kim, *Weak Interaction Singlet and Strong CP Invariance*, *Phys. Rev. Lett.* **43** (1979) 103.
- [6] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Can Confinement Ensure Natural CP Invariance of Strong Interactions?*, *Nucl. Phys.* **B166** (1980) 493.
- [7] M. Dine, W. Fischler and M. Srednicki, *A Simple Solution to the Strong CP Problem with a Harmless Axion*, *Phys. Lett.* **B104** (1981) 199.
- [8] A.R. Zhitnitsky, *On Possible Suppression of the Axion Hadron Interactions. (In Russian)*, *Sov. J. Nucl. Phys.* **31** (1980) 260.
- [9] I.G. Irastorza and J. Redondo, *New experimental approaches in the search for axion-like particles*, [1801.08127](#).
- [10] P.W. Graham, I.G. Irastorza, S.K. Lamoreaux, A. Lindner and K.A. van Bibber, *Experimental Searches for the Axion and Axion-Like Particles*, *Ann. Rev. Nucl. Part. Sci.* **65** (2015) 485 [[1602.00039](#)].
- [11] CAST collaboration, *New CAST Limit on the Axion-Photon Interaction*, *Nature Phys.* **13** (2017) 584 [[1705.02290](#)].
- [12] O.e.a. Straniero, *The RGB tip of galactic globular clusters and the revision of the bound of the axion-electron coupling*, [2010.03833](#).
- [13] B. Melendez, M.M. Bertolami and L. Althaus, *Revisiting the Impact of Axions in the Cooling of White Dwarfs*, [1210.0263](#).
- [14] HESS collaboration, *A Low level of extragalactic background light as revealed by gamma-rays from blazars*, *Nature* **440** (2006) 1018 [[astro-ph/0508073](#)].
- [15] MAGIC collaboration, *Very-High-Energy Gamma Rays from a Distant Quasar: How Transparent Is the Universe?*, *Science* **320** (2008) 1752 [[0807.2822](#)].
- [16] I. Collaboration, *Physics potential of the international axion observatory (IAXO)*, *Journal of Cosmology and Astroparticle Physics* **2019** (2019) 047.
- [17] CTA CONSORTIUM, MST TEAM collaboration, *Status of the Medium-Sized Telescope for the Cherenkov Telescope Array*, *PoS ICRC2015* (2016) 959 [[1509.01361](#)].
- [18] BABYIAXO collaboration, *Conceptual Design of BabyIAXO, the intermediate stage towards the International Axion Observatory*, [2010.12076](#).
- [19] J. Galán, “REST v2.0 : A data analysis and simulation framework for micro-patterned readout detectors.” talk given at the 8th Symposium on Large TPCs for Rare Event detection.