

Getting to understand more and more a (simple?) model of stochastic gene expression

Romain Yvinec

► To cite this version:


Romain Yvinec. Getting to understand more and more a (simple?) model of stochastic gene expression. 7th International Congress on Industrial and Applied Mathematics (ICIAM 2011), Jul 2011, Vancouver, Canada. hal-03114889


HAL Id: hal-03114889 https://hal.science/hal-03114889


Submitted on 7 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Getting to understand more and more a (simple?) model of stochastic gene expression Self-Regulation Nonlinear Regulation **Two different sources of noise** White noise **Yvinec Romain (ICJ)** Using Sample paths simulation he presence of noise can drastically alter regions of bistability The reduced 1-The full 3-dimensional dimensional stochastic deterministic model *model* has been analytically BURSTING is well known. characterized when either *Mono/Bistable behaviour* DIFFUSION one source of noise is *•in the positive regulation;* Monostable/Limit cycle in DNA present: *-Uni/Bimodal asymptotic* the negative regulation. 60 $\blacktriangleright 0$ Inducible or Repressible density in the positive **Questions that remains: Questions that remains:** $\sim \sim \sim$ regulation; **mRNA** →Transient Behaviour? Adiabatic elimination when noise Noise-induced bi-stability →Difference between the *-Unimodal asymptotic* is present? Regulation density in the negative f(E Oscillation property when noise is two source of noise? regulation. present in the negative regulation? Intermediate Time-dependent densities are BURSTING different for the two sources of Effector noise. The bursting model presents higher variability. **S** Protein bf(X)The stochastic limit cycle may be identified through a








CONCLUSION:


The two sources of noise have distinct transient properties as well as different convergence rates toward equilibrium; Memory and switch property also differs;
Noisy and bistable regime may occur transiently;
Key scaling parameters has been identified;
Oscillations and limit cycle property is being characterized.

Interplay between the two source of noise is still mysterious;

The bursting noise gives rise to higher or slower **autocorrelation time** depending on the respective value of the mean production rate and the degradation rate.




The diffusion model **switch** more an faster, specifically due to a nonrefractory time in the bursting model while going from a high to a low valu (the bursting noise is asymmetric)



σ=1.41 σ=2

When both noise are present, the strength of the noise governs mainly the **convergence rate** of the **mean**. For the **variance**, we can observe non-monotonic behaviour and more complex dependencies of the two source of noise.

Same for the **auto-correlation time**



Interpretation in terms of biology: possible interpretation as Intrinsic and Extrinsic noise?
 Adiabatic reduction not fully understood. Will some properties such bistable regime be transmitted?
 Analytical results....

<u>References:</u>

Molecular distributions in gene regulatory dynamics. Mackey MC, Tyran-Kamińska M, Yvinec R. J Theor Biol. 2011 274(1):84-96 On the Poincaré–Hill cycle map of rotational random walk: locating the stochastic limit cycle in a reversible Schnakenberg model, Vellela M. and Qian H., Proc. R. Soc. 2010 vol. 466 no. 2115 771-78