



HAL
open science

Continuity of the dual Haar measure

Jean N Renault

► **To cite this version:**

| Jean N Renault. Continuity of the dual Haar measure. 2021, 10.5802/crmath.183 . hal-03116966

HAL Id: hal-03116966

<https://hal.science/hal-03116966>

Submitted on 20 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONTINUITY OF THE DUAL HAAR MEASURE

JEAN N. RENAULT

ABSTRACT. Given a locally compact group bundle, we show that the system of the Plancherel weights of their C^* -algebras is lower semi-continuous. As a corollary, we obtain that the dual Haar system of a continuous Haar system of a locally compact abelian group bundle is also continuous.

Let G be a locally compact abelian group with Haar measure λ . The dual Haar measure $\hat{\lambda}$ on the dual group \hat{G} is the Haar measure on \hat{G} which makes the Fourier transform defined by

$$\mathcal{F}(f)(\chi) = \int f(\gamma) \overline{\chi(\gamma)} d\lambda(\gamma)$$

for f in the space $C_c(G)$ of complex-valued continuous functions with compact support on G , an isometry from $L^2(G, \lambda)$ to $L^2(\hat{G}, \hat{\lambda})$. Suppose now that $p : G \rightarrow X$ is a locally compact abelian group bundle. Here, we mean that G and X are locally compact Hausdorff spaces, p is a continuous surjection and the fibres $G_x = p^{-1}(x)$ are abelian groups. We also require that the multiplication, as a map from $G \times_X G$ to G , and the inverse, as a map from G to G are continuous. A Haar system for G is a family of measures $(\lambda^x)_{x \in X}$, where for all $x \in X$, λ^x is a Haar measure of G_x ; it is said to be continuous if for all $f \in C_c(G)$, the function $x \mapsto \int f d\lambda^x$ is continuous. Then we can form the dual group bundle $\hat{p} : \hat{G} \rightarrow X$. As a set, \hat{G} is the disjoint union of the dual groups \hat{G}_x . Moreover, it can be identified with the spectrum of the commutative C^* -algebra $C^*(G)$, where G is viewed as a locally compact groupoid with Haar system $(\lambda^x)_{x \in X}$. Hence it is endowed with a locally compact topology. It can be checked ([12, Corollary 3.4]) that $\hat{p} : \hat{G} \rightarrow X$ is indeed a locally compact abelian group bundle in the above sense. Of course, one expects that the dual Haar system $(\hat{\lambda}^x)_{x \in X}$ is continuous. This is stated as Proposition 3.6 of [12]. However it was recently pointed to the authors by Henrik Kreidler that their proof is defective. This note corrects this and gives a more general result, based on the fact that the dual Haar measure $\hat{\lambda}$ on the dual group \hat{G} of a locally compact abelian group G , viewed as its Haar weight and defined as the canonical weight of $L^\infty(\hat{G})$ acting on $L^2(\hat{G}, \hat{\lambda})$, corresponds under the Fourier transform to the Plancherel weight of G , defined as the canonical weight of the von Neumann algebra $VN(G)$ of G acting on $L^2(G, \lambda)$. Therefore one can consider a locally compact group bundle $p : G \rightarrow X$ where the fibres G_x are no longer abelian. Our main result is Proposition 2.3 which says that the Plancherel weight of G_x varies continuously in a suitable sense. This lead us to the definition of a lower semi-continuous $C_0(X)$ -weight on a $C_0(X)$ - C^* -algebra which we illustrate by three examples.

The first section recalls the construction of the canonical weight of a left Hilbert algebra and the properties which are needed to prove the crucial Corollary 1.6. In the second section, we consider the case of a locally compact group bundle and prove our main result. This example motivates the definition of a $C_0(X)$ -weight of a $C_0(X)$ - C^* -algebra, given in the third section. The fact that the Plancherel $C_0(X)$ -weight of the C^* -algebra of a group bundle is densely defined and lower semi-continuous gives Proposition 3.6 of [12].

1. THE PLANCHEREL WEIGHT OF A LOCALLY COMPACT GROUP

We recall first some elements of Tomita-Takesaki's theory, using the standard notation from [16]. Given a left Hilbert algebra \mathcal{A} where the product, the involution and the scalar product are respectively denoted by ab , a^\sharp and $(a|b)$, we denote by \mathcal{H} the Hilbert space completion of \mathcal{A} and by $\pi : \mathcal{A} \rightarrow \mathcal{L}(\mathcal{H})$ the left representation. We denote by $\mathcal{M} = \pi(\mathcal{A})''$ the left von Neumann algebra of \mathcal{A} . We denote by S the closure of the involution $a \mapsto a^\sharp$ and by F its adjoint. The domain of S [resp. F] is denoted by \mathcal{D}^\sharp [resp. \mathcal{D}^b] and one writes $\xi^\sharp = S\xi$ [resp. $\eta^b = F\eta$] for $\xi \in \mathcal{D}^\sharp$ [resp. for $\eta \in \mathcal{D}^b$]. An element $\eta \in \mathcal{H}$ is called right bounded if there exists a bounded operator $\pi'(\eta)$ on \mathcal{H} such that $\pi'(\eta)a = \pi(a)\eta$ for all $a \in \mathcal{A}$. One then writes $\xi\eta = \pi'(\eta)\xi$ for all $\xi \in \mathcal{H}$. The set of right bounded elements is denoted by \mathcal{B}' . One shows that $\mathcal{A}' = \mathcal{B}' \cap \mathcal{D}^b$ with involution ξ^b is a right Hilbert algebra in the same Hilbert space \mathcal{H} . An element $\xi \in \mathcal{H}$ is called left bounded if there exists a bounded operator $\pi(\xi)$ on \mathcal{H} such that $\pi(\xi)\eta = \pi'(\eta)\xi$ for all $\eta \in \mathcal{A}'$. One then writes $\xi\eta = \pi(\xi)\eta$ for all $\eta \in \mathcal{H}$. If ξ is left bounded and η is right bounded, the notation is consistent: $\pi(\xi)\eta = \xi\eta = \pi'(\eta)\xi$. The set of left bounded elements is denoted by \mathcal{B} . We shall need the following well-known lemmas.

Lemma 1.1. *Let \mathcal{B} be the set of left bounded elements of \mathcal{H} . Then,*

- (i) \mathcal{B} is a linear subspace of \mathcal{H} containing \mathcal{A} .
- (ii) $\pi(\mathcal{B})$ is contained in the left von Neumann algebra \mathcal{M} .
- (iii) \mathcal{B} is stable under \mathcal{M} . More precisely, for $T \in \mathcal{M}$ and $\xi \in \mathcal{B}$, $\pi(T\xi) = T\pi(\xi)$.

Proof. See [2, Section 2]. □

Lemma 1.2. *Let $\xi, \xi' \in \mathcal{B}$ such that $\pi(\xi') = \pi(\xi)^*$. Then ξ belongs to \mathcal{D}^\sharp and $\xi' = \xi^\sharp$.*

Proof. For all $\eta_1, \eta_2 \in \mathcal{A}'$, one has:

$$\begin{aligned}
(\xi | \eta_1 \eta_2^b) &= (\xi | \pi'(\eta_2^b) \eta_1) \\
&= (\xi | \pi'(\eta_2)^* \eta_1) \\
&= (\pi'(\eta_2) \xi | \eta_1) \\
&= (\pi(\xi) \eta_2 | \eta_1) \\
&= (\pi(\xi')^* \eta_2 | \eta_1) \\
&= (\eta_2 | \pi(\xi') \eta_1) \\
&= (\eta_2 | \pi'(\eta_1) \xi') \\
&= (\pi'(\eta_1)^* \eta_2 | \xi') \\
&= (\pi'(\eta_1^b) \eta_2 | \xi') \\
&= (\eta_2 \eta_1^b | \xi')
\end{aligned}$$

According to [16, Lemma 3.4] and the comments which follow, this suffices to conclude. \square

Definition 1.3. The canonical weight τ of the left Hilbert algebra \mathcal{A} is the map $\tau : \mathcal{M}_+ \rightarrow [0, \infty]$ defined by

$$\tau(T) = \begin{cases} \|\xi\|^2 & \text{if } \exists \xi \in \mathcal{B} : T = \pi(\xi)^* \pi(\xi) \\ \infty & \text{otherwise} \end{cases}$$

Lemma 1.4. [2, Théorème 2.11] *The above canonical weight τ is well-defined and is a faithful, semi-finite, σ -weakly lower semi-continuous weight on \mathcal{M}_+ .*

The following lemma is certainly well-known but I did not find a reference for it.

Lemma 1.5. *For all $\xi \in \mathcal{B}$, one has*

$$\tau(\pi(\xi)\pi(\xi)^*) = \begin{cases} \|\xi^\sharp\|^2 & \text{if } \xi \in \mathcal{D}^\sharp \\ \infty & \text{otherwise} \end{cases}$$

Proof. Suppose that $\xi \in \mathcal{D}^\sharp$. Then ξ belongs to the full Hilbert algebra $\mathcal{A}'' = \mathcal{B} \cap \mathcal{D}^\sharp$ of \mathcal{A} . Therefore, ξ^\sharp is also left bounded and $\pi(\xi)^* = \pi(\xi^\sharp)$. Thus,

$$\tau(\pi(\xi)\pi(\xi)^*) = \tau(\pi(\xi^\sharp)^* \pi(\xi^\sharp)) = \|\xi^\sharp\|^2$$

Suppose now that the left handside is finite. Then there exists $\eta \in \mathcal{B}$ such that $\pi(\xi)\pi(\xi)^* = \pi(\eta)^* \pi(\eta)$. Let $\pi(\xi) = U|\pi(\xi)|$ [resp. $\pi(\eta) = V|\pi(\eta)|$] be the polar decomposition of $\pi(\xi)$ [resp. $\pi(\eta)$]. All these operators belong to the left von Neumann algebra \mathcal{M} . Our assumption implies that $|\pi(\eta)| = U|\pi(\xi)|U^*$. Therefore,

$$\begin{aligned}
(U^*V^*\pi(\eta))^* &= \pi(\eta)^*VU \\
&= |\pi(\eta)|V^*VU \\
&= |\pi(\eta)|U \\
&= U|\pi(\xi)| \\
&= \pi(\xi)
\end{aligned}$$

Moreover, according to Lemma 1.1, $U^*V^*\pi(\eta) = \pi(U^*V^*\eta)$ and $\zeta = U^*V^*\eta$ is left bounded. Since $\pi(\xi) = \pi(\zeta)^*$ we deduce from Lemma 1.2 that $\xi \in \mathcal{D}^\sharp$. \square

We consider now the left Hilbert algebra associated with the left regular representation of a locally compact group G endowed with a left Haar measure λ . Using the framework of [14, pages 56-58], we choose the left Hilbert algebra $\mathcal{A} = C_c(G)$, where the product is the usual convolution product, the involution, denoted by f^* instead of f^\sharp , is $f^*(\gamma) = \overline{f(\gamma^{-1})}$ and the scalar product is

$$(f | g) = \int f \bar{g} d\lambda^{-1}.$$

The left representation is denoted by L instead of π . It acts on the Hilbert space $\mathcal{H} = L^2(G, \lambda^{-1})$ by $L(f)g = f * g$ for $f, g \in C_c(G)$. The left von Neumann algebra \mathcal{M} is the group von Neumann algebra $VN(G)$. The canonical weight τ of this left Hilbert algebra is also called the Plancherel weight of the group G ([15, 17]). We also refer the reader to [6, Section 2] for a deep study of its properties. It satisfies

$$\tau(L(f^* * f)) = \|f\|^2 = \int |f|^2 d\lambda^{-1} = (f^* * f)(e)$$

for $f \in C_c(G)$. When G is abelian, we can use the Fourier transform to compute the Plancherel weight (see [17, VII.3]). The canonical weight $\hat{\tau}$ of the commutative Hilbert algebra $L^2(\hat{G}, \hat{\lambda}) \cap L^\infty(\hat{G})$ is the integral with respect to the dual Haar measure $\hat{\lambda}$ on $L^\infty(\hat{G})_+$. This weight is called the Haar weight of the dual group \hat{G} . The Fourier transform \mathcal{F} implements an isomorphism between the full left Hilbert algebra of the regular representation of G and the Hilbert algebra $L^2(\hat{G}, \hat{\lambda}) \cap L^\infty(\hat{G})$. Therefore, $\tau(T) = \hat{\tau}(\mathcal{F} \circ T \circ \mathcal{F}^{-1})$ for $T \in VN(G)$. One has in particular

$$\tau(L(a)) = \int \mathcal{G}(a) d\hat{\lambda},$$

for $a \in C^*(G)_+$ where $\mathcal{G} : C^*(G) \rightarrow C_0(\hat{G})$ is the Gelfand transform.

Let us express Lemma 1.5 in the case of the left regular representation of a locally compact group G with a left Haar measure λ .

Corollary 1.6. *For all left bounded element ξ of $L^2(G, \lambda^{-1})$, one has*

$$\tau(L(\xi)L(\xi)^*) = \int |\xi|^2 d\lambda.$$

Remark 1.7. One can give a more direct proof of this result since the equality $L(\xi)^*\eta = \xi^* * \eta$ can be established by usual integration techniques and [5, Lemme 3.1]. However, its natural framework is Tomita-Takesaki's theory.

2. THE PLANCHEREL WEIGHT OF A GROUP BUNDLE

We consider now the case of a locally compact group bundle $p : G \rightarrow X$. We assume that the groups G_x have a left Haar measure λ^x such that $x \mapsto \lambda^x$ is continuous. This is the particular case of a locally compact groupoid with Haar system where the range and source maps coincide. Thus, we can construct its C^* -algebra $C^*(G)$ as usual. If the groups G_x are abelian, it is a commutative C^* -algebra. Then the Gelfand transform identifies it with $C_0(\hat{G})$. The space \hat{G}

is the total space of the bundle $\hat{p} : \hat{G} \rightarrow X$, where the fibre \hat{G}_x is the dual group of G_x . For $a \in C^*(G)_+$ and $x \in X$, we define

$$\mathcal{T}(a)(x) \stackrel{\text{def}}{=} \int \mathcal{G}(a) d\widehat{\lambda}^x = \int \mathcal{G}_x(a_x) d\widehat{\lambda}^x = \tau_x \circ L_x(a_x),$$

where L_x is the left regular representation of G_x and τ_x is the Plancherel weight of G_x . The last expression is defined when the groups are non-abelian and we turn now to this case.

Lemma 2.1. *Let $G \rightarrow X$ be a locally compact group bundle with Haar system $\lambda = (\lambda^x)_{x \in X}$. Then,*

- (i) $x \mapsto C^*(G_x)$ is an upper semi-continuous field of C^* -algebras;
- (ii) its sectional algebra is $C^*(G)$.

Proof. See for example [10, Section 5]. □

Equivalently, this lemma says that $C^*(G)$ is a $C_0(X)$ - C^* -algebra. We shall view an element a of $C^*(G)$ as a continuous field $x \mapsto a_x$, where $a_x \in C^*(G_x)$. For $a \in C^*(G)_+$, we define as in the commutative case the function

$$\mathcal{T}(a) : x \mapsto \tau_x(L_x(a_x)),$$

where L_x is the left regular representation of $C^*(G_x)$ and τ_x is the Plancherel weight of G_x . Our main result will be that the function $x \mapsto \mathcal{T}(a)(x)$ is lower semi-continuous. We consider now weights on C^* -algebras rather than von Neumann algebras and refer the reader to [1] for the main definitions and results.

Lemma 2.2. *Let $\varphi : A_+ \rightarrow [0, \infty]$ be a weight on a C^* -algebra A . Assume that $(e_i)_{i \in I}$ is an approximate identity for A such that $\|e_i\| \leq 1$ for all $i \in I$. Let $a \in A_+$. Then,*

- (i) $\varphi(a^{1/2}e_i a^{1/2}) \leq \varphi(a)$;
- (ii) if φ is lower semi-continuous, then $\varphi(a) = \sup_i \varphi(a^{1/2}e_i a^{1/2})$.

Proof. For (i), we have $a^{1/2}e_i a^{1/2} \leq \|e_i\|a \leq a$. Since φ is increasing, we obtain the desired inequality. For (ii), since $a^{1/2}e_i a^{1/2}$ converges to a , we have

$$\varphi(a) \leq \liminf \varphi(a^{1/2}e_i a^{1/2}) \leq \sup \varphi(a^{1/2}e_i a^{1/2}) \leq \varphi(a)$$

□

Proposition 2.3. *Let $G \rightarrow X$ be a locally compact group bundle with Haar system $\lambda = (\lambda^x)_{x \in X}$. Given $a \in C^*(G)_+$, the function*

$$\mathcal{T}(a) : x \mapsto \tau_x(L_x(a_x))$$

is lower semi-continuous.

Proof. Proposition 2.10 of [11] gives the existence of an approximate unit (e_i) of $C^*(G)$ where each e_i is a finite sum of elements of the form $f * f^*$ with $f \in C_c(G)$; moreover, according to its construction, $\|e_i\|_I$ tends to 1, where $\|f\|_I = \max(\sup_x \int |f| d\lambda^x, \sup_x \int |f| d\lambda_x)$ and $\lambda_x = (\lambda^x)^{-1}$. Replacing e_i by $(1/\|e_i\|_I)e_i$, we can assume that $\|e_i\|_I = 1$ for all $i \in I$. We then have $\|e_i\| \leq \|e_i\|_I \leq 1$. For

all $x \in X$, the image $(e_i(x))$ of (e_i) in $C^*(G_x)$ satisfies the same assumptions. Since the Plancherel weight τ_x is σ -weakly lower semi-continuous, we have

$$\tau_x(L_x(a_x)) = \sup_i \tau_x(L_x(a_x^{1/2} e_i(x) a_x^{1/2})).$$

We will show that the function $x \mapsto \tau_x(L_x(a_x^{1/2} e_i(x) a_x^{1/2}))$ is lower semi-continuous. This will imply that the function $x \mapsto \tau_x(L_x(a_x))$ is lower semi-continuous as lower upper bound of a family of lower semi-continuous functions. It suffices to show that $x \mapsto \tau_x(L_x(a_x^{1/2} * (f * f^*)_x * a_x^{1/2}))$ is lower semi-continuous for all $f \in C_c(G)$. Let us fix $x \in X$. According to Lemma 1.1, f_x and $g_x = L_x(a_x^{1/2})f_x$ are left bounded elements of $L^2(G_x, \lambda_x)$. Thus we have according to Lemma 1.5 :

$$\tau_x(L_x(a_x^{1/2} * (f * f^*)_x * a_x^{1/2})) = \tau_x(L_x(g_x)L_x(g_x)^*) = \int |g_x(\gamma)|^2 d\lambda^x(\gamma).$$

Note that $g : x \mapsto g_x$ is an element of the $C_0(X)$ -Hilbert module $L^2(G, \lambda^{-1})$ (see [7, Section 2]). If the groups G_x are unimodular, then

$$\int |g(\gamma)|^2 d\lambda^x(\gamma) = \int |g(\gamma)|^2 d\lambda_x(\gamma)$$

depends continuously on x . If not, we first observe that for all $\rho \in C_c(G)$, ρg belongs to $L^2(G, \lambda^{-1})$. Let $D : G \rightarrow \mathbf{R}_+^*$ be the modular function of the group bundle G . It is a continuous homomorphism such that $\lambda^x = D\lambda_x$ for all $x \in X$ (see [8, Lemma 2.4]). There exists an increasing sequence (D_n) of continuous non-negative functions with compact support which converges pointwise to D . Then, for all $x \in X$, $\int D_n(\gamma)|g(\gamma)|^2 d\lambda_x(\gamma)$ is an increasing sequence which converges to

$$\int D(\gamma)|g(\gamma)|^2 d\lambda_x(\gamma) = \int |g(\gamma)|^2 d\lambda^x(\gamma).$$

Therefore, the function $x \mapsto \int |g(\gamma)|^2 d\lambda^x(\gamma)$ is lower semi-continuous as a limit of an increasing sequence of continuous functions. \square

3. $C_0(X)$ -WEIGHTS ON $C_0(X)$ - C^* -ALGEBRAS

In this section, given a topological space X , $LSC(X)_+$ denotes the convex cone of lower semi-continuous functions $f : X \rightarrow [0, \infty]$.

Definition 3.1. Let X be a locally compact Hausdorff space. A $C_0(X)$ -weight on a $C_0(X)$ - C^* -algebra A is a map

$$\Phi : A_+ \rightarrow LSC(X)_+$$

such that

- (i) $\Phi(a + b) = \Phi(a) + \Phi(b)$ for all $a, b \in A_+$;
- (ii) $\Phi(ha) = h\Phi(a)$ for all $h \in C_0(X)_+$ and $a \in A_+$.

It is called lower semi-continuous if $a_n \rightarrow a$ implies $\Phi(a) \leq \liminf \Phi(a_n)$ and densely defined if its domain, defined as

$$P = \{a \in A_+ : \Phi(a) \text{ is finite and continuous}\}$$

is dense in A_+ .

In order to include the first example, we need to modify slightly this definition. Given a topological space X , we denote by $C_b(X)$ the space of complex-valued bounded continuous functions on X . We say that a C^* -algebra A is a $C_b(X)$ -algebra if it is endowed with a nondegenerate morphism of $C_b(X)$ into the centre of the multiplier algebra of A . Then we define a $C_b(X)$ -weight by replacing $C_0(X)$ by $C_b(X)$ in the above definition.

Remark 3.2. The above definition of a $C_0(X)$ -weight on a $C_0(X)$ - C^* -algebra is motivated by the examples below; it is different from the definition of a C^* -valued weight given in [9].

Lemma 3.3. *The domain P of a $C_b(X)$ -weight or of $C_0(X)$ -weight is hereditary: if $0 \leq b \leq a$ and $a \in P$, then $b \in P$.*

Proof. See [4, Lemme 4.4.2.i]. □

Example 3.4. *The canonical center-valued trace* (cf [3, Section 3.4]). Let A be a C^* -algebra and let $X = \text{Prim}(A)$ its primitive ideal space endowed with the Jacobson topology. Through the Dauns-Hofmann theorem, we view A as a $C_b(X)$ -algebra. Given $a \in A_+$, according to [13, Proposition 4.4.9], one can define $\Phi(a)(P) = \text{Trace}(\pi(a))$ where P is a primitive ideal and π is any irreducible representation admitting P as kernel, and $\Phi(a)$ is lower semi-continuous on X . Condition (i) of the definition results from the additivity of the trace and condition (ii) from the irreducibility of π . Thus $\Phi : A_+ \rightarrow LSC(X)_+$ is a $C_b(X)$ -weight. Since the usual Trace is σ -weakly lower semi-continuous, Φ is lower semi-continuous. By definition, Φ is densely defined if and only if A is a continuous-trace C^* -algebra.

Example 3.5. *The Plancherel $C_0(X)$ -weight of a group bundle.* This is the example described in the previous section:

Theorem 3.6. *Let $p : G \rightarrow X$ be a locally compact group bundle endowed with a continuous Haar system. Then the map*

$$\mathcal{T} : C^*(G)_+ \rightarrow LSC(X)_+$$

such that $\mathcal{T}(a)(x) = \tau_x(L_x(a_x))$ for $a \in C^(G)_+$ and where τ_x is the Plancherel weight of G_x and L_x the left regular representation of G_x is a densely defined lower semi-continuous $C_0(X)$ -weight, which we call the Plancherel $C_0(X)$ -weight of the group bundle.*

Proof. The main point was to show that the range of this map is contained in $LSC(X)_+$, which was done in Proposition 2.3. The conditions (i) and (ii) of the definition are clear. It is densely defined since its domain contains the elements $f^* * f$ where $f \in C_c(G)$, whose linear span is dense in $C^*(G)$. The lower semi-continuity of $\tau_x \circ L_x$ for all $x \in X$ gives the lower semi-continuity of \mathcal{T} . □

Example 3.7. *The commutative case.* Let $\pi : Y \rightarrow X$ be a continuous, open and surjective map, where Y and X are locally compact Hausdorff spaces. For $x \in X$, let $Y_x = \pi^{-1}(x)$ be the fibre over x . Endowed with the fundamental family

of continuous sections $C_c(Y)$, $x \mapsto C_0(Y_x)$ is a continuous field of C^* -algebras. Its C^* -algebra of continuous sections is identified to $C_0(Y)$. Thus $C_0(Y)$ is a $C_0(X)$ - C^* -algebra. The following proposition gives a description of the densely defined lower semi-continuous $C_0(X)$ -weights of $C_0(Y)$.

Proposition 3.8. *In the above situation, we have natural one-to-one correspondence between*

- (i) *the continuous π -systems of measures, by which we mean families of Radon measures $\alpha = (\alpha_x)_{x \in X}$ on Y , where α_x is supported on $\pi^{-1}(x)$, and such that for all $f \in C_c(Y)$, the function $x \mapsto \int f d\alpha_x$ is continuous;*
- (ii) *the densely defined and lower semi-continuous $C_0(X)$ -weights of $C_0(Y)$.*

Proof. When X is reduced to a point, this is a well-known result given for example in the introduction of [1]. A crucial point which we shall use again below is that a densely defined weight on $C_0(Y)$ is necessarily finite on $C_c(Y)_+$ because the linear span of its domain of definition is a dense ideal, hence it contains the minimal dense ideal, called the Pedersen ideal, which in our case is $C_c(Y)$ ([13, 5.6.3.]). The proof of the general case is essentially the same. Suppose that $\alpha = (\alpha_x)_{x \in X}$ is a continuous π -system. For $f \in C_0(Y)_+$, we define $\Phi(f)(x) = \int f d\alpha_x$. There exists an increasing sequence (f_n) in $C_c(Y)_+$ converging uniformly to f . Therefore, $\Phi(f) = \sup_n \Phi(f_n)$ is lower semi-continuous. It is clear that the other properties of a $C_0(X)$ -weight are satisfied. Since its domain contains $C_c(Y)$, Φ is densely defined. The lower semi-continuity of the Radon measures gives the lower semi-continuity of Φ . Conversely, let Φ be a densely defined and lower semi-continuous $C_0(X)$ weight on $C_0(Y)$. For all $x \in X$, we can define $\varphi_x : C_0(Y_x)_+ \rightarrow [0, \infty]$ such that $\Phi(f)(x) = \varphi_x(f|_{Y_x})$ for all f . Then φ_x is a densely defined and lower semi-continuous weight on $C_0(Y_x)$. As we recalled at the beginning, there exists a unique Radon measure α_x such that $\varphi_x(f) = \int f d\alpha_x$ for all $f \in C_0(Y_x)$. By assumption, the linear span of the domain of definition P of Φ is dense. Since it is also an ideal, it contains the Pedersen ideal $C_c(Y)$ of $C_0(Y)$. This shows that $\alpha = (\alpha_x)_{x \in X}$ is a continuous π -system of measures. \square

Example 3.9. *The Haar $C_0(X)$ -weight of a groupoid.* This example is a particular case of the previous example. Let G be a locally compact groupoid. We use the range map $r : G \rightarrow G^{(0)}$ to turn the commutative C^* -algebra $C_0(G)$ into a $C_0(X)$ -algebra, where $X = G^{(0)}$. Then a continuous Haar system $(\lambda^x)_{x \in X}$ defines a densely defined and lower semi-continuous $C_0(X)$ -weight of $C_0(G)$, which in accordance with the group case, can be called the Haar $C_0(X)$ -weight of G . On the other hand, we cannot define a Plancherel $C_0(X)$ -weight for an arbitrary groupoid G since $C^*(G)$ is usually not a $C_0(X)$ -algebra.

We now have all the elements to prove the Proposition 3.6 in [12].

Corollary 3.10. *Let $p : G \rightarrow X$ a locally compact bundle of abelian groups, equipped with a continuous Haar system $\lambda = (\lambda^x)_{x \in X}$. Then the family of dual Haar measures $\hat{\lambda} = (\hat{\lambda}^x)_{x \in X}$ is a continuous Haar system for $\hat{p} : \hat{G} \rightarrow X$.*

Proof. From Theorem 3.6, we know that the Plancherel $C_0(X)$ -weight \mathcal{T} on $C^*(G)$ is lower semi-continuous and densely defined. Therefore $\Phi = \mathcal{T} \circ \mathcal{G}^{-1}$, where

$\mathcal{G} : C^*(G) \rightarrow C_0(\hat{G})$ is the Gelfand transform, is a lower semi-continuous and densely defined $C_0(X)$ -weight on $C_0(\hat{G})$. Moreover, for all $x \in X$, the Plancherel weight τ_x of G_x correspond to the Haar weight of \hat{G}_x , which is given by the Radon measure $\hat{\lambda}^x$. From Proposition 3.8, the \hat{p} -system $\hat{\lambda} = (\hat{\lambda}^x)_{x \in X}$ is continuous. \square

Acknowledgements. I thank Henrik Kreidler for drawing the erroneous proof of Proposition 3.6 in [12] to our attention, Michel Hilsun for discussions which led to the present proof of Corollary 3.10 and Dana Williams for sharing another proof and providing inspiring feedback and valuable comments.

REFERENCES

- [1] F. Combes: *Poids sur une C^* -algèbre*, J. Math. pures et appl., **47** (1968), 57–100.
- [2] F. Combes: *Poids associé à une algèbre hilbertienne à gauche*, Compositio Mathematica, **23** no 1 (1971), 49–77.
- [3] J. Dixmier: *Les algèbres de von Neumann*, Gauthier-Villars, 1969.
- [4] J. Dixmier: *Les C^* -algèbres et leurs représentations*, Gauthier-Villars, 1969.
- [5] P. Eymard: *L'algèbre de Fourier d'un groupe localement compact*, Bull. Soc. math. France, **92** (1964), 181–236.
- [6] U. Haagerup: *On the dual weights for crossed products of von Neumann algebras II*, Math. Scand. **43** (1978) 119–140.
- [7] M. Khoshkam and G. Skandalis: *Regular representation of groupoid C^* -algebras and applications to inverse semi-groups*, J. reine angew. Math. **546** (2002) 47–72.
- [8] M. Ionescu, A. Kumjian, J. Renault, A. Sims and D. Williams: *C^* -algebras of extensions of groupoids by group bundles*, arXiv 2001:01312.
- [9] J. Kustermans: *Regular C^* -valued weights*, J. Operator Theory, **44** No 1 (2000), 151–205.
- [10] N. P. Landsman and B. Ramazan: *Quantization of Poisson algebras associated to Lie algebroids*, Contemporary Mathematics **282** (2001), 159–192.
- [11] P. Muhly, J. Renault and D. Williams: *Equivalence and isomorphism for groupoid C^* -algebras*, J. Operator Theory, **17**(1987), 3–22.
- [12] P. Muhly, J. Renault and D. Williams: *Continuous trace groupoid C^* -algebras III*, Trans. Amer. Math. Soc. **348** (1996), 3621–3641.
- [13] G. Pedersen: *C^* -algebras and their automorphism groups*, Academic Press, 1979.
- [14] J. Renault: *A groupoid approach to C^* -algebras*, Lecture Notes in Mathematics, **793** (1980), Springer.
- [15] S. Stratila: *Modular theory in operator algebras*, Abacus Press, 1981.
- [16] M. Takesaki: *Tomita's theory of modular Hilbert algebras and its applications*, Lecture Notes in Mathematics, **128** (1970), Springer.
- [17] M. Takesaki: *Theory of operator algebras II*, Encyclopedia of Mathematical Sciences, Springer Verlag, 2002.

INSTITUT DENIS POISSON (UMR 7013), UNIVERSITÉ D'ORLÉANS ET CNRS, 45067 ORLÉANS CEDEX 2, FRANCE

Email address: jean.renault@univ-orleans.fr