



HAL
open science

Almost harmonic spinors

Nicolas Ginoux, Jean-François Grosjean

► **To cite this version:**

Nicolas Ginoux, Jean-François Grosjean. Almost harmonic spinors. *Comptes Rendus. Mathématique*, 2010, 348 (13-14), pp.811-814. 10.1016/j.crma.2010.06.010 . hal-03117246

HAL Id: hal-03117246

<https://hal.science/hal-03117246>

Submitted on 21 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Almost harmonic spinors

Nicolas Ginoux*, Jean-François Grosjean†

January 21, 2021

Abstract. We show that any closed spin manifold not diffeomorphic to the two-sphere admits a sequence of volume-one-Riemannian metrics for which the smallest non-zero Dirac eigenvalue tends to zero. As an application, we compare the Dirac spectrum with the conformal volume.

Spineurs presque harmoniques

Résumé. Nous montrons que, sur toute variété spinorielle compacte sans bord non diffeomorphe à la sphère de dimension deux, il existe une suite de métriques riemanniennes de volume un pour laquelle la plus petite valeur propre non nulle de l'opérateur de Dirac tend vers zéro. Comme application, nous comparons le spectre de l'opérateur de Dirac avec le volume conforme.

1 Introduction and statement

Let M^n be an $n(\geq 2)$ -dimensional closed spin manifold and denote by D_g the spin Dirac operator associated to a Riemannian metric g . We denote by $\lambda_1(D_g^2)$ and $\lambda_1^+(D_g^2)$ the smallest and the smallest positive eigenvalue of D_g^2 respectively. It is well-known that the product $\lambda_1^+(D_g^2)\text{Vol}(M^n, g)^{\frac{2}{n}}$ is scaling-invariant and bounded from below by a positive constant in any conformal class [1, Thm. 2.3]. One can ask whether the infimum of $\lambda_1^+(D_g^2)\text{Vol}(M, g)^{\frac{2}{n}}$ on the space of all Riemannian metrics remains positive. This holds true if M is the 2-sphere \mathbb{S}^2 since it has only one conformal class; alternatively, it follows from C. Bär's estimate [6] valid for any Riemannian metric g on \mathbb{S}^2 :

$$\lambda_1(D_g^2)\text{Area}(\mathbb{S}^2, g) \geq 4\pi. \quad (1)$$

In this respect \mathbb{S}^2 is the only exception:

Theorem 1.1 *For any $n(\geq 2)$ -dimensional closed spin manifold M^n not diffeomorphic to \mathbb{S}^2 there exists a sequence $(g_p)_{p \in \mathbb{N}}$ of Riemannian metrics on M^n such that $\lambda_1^+(D_{g_p}^2)\text{Vol}(M^n, g_p)^{\frac{2}{n}} \xrightarrow{p \rightarrow \infty} 0$.*

Therefore one can get the Dirac spectrum as close to 0 as one wants with fixed volume. Note however that Theorem 1.1 does not prove the existence of non-zero harmonic spinors, i.e., that 0 is a Dirac eigenvalue.

Theorem 1.1 is proved in Section 2. In Section 3 we apply it to compare the Dirac spectrum with the conformal volume.

Acknowledgment. This work started during a stay of the first-named author at the Institut Élie Cartan de Nancy, which he would like to thank for its hospitality and support. It is also a pleasure to thank Bernd Ammann and Emmanuel Humbert for fruitful discussions and their critical reading of the paper.

*NWF I - Mathematik, Universität Regensburg, D-93040 Regensburg, E-mail: nicolas.ginoux@mathematik.uni-regensburg.de

†Institut Élie Cartan (Mathématiques), Université Henri Poincaré Nancy I, B.P.239 F-54506 Vandoeuvre-Lès-Nancy Cedex, E-mail: grosjean@iecn.u-nancy.fr

2 Proof

The proof of Theorem 1.1 relies on a standard technique first used in the spinorial context by C. Bär [7] to show the existence of metrics with harmonic spinors. Namely we prove the result by gluing a model manifold admitting such a sequence and by studying the convergence of the spectrum on the connected sum. Thus the proof is two-step.

Lemma 2.1

- i) Theorem 1.1 holds true on the standard sphere \mathbb{S}^n for any $n \geq 3$.
- ii) Theorem 1.1 holds true on the 2-torus \mathbb{T}^2 endowed with any of its 4 spin structures.

Proof: Both statements follow from elementary arguments.

i) For any $n \geq 3$ there exists on \mathbb{S}^n a metric \tilde{g} with $\text{Ker}(D_{\tilde{g}}) \neq \{0\}$: for $n \equiv 3 \pmod{4}$ it is some Berger metric as shown by C. Bär [7, Cor. p.906], for $n = 2m \geq 4$ and $n \geq 5$ the existence of such a metric has been proved by L. Seeger [13] and M. Dahl [9, Cor. 4.2] respectively. Linear interpolation between the standard round metric and \tilde{g} provides a smooth one-parameter-family of Riemannian metrics $(g_t)_{0 \leq t \leq 1}$ with $g_0 = \text{can}$ and $g_1 = \tilde{g}$. Since the volume remains bounded and the Dirac spectrum depends continuously on the metric in the C^1 -topology, we obtain the result.

ii) For a real parameter $a > 1$ consider the 2-torus $\mathbb{T}^2 := \mathbb{R}^2/\Gamma$, where $\Gamma := \mathbb{Z} \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} \oplus \mathbb{Z} \cdot \begin{pmatrix} 0 \\ a \end{pmatrix}$, with induced flat metric g_a . It carries 4 spin structures, 3 of which can be deduced from each other by an orientation-preserving diffeomorphism of \mathbb{T}^2 (see e.g. [4]). Thus it suffices to prove the statement for two spin structures which cannot be obtained from each other by a diffeomorphism, for example for the spin structure inducing a trivial covering on both factors and for the spin structure inducing a trivial covering on the first factor and a non-trivial one on the second one. For the former spin structure the smallest positive eigenvalue of $D_{g_a}^2$ is $\frac{4\pi^2}{a^2}$ and for the latter one it is $\frac{\pi^2}{a^2}$. Since $\text{Area}(\mathbb{T}^2, g_a) = a$ we conclude that in both situations $\lambda_1^+(D_{g_a}^2) \text{Area}(\mathbb{T}^2, g_a) \xrightarrow{a \rightarrow \infty} 0$ (compare with [4, Sec. 3]). \square

In the second step we consider the dimensions $n = 2$ and $n \geq 3$ separately. In the latter case it only remains to know how the Dirac spectrum behaves under connected sum¹:

Theorem 2.2 (C. Bär [8]) *Let (N_1^n, g_1) and (N_2^n, g_2) be closed Riemannian spin manifolds of dimension $n \geq 3$. Let $L > 0$ and $\eta \geq 0$ with $\pm(L + \eta) \notin (\text{Spec}(D_{g_1}) \cup \text{Spec}(D_{g_2}))$.*

Then for any $\varepsilon > 0$, there exists a Riemannian metric \tilde{g} on the connected sum $\tilde{N}^n := N_1^n \sharp N_2^n$ such that the Dirac eigenvalues of $N_1^n \cup N_2^n$ and (\tilde{N}^n, \tilde{g}) in $]-L - \eta, L + \eta[$ differ at most by ε and that $\text{Vol}(\tilde{N}^n, \tilde{g}) \leq \text{Vol}(N_1^n, g_1) + \text{Vol}(N_2^n, g_2) + \varepsilon$.

Note that, as an easy consequence, Theorem 2.2 remains valid when replacing the eigenvalues of the Dirac operator by those of its square. Fix now any Riemannian metric g on M^n (with $n \geq 3$). If p is any positive integer, pick from Lemma 2.1 a Riemannian metric g_p of volume one on \mathbb{S}^n with $\lambda_1^+(D_{g_p}^2) \leq \frac{1}{p}$.

Setting $L := \lambda_1^+(D_{g_p}^2)$, $\varepsilon := \frac{\lambda_1^+(D_{g_p}^2)}{2}$ and choosing $\eta > 0$ with $L + \eta \notin (\text{Spec}(D_{g_p}^2) \cup \text{Spec}(D_g^2))$, Theorem 2.2 implies the existence of a Riemannian metric \tilde{g}_p on $\tilde{N}^n := M^n \sharp \mathbb{S}^n$ such that at least one eigenvalue of $D_{\tilde{g}_p}^2$ lies in the interval $[\frac{\lambda_1^+(D_{g_p}^2)}{2}, \frac{3\lambda_1^+(D_{g_p}^2)}{2}]$ and that $\text{Vol}(\tilde{N}^n, \tilde{g}_p) \leq \text{Vol}(M^n, g) + 1 + \frac{1}{2p}$. Since \tilde{N}^n is spin diffeomorphic to M^n we conclude the proof of Theorem 1.1 for $n \geq 3$.

In dimension $n = 2$ we perform an induction on the genus of the surface. On \mathbb{T}^2 Theorem 1.1 already holds true by Lemma 2.1. Assume it to hold true for any closed oriented surface $M^2(\gamma)$ of genus $\gamma > 0$ and consider a closed oriented surface $M^2(\gamma + 1)$ of genus $\gamma + 1$. The oriented surface can be obtained as the connected sum of some $M^2(\gamma)$ and \mathbb{T}^2 . Moreover, the spin structure induced on a circle bounding a compact oriented surface is always a non-trivial covering [11, p.91], in particular it itself bounds a disk. Therefore, every spin structure on $M^2(\gamma + 1)$ is induced by some spin structure on $M^2(\gamma)$ and some on \mathbb{T}^2 . It would remain to prove the analog of Theorem 2.2 for surgeries of codimension 2, at least for connected sums of surfaces. We conjecture this holds true, using arguments and techniques from [2]. Actually much less is needed here:

¹Thanks to Christian Bär for indicating to us the right reference.

²The argument given in the published version was wrong. Thanks also to Christian Bär.

Lemma 2.3 *Let (M_1, g_1) and (M_2, g_2) be any oriented closed Riemannian surfaces. Fix $L > 0$ and $\eta \geq 0$ with $\pm(L + \eta) \notin (\text{Spec}(D_{g_1}) \cup \text{Spec}(D_{g_2}))$.*

Then for any $\varepsilon > 0$ there exists a Riemannian metric \tilde{g} on $M_1 \sharp M_2$ such that, for any eigenvalue λ of D_{g_1} or D_{g_2} in $] -L - \eta, L + \eta[$, there exists an eigenvalue $\tilde{\lambda}$ of $D_{\tilde{g}}$ such that $|\tilde{\lambda} - \lambda| \leq \varepsilon$ and $\text{Area}(M_1 \sharp M_2, \tilde{g}) \leq \text{Area}(M_1, g_1) + \text{Area}(M_2, g_2) + \varepsilon$.

Proof: The proof relies on a classical cut-off procedure for eigenvectors of D_{g_1} and D_{g_2} . We want to show that $\dim(\text{Ker}(D_{g_1} - \lambda \text{Id})) + \dim(\text{Ker}(D_{g_2} - \lambda \text{Id})) \leq \dim\left(\bigoplus_{\mu \in [\lambda - \varepsilon, \lambda + \varepsilon]} \text{Ker}(D_{\tilde{g}} - \mu \text{Id})\right)$. Fix $p_i \in M_i$,

$i = 1, 2$, and some sufficiently small $\delta > 0$. Consider the connected sum $\tilde{M} := M_1 \sharp M_2$ obtained by gluing $M_1 \setminus B_{p_1}(\delta)$ and $M_2 \setminus B_{p_2}(\delta)$ along their boundary, where $B_p(r)$ denotes the open metric disc of center p and radius r . From [7, p.932] or [2, Sec. 3.1-3.2] there exists a smooth Riemannian metric \tilde{g}_δ on \tilde{M} which coincides with g_i on $M_i \setminus B_{p_i}(\sqrt{\delta})$ and such that $\text{Area}(M_1 \sharp M_2, \tilde{g}_\delta) \leq \text{Area}(M_1, g_1) + \text{Area}(M_2, g_2) + c \cdot \sqrt{\delta}$, where $c > 0$ is a constant depending only on the metrics g_1 and g_2 . In particular we may choose $\delta > 0$ such that $c \cdot \sqrt{\delta} < \varepsilon$. For $i = 1, 2$ define $\chi_i \in C(\tilde{M}, [0, 1])$ by $\chi_i|_{M_i \setminus B_{p_i}(\sqrt{\delta})} := 1$, $\chi_i|_{B_{p_i}(\delta)} := 0$ and $\chi_i(x) := 2 - 2 \frac{\ln(d(x, p_i))}{\ln(\delta)}$ otherwise, where $d(x, p)$ denotes the distance between x and p . Note that χ_1 and χ_2 are well-defined and continuous on the whole \tilde{M} and that they can be smoothed out at both $\partial B_{p_i}(\delta)$ and $\partial B_{p_i}(\sqrt{\delta})$ such that the L^2 -norm of their gradient changes arbitrarily little. We keep denoting the corresponding smooth functions by χ_1 and χ_2 . Consider the map

$$\begin{aligned} \Phi : \text{Ker}(D_{g_1} - \lambda \text{Id}) \oplus \text{Ker}(D_{g_2} - \lambda \text{Id}) &\longrightarrow \Gamma(\Sigma \tilde{M}) \\ (\varphi_1, \varphi_2) &\longmapsto \chi_1 \varphi_1 + \chi_2 \varphi_2, \end{aligned}$$

which is well-defined because of $\chi_i|_{B_{p_i}(\delta)} = 0$ and injective by the unique continuation property (each Dirac eigenvector vanishing on an open subset of a connected Riemannian spin manifold must vanish identically).

Now from the min-max principle it suffices to show that $\frac{\|(D_{\tilde{g}_\delta} - \lambda)\varphi\|_{L^2}}{\|\varphi\|_{L^2}} \leq \varepsilon$ for all $\varphi \in \text{Im}(\Phi) \setminus \{0\}$.

Since the subspaces $\Phi(\text{Ker}(D_{g_1} - \lambda \text{Id}))$ and $\Phi(\text{Ker}(D_{g_2} - \lambda \text{Id}))$ are L^2 -orthogonal to each other (for $\text{supp}(\chi_1) \cap \text{supp}(\chi_2)$ has zero measure), we can assume that $\varphi \in \Phi(\text{Ker}(D_{g_1} - \lambda \text{Id}))$ with $\|\varphi\|_{L^2} = 1$. Using the formula $D_g(f\varphi) = df \cdot \varphi + f D_g \varphi$, we compute:

$$\begin{aligned} \int_{\tilde{M}} |(D_{\tilde{g}_\delta} - \lambda)\varphi|^2 v_{\tilde{g}_\delta} &= \int_{M_1} |(D_{\tilde{g}} - \lambda)\chi_1 \varphi_1|^2 v_{\tilde{g}_\delta} \\ &= \int_{M_1} |d\chi_1|^2 |\varphi_1|^2 v_{\tilde{g}_\delta} \\ &\leq C \sup_{M_1} (|\varphi_1|^2) \int_{\delta}^{\sqrt{\delta}} \frac{4}{r^2 \ln(\delta)^2} r dr \\ &\leq \frac{C'}{\ln(\delta)}, \end{aligned}$$

where $C > 0$ is a constant depending only on the original metrics in the ring $B_{p_1}(\sqrt{\delta}) \setminus B_{p_1}(\delta)$ and $C' = 2C \cdot \sup_{\varphi_1 \in \text{Ker}(D_{g_1} - \lambda \text{Id})} (\sup_{M_1} (|\varphi_1|^2))$ (note that $C' < \infty$ since $\text{Ker}(D_{g_1} - \lambda \text{Id})$ is finite-dimensional). We

deduce that $\|(D_{\tilde{g}_\delta} - \lambda)\varphi\|_{L^2}^2 \xrightarrow{\delta \rightarrow 0} 0$ and the statement of Lemma 2.3. \square

The proof of Theorem 1.1 for $n = 2$ follows the lines of that for $n \geq 3$: given any Riemannian metric g on $M^2(\gamma)$ and a positive $p \in \mathbb{N}$, pick from Lemma 2.1 a Riemannian metric g_p of unit area on \mathbb{T}^2 with $\lambda_1^+(D_{g_p}^2) \leq \frac{1}{p}$, whatever the spin structure of \mathbb{T}^2 is. Lemma 2.3 ensures the existence of a Riemannian metric \tilde{g}_p on $M^2(\gamma + 1) = M^2(\gamma) \sharp \mathbb{T}^2$ such that at least one eigenvalue of $D_{\tilde{g}_p}^2$ lies in the interval $[\frac{\lambda_1^+(D_{g_p}^2)}{2}, \frac{3\lambda_1^+(D_{g_p}^2)}{2}]$ and that $\text{Area}(M^2(\gamma + 1), \tilde{g}_p) \leq \text{Area}(M^2(\gamma), g) + 1 + \frac{1}{2p}$. This proves the result for $\gamma + 1$ and concludes the proof of Theorem 1.1.

3 Application

This note was motivated by the study of the relationship between the Dirac spectrum and the so-called conformal volume, which is the conformal invariant defined for any closed Riemannian manifold (M^n, g)

by

$$V_c(M^n, [g]) := \inf_{N \in \mathbb{N}} \left(\inf_{\varphi \in \text{Imm}_c(M^n, \mathbb{S}^N)} \left(\sup_{\gamma \in \text{Conf}(\mathbb{S}^N)} (\text{Vol}(M^n, (\gamma \circ \varphi)^* \text{can})) \right) \right),$$

where $\text{Imm}_c(M^n, \mathbb{S}^N)$ denotes the set of conformal immersions $(M^n, g) \rightarrow (\mathbb{S}^N, \text{can})$ and $\text{Conf}(\mathbb{S}^N)$ the group of conformal diffeomorphisms of $(\mathbb{S}^N, \text{can})$. First introduced by P. Li and S.-T. Yau [12], it has been shown to be directly related to the Laplace spectrum since it provides an upper bound of the corresponding spectral invariant [12, 10]: $(0 <) \lambda_1(\Delta) \text{Vol}(M^n, g)^{\frac{2}{n}} \leq n V_c(M^n, [g])^{\frac{2}{n}}$. For the Dirac operator such a result cannot be expected because of $\sup_{\bar{g} \in [g]} (\lambda_1^+(D_{\bar{g}}^2) \text{Vol}(M^n, \bar{g})^{\frac{2}{n}}) = \infty$, see [5, Thm. 1.1].

However, one could reasonably conjecture that the conformal volume bounds $\lambda_1(D_g^2) \text{Vol}(M, g)^{\frac{2}{n}}$ from below, provided the possible eigenvalue 0 is left aside. For $M = \mathbb{S}^2$ this is the case because of (1) and $4\pi = \text{Area}_c(\mathbb{S}^2)$ (see [12]). It is hopeless for any other manifold:

Corollary 3.1 *For any $n(\geq 2)$ -dimensional closed Riemannian spin manifold (M^n, g) not diffeomorphic to \mathbb{S}^2 there exists no positive constant $c(M)$ (depending only on M) such that*

$$\lambda_1^+(D_g^2) \text{Vol}(M, g)^{\frac{2}{n}} \geq c(M) V_c(M^n, [g])^{\frac{2}{n}}.$$

Proof: It is elementary to show that [12, Fact 2]

$$V_c(M^n, [g]) \geq \text{Vol}(\mathbb{S}^n, \text{can}), \quad (2)$$

whose r.h.s. does not depend on the metric g . We conclude with Theorem 1.1. \square

Still there exists a subtle relationship between the Dirac spectrum and the conformal volume. Indeed by [1, Thm. 3.1 & 3.2] and [3, Thm. 1.1], $\inf_{\bar{g} \in [g]} (\lambda_1^+(D_{\bar{g}}^2) \text{Vol}(M^n, \bar{g})^{\frac{2}{n}}) \leq \frac{n^2}{4} \text{Vol}(\mathbb{S}^n, \text{can})^{\frac{2}{n}}$, hence combining with (2) one obtains

$$\inf_{\bar{g} \in [g]} (\lambda_1^+(D_{\bar{g}}^2) \text{Vol}(M^n, \bar{g})^{\frac{2}{n}}) \leq \frac{n^2}{4} V_c(M^n, [g])^{\frac{2}{n}}.$$

References

- [1] B. Ammann, *A spin-conformal lower bound of the first positive Dirac eigenvalue*, *Diff. Geom. Appl.* **18** (2003), no. 1, 21–32.
- [2] B. Ammann, M. Dahl, E. Humbert, *Surgery and harmonic spinors*, *Adv. Math.* **220** (2009), no. 2, 523–539.
- [3] B. Ammann, J.-F. Grosjean, E. Humbert, B. Morel, *A spinorial analogue of Aubin’s inequality*, *Math. Z.* **260** (2008), no. 1, 127–151.
- [4] B. Ammann, E. Humbert, *The first conformal Dirac eigenvalue on 2-dimensional tori*, *J. Geom. Phys.* **56** (2006), no. 4, 623–642.
- [5] B. Ammann, P. Jammes, *The supremum of conformally covariant eigenvalues in a conformal class*, [arXiv:0708.0529v1](https://arxiv.org/abs/0708.0529v1) (2007).
- [6] C. Bär, *Lower eigenvalue estimates for Dirac operators*, *Math. Ann.* **293** (1992), no. 1, 39–46.
- [7] ———, *Metrics with Harmonic Spinors*, *Geom. Funct. Anal.* **6** (1996), 899–942.
- [8] ———, *Harmonic spinors for twisted Dirac operators*, *Math. Ann.* **309** (1997), no. 2, 225–246.
- [9] M. Dahl, *On the space of metrics with invertible Dirac operator*, *Comment. Math. Helv.* **83** (2008), no. 2, 451–469.
- [10] A. El Soufi, S. Ilias, *Immersiones minimales, première valeur propre du laplacien et volume conforme*, *Math. Ann.* **275** (1986), no. 2, 257–267.
- [11] H.B. Lawson, M.-L. Michelsohn, *Spin geometry*, Princeton University Press, Princeton, 1989.
- [12] P. Li, S.-T. Yau, *A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces*, *Invent. Math.* **69** (1982), no. 2, 269–291.
- [13] L. Seeger, *Metriken mit harmonischen Spinoren auf geradedimensionalen Sphären*, Shaker Verlag, Aachen, 2001.