- 1 Title Psychological vulnerability of French university students during COVID-19 lockdown
- 2 **Authors:**
- 3 Sarah Le Vigouroux^{\$}; Aurélie Goncalves^{*}\$; Elodie Charbonnier
- 45 Affiliation
- 6 UNIV. NIMES, EA7352 CHROME / APSY-V, F-30021 Nîmes Cedex, France
- 7 \$ Co-first author

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

- Abstract: Even before the current pandemic, university students were known to have high levels of hassles, anxiety, and depressive symptoms. In all probability, the advent of COVID-19 has substantially raised these levels. The present study measured the emotional state of university students during lockdown, and identified the relevant situational and psychological factors. To this end, 1297 French university students were assessed during lockdown, which lasted from 16 March to 11 May in France. Situational factors included the belief that lockdown was compromising their future job prospects, COVID-19 symptoms, and health concerns. Psychological factors included students' implicit theory of emotion, and the coping strategies they used during lockdown. We explored the extent to which these factors were associated with levels of hassles, anxiety, and depressive symptoms. Results indicated that students' belief that lockdown was compromising their future job prospects was positively related to hassles (r =.34, p < .001), while concerns about their own health and that of relatives were positively associated with anxiety (r = .37 and .34; p < .001). In addition, use of the positive reframing coping strategy mediated the effect of students' implicit theory of emotion on their depressive symptoms. Even though the national lockdown is now over, the pandemic will continue to have a major impact on university students over the coming months. It is therefore essential for universities to provide them with adequate psychological support.
- 26 **Keywords:** coronavirus; university students; depression; anxiety; hassles; lockdown;
- 27 pandemic

1. Introduction

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

On 12 March 2020, the French Government introduced a series of unprecedented measures designed to combat the spread of coronavirus (COVID-19), and on 16 March, the entire population of France was placed under lockdown. Universities were immediately closed, leading to urgent changes in learning habits and exams, generating considerable uncertainty, and massively reducing social contacts, especially for students living alone. Even before the pandemic, university students had been identified as having poorer mental health than their nonstudent peers (Stanley & Manthorpe, 2001). For example, a study conducted in 21 countries worldwide found that 18.5% of university students had had a least one major depressive episode, and 16.7% displayed a generalized anxiety disorder (Auerbach et al., 2018). Ibrahim et al. (2013)'s literature review including 24 studies found that the prevalence of depression among university students was 30.6%. In addition, a recent systematic review including 48 articles from different parts of the world showed a prevalence of 26.1% for depressive symptoms and 24.5% for anxiety among 56,816 students (de Paula et al., 2020). However, both reviews highlighted substantial disparities between countries and studies, in terms of education level, subject, or measures used. For example, 30% of Japanese students were found to have probable anxiety and depressive symptoms (Matsudaira et al., 2009), compared with only 13% of students in Lithuania (Bunevicius et al., 2008). Despite these disparities in prevalence, there is a consensus that university students worldwide are vulnerable to anxiety and depressive symptoms.

Ordinarily, students' poor mental health stems partly from the need to deal with an accumulation of hassles, in the shape of university pressure, schedule changes, material tasks, paid work activities, financial difficulties, and isolation (Réveillère et al., 2001). Coping strategies therefore play a central role. These strategies, and thus emotional experience, are influenced by beliefs about emotions (Tamir et al., 2007). Some individuals see emotions as the direct and uncontrollable consequences of events or contexts (i.e., static implicit theory).

Others, however, consider them to be malleable, and have a clearer sense of being able to act upon their emotional experience (i.e., dynamic implicit theory). Endorsement of a dynamic implicit theory of emotion is associated with less anxiety (De Castella et al., 2014), through better use of strategies such as cognitive reappraisal (Tamir et al., 2007). We can therefore hypothesize that implicit theory of emotion played a role in the coping strategies used by university students during COVID-19 lockdown, and thus in their emotional state. In Europe, there have only been four studies-all descriptive-of the impact of COVID-19 on students. These highlighted high levels of anxiety and depressive symptoms (Essadek & Rabeyron, 2020; Husky et al., 2020; Kaparounaki et al., 2020; Odriozola-González et al., 2020). To date, the factors involved in the expression of these symptoms during lockdown have mostly been overlooked. More specifically, no studies have evaluated the reciprocal influences of situational factors (e.g., COVID-19 symptoms, health concerns), and psychological factors (e.g., coping, implicit theory of emotions), on students' emotional states (e.g., anxiety, depression, hassles). The purpose of the present study was thus to assess the emotional state of French university students during lockdown, and the psychological and situational factors involved. More specifically, the objectives of the present study were to (a) investigate the links between

situational factors and students' emotional state, (b) investigate the links between psychological factors and their emotional state, and (c) explore the possible reciprocal influence of dispositional and situational factors on their emotional state.

2. Method

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

2.1. Sample and procedure

Data were collected anonymously between 23 April and 8 May (i.e., after 1 month of lockdown), via an online survey designed with Qualtrics software. A link to the survey was sent by e-mail to teachers in different faculties (Sciences, Sport Sciences, Psychology, Art, etc.) at four French universities (Nimes, Lorraine, Strasbourg, and Catholic University of the West), who forwarded it to their students, using either the university's mailing lists or the students'

social media groups. The link to the survey was also distributed via social media by the students who responded, thereby allowing students from other universities to respond. A total of 2153 people initially clicked on the link, but 11 refused to give their consent and were therefore not given access to the survey. Another 11 were excluded, because they mentioned that they were not students, while 833 people started but did not complete all the questionnaire. Finally, one person was excluded from the study by the investigators because of inconsistent responses. Our final sample therefore comprised 1297 students from different French universities, mainly in the south east, north west and north east (region particularly badly affected by COVID-19). Participants were aged 17-75 years (M = 21.27, SD = 4.72). Participants' characteristics are set out in Table 1. About 38.32% of students reported that, on average, more than two relatives (M = 2.43, SD = 2.08) in their environment had exhibited COVID-19 symptoms.

INSERT TABLE 1 ABOUT HERE

2.2. Measures

2.2.1. Hassles

We measured students' level of hassles with items taken from the Revised University Student Hassles Scale (RUSHS; Pett & Johnson, 2005). We included items (N = 22) belonging to the Time pressures, Friendships, and Physical appearance subscales in the questionnaire, as these appeared to be the most relevant in the specific context of lockdown. Participants indicated the severity of each type of hassle since lockdown on a 5-point scale ranging from 1 ($Not\ at\ all\ severe$) to 5 ($Extremely\ severe$). Higher scores reflected higher levels of hassles. Cronbach's alphas were satisfactory: .89 for Time pressures, .74 for Friendships, .80 for Physical appearance, and .88 for Total hassles.

2.2.2. Anxiety and depressive symptoms

Anxiety and depressive symptoms were assessed using a French version of the Hospital Anxiety and Depression Scale (HADS; Lepine et al., 1985). This 14-item self-report questionnaire assesses anxiety symptoms (7 items) and depressive symptoms (7 items) with

labels varying from one item to the next. Scores range from 0 to 21 for each dimension, with higher scores reflecting higher levels of anxiety or depressive symptoms.

2.2.3. Implicit theory of emotions

Students' implicit theory of emotion was assessed using a French translation of the Implicit Theories of Emotion Scale (Tamir et al., 2007), based on the procedure used by Vallerand (1989). This 4-item self-report measure assesses dynamic implicit theory (2 items; r = .50, p < .001) and static implicit theory (2 items; r = .68, p < .001). Scores on the latter were reversed to calculate a composite dynamic implicit emotion score ($\alpha = .80$). Participants answered on a 6-point Likert scale ranging from 1 (*Strongly disagree*) to 6 (*Strongly agree*).

2.2.4. Strategies used to cope with lockdown

Coping strategies were assessed using a French version of the situational version of the Brief-COPE (Muller & Spitz, 2003). Students were instructed to refer to a stressful situation related to lockdown. This self-report scale assesses 14 coping strategies (2 items per strategy): active coping, planning, instrumental support, use of emotional support, venting, behavioral disengagement, self-distraction, self-blame, positive reframing, humor, denial, acceptance, religion, and substance use. Participants rated each of the 28 items on a 4-point Likert scale ranging from *Never* to *Always*. Higher scores reflected higher levels of strategy use.

2.2.5. Situational factors

We considered six situational factors: (a) the extent to which participants felt that lockdown was compromising their future job prospects (scale ranging from 0 to 100); (b) the extent to which university studies were essential to participants (scale ranging from 0 to 100); (c) the presence or absence of COVID symptoms; (d) the presence or absence of COVID symptoms in their relatives; (e) participants' level of concern about their health owing to the COVID-19 crisis (scale ranging from 0 to 100); and (f) participants' level of concern about their relatives' health owing to the COVID-19 crisis (scale ranging from 0 to 100).

2.3. Statistical analyses

First, correlations were calculated to examine the associations between all numerical variables (i.e., essential nature of university studies, belief that lockdown was compromising students' future job prospects, concerns about students' own health and that of their relatives, anxiety and depressive symptoms, hassles, implicit theory of emotion, and coping strategies). Second, mean comparisons (with Student's *t* tests or analyses of variance, ANOVAs) were conducted to examine differences in psychological experiences as a function of categorical variables (i.e., paid work activity, lockdown with parents, COVID-19 symptoms). Finally, a network analysis based on partial correlations was carried out to study all the relationships.

3. Results

3.1. Links between emotional state and situational factors

Our first objective was to investigate the links between the situational factors and hassles, anxiety, and depressive symptoms. Results showed that 42.48% of participants had possible depressive symptoms (HADS-D \geq 8), and 33.38% had probable anxiety (HADS-A \geq 11). Our correlational analyses (Table 2) showed that the more university students worried about their health, the more they worried about the health of their loved ones (r = .50, p < .001). In addition, these two variables were positively associated with hassles, depressive symptoms, and above all anxiety (r = .37, p < .001 and r = .34, p < .001). Similarly, the more university students considered that lockdown was compromising their future job prospects, the higher their levels of hassles, anxiety, and depressive symptoms. However, the notion that university studies are essential was not related to psychological issues, and was only weakly related to the belief that lockdown was compromising students' future job prospects. Finally, university students who had symptoms of COVID-19, and/or who had relatives with COVID-19 symptoms, had higher levels of hassles, anxiety and depressive symptoms, compared with participants who had no symptoms (Table 3). However, they were no more concerned about either their own health

or that of their relatives. In addition, differences in emotional state were not explained either by paid work activity or by lockdown with parents.

INSERT TABLES 2 & 3 ABOUT HERE

3.2. Links between emotional state and psychological factors

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

Our second objective was to evaluate the associations between psychological factors (coping strategies and implicit theory of emotion) and students' emotional state. The first observation yielded by the descriptive data (Table 2) was that participants had quite low mean levels of use of coping strategies, except for acceptance and positive reframing. The five coping strategies most commonly used by our participants were acceptance, positive reframing, planning, self-distraction, and venting. Our correlational analyses (Table 2) showed that hassles, anxiety, and depressive symptoms were related to all the coping strategies, except for religion, venting, and self-distraction. More specifically, hassles, anxiety, and depressive symptoms were positively correlated with behavioral disengagement (r = .30, r = .32, and r = .30). .41; ps < .001), self-blame (r = .46, r = .51, and r = .44; ps < .001), and were negatively correlated with acceptance (r = -.30, r = -.41, and r = -.46; ps < .001), and positive reframing (r = -.25, r = -.37, and r = -.50; ps < .001). Similar associations were observed between these strategies and implicit theory of emotion (behavioral disengagement: r = -.22; self-blame: r = -.19; acceptance: r = .23; and positive reframing r = .32; all ps < .001). In other words, participants who above all endorsed an entity theory of emotion (i.e., belief that even with effort, it is impossible to change the emotions one feels), exhibited poorer adjustment to lockdown and higher levels of stress, anxiety, and depressive symptoms.

3.3. Reciprocal influence of dispositional and situational factors on emotional state

Finally, our third objective was to explore the reciprocal influence of dispositional and situational factors on students' emotional state. A network analysis based on partial correlations (Fig. 1) highlighted the centrality of anxiety and depressive symptoms in the psychological

functioning of university students during lockdown. Our results indicated that the belief that lockdown was compromising their future job prospects was related to more hassles, while concerns about their own and relatives' health were related to anxiety. As expected, implicit theory of emotion was indirectly associated with depressive symptoms, via positive reframing.

INSERT FIGURE 1 ABOUT HERE

4. Discussion

According to UNESCO, on 1 April 2020, schools and universities were closed in 185 countries around the world, affecting 1 542 412 000 learners (i.e., nearly 89.4% of the world's enrolled learners). In France, a total lockdown was imposed on 16 March, and all universities were closed. On 18 May, the lockdown started to be lifted, but French universities remained closed to students until September 2020. Unfortunately, even if the virus is partially under control, it is likely to have a lasting impact on universities. Since universities reopened, social distancing rules and the wearing of masks have led to many changes in the way they operate. To reduce the number of students on campus, hybrid learning or e-learning is being heavily promoted. In addition, the threat of a further spread of the pandemic and full closure of universities continues to hang over students. By 14 September, several French universities (e.g., Montpellier School of Medicine, Political Science Institute in Reims) had already had to close again, owing to the high prevalence of the virus (*Le Monde*¹). In this context, identifying the emotional state of French university students during the spring lockdown, as well as the psychological and situational factors involved, could help the authorities anticipate potential difficulties during the 2020-2021 academic year.

¹ Le Monde (15 September 2020). Covid-19: pour les étudiants, rentrée universitaire, premiers clusters et séances de sensibilisation.

Our results showed that during lockdown, 42.48% of our participants displayed possible depressive symptoms, whereas depressive symptoms usually affect about 30% of students in France (Boujut et al., 2009). This also seems to have been the case in other countries. For instance, a rate of 31.9% was found in Nigerian students during their lockdown (Ojewale, 2020), whereas the pre-COVID rate was 12.7% (Seun-Fadipe & Mosaku, 2017). Concerning anxiety, our results showed that during lockdown, 33.38% of our participants had probable anxiety. This is also a higher rate than the one that is usually observed for students in France (24.2%; Balayssac et al., 2018). In addition, this rate is higher than the rate (22.5%) for French nonstudent youth (18-24 years) 2 weeks after the start of lockdown (Chan-Chee et al., 2020). These results reflect the high levels of anxiety among French university students during lockdown. Taken together, our results suggest that students, known to constitute a vulnerable population (see systematic reviews by Ibrahim et al., 2013; de Paula et al., 2020), could become even more vulnerable during the 2020-2021 academic year, meaning that universities will have to be extremely vigilant. This vigilance will be all the more necessary, given that studies that have tracked the long-term consequences of previous pandemics suggest that psychological problems (including anxiety and depressive symptoms) may persist for months and even years after the event (e.g., Bonanno et al., 2008; Lee et al., 2007; Liu et al., 2012).

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

Our results showed that anxiety and depressive symptoms were strongly related to the types of coping strategies used during lockdown, especially avoidance strategies (i.e., self-distraction, denial, substance use, behavioral disengagement, venting, and self-blaming). This finding is consistent with studies conducted during the lockdown in the United Kingdom (Dawson & Golijani-Moghaddam, 2020), but also prior to the pandemic (Garnefski et al., 2002; Whatley et al., 1998). In addition, it is interesting to note that participants' mean scores were quite low for most of the coping strategies we measured. According to Dawson and Golijani-Moghaddam (2020), the isolating and restrictive pandemic context may mean that some individuals can no longer successfully use their normal ways of coping. It may be that our

participants had difficulty finding ways to cope with lockdown, and therefore resorted less to coping strategies. However, it is also important to point out that approach coping, and more specifically acceptance and positive reframing, were the strategies our participants used the most, and these strategies played a generally protective role against anxiety and depressive symptoms. In summary, our study showed that university students had difficulty diversifying their coping strategies during lockdown, but were nonetheless able to implement several effective strategies, such as positive reframing, planning, and acceptance. It is possible that those who had difficulty implementing these strategies were also those who expressed the highest levels of anxiety and depressive symptoms.

Beyond the importance of selecting and using effective coping strategies for reducing the anxiety associated with the pandemic and lockdown, our study highlighted the importance of students' beliefs about the capacity for emotional regulation, and more specifically their implicit theory of emotions. Our results showed that the more students adhered to an implicit static theory of emotions (i.e. belief that even with effort, it is impossible to change the emotions they feel), the poorer their adjustment to lockdown and the higher their levels of hassles, anxiety and depressive symptoms. The links between students' adjustment and their implicit theory of emotion are consistent with previous studies (e.g., Tamir et al., 2007), and support the idea that interventions aimed at developing students' emotion regulation skills could help to prevent them developing anxiety and depressive symptoms, and improve their ability to cope with stress. To this end, it seems appropriate to encourage their adhesion to a dynamic theory of emotion.

Our results also showed that situational variables played a central role in students' emotional state during lockdown. More specifically, the belief that lockdown was compromising their future prospects was positively associated with hassles, anxiety, and depressive symptoms. This finding is consistent with a previous study (Hasan & Bao, 2020) that showed that fear of losing an academic year was the concern that most exacerbated students' psychological anxiety in Bangladesh during lockdown. In the same way, our results

showed that the more students worried about their own health that of their loved ones, the higher their levels of hassles, anxiety and depressive symptoms. All these associations can be explained by the uncertainty (about health and the future) generated by COVID-19, a central concept in the fear of this virus (Schimmenti et al., 2020). In support of this argument, Satici et al. (2020) demonstrated that the inability to tolerate uncertainty in the COVID-19 pandemic can trigger fear of virus and impact negatively on psychological wellbeing.

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

COVID-19 has brought about a digital revolution in higher education (e.g., use of videoconferencing, digital open books, online exams, interaction in virtual environments; Strielkowski, 2020). Although e-learning was essential for maintaining education during lockdown, it was not without consequences (e.g., high levels of stress and strain), as illustrated by the global survey of the International Association of Universities (IAU, 2020). According to Baticulon et al. (2020), there can be a number of obstacles to e-learning, including technological, personal, family, institutional, or community barriers. In their study, only 41% of students in the Philippines thought they could physically and mentally participate in this type of education. In addition, 25% of college students in China suffered from severe anxiety due to e-learning crack-up (Cao et al., 2020), and anxiety also appears to be related to a lack of enjoyment in the classroom (Dewaele et al., 2019). Despite the reopening of most universities around the world in September 2020, social distancing rules impose the use of distance and/or hybrid learning, for which students are poorly prepared (especially first-year undergraduates). Some students may be overwhelmed by the increased flow of information and the lack of a regular schedule, which can lead to dropout and university failure. This context may well increase hassles about future job prospects, especially among students for whom university education is essential, further increasing their levels of hassles stress and anxiety. All this suggests that university students, who are already known to be a vulnerable population, may be particularly at risk during the 2020-2021 academic year.

It is therefore essential that universities' preventive medicine departments provide adequate psychological support for students. Based on the results of our partial correlation-based network analysis, psychological interventions aimed at reducing depressive symptoms could focus on promoting the use of positive reframing and acceptance, as well as discouraging the use of behavioral disengagement. To reduce hassles and anxiety, it also seems appropriate to limit the use of self-blame, and to analyze students' concerns about their future job prospects, their health, and that of their loved ones. To limit contact and improve coping strategies, *online self-help* interventions (e.g., web page, Facebook page, e-learning) could be used more. A pilot study conducted among first-year medical students showed that a stress management intervention based exclusively on Facebook (including educational, information and stress management resources based primarily on original videos) improved their coping strategies (George et al., 2013). As a precautionary measure, at the beginning of the academic year, universities should provide information about COVID-19 and its risks, hybrid education, hassles, anxiety and depressive symptoms, and how to manage it all.

The present results need to be interpreted with caution. First, although our sample size was large, we cannot be sure that it was representative of all students in France. Although our survey was made available to students in several universities, there was a low response rate-possibly because students were more interested in learning about their courses, and less available to take part in a survey. Nevertheless, students from different subjects and different levels of education were included, leading us to believe that our results reflect French students' emotional state during lockdown rather well. Second, these results exclusively concerned French university students, thus limiting their generalizability. However, the few studies conducted in other countries thus far have also reported increased anxiety and depressive symptoms among university students during lockdown. The impact of fear of COVID-19, academic failure and coping strategies on these symptoms has also been highlighted in research

in other countries. These data suggest that our results could also inform the dynamics in other countries, although further research is warranted at this stage. Third, the cross-sectional design of our study did not allow us to state that our participants' symptoms were lower before lockdown. Based on previous studies demonstrating that the psychological consequences of pandemics can last for several months, it is essential for future studies to conduct longitudinal assessments of students' emotional state throughout the 2020-2021 academic year.

5. Conclusion

In conclusion, our study highlighted high levels of hassle, probable anxiety, and/or probable depressive symptoms among university students during France's COVID-19 lockdown. Results showed that the students most at risk were those who thought that lockdown would have a detrimental effect on their future job prospects, worried about their health and that of their loved ones, believed that emotions are fixed, favored the use of avoidance strategies (e.g., behavioral disengagement, denial), and made little use of positive reframing or acceptance. These results are all the more important given that the threat of a so-called second wave and closure of the universities, and even a new lockdown, may maintain or even increase students' concerns for their health and that of their loved ones, as well as their anxiety.

320

321

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

6. References

- 322 Auerbach, R. P., Mortier, P., Bruffaerts, R., Alonso, J., Benjet, C., Cuijpers, P., 323 Demyttenaere, K., Ebert, D. D., Green, J. G., Hasking, P., Murray, E., Nock, M. K., Pinder-Amaker, S., Sampson, N. A., Stein, D. J., Vilagut, G., Zaslavsky, A. M., & 324 325 Kessler, R. C. (2018). WHO world mental health surveys international college student 326 project: Prevalence and distribution of mental disorders. Journal of Abnormal 327 Psychology, 127(7), 623–638. https://doi.org/10.1037/abn0000362 328 Balayssac, D., Pereira, B., Darfeuille, M., Cuq, P., Vernhet, L., Collin, A., Vennat, B., & Authier, N. (2018). Use of psychotropic medications and illegal drugs, and related 329 330 consequences among French pharmacy students - SCEP study: A nationwide cross-331 sectional study. Frontiers in Pharmacology, 9(July), 1–12. https://doi.org/10.3389/fphar.2018.00725
- 332
- 333 Baticulon, R. E., Rose Alberto, N. I., Beatriz Baron, M. C., Earl Mabulay, R. C., Gabriel

- Rizada, L. T., Jenkin Sy, J., Jan Tiu, C. S., Clarion, C. A., & Carlo Reyes, J. B. (2020).
 Barriers to online learning in the time of COVID-19: A national survey of medical students in the Philippines. *MedRxiv*, 2020.07.16.20155747.
- 337 https://doi.org/10.1101/2020.07.16.20155747

362

363

364

365

366

367

368

369

370

- Bonanno, G., Ho, S. M., Wong, C. P., Chan, J. C., Kwong, R. S., Cheung, C. K., & Wong, V. C. (2008). Psychological resilience and dysfunction among hospitalized survivors of the SARS epidemic in Hong Kong: A latent class approach. *Health Psychology*, *27*(5), 659-667. https://doi.org/10.1037/0278-6133.27.5.659
- Boujut, E., Koleck, M., Bruchon-Schweitzer, M., & Bourgeois, M. L. (2009). Mental health among students: A study among a cohort of freshmen. *Annales Medico-Psychologiques*, 167(9), 662–668. https://doi.org/10.1016/j.amp.2008.05.020
- Bunevicius, A., Katkute, A., & Bunevicius, R. (2008). Symptoms of anxiety and depression in medical students and in humanities students: Relationship with big-five personality dimensions and vulnerability to stress. *International Journal of Social Psychiatry*, 54(6), 494–501. https://doi.org/10.1177/0020764008090843
- Cao, W., Fang, Z., Hou, G., Han, M., Xu, X., Dong, J., & Zheng, J. (2020). The psychological
 impact of the COVID-19 epidemic on college students in China. *Psychiatry Research*,
 287. https://doi.org/10.1016/j.psychres.2020.112934
- Chan-Chee, C., Léon, C., Lasbeur, L., Lecrique, J., Raude, J., Arwidson, P., & du Roscoät, E. (2020). La santé mentale des Français face au Covid-19: Prévalences, évolutions et déterminants de l'anxiété au cours des deux premières semaines de confinement (Enquête CoviPrev 23-25 mars et 30 mars-1er avril 2020). Bulletin Épidémiologique Hebdomadaire, 13, 260–269.
- Dawson, D. L., & Golijani-Moghaddam, N. (2020). COVID-19: Psychological flexibility, coping, mental health, and wellbeing in the UK during the pandemic. *Journal of Contextual Behavioral Science*, 17(June), 126–134. https://doi.org/10.1016/j.jcbs.2020.07.010
 - de Castella, K., Goldin, P., Jazaieri, H., Ziv, M., Heimberg, R. G., & Gross, J. J. (2014). Emotion beliefs in social anxiety disorder: Associations with stress, anxiety, and wellbeing. *Australian Journal of Psychology*, 66(2), 139–148. https://doi.org/10.1111/ajpy.12053
 - de Paula, W., Breguez, G. S., Machado, E. L., & Meireles, A. L. (2020). Prevalence of anxiety, depression, and suicidal ideation symptoms among university students: A systematic review. *Brazilian Journal of Health Review*, *3*(4), 8739–8756. https://doi.org/10.34119/bjhrv3n4-119
 - Dewaele, J., Magdalena, A. F., & Saito, K. (2019). The effect of perception of teacher characteristics on Spanish EFL learners' anxiety and enjoyment. *Modern Language Journal*, 103(2), 412–427. https://doi.org/10.1111/modl.12555
- Essadek, A., & Rabeyron, T. (2020). Mental health of French students during the Covid-19 pandemic. *Journal of Affective Disorders*.

 https://doi.org/https://doi.org/10.1016/j.jad.2020.08.042
- Garnefski, N., Legerstee, J., Kraaij, V., Van Den Kommer, T., & Teerds, J. (2002). Cognitive coping strategies and symptoms of depression and anxiety: A comparison between adolescents and adults. *Journal of Adolescence*. https://doi.org/10.1006/jado.2002.0507
- George, D. R., Dellasega, C., Whitehead, M. M., & Bordon, A. (2013). Facebook-based stress management resources for first-year medical students: A multi-method evaluation.

 Computers in Human Behavior. https://doi.org/10.1016/j.chb.2012.12.008
- Hasan, N., & Bao, Y. (2020). Impact of "e-Learning crack-up" perception on psychological distress among college students during COVID-19 pandemic: A mediating role of "fear of academic year loss". *Children and Youth Services Review*, 118(July), 105355.
- 384 https://doi.org/10.1016/j.childyouth.2020.105355

- Husky, M. M., Kovess-Masfety, V., & Swendsen, J. D. (2020). Stress and anxiety among university students in France during Covid-19 mandatory confinement. *Comprehensive Psychiatry*, *102*, 152191. https://doi.org/10.1016/j.comppsych.2020.152191
- 388 IAU. (2020). Covid-19: Higher education challenges and responses. The Global Voice of 389 Higher Education. https://www.iau-aiu.net/COVID-19-Higher-Education-challenges-390 and-responses
- Ibrahim, A. K., Kelly, S. J., Adams, C. E., & Glazebrook, C. (2013). A systematic review of
 studies of depression prevalence in university students. *Journal of Psychiatric Research*, 47(3), 391–400. https://doi.org/10.1016/j.jpsychires.2012.11.015
- Kaparounaki, C. K., Patsali, M. E., Mousa, D.-P. V, Papadopoulou, E. V. K., Papadopoulou,
 K. K. K., & Fountoulakis, K. N. (2020). University students' mental health amidst the
 COVID-19 quarantine in Greece. *Psychiatry Research*, *290*, 113111.
 https://doi.org/10.1016/j.psychres.2020.113111
- Lee, A. M., Wong, J. G. W. S., McAlonan, G. M., Cheung, V., Cheung, C., Sham, P. C., Chu, N. M., Wong, P. C., Tsang, K. W. T., & Chua, S. E. (2007). Stress and psychological distress among SARS survivors 1 year after the outbreak. *Canadian Journal of Psychiatry*. https://doi.org/10.1177/070674370705200405
- 402 Lepine, J. P., Godchau, M., & Brun, P. (1985). Anxiety and depression in inpatients. In *The Lancet*. https://doi.org/10.1016/S0140-6736(85)92589-9
- Liu, X., Kakade, M., Fuller, C. J., Fan, B., Fang, Y., Kong, J., Guan, Z., & Wu, P. (2012).

 Depression after exposure to stressful events: Lessons learned from the severe acute respiratory syndrome epidemic. *Comprehensive Psychiatry*.

 https://doi.org/10.1016/j.comppsych.2011.02.003

- Matsudaira, T., Igarashi, H., Kikuchi, H., Kano, R., Mitoma, H., Ohuchi, K., & Kitamura, T. (2009). Factor structure of the Hospital Anxiety and Depression Scale in Japanese psychiatric outpatient and student populations. *Health and Quality of Life Outcomes*, 7, 1–9. https://doi.org/10.1186/1477-7525-7-42
 - Muller, L., & Spitz, E. (2003). Évaluation multidimensionnelle du coping: Validation du Brief COPE sur une population française. *Encephale*, *29*(6 I), 507–518.
- Odriozola-González, P., Planchuelo-Gómez, Á., Irurtia, M. J., & de Luis-García, R. (2020).
 Psychological effects of the COVID-19 outbreak and lockdown among students and workers of a Spanish university. *Psychiatry Research*, 290, 113108.
 https://doi.org/10.1016/j.psychres.2020.113108
- Ojewale, L. Y. (2020). Psychological state and family functioning of University of Ibadan students during the COVID-19 lockdown. https://doi.org/10.1101/2020.07.09.20149997
- Pett, M. A., & Johnson, M. J. M. (2005). Development and psychometric evaluation of the
 Revised University Student Hassles Scale. *Educational and Psychological Measurement*, 65(6), 984–1010. https://doi.org/10.1177/0013164405275661
- 423 Réveillère, C., Nandrino, J. ., Sailly, F., Mercier, C., & Moreel, V. (2001). Étude des tracas 424 quotidiens des étudiants: Liens avec la santé perçue. *Annales Médico-Psychologiques*, 425 *Revue Psychiatrique*, 159(6), 460–465. https://doi.org/10.1016/S0003-4487(01)00070-1
- Sarmento, M. (2015). A "mental health profile" of higher education students. *Procedia* Social and Behavioral Sciences, 191, 12–20.
 https://doi.org/10.1016/j.sbspro.2015.04.606
- Satici, B., Saricali, M., Satici, S. A., & Griffiths, M. D. (2020). Intolerance of uncertainty and mental wellbeing: Serial mediation by rumination and fear of COVID-19. *International Journal of Mental Health and Addiction*. https://doi.org/10.1007/s11469-020-00305-0
- Schimmenti, A., Billieux, J., & Starcevic, V. (2020). The four horsemen of fear during the COVID pandemic. *Clinical Neuropsychiatry*, *17*(2), 41–45.
- Seun-Fadipe, C. T., & Mosaku, K. S. (2017). Sleep quality and psychological distress among undergraduate students of a Nigerian university. *Sleep Health*, *3*(3), 190–194.

436	https://doi.org/10.1016/j.sleh.2017.02.004
437	Stanley, N., & Manthorpe, J. (2001). Responding to students' mental health needs:
438	Impermeable systems and diverse users. <i>Journal of Mental Health</i> , 10(1), 41–52.
439	Strielkowski, W. (2020). COVID-19 pandemic and the digital revolution in academia and
440	higher education. Preprints. https://doi.org/10.20944/preprints202004.0290.v1
441	Tamir, M., John, O. P., Srivastava, S., & Gross, J. J. (2007). Implicit theories of emotion:
442	Affective and social outcomes across a major life transition. Journal of Personality and
443	Social Psychology, 92, 731–744.
444	Vallerand, R. J. (1989). Vers une méthodologie de validation transculturelle de questionnaires
445	psychologiques: Implications pour la recherche en langue française. Revue Canadienne
446	de Psychologie, 30, 662–680.
447	Whatley, S. L., Foreman, A. C., & Richards, S. (1998). The relationship of coping style to
448	dysphoria, anxiety, and anger. Psychological Reports.
449	https://doi.org/10.2466/pr0.1998.83.3.783
450	

Table 1 Characteristics of survey respondents (N = 1297)

Characteristics	Number	%
Sex		
Female	1009	77.79
Male	268	20.66
Other	20	1.54
University		
Nîmes	560	43.14
Lorraine	370	28.51
Strasbourg	212	16.33
UCO Angers	86	6.63
Others	70	5.39
Level		
Undergraduate	1130	86.86
First year	488	37.51
Second year	314	24.14
Third year	328	25.21
Master's	156	11.99
Fourth year	82	6.30
Fifth year	74	5.69
PhD	11	0.85
Undefined	4	0.31
Subject	·	0.01
Psychology	473	36.84
Sports science	203	15.81
Languages	159	12.38
History/Geography	99	7.71
Sciences	82	6.39
Art/Design	56	4.36
Law/Economics/Management	48	3.74
French Literature	39	3.04
Lockdown with parents	33	3.04
Yes	914	70.25
No	387	29.75
Paid work activity during lockdown	367	29.73
Yes	219	16.86
Contact with the public	45	13.42
	43 174	3.47
No contact with the public No		
	1078	83.11
Positive COVID-19 test or symptoms	177	12.00
Yes	177	13.60
No	1124	86.40
Relative with positive COVID-19 test or		
symptoms	407	20.20
Yes	497	38.20
No	804	61.80
Anxiety	506	45.05
No symptoms (≤ 7)	596	45.95
Possible anxiety (8-10)	268	20.66
Probable anxiety (≥ 11)	433	33.38
Depression		
No symptoms (≤ 7)	759	58.52
Possible depression (8-10)	275	21.20
Probable depression (≥ 11)	263	20.28

Table 2 Descriptive data (mean, standard deviation, minimum and maximum) and correlations between variables of interest

	Mean (SD)	Min-	1	2	3	4	5	6	7	8	
		Max	r p	r p	r p	r p	r p	r p	r p	r p	
1. Essential nature of university studies	79.50 (20.60)	0-100	_								
Belief that lockdown is compromising future job prospects	55.77 (30.28)	0-100	.09 .002								
3. Concerns about own health	32.14 (27.92)	0-100	.17 < .001	.18 < .001							
4. Concerns about relatives' health	67.84 (27.15)	0-100	.17 < .001	.21 < .001	.50 < .001						
5. Anxiety	8.76 (4.69)	0-21	.08 .003	.27 < .001	.37 < .001	.34 < .001					
6. Depressive symptoms	7.01 (4.02)	0-19	.00 .892	.31 < .001	.19 < .001	.15 < .001	.60 < .001				
7. Hassles	3.09 (.77)	1-5	.08 .006	.34 < .001	.24 < .001	.25 < .001	.57 < .001	.54 < .001			
8. Implicit theories of emotions	15.08 (4.46)	4-24	.06 .021	05 .09	03 .33	06 .024	21 < .001	30 < .001	17 < .001		
Coping											
9. Venting	4.24 (1.74)	2-8	.04 .157	.01 .72	.12 < .001	.08 .004	.08 .006	08 .005	.05 .065	.10 < .001	
10. Using emotional support	4.02 (1.81)	2-8	.06 .04	.13 < .001	.23 < .001	.19 < .001	.37 < .001	.19 < .001	.26 < .001	05 .06	
11. Using instrumental support	3.77 (1.68)	2-8	.06 .038	.06 .044	.23 < .001	.18 < .001	.22 < .001	.05 .087	.16 < .001	.02 .502	
12. Planning	4.48 (1.78)	2-8	.14 < .001	09 < .001	.04 .168	.03 .358	05 .076	20 < .001	06 .04	.18 < .001	
13. Active coping	3.75 (1.51)	2-8	.16 < .001	08 .004	.08 .006	.06 .031	05 .071	25 < .001	09 < .001	.19 < .001	
14. Positive reframing	5.23 (1.81)	2-8	.04 .137	21 < .001	16 < .001	09 .001	37 < .001	50 < .001	25 < .001	.32 < .001	
15. Humor	3.69 (1.72)	2-8	09 .002	12 < .001	24 < .001	20 < .001	31 < .001	25 < .001	16 < .001	.19 < .001	
16. Acceptance	6.13 (1.65)	2-8	.01 .709	26 < .001	21 < .001	16 < .001	41 < .001	46 < .001	30 < .001	.23 < .001	
17. Denial	3.02 (1.37)	2-8	.05 .099	.17 < .001	.16 < .001	.13 < .001	.27 < .001	.28 < .001	.27 < .001	12 < .001	
18. Self-blame	3.85 (1.71)	2-8	.00 .958	.23 < .001	.16 < .001	.17 < .001	.51 < .001	.44 < .001	.46 < .001	19 < .001	
19. Behavioral disengagement	3.76 (1.59)	2-8	08 .002	.15 < .001	.08 .004	.04 .112	.32 < .001	.41 < .001	.30 < .001	22 < .001	
20. Substance use	2.62 (1.39)	2-8	07 .011	.17 < .001	.07 .008	.04 .15	.24 < .001	.23 < .001	.21 < .001	04 .162	
21. Self-distraction	4.82 (1.60)	2-8	.10 < .001	.00 .972	.11 < .001	.13 < .001	.10 < .001	06 .04	.07 .013	.05 .083	
22. Religion	2.82 (1.52)	2-8	02 .428	03 .324	.02 .387	.00 .928	.03 .26	02 .398	.03 .215	.15 < .001	

454

Table 3. Means comparisons (standard deviation) of psychological experience of lockdown as a function of living conditions (paid work

456 activity, lockdown with parents, symptoms and university)


	D	£:	l et de						COVID-19 symptoms													
	Professional activity			Lockdown with parents			Student				Relative				University							
	Yes	No	t	p	Yes	No	t	р	Yes	No	t	р	Yes	No	t	р	Strasbourg	Lorraine	Nîmes	UCO Angers	F	р
Lockdown is	57.76	55.36	1.07	.285	55.55	56.28	20	COF	60.85	54.96	-2.41 .016	57.22	54.86	4.27	172	56.18	55.19	55.41	57.44	4.6	022	
compromising my future job prospects	(28.11)	(3.70)	1.07		(29.55)	(31.98)	39 .69	.695	(32.54)	(29.84)		.016	(30.62)	(30.05)	-1.37	.172	(29.11)	(31.44)	(3.54)	(25.9)	.16	.922
Concerns about personal	31.71	32.22	25	225	31.9	32.69	47 .642	6.42	32.43	32.09	15 .883	001	30.67	33.05	1.49	127	31.08	31.97	33.39	28.14	1.05 .3	200
health	(26.60)	(28.20)	25	.805	(27.64)	(28.62)		.642	(29.53)	(27.68)		.881	(28.31)	(27.66)		.137	(27.13)	(29.47)	(28.23)	(23.22)		.368
Concerns about relatives'	69.97	67.40	1 20	.203	68.17	67.04	.68	.495	68.80	67.68	51 .611	611	69.13	67.03	-1.36	.175	67.5	69.17	68.59	60.86	2.32 .0	.073
health	(25.39)	(27.49)	1.28	.203	(26.74)	(28.13)		.495	(28.98)	(26.86)		.011	(27.31)	(27.04)		.1/3	(25.45)	(27.79)	(27.72)	(24.42)		.075
Anvioty	8.75	8.76	03 .	.979	8.63	9.06	1 5	-1.5 .133	9.72	8.61	-2.95 .	.003	9.31	8.42	-3.31	<.001	8.36	8.85	9.02	7.29	3.97 .	.008
Anxiety	(4.83)	(4.66)	03	.373	(4.61)	(4.86)	-1.5		(4.91)	(4.64)		.003	(4.87)	(4.54)		.51 <.001	(4.57)	(4.98)	(4.69)	(3.18)		.006
Depressive symptoms	7.05	7	.16	.876	6.97	7.09	46	.646	8.01	6.85	-3.59	<.001	7.44	6.74	-3.06	.002	6.39	7.38	7.01	6.69	2.94	.032
Depressive symptoms	(4.03)	(4.02)	.10	.670	(3.93)	(4.24)	40	.040	(4.38)	(3.94)		<.001	(4.28)	(3.83)	-3.00	.002	(3.93)	(4.33)	(.88)	(3.51)		.032
Hassles	3.14	3.08	1.16	.248	3.07	3.12	1 1 5	.252	3.20	3.07	-2.19	.028	3.17	3.03	-3.2	.2 .001	2.97	3.07	3.16	2.92	4.49	.004
Паззіез	(.77)	(.77)	1.10	.240	(.77)	(.78)	-1.15	.232	(.75)	(.77)	-2.19		(.77)	(.76)	-3.2		(.77)	(.80)	(.75)	(.71)		.004

457 458

Note. For the dichotomous variables, we ran Student's t tests for independent groups. For the university variable including four modalities, we ran

an ANOVA.

Figure 1: Partial correlations network



461

460

- 1 Lockdown is compromising my future job prospects
- 2 Concerns about personal health
- 3 Concerns about relatives' health
- 4 Anxiety symptoms
- 5 Depressive symptom
- 6 Hassles
- 7 Implicit theories dynamics of emotion
- 8 Coping: Acceptance
- 9 Coping: Positive reframing
- 10 Coping: Self-blame

11 Coping: Behavioural disengagement

462 463 464

Note. Solid lines correspond to positive partial correlations, and dashed lines to negative partial correlations. The thicker the line, the stronger the partial correlation.