



**HAL**  
open science

# Comparative studies of egg parasitoids of the pine processionary moth (*Thaumetopoea pityocampa*, Den. & Schiff.) in historic and expansion areas in France and Bulgaria

Georgi Georgiev, Jérôme Rousselet, Mathieu Laparie, Christelle Robinet, Margarita Georgieva, Gergana Zaemdzhikova, Alain Roques, Alexis Bernard, Laura Poitou, Maurane Buradino, et al.

## ► To cite this version:

Georgi Georgiev, Jérôme Rousselet, Mathieu Laparie, Christelle Robinet, Margarita Georgieva, et al.. Comparative studies of egg parasitoids of the pine processionary moth (*Thaumetopoea pityocampa*, Den. & Schiff.) in historic and expansion areas in France and Bulgaria. *Forestry*, 2021, 94 (2), pp.324-331. 10.1093/forestry/cpaa022 . hal-03123290

**HAL Id: hal-03123290**

**<https://hal.science/hal-03123290v1>**

Submitted on 9 Oct 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Comparative studies of egg parasitoids of the pine processionary moth (*Thaumetopoea pityocampa*, Den. & Schiff) in historic and expansion areas in France and Bulgaria

Georgi Georgiev<sup>1\*</sup>, Jérôme Rousselet<sup>2</sup>, Mathieu Laparie<sup>2</sup>, Christelle Robinet<sup>2</sup>, Margarita Georgieva<sup>1</sup>, Gergana Zaemdzhikova<sup>1</sup>, Alain Roques<sup>2</sup>, Alexis Bernard<sup>2</sup>, Laura Poitou<sup>2</sup>, Maurane Buradino<sup>3</sup>, Carole Kerdelhue<sup>4</sup>, Jean-Pierre Rossi<sup>4</sup>, Maria Matova<sup>1</sup>, Peter Boyadzhiev<sup>5</sup> and Plamen Mirchev<sup>1</sup>

<sup>1</sup> Forest Research Institute, Bulgarian Academy of Sciences, 1756 Sofia, Bulgaria

<sup>2</sup> L'Unité de Recherche de Zoologie Forestière (URZF), INRA, 45075 Orléans, France

<sup>3</sup> Entomologie et Forêt Méditerranéenne (UEFM), INRA, UE 0348, Avignon, France

<sup>4</sup> Centre de Biologie pour la Gestion des Populations (CBGP), INRA, 1062 UMR, Montpellier, France

<sup>5</sup> Department of Zoology, University of Plovdiv, 4000 Plovdiv, Bulgaria

\*Corresponding author E-mail: ggeorgiev.fri@gmail.com

To investigate enemy pressure across the range of the pine processionary moth (*Thaumetopoea pityocampa*, Den. & Schiff) as it spreads northwards in Europe because of climate change, a survey of 206 egg-batches from historical and newly colonized areas at both the westernmost and the easternmost ends of the front was carried out in 2016-2018. Out of them, 97 egg-batches were collected from nine locations in three regions in France, and 109 from four localities in two regions in Bulgaria, both within historical and newly colonized ranges of the pest. The average number of eggs per batch collected in Bulgaria ( $226.6 \pm 43.2$ ) was higher than that in France ( $194.3 \pm 50.1$ ). However, the hatching percentage was higher in French samples, varying from 69.8% to 95.7%, compared to a lower range between 49.8% and 85.2% in Bulgarian samples. Four primary parasitoids (*Ooencyrtus pityocampae*, *Baryscapus servadeii*, *Anastatus bifasciatus*, *Trichogramma* sp.) and a hyperparasitoid (*B. transversalis*, found only in two regions of the historical range in Bulgaria) were identified. The oligophagous species *B. servadeii* was present in all sites within the historical range of the pest. In newly colonized areas, the impact of primary parasitoids on the host was significantly lower suggesting that they lag behind the range expansion of pine processionary moth. In France, the most abundant species in the *T. pityocampa* parasitoid complex were *B. servadeii* in Ré Island and Orléans (97.3% and 87.4%, respectively), and *Trichogramma* sp. (99.7%) in a newly colonized locality in Fréhel. *Ooencyrtus pityocampae* prevailed in three out of four Bulgarian localities (72.1% in Sandanski, 89.7% in Maglizh and 65.7% in Sladak kladenets), whereas *B. servadeii* was predominant in Gega (75.4%). Mortality of *B. servadeii* and *O. pityocampae* caused by the hyperparasitoid *B. transversalis* amounted to 4.8-6.2%. The impact of predators on the pine processionary moth in the egg stage was negligible at most sites, reaching 12.5% in only one site (Maglizh).

**Key words:** *Thaumetopoea pityocampa*, range expansion, survival, egg parasitoids, France, Bulgaria

## Introduction

The pine processionary moth, *Thaumetopoea pityocampa* (Denis & Schiffmüller, 1775) (Lepidoptera: Notodontidae) is one of the most important forest pests in the Mediterranean region. Due to climate change, and particularly winter warming (Battisti *et al.*, 2005, 2006; Robinet *et al.*, 2007), an altitudinal and latitudinal expansion of its range has been observed over the last few decades (Hoch *et al.*, 2005; Imber, 2012; Mirchev *et al.*, 2011a; Netherer and Schopf, 2010;

Roques *et al.*, 2005; Toffolo *et al.*, 2006). In its current range, *T. pityocampa* lives in different habitats both in its historical range and newly colonized areas.

In France, five egg parasitoids of pine processionary moth have been recorded: *Baryscapus servadeii* (Domenichini, 1965) (Hymenoptera: Eulophidae), *Ooencyrtus pityocampae* (Mercet, 1921) (Hymenoptera: Encyrtidae), *Anastatus bifasciatus* (Geoffroy, 1785), *Eupelmus* sp. (Hymenoptera: Eupelmidae) and *Trichogramma* sp. (Hymenoptera: Trichogrammatidae) (Auger-Rozenberg *et al.*, 2015). In addition, *Baryscapus transversalis* Graham, 1991, *Pediobius bruchicida* (Rondani, 1872) (Hymenoptera: Eulophidae) and *Eupelmus (Macroneura) vesicularis* (Retzius, 1783) (Hymenoptera: Eulophidae), have been established in Bulgaria (Mirchev *et al.*, 2011b; Tsankov *et al.*, 1996a). The performance and distribution of the pine processionary moth and its insect enemies are highly dependent on climatic conditions. Higher temperatures in winter favour larval survival of the pine processionary moth (Battisti *et al.*, 2005), whereas summer temperatures above 32° C are considered lethal for the development of *T. pityocampa* eggs and young larvae (Robinet *et al.*, 2015). The climatic conditions in a particular biotope comprehensively influence the variety of egg parasitoid species as well. Indeed, Masutti (1964) reported that *O. pityocampae* is sensitive to high temperatures and biotopes with prolonged periods above 30° C are not suitable for its survival. In contrast, another hymenopteran species, *B. servadeii*, is more tolerant and develops successfully even at high temperatures. The parasitoid complex in newly colonized areas may also depend on the pace of the range expansion of *T. pityocampa*. The egg parasitoid assemblage of *T. pityocampa* is an indicator of two different types of pest expansion in new habitats: *i*) when *T. pityocampa* penetrates into new regions, relatively remote from the primary habitats where the specific *B. servadeii* is absent (Mirchev *et al.*, 2017); *ii*) when the pine processionary moth gradually enlarges its range in adjacent stands at the rate of 2-3 km per year, the host is followed by *B. servadeii* (Mirchev *et al.*, 2018).

The aim of this study was to compare the egg parasitoids complex of *T. pityocampa* in historic and expansion areas in France and Bulgaria, the westernmost and easternmost countries where the northward expansion of the pest is observed.

## Materials and methods

During the period 2016-2017, a total of 206 egg-batches of *T. pityocampa* were collected from France and Bulgaria between the end of August and November (Table 1). In Bulgaria, all samples were collected in Austrian pine (*Pinus nigra* Arn.) plantations. In France, samples were collected in *P. nigra*, maritime pine (*Pinus pinaster* Aiton) and Monterey pine (*Pinus radiata* D.Don) stands (Table 1).

### Table 1.

In France, 97 egg-batches were collected in nine localities from three regions corresponding to different colonization dates (Fig. 1):

- Ré Island, a small island in the Atlantic Ocean situated about 4 km away from mainland France, which is within the historical range (there is data from the 1970s);

- Orléans, in central France, which has been colonized in the 1990s;
- Fréhel, in Brittany, which has been newly colonised in the 2010s.

**Fig. 1.**

In Bulgaria, 109 egg-batches were collected in two ‘historical’ and two ‘newly colonized’ localities from two distinct regions (Fig. 2):

- Southwest Bulgaria - Struma valley (town of Sandanski) and Ograzhden Mt. (Gega vill.). In that area, the pine processionary moth has been present since 1949, considered historical range.
- Central Bulgaria - in the land of Sladak kladenets vill. In Sredna gora Mt. (the pest was first recorded in 2012) and in the region of Maglizh (newly colonized by *T. pityocampa* in 2015).

**Fig. 2.**

Upon collection, all egg batches were transported to the laboratory of entomology of Forest Research Institute in Sofia. The scales of egg-masses were removed, and the samples were analysed according to the protocol described in Tsankov *et al.* (1996). The egg-batches were put singly in test tubes, covered with cotton stoppers, and kept at room temperature (20-22° C).

During the laboratory studies, the samples were checked five times and the parasitoids that emerged were separated in plastic capsules for identification. During November-December 2018, the eggs without exit holes were dissected, and the meconia and remains of emerged or dead insects were determined under a stereomicroscope (40×). Parasitoids that emerged before collection were determined by their meconia and remains according to Schmidt and Kitt (1994), Tanzen and Schmidt (1995), Schmidt *et al.* (1997a) and Tsankov *et al.* (1996, 1998).

The data was analysed using descriptive statistic of MS Excel 2013. The Clopper-Pearson method (Hollander and Wolfe, 1973) was used to compare relative share of the specific *B. servadeii* in parasitoid assemblages of *T. pityocampa* in old and newly colonized areas.

## **Results**

### **Viability of the pine processionary moth in the egg stage**

The number of eggs in *T. pityocampa* batches varied from 185.7±47.7 to 232.2±43.5 in France and from 216.4±41.0 to 251.3±31.3 in Bulgaria (Table 2). In Bulgaria, the average number of eggs in a batch (226.6±43.2) was higher than that obtained from France (194.3±50.1). The highest average egg production per female (251.3±31.3) and the smallest dispersion between samples (204-299 eggs/batch) was recorded in Maglizh. In this locality, the proportion of hatched larvae was almost half (49.8%) of all analysed eggs. A significant share (41.5%) of parasitized and destroyed by predators eggs was registered. In the rest of the localities, the rate of parasitized and destroyed eggs varied between 10.3% (Sladak kladenets) and 19.2% (Sandanski).

The lowest average number of eggs per batch (185.7±47.7) and a large dispersion between samples (55-278) were recorded in the recently infested by the pine processionary moth region of Fréhel. The highest proportion of hatched larvae was established in this region (95.7%), and the rate of eggs parasitized and destroyed by predators was only 2.9%. In the rest of the French

localities the impact of parasitoids and predators reached up to 13.6% (Ré Island) and 24.5% (Orléans) (Table 2).

### **Table 2.**

Another factor that also affected the rate of successful hatching was the abortion of eggs, which varied between 1.4% (Fréhel) and 10.8% (Sandanski) (Table 2). The number of dead larvae in the eggs (without exit holes) varied between 0.7% and 2.5% in France, and between 2.1% and 4.7% in Bulgaria. The proportion of sterile, non-fertilized eggs recorded in all samples varied between 0.6% (Fréhel) and 5.0% (Sandanski).

The predatory effect in most localities was negligible (0.02-2.5%) with the exception of Maglizh in Bulgaria (12.5%) (Table 2).

### **Complex of egg parasitoids in different localities**

In this study, the egg parasitoids were the main factor causing pine processionary moth mortality. The mortality rate in France varied between 2.9% (Fréhel) and 24.4% (Orléans) (Table 2). In Bulgaria, the parasitoids affected between 7.8% (Sladak kladenets) and 29.0% (Maglizh) of host eggs.

Four primary egg parasitoids were identified in both France and Bulgaria: *Ooecyrtus pityocampae*, *Baryscapus servadeii*, *Anastatus bifasciatus*, *Trichogramma* sp. (Table 3).

### **Table 3.**

In a recently colonized area in France (Fréhel), only *Trichogramma* sp. and a single specimen of *O. pityocampae* represented the complex of parasitoids (Table 3). In areas of historical range (Orléans and Ré Island), *B. servadeii* was dominant with 84.7% and 97.3%, respectively. A minor presence of *O. pityocampae* and *Trichogramma* sp. was also recorded in them, whereas in Orléans, four specimens of *A. bifasciatus* were detected.

In the historical range of *T. pityocampa* in Bulgaria, the most numerous was the polyphagous *O. pityocampae* in Sandanski (72.1%) and the oligophagous *B. servadeii* in Gega (75.4%). In the newly colonized sites, the polyphagous parasitoids prevailed. In Sladak kladenets, only *O. pityocampae* and *A. bifasciatus* were identified, with a significant prevalence of the former species. In Maglizh, four primary parasitoids were reared from host eggs but the specific species *B. servadeii* occupied only 2.1% of the parasitoid complex.

In general, *B. servadeii* is common at all sites where the pine processionary moth has occurred for a long time (Ré Island, Orléans, Sandanski, Gega) and it is insignificant in all sites where the pest has penetrated recently (Fréhel, Sladak kladenets, Maglizh) (Table 3). This regularity of *B. servadeii* distribution has been confirmed in all studied areas. The Clopper-Pearson method determined the confidence interval [59, 100] of the percentage of cases in which the rule is confirmed with at a confidence level of 0.95. This suggests that the spread of *B. servadeii* is related to the old habitats of pine processionary moth and exceptions to this rule are rare (maximum 31%).

The hyperparasitoid *B. transversalis* was found only in Bulgaria at Sandanski and Gega, both within the historical range of the pest.

### **Bioecological characteristics of egg parasitoids**

A difference in the phenology of *O. pityocampae* was observed in France and Bulgaria. In samples from Ré Island and Orléans, most of the individuals emerged after collection of egg batches – 77.8% and 67.0%, respectively (Table 4). In contrast, in three Bulgarian localities (Gega, Sladak kladenets and Maglizh), the predominant part of *O. pityocampae* adults (60.6-73.6%) had emerged before collection of egg batches, and only in Sandanski, most of them (55.8%) emerged after collection. The proportion of established dead *O. pityocampae* adults in host's eggs was the lowest in Orléans (3.3%) whereas it ranged from 10.2 to 32.2% in Bulgarian localities. The average survival rate of the parasitoid in French and Bulgarian localities was 96.0% and 85.4%, respectively.

#### **Table 4.**

High average survival rate of *B. servadeii* 97.3% was recorded for both countries, and the mortality was low in both French and Bulgarian sites (1.5-3.4%). There was no difference in phenology for this parasitoid in both countries.

The polyphagous *A. bifasciatus* was established in one locality in France (Orléans) and all studied localities in Bulgaria (Table 3). In Bulgaria, the majority of adults emerged after hibernation.

The polyembryonic *Trichogramma* sp. was found in 9.5-13.2% of *T. pityocampa* egg batches from the historical range localities in France and Bulgaria, and in 38.5-39.1% from the newly colonized areas (Table 4). A high mortality (up to 70.0%) of adults was observed in the eggs. The hyperparasitoid *B. transversalis* parasitized 4.8-6.2% of *B. servadeii* and *O. pityocampae*. A high mortality (16.7-21.4%) of the hyperparasitoid was recorded in Sandanski and Gega (Table 4).

In this study, all emerging adults of *B. servadeii* and *O. pityocampae* were female, unlike *A. bifasciatus* where the specimens recorded were only male.

### **Discussion**

Female moths of *T. pityocampa* usually lay only one egg batch (Douma-Petridou, 1990), and therefore the average number of eggs in batches corresponds to the fertility of the species. The average number of eggs per batch collected in Bulgaria (226.6) was higher than that collected in France (194.3). The fertility of *T. pityocampa* depends on climatic conditions, host plant quality and population dynamics (Masutti and Battisti, 1990). It is higher in the northern parts of its range (Mirchev, 2005) and increases with altitude (Özkazans, 1987; Tiberi and Roversi, 1987). In contrast, Masutti and Battisti (1990) reported lower fertility in colder areas. The biochemical composition of the host plant is one of the significant environmental factors affecting *T. pityocampa* (Carvalho *et al.*, 1999; Tiberi *et al.*, 1999, Moura *et al.*, 1999; Roussis *et al.*, 1994). In mixed stands of *Pinus radiata* and *Pinus pinaster*, the pine processionary moth prefers the former

species on which caterpillars grow faster, reach a larger size and pupate earlier (Buxton, 1990). The parasitoid assemblage of *T. pityocampa* and its impact on the pest may also be affected by the identity of the host plant. According to Mirchev (2005), the rate of parasitism in different pine species varies greatly between years and localities: in Turkish pine (*Pinus brutia* Tenore) stands in Turkey, the average parasitism in Köyceğiz was 9.8%, but in Iskenderun - 34.9%; in Bulgaria, in *P. nigra* plantations, the parasitism varied from 4.4% (Kurtovo) to 43.7% (Ploski), and in Scots pine (*Pinus sylvestris* L.) - from 0% (Kyustendil) to 23.5% (Satovcha); in Greece, the lowest and highest rates of parasitism in Aleppo pine (*Pinus halepensis* Mill.) stands were 13.3% (Fteri-Egio) and 39.6% (Hydra Island), respectively.

Egg parasitoids are considered the most important regulator of *T. pityocampa* egg stage in both Bulgaria and France, that was found in many previous studies (Battisti, 1989; Biliotti, 1958; Biliotti *et al.*, 1962; Cadahia and Cuevas, 1964; Ceballos, 1969; Demolin and Delmos, 1967; Harapin, 1986; Jamaâ *et al.*, 1996; El-Yousfi, 1989; Kitt and Schmidt, 1993; Tiberi and Roversi, 1987; Tsankov, 1990; Szczepański and Tzankov, 1967). Percent parasitism is an indirect indicator of parasitoid effects on host populations, although life table studies that consider stage-specific effects of parasitoids and other mortality factors provide a more accurate picture of the role of different parasitoids (Van Driesche *et al.*, 1991). In the present study, a higher rate of parasitized eggs was detected at the localities of historical range of the species in France - Ré Island and Orléans, compared to a newly colonized one, Fréhel. In Bulgaria, the host mortality caused by parasitoids in historical localities (Sandanski and Gega) was 19.2-17.3%. In one of the new localities, Sladak kladenets, a typical low parasitism rate was recorded. In contrast, an unusually high parasitism rate (29.0%) was established in the second newly colonized locality, Maglizh. The high value of parasitism in Maglizh is due to the polyphagous *O. pityocampae* which is associated with a wide range of hemipteran and lepidopteran hosts (Noyes, 2018). As this parasitoid is less specialized and adapted to *T. pityocampa* (the parasitoid's emergence occurs in late autumn or spring), survival on other hosts is probably responsible for its high impact on the pest.

*Baryscapus servadeii* and *O. pityocampae* are widespread across the distribution range of *T. pityocampa* (Mirchev, 2005). The oligophagous *B. servadeii* parasitizes not only pine processionary moth, but also two other species of *Thaumetopoea* genus - *T. bonjeani* (Demolin, 1988) and *T. wilkinsoni* (Halperin, 1990). The number of *B. servadeii* and *O. pityocampae* may be limited by different adaptability to environmental conditions. *Baryscapus servadeii* is more adaptive to thermal stressors (Masutti, 1964). It successfully infests *T. pityocampa* eggs from their deposition to larval hatching whereas *O. pityocampae* cannot develop after 32 days of host incubation period (Halperin, 1990). The number of egg parasitoids of *T. pityocampa* is also influenced by the presence of scales on egg batches (Biliotti, 1958). This is obviously much more relevant for small-sized parasitoids such as *Trichogramma* sp. since females cannot overcome the scales and can only parasitize uncovered eggs in the ends of batches (Tsankov, 1990).

The hyperparasitoid *B. transversalis* was found in Bulgarian samples. It is distributed in Southeast Europe: Albania, Bulgaria, Bosnia and Herzegovina, Greece and Turkey (Noyes, 2018). Another representative of the genus, *Baryscapus endemus* (Walker, 1839) (syn. *Tetrastichus tibialis* Kurdjumov, 1913) was reported as a secondary parasitoid in France (Biliotti, 1958). It was

also found on *T. pityocampa* in Spain (Cadahia and Cuevas, 1964) and North Macedonia (Roques *et al.*, 2015). *B. endemus* is a polyphage of many insect hosts from Coleoptera, Hemiptera, Hymenoptera and Lepidoptera orders (Noyes, 2018).

Abiotic factors (high temperature, low humidity, etc.) are the main causes for embryonal mortality of *T. pityocampa* (Mirchev, 2005). When the egg batches are exposed to direct sunlight, their temperature might rise by 14° C (Milani, 1990). Sterile, unfertilized or empty eggs in *T. pityocampa* batches were reported in Greece (Bellin *et al.*, 1990), Morocco (Schmidt *et al.*, 1997b), Portugal (Mirchev and Tsankov, 2000) and other countries.

The proportion of eggs destroyed by predators is usually negligible, but in Maglitzh, the regulating effect reached high values (>10%). In Bulgaria, *Ephippiger ephippiger* (Fiebig, 1784) (Orthoptera: Bradyporidae) and *Pterolepis germanica* (Herrich-Schäffer, 1840) (Orthoptera: Tettigoniidae) were observed to destroy *T. pityocampa* egg batches (Tsankov *et al.*, 1996b). The first species was reported as a predator of the pine processionary moth in France as well (Demolin and Delmos, 1967). Halperin (1990) reported *Monomorium dentigerum* (Roger, 1862) (Hymenoptera: Formicidae) and tetigonid species as predators of *T. wilkinsoni* in Israel, and *Monomorium destructor* (Jerdon, 1851) (syn. *Myrmica gracillima* Smith, F., 1861) in Cyprus.

In conclusion, parasitoids were the most important regulating factor of *T. pityocampa* at the egg stage in both France and Bulgaria. In newly colonized areas, the impact of the parasitoids on the host is significantly lower. This suggests that parasitoids lag behind the expansion of the pine processionary moth. In one of the most important species, *O. pityocampae*, the temperature conditions and the presence of alternative hosts are essential factors for its effectiveness. The hyperparasitoid *B. transversalis* caused relatively low mortality (4.8-6.2%) of primary parasitoids at sites within the historical range of *T. pityocampa* in Bulgaria.

## Acknowledgements

This study was supported by RILA Project DTNS France 01/8/09.05.2017 ‘Standardizing Automated large-scale Monitoring’ to understand atypic phenologies at both ends (France and Bulgaria) of the climate-driven expansion front of a European urticating pest, the Pine Processionary Moth (SAM3PM) financed by the National Scientific Fund. The research was funded by the French Ministry of Europe and Foreign Affairs (MEAE) and the French Ministry of Higher Education, Research and Innovation (MESRI).

## Funding

This work was supported by the Bulgarian National Science Fund (RILA Project DTNS France 01/8/09.05.2017), the French Ministry of Europe and Foreign Affairs (MEAE) and the French Ministry of Higher Education, Research and Innovation (MESRI).

## References



- Auger-Rozenberg, M.-A., Barbaro, L., Battisti, A., Blache, S., Charbonnier, Y., Denux, O., Garcia, J., Goussard, F., Imbert, C.-E., Kerdelhué, C., Roques, A., Torres-Leguizamon, M. and Vetillard, F. 2015 Ecological responses of parasitoids, predators and associated insect communities to the climate-driven expansion of the pine processionary moth. In *Processionary Moths and Climate Change: An Update*. A. Roques (ed.) Springer, Netherlands, pp. 311-357.
- Battisti, A. 1989 Field studies on the behaviour of two egg parasitoids of the pine processionary moth *Thaumetopoea pityocampa*. *Entomophaga* **34**, 29-38.
- Battisti, A., Colazza, S., Roversi, P. F. and Tiberi, R. 1988 Alternative hosts of *Ooencyrtus pityocampae* Mercet (Hymenoptera: Encyrtidae) in Italy. *Redia* **71**, 321-328.
- Battisti, A., Stastny, M., Netherer, S., Robinet, C., Schopf, A., Roques, A. and Larsson, S. 2005 Expansion of geographic range in the pine processionary moth caused by increased winter temperatures. *Ecol. Appl.* **15**, 2084-2096.
- Battisti, A., Stastny, M., Buffo, E. and Larsson, S. 2006 A rapid altitudinal range expansion in the pine processionary moth produced by the 2003 climatic anomaly. *Global Change Biol.* **12**, 662-671.
- Bellin, S., Schmidt, G.H. and Douma-Petridou, E. 1990 Structure, ooparasitoid spectrum and rate of parasitism of egg-batches of *Thaumetopoea pityocampa* (Den. & Schiff.) (Lep., Thaumetopoeidae) in Greece. *J. Appl. Entomol.* **110**, 113-120.
- Biliotti, E. 1958 Les parasites et prédateurs de *Thaumetopoea pityocampa* Schiff. (Lepidoptera). *Entomophaga* **3**, 23-34.
- Biliotti, E., Vago, C. and Halperin, J. 1962 Une virose de *Thaumetopoea wilkinsoni* et ses rapports avec la polyédrie cytoplasmatique de *T. pityocampa*. *C. r. hebd. séances Acad. sci.* **225**, 1039-1041.
- Buxton, R.D. 1990 The influence of host tree species on timing of pupation of *Thaumetopoea pityocampa* Schiff. (Lep., Thaumetopoeidae) and its exposure to parasitism by *Phryxe caudata* Rond. (Dipt., Larvaevoridae). *J. Appl. Entomol.* **109**, 302-310.
- Cadahia, D. and Cuevas, P. 1964 Tratamientos de la 'Processionaria del Pino', en la campaña 1963/64. *Bol. Serv. Plagas Forestales* **7**, 12-22.
- Carvalho, L., Mateus, E.P., Farrall, M.H. and Paiva, M.R. 1999 Caracterização do ataque de *Thaumetopoea pityocampa* num povoamento de *Pinus pinaster* por SPME e GC-MS. *Revista de Biol. (Lisboa)* **17**, 283-287.
- Ceballos, P. 1969 Predadores de *Thaumetopoea pityocampa* y posibilidades de utilizacion. *Bol. Serv. Plagas Forestales* **12**, 35-38.
- Demolin, J. 1988 Intensification de la protection phytosanitaire des forêts. Algérie 1986-1987. La processionnaire du cèdre: *Thaumetopoea bonjeani*. *FAO Rapport scientifique et iconographique*, Rome, Italy, 21 pp.
- Demolin, G. and Delmos, J.C. 1967 Les Ephippigeres (Orthopteres, Tettigonidae) prédateurs occasionnels, mails importants de, *Thaumetopoea pityocampa* Schiff. *Entomophaga* **12**, 399-401.
- Douma-Petridou, E. 1990 European *Thaumetopoea* species. (Lep., Thaumetopoeidae): Characteristics and life-cycles. In *Proceeding Thaumetopoea Symposium*. G.H. Schmidt (ed.). Department of Zoology-Entomology, University of Hannover, pp. 12-18.

- El-Yousfi, M. 1989 La processionaria del cedro, *Thaumetopoea bonjeani* (Powell). *Bol. San. Veg., Plagas* **15**, 43-56.
- Halperin, J. 1990 Natural enemies of *Thaumetopoea* spp. (Lep., Thaumetopoeidae) in Israel. *J. Appl. Entomol.* **109**, 425-435.
- Harapin, M. 1986 A note on the parasite complex of *Thaumetopoea pityocampa* in Yugoslavia. *EPPO Bulletin* **16**, 625-626.
- Hoch, G., Netherer, S., Pennerstorfer, J., Baier, P., Battisti, A. and Schopf, A. 2005 Range expansion of the pine processionary moth in Europe II: Activity and survival of *Thaumetopoea pityocampa* (Lep.: Thaumetopoeidae) during the winter months in an Alpine valley. In *Proceedings, 16th U.S. Department of Agriculture interagency research forum on gypsy moth and other invasive species*. K.W. Gottschalk (ed.), p. 42.
- Hollander, M. and Wolfe, D.A. 1973 *Nonparametric statistical methods*. John Wiley & Sons. New York, 503 pp.
- Imber, C-E. 2012 Expansion d'un ravageur forestier sous l'effet du Rechauffement climatique: la processionnaire du pin affecte-t-elle la biodiversité entomologique dans les zones nouvellement colonisées? *Thèse pour obtenir le grade de: Docteur de l'université d'Orleans*, 197 pp.
- Jamaâ, B., Graf, P. and Bourarach, K. 1996 Factors of natural death of the forest caterpillar: *Thaumetopoea pityocampa* Schiff. (Lepidoptera: Thaumetopoeidae) at the egg stage in Morocco. *Arab J. Pl. Protect.* **14** (2), 86-90.
- Kitt, J. and Schmidt, G.H. 1993 Parasitism of egg-batches of the pine processionary moth *Thaumetopoea wilkinsoni* Tams (Lep., Thaumetopoeidae) in the mountains of Lahav (Israel). *J. Appl. Entomol.* **115**, 484-498.
- Masutti, L. 1964 Ricerche sui parassiti oofagi della *Thaumetopoea pityocampa* (Schiff.). *Annali del Centro di Economia Montana delle Venezie* **4**, 205-271.
- Masutti, L. and Battisti, A. 1990 *Thaumetopoea pityocampa* (Den. & Schiff.) in Italy. Bionomics and perspectives of integrated control. *J. Appl. Entomol.* **110**, 229-234.
- Milani, R. 1990 The temperature of the egg masses of *Thaumetopoea pityocampa* (Den. & Schiff.) (Lepidoptera, Thaumetopoeidae). *Redia* **73**, 149-161.
- Mirchev, P. 2005 Egg parasitoids on pine processionary moth *Thaumetopoea pityocampa* (Den. & Schiff.) (Lepidoptera: Thaumetopoeidae) in countries of Balkan Peninsula. *DSc Thesis*, Forest Research Institute, Sofia, 276 pp.
- Mirchev, P. and Tsankov, G. 2000 Parasitism of egg-batches of pine processionary moth *Thaumetopoea pityocampa* (Den. et Schiff.) (Lepidoptera: Thaumetopoeidae) collected in Portugal. *Forest Sci.* **4**, 65-71.
- Mirchev, P., Georgiev, G. and Matova, M. 2011a Prerequisites for expansion of pine processionary moth *Thaumetopoea pityocampa* (Den. & Schiff.) in Bulgaria. *J. Balkan Ecol.* **14**, 117-130.
- Mirchev, P., Tsankov, G., Georgiev, G. and Boyadzhiev, P. 2011b *Pediobius bruchicida* (Rondani) (Hymenoptera: Eulophidae) – an egg parasitoid of pine processionary moth, *Thaumetopoea pityocampa* (Denis & Schiffermuller) (Lepidoptera: Notodontidae) and a new species for Bulgarian fauna. *Acta Zool. Bulg.* **63**, 319-322.

- Mirchev, P., Georgiev, G. and Tsankov, G. 2017 Long-term studies on egg parasitoids of pine processionary moth (*Thaumetopoea pityocampa*) in a new locality in Bulgaria. *J. Entomol. Res. Soc.* **19** (3), 15-25.
- Mirchev, P., Georgiev, G., Georgieva, M., Matova, M. and Zaemdzhikova, G. 2018 Enlargement of the pine processionary moth (*Thaumetopoea pityocampa*) range in Bulgaria. *Forest Review* **48**, 4-7.
- Moura, E.B., Rodrigues, M.T. and Paiva, M.R. 1999 Estudo da distribuição espacial da processionária (*Thaumetopoea pityocampa*) num povoamento de *Pinus pinaster*. In Actas do 3º Encontro Nacional de Ecologia, Faro, Portugal, 3-5 de Dezembro de 1998. *Revista de Biol. (Lisboa)* **17**, 203-209.
- Özkazans, O. 1987 Cam keseböceği (*Thaumetopoea pityocampa* Schiff.) (Lep.: Thaumetopoeidae) 'nin yumurta birakama dayanislari üzerinde incelemeler. In *Türkiye I. Entomoloji Kongresi Bildirileri*, 13-16 Ekim 1987, Ege Üniversitesi, Bornova, Izmir. Bornova/Izmir, Turkey; Ege Üniversitesi Atatürk Kültür Merkezi, pp. 727-735.
- Netherer, S. and Schopf, A. 2010 Potential effects of climate change on insect herbivores in European forests-General aspects and the pine processionary moth as specific example. *Forest Ecol. Manag.* **259**, 831-838.
- Nicholls, J.A., Fuentes-Utrilla, P., Hayward, A., Melika, G., Csoka, G., Nieves-Aldrey, J.-L., Pujade-Villar, J., Tavakoli, M., Schönrogge, K. and Stone, G.N. 2010 Community impacts of anthropogenic disturbance, natural enemies exploit multiple routes in pursuit of invading herbivore hosts. *BMC Evol. Biol.* **10**, 322.
- Noyes, J.S. 2018 Universal Chalcidoidea Database. World Wide Web electronic publication. <http://www.nhm.ac.uk/chalcidoids>, Database last updated March 2018.
- Robinet, C., Baier, P., Pennerstorfer, J., Schopf, A. and Roques, A. 2007 Modelling the effects of climate change on the potential feeding activity of *Thaumetopoea pityocampa* (Den. & Schiff.) (Lep. Notontidae) in France. *Global Ecol. and Biogeogr.* **16**, 460-471.
- Roques, A., Rousset, J., Robinet, C., Goussard, F. and Battisti, A. 2005 *Range expansion of the pine processionary moth in Europe. I. Mechanisms underlying PPM expansion in France in relation to global warming*. In Proceedings, 16th U.S. Department of Agriculture interagency research forum on gypsy moth and other invasive species, 2005. K.W.Gottschalk (ed.) USDA Forest Service, General Technical Report NE-337, pp. 71-72.
- Roques, A., Rousset, J., Avci, M., Avtzis, D.N., Basso, A., Battisti, A., Ben Jamaa, M.L., Bensidi, A., Berardi, L., Berretima, W., Branco, M., Chakali, G., Çota, E., Dautbašić, M., Delb, H., El Alaoui El Fels, M.A., El Mercht, S., El Mokhefi, M., Forster, B., Garcia, J., Georgiev, G., Glavendekić, M.M., Goussard, F., Halbig, P., Henke, L., Hernández, R., Hódar, J.A., İpekdağ, K., Jurc, M., Klimetzek, D., Laparie, M., Larsson, S., Mateus, E., Matošević, D., Meier, F., Mendel, Z., Meurisse, N., Mihajlović, L., Mirchev, P., Nasceski, S., Nussbaumer, C., Paiva, M.-R., Papazova, I., Pino, J., Podlesnik, J., Poirot, J., Protasov, A., Rahim, N., Peña, G.S., Santos, H., Sauvard, D., Schopf, A., Simonato, M., Tsankov, G., Wagenhoff, E., Yart, A., Zamora, R., Zamoum, M. and Robinet, C. 2015 Climate Warming and Past and Present Distribution of the Processionary Moths (*Thaumetopoea* spp.) in Europe, Asia Minor and North Africa. In *Processionary Moths and Climate Change: An Update*. A. Roques (ed.). Springer, pp. 81-161.

- Roussis, V., Petrakis, V.P., Ortiz, A. and Mazomenos, E.B. 1994 Pine secondary metabolites as perceived by the mother *Thaumetopoea pityocampa* (Den. & Schiff.) and experienced by the offspring. In *5<sup>th</sup> European Congress of Entomology*, Univ. of York, UK, 29 August-2 September 1994, p. 134.
- Schmidt, G.H. and Kitt, J. 1994 Identification by meconia of two egg parasitoids of *Thaumetopoea wilkinsoni*. *Phytoparasitica* **22**, 39-41.
- Schmidt, G. H., Tsankov, G. and Mirchev, P. 1997a Notes on the egg parasitoids of *Thaumetopoea pityocampa* (Den. & Schiff.) (Insecta Lepidoptera Thaumetopoeidae) collected on the Greek island Hydra. *Boll. Zool. Agraria e di Bachic., Ser. II* **29**, 91-99.
- Schmidt, G.H., Mirchev, P. and Tsankov, G. 1997b The egg parasitoids of *Thaumetopoea pityocampa* in the Atlas mountains near Marrakech (Morocco). *Phytoparasitica* **25**, 275-281.
- Szczepański, H. and Tsankov, G. 1967 Materialy do znajności pasożytów jaj korowódki śródziemnomorskiej *Thaumetopoea pityocampa* Schiff. (Lepidoptera, Thaumetopoeidae) w Bulgarii. *Sylwan* **4**, 37-43.
- Tanzen, E. and Schmidt, G.H. 1995 Identification by meconia of four species egg parasitoids *Thaumetopoea pityocampa* (Den. & Schiff.) (Insecta Lepidoptera Thaumetopoeidae). *Boll. Zool. Agrar. Bachic., Ser. II* **27**, 61-70.
- Tiberi, R. and Roversi, F.P. 1987 I parassitoidi oofagi di *Thaumetopoea pityocampa* (Den. & Schiff.) su *Pinus halepensis* Mill. Nel Gargano, Puglia (Hymenoptera, Chalcidoidea; Lepidoptera, Thaumetopoeidae). *Redia* **70**, 1-19.
- Tiberi, R., Niccoli, A., Curini, M., Epifano, F., Marcotullio, M.C. and Rosati, O. 1999 The role of the monoterpene composition in *Pinus* spp., needles in host selection by the pine processionary caterpillar, *Thaumetopoea pityocampa*. *Phytoparasitica* **27**, 263-272.
- Toffolo, E.P., Bernardinelli, I., Stergulc, F. and Battisti, A. 2006 Climate change and expansion of the Pine Processionary Moth, *Thaumetopoea pityocampa*, in Northern Italy. In *IUFRO Working Party 7.03.10 Proceedings of the Workshop 2006*, Gmunden/Austria, pp. 331-340.
- Tsankov, G. 1990 Egg parasitoids of the pine processionary moth, *Thaumetopoea pityocampa* (Den. & Schiff.) (Lep., Thaumetopoeidae) in Bulgaria. *J. Appl. Entomol.* **110**, 7-13.
- Tsankov, G., Schmidt, G.H. and Mirchev, P. 1996a Parasitism of egg-batches of the pine processionary moth *Thaumetopoea pityocampa* (Den. & Schiff.) (Lep., Thaumetopoeidae) in various regions of Bulgaria. *J. Appl. Entomol.* **120**, 93-105.
- Tsankov, G., Schmidt, G.H. and Mirchev, P. 1996b Structure and parasitism of egg-batches of a processionary moth population different from *Thaumetopoea pityocampa* (Den. & Schiff.) (Lep. Thaumetopoeidae) found in Bulgaria. *Boll. Zool. Agrar. Bachic., Ser. II* **28**, 195-207.
- Tsankov, G., Schmidt, G.H. and Mirchev, P. 1998 Distribution of egg parasitoids of the pine processionary moth *Thaumetopoea pityocampa* (Den. et Schiff.) (Lep., Thaumetopoeidae) in the southwestern region of Bulgaria. *Forest Sci.* **3/4**, 5-17.
- Van Driesche, R.G., Bellows Jr, T.S., Elkinton, J.S., Gould, J.R. and Ferro, D.N. 1991 The meaning of percentage parasitism revisited: solutions to the problem of accurately estimating total losses from parasitism. *Env. Entomol.* **20**, 1-7.

Table 1. Main characteristics of sample plots in France and Bulgaria

Region	Range	Locality	Coordinates	Altitude, m	Date of collection	Egg batches	Host tree
<b>France</b>							
Ré Island	Historical	Portes-en-Ré	N46°15'11.3" W01°30'54.0"	14	21.11.2017	15	<i>Pinus nigra</i>
		Ars-en-Ré	N46°13'15.1" W01°31'49.9"	0	21.11.2017	2	<i>Pinus pinaster</i>
		St.-Martin-en-Ré	N46°11'13.4" W01°22'06.8"	14	21.11.2017	4	<i>Pinus pinaster</i>
Fréhel	Newly colonized	Camp de Fréhel	N48°39'15.9" W02°21'22.5"	14	22.11.2017	40	<i>Pinus radiata</i>
		Camp de Fréhel	N48°39'15.9" W02°21'22.5"	14	22.11.2017	14	<i>Pinus pinaster</i>
		Plurien	N48°38'02.5" W02°24'37.0"	8	22.11.2017	10	<i>Pinus pinaster</i>
Orléans	Newly colonized	Saran	N47°56'58.5" E01°50'16.0"	121	23.11.2017	8	<i>Pinus nigra</i>
		Orléans	N47°49'22.3" E01°55'29.5"	108	23.11.2017	2	<i>Pinus nigra</i>
		Châteaudun	N48°03'34.4" E01°23'39.0"	127	23.11.2017	2	<i>Pinus nigra</i>
<b>Bulgaria</b>							
Sandanski	Historical	Sandanski	N41°34'22.0" E23°17'14.9"	325	20.09.2017	22	<i>Pinus nigra</i>
Ograzhden Mt.	Historical	Gega	N41°27'09.7" E22°59'54.0"	856	31.08.2017	38	<i>Pinus nigra</i>
St. Zagora	Newly colonized	Sladak kladenets	N42°24'19.8" E25°22'58.5"	403	28.10.2016	36	<i>Pinus nigra</i>
Maglzh	Newly colonized	Maglzh	N42°36'08.6" E25°30'31.9"	374	03.10.2017	13	<i>Pinus nigra</i>

Table 2. Viability of *Thaumetopoea pityocampa* in egg stage in different localities

Parameters	France				Bulgaria		
	Ré Island	Fréhel	Orléans	Sandanski	Gega	Sladak kladenets	Maglizh
Generation of PPM	2017	2017	2017	2017	2017	2016	2017
Total number (n) of egg batches	21	64	12	22	38	36	13
Larvae died without opening, n (%)	30 (0.7)	84 (0.7)	70 (2.5)	232 (4.7)	242 (2.9)	173 (2.1)	112 (3.4)
Larvae died with opening, n (%)	9 (0.2)	13 (0.1)	3 (0.1)	49 (1.0)	115 (1.4)	26 (0.3)	35 (1.1)
Undeveloped eggs with dried-up yolk, n (%)	76 (1.7)	74 (0.6)	72 (2.6)	245 (5.0)	226 (2.7)	149 (1.8)	97 (3.0)
Eggs totally empty, without any remains, n (%)	7 (0.2)	-	11 (0.4)	4 (0.1)	99 (1.2)	26 (0.3)	41 (1.3)
Total number of aborted eggs, n (%)	122 (2.8)	171 (1.4)	156 (5.6)	530 (10.8)	682 (8.3)	374 (4.5)	285 (8.7)
Number of larvae hatched, n (%)	3642 (83.6)	11198 (95.7)	1947 (69.8)	3452 (70.0)	6120 (74.4)	7052 (85.2)	1626 (49.8)
Impact of parasitoids, n (%)	593 (13.6)	332 (2.9)	681 (24.4)	947 (19.2)	1392 (16.9)	650 (7.8)	947 (29.0)
Impact of predators, n (%)	1 (0.02)	2 (0.02)	3 (0.1)	-	32 (0.4)	203 (2.5)	409 (12.5)
Total number of eggs, n (%)	4358 (100)	11703 (100)	2787 (100)	4929 (100)	8226 (100)	8279 (100)	3267 (100)
Range of eggs per egg batches	106-287	55-278	172-305	105-284	94-277	92-306	204-299
Mean eggs per egg batches±Std	207.5±50.5	185.7±47.7	232.2±43.5	224.0±41.8	216.4±41.0	230.0±48.8	251.3±31.3

Table 3. Percentage of egg parasitoids of *Thaumetopoea pityocampa* in different localities

Parasitoids	France						Bulgaria							
	Ré Island		Fréhel		Orléans		Sandanski		Gega		Sladak kladenets		Maglizeh	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%
<i>Ooencyrtus pityocampae</i>	9	1.6	1	0.3	91	14.7	457	72.1	130	13.3	236	65.7	549	89.7
<i>Baryscapus servadeii</i>	550	97.3	-	-	524	84.7	130	20.5	737	75.4	-	-	13	2.1
<i>Baryscapus transversalis</i>	-	-	-	-	-	-	28	4.4	54	5.5	-	-	-	-
<i>Anastatus bifasciatus</i>	-	-	-	-	4	0.6	19	3.0	36	3.7	123	34.3	30	4.9
<i>Trichogramma</i> sp.	6	1.1	331	99.7	-	-	-	-	21	2.1	-	-	20	3.3
<b>Total</b>	<b>565</b>	<b>100.0</b>	<b>332</b>	<b>100.0</b>	<b>619</b>	<b>100.0</b>	<b>634</b>	<b>100.0</b>	<b>978</b>	<b>100.0</b>	<b>359</b>	<b>100.0</b>	<b>612</b>	<b>100.0</b>

Table 4. Characteristics of egg parasitoids of *T. pityocampa* in different localities

Species and parameters	France				Bulgaria		
	Ré Island	Fréhel	Orléans	Sandanski	Gega	Sladak kladenets	Maglizh
<b><i>Ooencyrtus pityocampae</i>, n</b>	<b>9</b>	<b>1</b>	<b>91</b>	<b>457</b>	<b>130</b>	<b>236</b>	<b>549</b>
Emerged before collection of egg batches, n (%)	2 (22.2)	-	27 (29.7)	149 (32.6)	79 (60.8)	143 (60.6)	404 (73.6)
Emerged after collection of egg batches, n (%)	7 (77.8)	-	61 (67.0)	255 (55.8)	36 (27.7)	17 (7.2)	89 (16.2)
Adults died in eggs, n (%)	-	1 (100.0)	3 (3.3)	53 (11.6)	15 (11.5)	76 (32.2)	56 (10.2)
Egg batches with <i>O. pityocampae</i> , n (%)	3 (14.3)	1 (1.6)	7 (58.3)	21 (95.5)	16 (42.1)	22 (61.1)	12 (92.3)
<b><i>Baryscapus servadeii</i>, n</b>	<b>550</b>	<b>-</b>	<b>524</b>	<b>130</b>	<b>737</b>	<b>-</b>	<b>13</b>
Emerged before collection of egg batches, n (%)	113 (20.5)	-	100 (19.1)	2 (1.5)	374 (50.7)	-	3 (23.1)
Emerged after collection of egg batches, n (%)	429 (78.0)	-	406 (77.5)	126 (97.0)	339 (46.0)	-	10 (76.9)
Adults died in eggs, n (%)	8 (1.5)	-	18 (3.4)	2 (1.5)	24 (3.3)	-	-
Egg batches with <i>B. servadeii</i> , n (%)	12 (57.1)	-	9 (75.0)	8 (36.4)	29 (76.3)	-	4 (30.8)
<b><i>Baryscapus transversalis</i>, n</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>28</b>	<b>54</b>	<b>-</b>	<b>-</b>
Emerged after collection of egg batches, ♀, n (%)	-	-	-	12 (42.9)	24 (44.4)	-	-
Emerged after collection of egg batches, ♂, n (%)	-	-	-	10 (35.7)	21 (38.9)	-	-
Adults died in eggs, n (%)	-	-	-	6 (21.4)	9 (16.7)	-	-
Egg batches with <i>B. transversalis</i> , n (%)	-	-	-	5 (22.7)	13 (34.2)	-	-
<b><i>Anastatus bifasciatus</i>, n</b>	<b>-</b>	<b>-</b>	<b>4</b>	<b>19</b>	<b>36</b>	<b>123</b>	<b>30</b>
Emerged before collection of egg batches, n (%)	-	-	2 (50.0)	1 (5.3)	19 (52.8)	97 (78.9)	25 (83.3)
Emerged after collection of egg batches, n (%)	-	-	2 (50.0)	18 (94.7)	17 (47.2)	17 (13.8)	2 (6.7)
Adults died in eggs, n (%)	-	-	-	-	-	9 (7.3)	3 (10.0)
Egg batches with <i>A. bifasciatus</i> , n (%)	-	-	1 (8.3)	1 (4.5)	5 (13.2)	14 (38.9)	5 (38.5)
<b><i>Trichogramma</i> sp., n</b>	<b>6</b>	<b>331</b>	<b>-</b>	<b>-</b>	<b>21</b>	<b>-</b>	<b>20</b>
Emerged before collection of egg batches, n (%)	5 (83.3)	147 (44.4)	-	-	10 (47.6)	-	6 (30.0)
Emerged after collection of egg batches, n (%)	-	44 (13.3)	-	-	6 (28.6)	-	-
Adults died in eggs, n (%)	1 (16.7)	140 (42.3)	-	-	5 (23.8)	-	14 (70.0)
Egg batches with <i>Trichogramma</i> sp., n (%)	2 (9.5)	25 (39.1)	-	-	5 (13.2)	-	5 (38.5)
<b>Undetermined larvae of parasitoids, n</b>	<b>28</b>	<b>-</b>	<b>68</b>	<b>313</b>	<b>411</b>	<b>291</b>	<b>335</b>



