



HAL
open science

Formulae for two-variable Green functions

François Digne, Jean Michel

► **To cite this version:**

| François Digne, Jean Michel. Formulae for two-variable Green functions. 2021. hal-03127834v2

HAL Id: hal-03127834

<https://hal.science/hal-03127834v2>

Preprint submitted on 19 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FORMULAE FOR TWO-VARIABLE GREEN FUNCTIONS

FRANÇOIS DIGNE AND JEAN MICHEL

ABSTRACT. Based on results of Digne-Michel-Lehrer (2003) we give two formulae for two-variable Green functions attached to Lusztig induction in a finite reductive group. We present applications to explicit computation of these Green functions, to conjectures of Malle and Rotilio, and to scalar products between Lusztig inductions of Gelfand-Graev characters.

Let \mathbf{G} be a connected reductive group with Frobenius root F ; that is, some power F^δ is a Frobenius endomorphism attached to an \mathbb{F}_{q^δ} -structure on \mathbf{G} , where q^δ is a power of a prime p . Let \mathbf{L} be an F -stable Levi subgroup of a (non-necessarily F -stable) parabolic subgroup \mathbf{P} of \mathbf{G} . Let \mathbf{U} be the unipotent radical of \mathbf{P} and let $\mathbf{X}_{\mathbf{U}} = \{g\mathbf{U} \in \mathbf{G}/\mathbf{U} \mid g^{-1}Fg \in \mathbf{U} \cdot F\mathbf{U}\}$ be the variety used to define the Lusztig induction and restriction functors $R_{\mathbf{L}}^{\mathbf{G}}$ and $*R_{\mathbf{L}}^{\mathbf{G}}$. For $u \in \mathbf{G}^F, v \in \mathbf{L}^F$ unipotent elements, the two-variable Green function is defined as

$$Q_{\mathbf{L}}^{\mathbf{G}}(u, v) = \text{Trace}((u, v) \mid \sum_i (-1)^i H_c^i(\mathbf{X}_{\mathbf{U}})).$$

In this paper, using the results of [5], we give two different formulae for two-variable Green functions, and some consequences of these, including proving some conjectures of [15].

The two-variables Green functions occur in the character formulae for Lusztig induction and restriction. In particular, for unipotent elements these formulae read

Proposition 1. (*See for example [8, 10.1.2]*)

- If u is a unipotent element of \mathbf{G}^F , and ψ a class function on \mathbf{L}^F , we have

$$R_{\mathbf{L}}^{\mathbf{G}}(\psi)(u) = |\mathbf{L}^F| \langle \psi, Q_{\mathbf{L}}^{\mathbf{G}}(u, -) \rangle_{\mathbf{L}^F}.$$

- If v is a unipotent element of \mathbf{L}^F , and χ a class function on \mathbf{G}^F , we have

$$*R_{\mathbf{L}}^{\mathbf{G}}(\chi)(v) = |\mathbf{L}^F| \langle \chi, Q_{\mathbf{L}}^{\mathbf{G}}(-, v^{-1}) \rangle_{\mathbf{G}^F}.$$

TWO FORMULAE FOR TWO-VARIABLE GREEN FUNCTIONS

For an element u in a group G we denote by u^G the G -conjugacy class of u .

Proposition 2. *Assume either the centre $Z\mathbf{G}$ of \mathbf{G} is connected and $q > 2$, or q is large enough (depending just on the Dynkin diagram of \mathbf{G}). Then for u regular, $Q_{\mathbf{L}}^{\mathbf{G}}(u, -)$ vanishes outside a unique regular unipotent class of \mathbf{L}^F . For v in that class, we have $Q_{\mathbf{L}}^{\mathbf{G}}(u, v) = |v^{\mathbf{L}^F}|^{-1}$.*

Proof (Rotilio). Let $\gamma_u^{\mathbf{G}}$ be the normalized characteristic function of the \mathbf{G}^F -conjugacy class of u ; that is, the function equal to 0 outside the class of u and to $|C_{\mathbf{G}}(u)^F|^{-1}$ on that class. For $v' \in \mathbf{L}^F$ unipotent, Proposition 1 gives

$$*R_{\mathbf{L}}^{\mathbf{G}}(\gamma_u^{\mathbf{G}})(v') = |\mathbf{L}^F| \langle \gamma_u^{\mathbf{G}}, Q_{\mathbf{L}}^{\mathbf{G}}(-, v'^{-1}) \rangle_{\mathbf{G}^F} = |\mathbf{L}^F| Q_{\mathbf{L}}^{\mathbf{G}}(u, v'^{-1}).$$

Now, by [1, Theorem 15.2], there exists $v \in \mathbf{L}^F$ such that the left-hand side is equal to $\gamma_v^{\mathbf{L}}(v')$. In [1, Theorem 15.2] q is assumed large enough so that the Mackey formula holds and that p and q are such that the results of [10] hold (that is p almost good and q larger than some number depending on the Dynkin diagram of \mathbf{G}). The Mackey formula is known to hold if $q > 2$ by [2] and the results of [10] hold without condition on q if $Z\mathbf{G}$ is connected by [17] or [18, Theorem 4.2] (the assumption that p is almost good can be removed, see [13, section 89] and [12]). The proposition follows. \square

As in [5], we consider irreducible \mathbf{G} -equivariant local systems on unipotent classes. These local systems are partitioned into “blocks” parametrised by cuspidal pairs formed by a Levi subgroup and a cuspidal local system supported on a unipotent class of that Levi subgroup. Let us call $Q_{wF}^{\mathbf{L}, \mathcal{I}}$ the function defined in [5, 3.1(iii)] relative to an F -stable block \mathcal{I} of unipotently supported local systems on \mathbf{L} and to $wF \in W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})F$ where $(\mathbf{L}_{\mathcal{I}}, \iota_{\mathcal{I}})$ is the cuspidal datum of \mathcal{I} (for a Levi \mathbf{L} of a reductive group \mathbf{G} , we set $W_{\mathbf{G}}(\mathbf{L}) = N_{\mathbf{G}}(\mathbf{L})/\mathbf{L}$).

Proposition 3. *Assume either $Z\mathbf{G}$ is connected or q is large enough (depending just on the Dynkin diagram of \mathbf{G}). For u a unipotent element of \mathbf{G}^F and v a unipotent element of \mathbf{L}^F , we have*

$$Q_{\mathbf{L}}^{\mathbf{G}}(u, v) = |\mathbf{L}^F|^{-1} \sum_{\mathcal{I}} \sum_{w \in W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})} \frac{|Z^0(\mathbf{L}_{\mathcal{I}})^{wF}|}{|W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})|} \overline{Q_{wF}^{\mathbf{G}, \mathcal{I}_{\mathbf{G}}}}(u) Q_{wF}^{\mathbf{L}, \mathcal{I}}(v),$$

where \mathcal{I} runs over the F -stable blocks of \mathbf{L} and where $\mathcal{I}_{\mathbf{G}}$ is the block of \mathbf{G} with same cuspidal data as \mathcal{I} .

The part of the above sum for \mathcal{I} the principal block is the same formula as [7, Corollaire 4.4].

Proof. Proposition 1 applied with $\psi = Q_{wF}^{\mathbf{L}, \mathcal{I}}$ gives, if we write $R_{\mathbf{L}^F}^{\mathbf{G}^F}$ instead of $R_{\mathbf{L}}^{\mathbf{G}}$ to keep track of the Frobenius,

$$\langle Q_{wF}^{\mathbf{L}, \mathcal{I}}, Q_{\mathbf{L}}^{\mathbf{G}}(u, -) \rangle_{\mathbf{L}^F} = |\mathbf{L}^F|^{-1} R_{\mathbf{L}^F}^{\mathbf{G}^F}(Q_{wF}^{\mathbf{L}, \mathcal{I}})(u)$$

Now we have by [5, Proposition 3.2] $Q_{wF}^{\mathbf{L}, \mathcal{I}} = R_{\mathbf{L}_{\mathcal{I}}^{wF}}^{\mathbf{L}^F} \tilde{\chi}_{\iota_{\mathcal{I}}, wF}$ where $\tilde{\chi}_{\iota_{\mathcal{I}}, wF}$ is $q^{c_{\mathcal{I}}}$ times the characteristic function of $(\iota_{\mathcal{I}}, wF)$, a class function on $\mathbf{L}_{\mathcal{I}}^{wF}$. Here, as in [5, above Remark 2.1], for an irreducible \mathbf{G} -equivariant local system ι , we denote by C_{ι} the unipotent \mathbf{G} -conjugacy class which is the support of ι , and if $(\mathbf{L}_{\mathcal{I}}, \iota_{\mathcal{I}})$ is the cuspidal datum of ι we set $c_{\mathcal{I}} = \frac{1}{2}(\text{codim } C_{\iota} - \dim Z(\mathbf{L}_{\mathcal{I}}))$. In [5, Proposition 3.2] the assumptions on p and q come from [10] but by the same considerations than at the end of the proof of Proposition 2 it is sufficient to assume $Z\mathbf{G}$ connected or q large enough.

By the transitivity of Lusztig induction we get

$$\langle Q_{wF}^{\mathbf{L}, \mathcal{I}}, Q_{\mathbf{L}}^{\mathbf{G}}(u, -) \rangle_{\mathbf{L}^F} = |\mathbf{L}^F|^{-1} R_{\mathbf{L}_{\mathcal{I}}^{wF}}^{\mathbf{G}^F}(\tilde{\chi}_{\iota_{\mathcal{I}}, wF})(u) = |\mathbf{L}^F|^{-1} Q_{wF}^{\mathbf{G}, \mathcal{I}_{\mathbf{G}}}(u),$$

where $\mathcal{I}_{\mathbf{G}}$ is the block of \mathbf{G} with same cuspidal data as \mathcal{I} . Using the orthogonality of the Green functions $Q_{wF}^{\mathbf{L}, \mathcal{I}}$, see [5, Corollary 3.5] (where the assumption is that p is almost good which comes from [9], so can be removed now by [12]) and the fact they form a basis of unipotently supported class functions on \mathbf{L}^F , indexed by the $W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})$ -conjugacy classes of $W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})F$, we get the proposition. \square

Proposition 3 gives a convenient formula to compute automatically two-variable Green functions. Table 1 gives an example, computed with the package `Chevie` (see [16]).

We denote by \mathcal{Y}_ι the characteristic function of the F -stable local system ι , and by $A(u)$ the group of components of the centralizer of a unipotent element u .

Proposition 4. *Assume either $Z\mathbf{G}$ is connected or q is large enough (depending just on the Dynkin diagram of \mathbf{G}). Let $R_{\iota,\gamma}$ be the polynomials which appear in [5, Lemma 6.9]. Then*

$$Q_{\mathbf{L}}^{\mathbf{G}}(u, v) = |v^{\mathbf{L}^F}|^{-1} |A(v)|^{-1} \sum_{\mathcal{I}} \sum_{\iota \in \mathcal{I}_{\mathbf{G}^F}, \gamma \in \mathcal{I}^F} \overline{\mathcal{Y}_\iota(u)} \mathcal{Y}_\gamma(v) R_{\iota,\gamma} q^{c_\iota - c_\gamma},$$

where $c_\iota = \frac{1}{2}(\text{codim } C_\iota - \dim Z(\mathbf{L}_{\mathcal{I}}))$.

Proof. For a block \mathcal{I} of \mathbf{L} and $\iota \in \mathcal{I}^F$, let \tilde{Q}_ι be the function of [5, (4.1)]. Then by [5, (4.4)] applied respectively in \mathbf{G} and \mathbf{L} we have

$$Q_{wF}^{\mathbf{G}, \mathcal{I}_{\mathbf{G}}} (u) = \sum_{\iota \in \mathcal{I}_{\mathbf{G}^F}} \tilde{Q}_\iota(wF) \tilde{\mathcal{Y}}_\iota(u) \quad \text{and} \quad Q_{wF}^{\mathbf{L}, \mathcal{I}} (v) = \sum_{\kappa \in \mathcal{I}^F} \tilde{Q}_\kappa(wF) \tilde{\mathcal{Y}}_\kappa(v)$$

where $\tilde{\mathcal{Y}}_\iota = q^{c_\iota} \mathcal{Y}_\iota$. Thus, using the notation $\mathcal{Z}_{\mathbf{L}_{\mathcal{I}}}$ as in [5, 3.3] to denote the function $wF \mapsto |Z^0(\mathbf{L}_{\mathcal{I}})^{wF}|$ on $W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})F$, the term relative to a block \mathcal{I} in the formula of Proposition 3 can be written

$$|\mathbf{L}^F|^{-1} \langle \mathcal{Z}_{\mathbf{L}_{\mathcal{I}}} \sum_{\kappa \in \mathcal{I}^F} \tilde{Q}_\kappa \tilde{\mathcal{Y}}_\kappa(v), \sum_{\iota \in \mathcal{I}_{\mathbf{G}^F}} \tilde{\mathcal{Y}}_\iota(u) \text{Res}_{W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})F}^{W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})F} \tilde{Q}_\iota \rangle_{W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})F}.$$

Applying now [5, Lemma 6.9] this is equal to

$$|\mathbf{L}^F|^{-1} \langle \mathcal{Z}_{\mathbf{L}_{\mathcal{I}}} \sum_{\kappa \in \mathcal{I}^F} \tilde{Q}_\kappa \tilde{\mathcal{Y}}_\kappa(v), \sum_{\iota \in \mathcal{I}_{\mathbf{G}^F}, \gamma \in \mathcal{I}^F} \tilde{\mathcal{Y}}_\iota(u) R_{\iota,\gamma} \tilde{Q}_\gamma \rangle_{W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})F},$$

We use now [5, Corollary 5.2] which says that, $\langle \tilde{Q}_\gamma, \mathcal{Z}_{\mathbf{L}_{\mathcal{I}}} \tilde{Q}_\kappa \rangle_{W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})F} = 0$ unless $C_\gamma = C_\kappa$ and in this last case is equal to

$$|A(v)|^{-1} \sum_{a \in A(v)} |C_{\mathbf{L}}^0(v_a)^F| q^{-2c_\gamma} \mathcal{Y}_\gamma(v_a) \overline{\mathcal{Y}_\kappa(v_a)}$$

Thus the previous sum becomes

$$|\mathbf{L}^F|^{-1} \sum_{\iota \in \mathcal{I}_{\mathbf{G}^F}, \gamma \in \mathcal{I}^F} \overline{\tilde{\mathcal{Y}}_\iota(u)} R_{\iota,\gamma} |A(v)|^{-1} \sum_{a \in A(v)} |C_{\mathbf{L}}^0(v_a)^F| q^{-2c_\gamma} \mathcal{Y}_\gamma(v_a) \sum_{\kappa \in \mathcal{I}^F} \overline{\mathcal{Y}_\kappa(v_a)} \tilde{\mathcal{Y}}_\kappa(v).$$

But by [5, (4.5)] we have $\sum_{\kappa} \overline{\mathcal{Y}_\kappa(v_a)} \tilde{\mathcal{Y}}_\kappa(v) = \begin{cases} q^{c_\kappa} |A(v)|^F & \text{if } v_a = v \\ 0 & \text{otherwise} \end{cases}$, where κ runs

over all local systems. Thus, summing over all the blocks, we get the formula in the statement. \square

Corollary 5. *Assume either $Z\mathbf{G}$ is connected or q is large enough (depending just on the Dynkin diagram of \mathbf{G}). Then for any unipotent elements $u \in \mathbf{G}^F$ and $v \in \mathbf{L}^F$ we have:*

- (i) $Q_{\mathbf{L}}^{\mathbf{G}}(u, v)$ vanishes unless $v^{\mathbf{G}} \subseteq \overline{u^{\mathbf{G}}} \subseteq \text{Ind}_{\mathbf{L}}^{\mathbf{G}}(v^{\mathbf{L}})$, where $\text{Ind}_{\mathbf{L}}^{\mathbf{G}}(v^{\mathbf{L}})$ is the induced class in the sense of [14].
- (ii) $|v^{\mathbf{L}^F}| |A(v)| Q_{\mathbf{L}}^{\mathbf{G}}(u, v)$ is an integer and is a polynomial in q with integral coefficients.

Proof. For (i), we use [5, Lemma 6.9(i)] which states that $R_{\iota,\gamma} = 0$ unless $C_\gamma \subseteq \overline{C_\iota} \subseteq \text{Ind}_{\mathbf{L}}^{\mathbf{G}}(C_\gamma)$. Since $\tilde{\mathcal{Y}}_\kappa(v)$ vanishes unless $C_\kappa \ni v$, the only non-zero terms in the formula of proposition 4 have $C_\gamma \ni v$, whence the result since $\tilde{\mathcal{Y}}_\iota(u)$ vanishes unless $C_\iota \ni u$.

For (ii), we start with

Lemma 6. $q^{c_\iota - c_\gamma} R_{\iota,\gamma}$ is a polynomial in q with integral coefficients.

Proof. The defining equation of the matrix $\tilde{R} = \{q^{c_\iota - c_\gamma} R_{\iota,\gamma}\}_{\iota,\gamma}$ reads (see the proof of [5, Lemma 6.9(i)]):

$$\tilde{R} = P_{\mathbf{G}} C_{\mathbf{G}} I C_{\mathbf{L}}^{-1} P_{\mathbf{L}}^{-1}$$

where $C_{\mathbf{G}}$ is the diagonal matrix with diagonal coefficients q^{c_ι} for $\iota \in \mathcal{I}_{\mathbf{G}}$, and $C_{\mathbf{L}}$ is the similar matrix for \mathbf{L} and \mathcal{I} , where $P_{\mathbf{G}}$ is the matrix with coefficients $\{P_{\iota,\iota'}\}_{\iota,\iota' \in \mathcal{I}_{\mathbf{G}}}$ where these polynomials are those defined in [11, 6.5], and $P_{\mathbf{L}}$ is the similar matrix for \mathbf{L} and \mathcal{I} , and finally I is the matrix with coefficients

$$I_{\iota,\gamma} = \langle \text{Ind}_{W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})F}^{W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})F} \tilde{\varphi}_\gamma, \tilde{\varphi}_\iota \rangle_{W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})F}$$

where $\tilde{\varphi}_\gamma$ is the character of $W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})F$ which corresponds by the generalised Springer correspondence to γ (and similarly for $\tilde{\varphi}_\iota$). Since $P_{\mathbf{L}}$ and $P_{\mathbf{G}}$ are unitriangular matrices with coefficients integral polynomials in q , thus $P_{\mathbf{L}}^{-1}$ also, it suffices to prove that $C_{\mathbf{G}} I C_{\mathbf{L}}^{-1}$ has coefficients polynomial in q , or equivalently that

$$\text{if } \langle \text{Ind}_{W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})F}^{W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})F} \tilde{\varphi}_\gamma, \tilde{\varphi}_\iota \rangle_{W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})F} \neq 0, \text{ then } c_\iota - c_\gamma \geq 0.$$

We now use [5, Proposition 2.3(ii)] which says that the non-vanishing above implies $C_\gamma \subseteq \overline{C_\iota} \subseteq \text{Ind}_{\mathbf{L}}^{\mathbf{G}}(C_\gamma)$. We now use that, according to the definitions, $c_\iota - c_\gamma = \dim \mathcal{B}_u^{\mathbf{G}} - \dim \mathcal{B}_v^{\mathbf{L}}$ where $\mathcal{B}_u^{\mathbf{G}}$ is the variety of Borel subgroups of \mathbf{G} containing an element u of the support of ι , and where $\mathcal{B}_v^{\mathbf{L}}$ is the variety of Borel subgroups of \mathbf{L} containing an element v of the support of γ . Now the lemma follows from the fact that by [14, Theorem 1.3 (b)] we have $\dim \mathcal{B}_u^{\mathbf{G}} = \dim \mathcal{B}_v^{\mathbf{L}}$ if u is an element of $\text{Ind}_{\mathbf{L}}^{\mathbf{G}}(C_\gamma)$, and that $\dim \mathcal{B}_u^{\mathbf{G}}$ is greater for $u \in \overline{\text{Ind}_{\mathbf{L}}^{\mathbf{G}}(C_\gamma)} - \text{Ind}_{\mathbf{L}}^{\mathbf{G}}(C_\gamma)$. \square

Now (ii) results from the lemma: since the $\tilde{\mathcal{Y}}$ have values algebraic integers, by Proposition 4 the expression in (ii) is a polynomial in q with coefficients algebraic integers. But, since $|\mathbf{L}^F| Q_{\mathbf{L}}^{\mathbf{G}}(u, v)$ is a Lefschetz number (see for example [8, 8.1.3]), the expression in (ii) is a rational number; since this is true for an infinite number of integral values of q the expression in (ii) is a polynomial with integral coefficients. \square

SCALAR PRODUCTS OF INDUCED GELFAND-GRAEV CHARACTERS

The pretext for this section is as follows: in [2, Remark 3.10] is pointed the problem of computing $\langle R_{\mathbf{L}}^{\mathbf{G}} \Gamma_\iota, R_{\mathbf{L}}^{\mathbf{G}} \Gamma_\iota \rangle_{\mathbf{G}^F}$ when (\mathbf{G}, F) is simply connected of type 2E_6 , when \mathbf{L} is of type $A_2 \times A_2$, and when ι corresponds to a faithful character of $Z(\mathbf{L})/Z^0(\mathbf{L})$, and checking that the value is the same as given by the Mackey formula. We show now various ways to do this computation, where in this section we assume p and q large enough for all the results of [5] to hold (in particular, we assume p good for \mathbf{G} , thus not solving the problem of loc. cit. where we need $q = 2$).

Let $Z = Z(\mathbf{G})$, and let Γ_z be the Gelfand-Graev character parameterized by $z \in H^1(F, Z)$, see for instance [3, Definition 2.7]. Let u_z be a representative of the regular unipotent class parametrized by z . As in [11, 7.5 (a)] for ι an F -stable local system on the regular unipotent class we define $\Gamma_\iota = c \sum_{z \in H^1(F, Z)} \mathcal{Y}_\iota(u_z) \Gamma_z$ where $c = \frac{|Z/Z^0|}{|H^1(F, Z)|}$.

Note that the cardinality $|C_{\mathbf{G}}(u_z)^F|$ is independent of z ; actually it is equal to $|Z(\mathbf{G})^F|q^{\text{rkss}\mathbf{G}}$ (see [1, 15.5]). Thus we will denote this cardinality $|C_{\mathbf{G}^F}(u)|$ where $u \in \mathbf{G}^F$ is any regular unipotent element. There exists a character ζ of $H^1(F, Z)$ and a root of unity b_ι (see [4, above 1.5]) such that $\mathcal{Y}_\iota(u_z) = b_\iota \zeta(z)$. With these notations, we have

Proposition 7. *We have $\Gamma_\iota = \eta_{\mathbf{G}} \sigma_\zeta^{-1} c |C_{\mathbf{G}^F}(u)| D\mathcal{Y}_\iota$ where $\eta_{\mathbf{G}}$ and σ_ζ are defined as in [4, 2.5],*

Proof. This proposition could be deduced from [6, Theorem 2.8] using [4, Theorem 2.7]. We give here a more elementary proof.

With the notations of [3, (3.5')] we have $D\Gamma_z = \sum_{z' \in H^1(F, Z)} c_{z, z'} \gamma_{z'}$. By [4, lemma 2.3] we have $c_{z, z'} = c_{zz'^{-1}, 1}$ and $\sum_{z \in H^1(F, Z)} \zeta(z) c_{z, 1} = \eta_{\mathbf{G}} \sigma_\zeta^{-1}$. It follows that

$$\begin{aligned} c^{-1} b_\iota^{-1} D\Gamma_\iota &= \sum_{z \in H^1(F, Z)} \zeta(z) D\Gamma_z = \sum_{z, z' \in H^1(F, Z)} \zeta(z) c_{z, z'} \gamma_{z'} \\ &= \sum_{z, z' \in H^1(F, Z)} c_{zz'^{-1}, 1} \zeta(z) \gamma_{z'} \\ &= \sum_{z' \in H^1(F, Z)} \zeta(z') \gamma_{z'} \sum_{z'' \in H^1(F, Z)} c_{z'', 1} \zeta(z'') \\ &= \eta_{\mathbf{G}} \sigma_\zeta^{-1} \sum_{z' \in H^1(F, Z)} \zeta(z') \gamma_{z'} = \eta_{\mathbf{G}} \sigma_\zeta^{-1} b_\iota^{-1} |C_{\mathbf{G}^F}(u)| \mathcal{Y}_\iota \end{aligned}$$

□

Proposition 8. *If ι is a local system supported on the regular unipotent class of \mathbf{L} and \mathcal{I} denotes its block, we have*

$$\langle R_{\mathbf{L}}^{\mathbf{G}} \Gamma_\iota^{\mathbf{L}}, R_{\mathbf{L}}^{\mathbf{G}} \Gamma_\iota^{\mathbf{L}} \rangle_{\mathbf{G}^F} = \frac{|Z(\mathbf{L})|}{|Z^0(\mathbf{L})|^2} \sum_{w \in W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})} \frac{|Z^0(\mathbf{L}_{\mathcal{I}})^{wF} |W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})|}{|W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})|^2} \frac{|(wF)^{W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})} \cap W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})|}{|(wF)^{W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})}|}.$$

Note that in a given block \mathcal{I} there is at most one local system supported by the regular unipotent class (see [4, Corollary 1.10]).

Proof. When ι is supported by the regular unipotent class we have $\tilde{Q}_\iota = 1$, see the beginning of section 7, bottom of page 130 in [5]. Using this in the last formula of the proof of [5, Proposition 6.1], we get that $\Gamma_\iota^{\mathbf{L}}$ is up to a root of unity equal to $|A(C_\iota)| |W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})|^{-1} \sum_{w \in W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})} |Z^0(\mathbf{L}_{\mathcal{I}})^{wF}| Q_{wF}^{\mathbf{L}, \mathcal{I}}$. Since $R_{\mathbf{L}}^{\mathbf{G}} Q_{wF}^{\mathbf{L}, \mathcal{I}} = Q_{wF}^{\mathbf{G}, \mathcal{I}_{\mathbf{G}}}$, we get

$$\langle R_{\mathbf{L}}^{\mathbf{G}} \Gamma_\iota, R_{\mathbf{L}}^{\mathbf{G}} \Gamma_\iota \rangle_{\mathbf{G}^F} = |A(C_\iota)|^2 |W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})|^{-2} \sum_{w, w' \in W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})} |Z^0(\mathbf{L}_{\mathcal{I}})^{wF}| |Z^0(\mathbf{L}_{\mathcal{I}})^{w'F}| \langle Q_{wF}^{\mathbf{G}, \mathcal{I}_{\mathbf{G}}}, Q_{w'F}^{\mathbf{G}, \mathcal{I}_{\mathbf{G}}} \rangle_{\mathbf{G}^F}.$$

By [5, 3.5] the last scalar product is zero unless wF and $w'F$ are conjugate in $W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})$, and is equal to $|C_{W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})}(wF)|/|Z^0(\mathbf{L}_{\mathcal{I}})^{wF}|$ otherwise. We get

$$\begin{aligned} \langle R_{\mathbf{L}}^{\mathbf{G}}\Gamma_{\iota}, R_{\mathbf{L}}^{\mathbf{G}}\Gamma_{\iota'} \rangle_{\mathbf{G}^F} &= |A(C_{\iota})|^2 |W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})|^{-2} \\ &\quad \sum_{w \in W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})} |Z^0(\mathbf{L}_{\mathcal{I}})^{wF}| |C_{W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})}(wF)| |(wF)^{W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})} \cap W_{\mathbf{L}}(\mathbf{L}_{\mathcal{I}})|, \end{aligned}$$

which gives the formula of the proposition since $A(C_{\iota}) = Z(\mathbf{L})/Z(\mathbf{L})^0$. \square

Corollary 9. *Let ι and ι' be local systems supported on the regular unipotent class of \mathbf{G} , and $\mathcal{I}, \mathcal{I}'$ be their respective blocks: then*

$$\begin{aligned} \text{(i)} \quad \langle \Gamma_{\iota}^{\mathbf{G}}, \Gamma_{\iota'}^{\mathbf{G}} \rangle_{\mathbf{G}^F} &= \begin{cases} 0 & \text{if } \iota \neq \iota', \\ \left| \frac{Z(\mathbf{G})}{Z^0(\mathbf{G})} \right|^2 |Z^0(\mathbf{G})^F| q^{\dim Z(\mathbf{L}_{\mathcal{I}}) - \dim Z(\mathbf{G})} & \text{if } \iota = \iota'. \end{cases} \\ \text{(ii)} \quad \langle \mathcal{Y}_{\iota}, \mathcal{Y}_{\iota'} \rangle_{\mathbf{G}^F} &= \begin{cases} 0 & \text{if } \iota \neq \iota', \\ q^{-\text{rk}_{\text{ss}} \mathbf{G}} |Z^0(\mathbf{G})^F|^{-1} & \text{if } \iota = \iota'. \end{cases} \end{aligned}$$

Proof. The functions $Q_{wF}^{\mathbf{G}, \mathcal{I}_{\mathbf{G}}}$ and $Q_{w'F}^{\mathbf{G}, \mathcal{I}'_{\mathbf{G}}}$ are orthogonal to each other when $\mathcal{I}_{\mathbf{G}} \neq \mathcal{I}'_{\mathbf{G}}$ (see [9, V, 24.3.6] where the orthogonality is stated for the functions \mathcal{X}_i). Since there is a unique ι in a given block supported on the regular unipotent class, we get the orthogonality in (i). In the case $\iota = \iota'$ in (i), the specialization $\mathbf{L} = \mathbf{G}$ in Proposition 8 is $\langle \Gamma_{\iota}^{\mathbf{G}}, \Gamma_{\iota'}^{\mathbf{G}} \rangle_{\mathbf{G}^F} = \left| \frac{Z(\mathbf{G})}{Z^0(\mathbf{G})} \right|^2 \sum_{w \in W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})} \frac{|Z^0(\mathbf{L}_{\mathcal{I}})^{wF}|}{|W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})|}$. By [5, Corollary 5.2], where we use that $\tilde{Q}_{\iota} = 1$ when ι has regular support, we have $\sum_{w \in W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})} \frac{|Z^0(\mathbf{L}_{\mathcal{I}})^{wF}|}{|W_{\mathbf{G}}(\mathbf{L}_{\mathcal{I}})|} = q^{-2c_{\iota}} |C_{\mathbf{G}}(u)^{0F}|$. Whence $\langle \Gamma_{\iota}^{\mathbf{G}}, \Gamma_{\iota'}^{\mathbf{G}} \rangle_{\mathbf{G}^F} = \left| \frac{Z(\mathbf{G})}{Z^0(\mathbf{G})} \right|^2 q^{-\text{rk} \mathbf{G} + \dim Z(\mathbf{L}_{\mathcal{I}})} |C_{\mathbf{G}}^0(u)^F|$. Using $|C_{\mathbf{G}}^0(u)^F| = q^{\text{rk}_{\text{ss}} \mathbf{G}} |Z^0(\mathbf{G})^F|$, we get (i).

For (ii), we apply Proposition 7 in (i), using that D is an isometry and that $\sigma_{\zeta} \bar{\sigma}_{\zeta} = q^{\text{rk}_{\text{ss}} \mathbf{L}_{\mathcal{I}}}$ by [4, proposition 2.5]. \square

A particular case of Proposition 8 is

Corollary 10. *If (\mathbf{L}, ι) is a cuspidal pair, that is $\mathbf{L} = \mathbf{L}_{\mathcal{I}}$, then*

$$\langle R_{\mathbf{L}}^{\mathbf{G}}\Gamma_{\iota}^{\mathbf{L}}, R_{\mathbf{L}}^{\mathbf{G}}\Gamma_{\iota}^{\mathbf{L}} \rangle_{\mathbf{G}^F} = \left| \frac{Z(\mathbf{L})}{Z^0(\mathbf{L})} \right|^2 |W_{\mathbf{G}}(\mathbf{L})| |Z(\mathbf{L})^{0F}|.$$

We remark that this coincides with the value predicted by the Mackey formula

$$\langle R_{\mathbf{L}}^{\mathbf{G}}\Gamma_{\iota}^{\mathbf{L}}, R_{\mathbf{L}}^{\mathbf{G}}\Gamma_{\iota}^{\mathbf{L}} \rangle_{\mathbf{G}^F} = \sum_{x \in \mathbf{L}^F \backslash \mathcal{S}(\mathbf{L}, \mathbf{L}) / \mathbf{L}^F} \langle {}^*R_{\mathbf{L} \cap {}^x\mathbf{L}}^{\mathbf{L}}(\Gamma_{\iota}^{\mathbf{L}}), {}^*R_{\mathbf{L} \cap {}^x\mathbf{L}}^{{}^x\mathbf{L}}({}^x\Gamma_{\iota}^{\mathbf{L}}) \rangle_{(\mathbf{L} \cap {}^x\mathbf{L})^F}$$

Indeed, since the block \mathcal{I} which contains the local system ζ is reduced to the unique cuspidal local system (C, ζ) where C is the regular class of \mathbf{L} , all terms in the Mackey formula where $\mathbf{L} \cap {}^x\mathbf{L} \neq \mathbf{L}$ vanish. Thus the Mackey formula reduces to

$$\langle R_{\mathbf{L}}^{\mathbf{G}}\Gamma_{\iota}^{\mathbf{L}}, R_{\mathbf{L}}^{\mathbf{G}}\Gamma_{\iota}^{\mathbf{L}} \rangle_{\mathbf{G}^F} = \sum_{x \in W_{\mathbf{G}}(\mathbf{L})} \langle \Gamma_{\iota}^{\mathbf{L}}, {}^x\Gamma_{\iota}^{\mathbf{L}} \rangle_{\mathbf{L}^F}$$

and any x in $W_{\mathbf{L}}(\mathbf{L})$ acts trivially on $H^1(F, Z(\mathbf{L}))$ since, the map $h_{\mathbf{L}}$ being surjective, any element of $H^1(F, Z(\mathbf{L}))$ is represented by an element of $H^1(F, Z(\mathbf{G}))$; thus all the terms in the sum are equal, and we get the same result as Corollary 10 by applying Corollary 10 in the case $\mathbf{G} = \mathbf{L}$.

Another method for computing $\langle R_{\mathbf{L}}^{\mathbf{G}}D\Gamma_i, R_{\mathbf{L}}^{\mathbf{G}}D\Gamma_i \rangle_{\mathbf{G}^F}$ would be to use Proposition 1 and the values of the two-variable Green functions.

We give these values in the following table in the particular case of 2E_6 for the F -stable standard Levi subgroup of type $A_2 \times A_2$, for $q \equiv -1 \pmod{3}$, so that F acts trivially on $Z(\mathbf{G})/Z^0(\mathbf{G})$. This table has been computed in **Chevie** using Proposition 3. The method is to compute the one-variable Green functions which appear in the right-hand side sum by the Lusztig-Shoji algorithm; note that even though the characteristic function of cuspidal character sheaves are known only up to a root of unity, this ambiguity disappears when doing the sum, since such a scalar appears multiplied by its complex conjugate. However the Lusztig-Shoji algorithm depends also on the knowledge that when ι , with support the class of the unipotent element u , is parameterized by (u, χ) where $\chi \in \text{Irr}(A(u))$ then for the unipotent element $u_a \in \mathbf{G}^F$ parameterized by $a \in H^1(F, A(u))$ we have $\mathcal{Y}_\iota(u_a) = \chi(a)$. We assume that this hold. This is known when ι is in the principal block, but not for the two blocks with cuspidal datum supported on the Levi subgroup of type $A_2 \times A_2$.

Note that the table shows that the values of $|v^{\mathbf{L}^F}|Q_{\mathbf{L}}^{\mathbf{G}}(u, v)$ are not in general polynomials with integral coefficients but may have denominators equal to $|A(v)|$.

TABLE 1. Values of $|v^{\mathbf{L}^F}|Q_{\mathbf{L}}^{\mathbf{G}}(u, v)$ for $\mathbf{G} = {}^2E_6(q)$ simply connected and $\mathbf{L} = A_2(q^2)(q-1)^2$, for $q \equiv -1 \pmod{3}$.

$v \setminus u$	E_6	$E_{6(\zeta_3)}$	$E_{6(\zeta_3^2)}$	$E_6(a_1)$	$E_{6(a_1)(\zeta_3)}$	$E_{6(a_1)(\zeta_3^2)}$	D_5	$E_6(a_3)$
111, 111	0	0	0	0	0	0	0	0
21, 21	0	0	0	0	0	0	1	$4q+1$
3, 3	1	0	0	$(4q+1)/3$	$\Phi_2/3$	$\Phi_2/3$	$2q\Phi_2/3$	$(7q^2+2q-2)q/3$
$3, 3_{(\zeta_3)}$	0	1	0	$\Phi_2/3$	$(4q+1)/3$	$\Phi_2/3$	$2q\Phi_2/3$	$(q-2)q\Phi_2/3$
$3, 3_{(\zeta_3^2)}$	0	0	1	$\Phi_2/3$	$\Phi_2/3$	$(4q+1)/3$	$2q\Phi_2/3$	$(q-2)q\Phi_2/3$

$v \setminus u$	$E_6(a_3)_{(-\zeta_3^2)}$	$E_6(a_3)_{(\zeta_3)}$	$E_6(a_3)_{(-1)}$	$E_6(a_3)_{(\zeta_3^2)}$	$E_6(a_3)_{(-\zeta_3)}$
111, 111	0	0	0	0	0
21, 21	$2q+1$	$4q+1$	$2q+1$	$4q+1$	$2q+1$
3, 3	$q^2\Phi_2$	$(q-2)q\Phi_2/3$	$(3q+2)q^2$	$(q-2)q\Phi_2/3$	$q^2\Phi_2$
$3, 3_{(\zeta_3)}$	$(3q+2)q^2$	$(q-2)q\Phi_2/3$	$q^2\Phi_2$	$(7q^2+2q-2)q/3$	$q^2\Phi_2$
$3, 3_{(\zeta_3^2)}$	$q^2\Phi_2$	$(7q^2+2q-2)q/3$	$q^2\Phi_2$	$(q-2)q\Phi_2/3$	$(3q+2)q^2$

$v \setminus u$	A_5	$A_5(\zeta_3)$	$A_5(\zeta_3^2)$	$D_5(a_1)$
111, 111	0	0	0	0
21, 21	$(-2q-1)\Phi_2$	$(-2q-1)\Phi_2$	$(-2q-1)\Phi_2$	$3q+1$
3, 3	$q\Phi_2\Phi_3/3$	$(-5q^2-2q+1)q\Phi_2/3$	$(-5q^2-2q+1)q\Phi_2/3$	$q\Phi_1\Phi_2/3$
$3, 3_{(\zeta_3)}$	$(-5q^2-2q+1)q\Phi_2/3$	$q\Phi_2\Phi_3/3$	$(-5q^2-2q+1)q\Phi_2/3$	$q\Phi_1\Phi_2/3$
$3, 3_{(\zeta_3^2)}$	$(-5q^2-2q+1)q\Phi_2/3$	$(-5q^2-2q+1)q\Phi_2/3$	$q\Phi_2\Phi_3/3$	$q\Phi_1\Phi_2/3$

$v \setminus u$	A_4+A_1	D_4	A_4	$D_4(a_1)_{(111)}$
111, 111	0	1	0	$4q+1$
21, 21	$\Phi_2\Phi_3$	$3q\Phi_2\Phi_6$	$(3q^3+q^2+q+1)\Phi_2$	$(8q^3+2q^2+4q-2)q\Phi_2$
3, 3	$(2q+1)q^3\Phi_2/3$	$q\Phi_1\Phi_2^2\Phi_6/3$	$q^4\Phi_2^2$	$(4q+1)q^4\Phi_2^2/3$
$3, 3_{(\zeta_3)}$	$(2q+1)q^3\Phi_2/3$	$q\Phi_1\Phi_2^2\Phi_6/3$	$q^4\Phi_2^2$	$(4q+1)q^4\Phi_2^2/3$
$3, 3_{(\zeta_3^2)}$	$(2q+1)q^3\Phi_2/3$	$q\Phi_1\Phi_2^2\Phi_6/3$	$q^4\Phi_2^2$	$(4q+1)q^4\Phi_2^2/3$

$v \setminus u$	$D_4(a_1)_{(21)}$	$D_4(a_1)$	A_3+A_1	A_3
111, 111	$2q+1$	Φ_2	$(-2q-1)\Phi_2$	$(3q^3+2q+1)\Phi_2$
21, 21	$(8q^3+6q^2+2q+2)q^2$	$(2q+1)q\Phi_2\Phi_6$	$(-4q^3-q^2-2q+1)q\Phi_2^2$	$(3q^3-q^2+2q-1)q\Phi_2^2\Phi_4$
3, 3	$(2q+1)q^4\Phi_1\Phi_2/3$	$q^4\Phi_2\Phi_6/3$	$(-2q-1)q^5\Phi_2^2/3$	0
$3, 3_{(\zeta_3)}$	$(2q+1)q^4\Phi_1\Phi_2/3$	$q^4\Phi_2\Phi_6/3$	$(-2q-1)q^5\Phi_2^2/3$	0
$3, 3_{(\zeta_3^2)}$	$(2q+1)q^4\Phi_1\Phi_2/3$	$q^4\Phi_2\Phi_6/3$	$(-2q-1)q^5\Phi_2^2/3$	0

$v \setminus u$	$2A_2 + A_1$	$2A_2 + A_1(\zeta_3)$	$2A_2 + A_1(\zeta_3^2)$	$2A_2$
111, 111	$\Phi_2 \Phi_3$	$\Phi_2 \Phi_3$	$\Phi_2 \Phi_3$	$\Phi_2^2 \Phi_3 \Phi_6$
21, 21	$(2q^3 + 2q^2 + 4q + 1)q^3 \Phi_2$	$(2q^3 + 2q^2 + 4q + 1)q^3 \Phi_2$	$(2q^3 + 2q^2 + 4q + 1)q^3 \Phi_2$	$3q^4 \Phi_2^2 \Phi_3 \Phi_6$
3, 3	$q^6 \Phi_2 \Phi_3$	0	0	$q^6 \Phi_2^2 \Phi_3 \Phi_6$
$3, 3(\zeta_3)$	0	$q^6 \Phi_2 \Phi_3$	0	0
$3, 3(\zeta_3^2)$	0	0	$q^6 \Phi_2 \Phi_3$	0

$v \setminus u$	$2A_2(\zeta_3)$	$2A_2(\zeta_3^2)$	$A_2 + 2A_1$	$A_2 + A_1$
111, 111	$\Phi_2^2 \Phi_3 \Phi_6$	$\Phi_2^2 \Phi_3 \Phi_6$	$(2q^4 + q^3 + q^2 + q + 1)\Phi_2$	$\Phi_2^2 \Phi_3 \Phi_6$
21, 21	$3q^4 \Phi_2^2 \Phi_3 \Phi_6$	$3q^4 \Phi_2^2 \Phi_3 \Phi_6$	$(q^3 + 2q + 1)q^5 \Phi_2$	$(3q^2 + 2q + 1)q^4 \Phi_2^2 \Phi_6$
3, 3	0	0	0	0
$3, 3(\zeta_3)$	$q^6 \Phi_2^2 \Phi_3 \Phi_6$	0	0	0
$3, 3(\zeta_3^2)$	0	$q^6 \Phi_2^2 \Phi_3 \Phi_6$	0	0

$v \setminus u$	$A_2(11)$	A_2
111, 111	$(3q^5 + q^2 + q + 1)\Phi_2^2 \Phi_6$	$5q^9 + 3q^8 + 4q^7 + 4q^6 + 5q^5 + 4q^4 + 2q^3 + 2q^2 + 2q + 1$
21, 21	$(4q^2 + q + 1)q^4 \Phi_2^3 \Phi_6^2$	$(2q + 1)q^4 \Phi_1 \Phi_2 \Phi_3 \Phi_4 \Phi_6$
3, 3	0	0
$3, 3(\zeta_3)$	0	0
$3, 3(\zeta_3^2)$	0	0

$v \setminus u$	$3A_1$	$2A_1$
111, 111	$(-3q^9 - 3q^8 - 3q^6 - 2q^4 - q^3 - q^2 - q - 1)\Phi_2$	$(2q^8 + q^6 + q^5 + q^4 + 1)\Phi_2^2 \Phi_3 \Phi_6$
21, 21	$(-2q^2 - 1)q^7 \Phi_2^3 \Phi_6$	$q^9 \Phi_2^3 \Phi_3 \Phi_4 \Phi_6$
3, 3	0	0
$3, 3(\zeta_3)$	0	0
$3, 3(\zeta_3^2)$	0	0

$v \setminus u$	A_1	1
111, 111	$(2q^{10} + q^9 + q^8 + q^7 + 2q^6 + q^5 + q^4 + q^2 + q + 1)\Phi_2^3 \Phi_6 \Phi_{10}$	$\Phi_2^4 \Phi_3 \Phi_4 \Phi_6^2 \Phi_8 \Phi_{10} \Phi_{12} \Phi_{18}$
21, 21	0	0
3, 3	0	0
$3, 3(\zeta_3)$	0	0
$3, 3(\zeta_3^2)$	0	0

REFERENCES

- [1] C. Bonnafé, Actions of relative Weyl groups II *J. Group Theory* **8**(2005) 351–387
- [2] C. Bonnafé and J. Michel, A computational proof of the Mackey formula for $q > 2$, *J. Algebra* **327**(2011) 506–526.
- [3] F. Digne, G. Lehrer and J. Michel, The characters of the group of rational points of a reductive group with non-connected centre, *J. Crelle* **425**(1992) 155–192.
- [4] F. Digne, G. Lehrer and J. Michel, On Gel'fand-Graev characters of reductive groups with disconnected centre, *J. Crelle* **491**(1997), 131–147.
- [5] F. Digne, G. Lehrer and J. Michel, The space of unipotently supported class functions on a reductive group, *J. of Algebra* **260**(2003), 111–137.
- [6] F. Digne, G. Lehrer and J. Michel, On character sheaves and characters of reductive groups at unipotent elements, *Pure Appl. Math. Quart.* **10** (2014) 459–512.
- [7] F. Digne, and J. Michel, Foncteurs de Lusztig et caractères des groupes linéaires et unitaires sur un corps fini, *J. Algebra* **107** (1987) 217–255.
- [8] F. Digne, and J. Michel, Representations of finite groups of Lie type, *LMS student books* **95** (2020) CUP.
- [9] G. Lusztig, Character Sheaves V, *Adv. Math.* **61** (1986) 103–155
- [10] G. Lusztig, Green functions and character sheaves *Ann. of Math.* **131** (1990) 355–408

- [11] G. Lusztig, A unipotent support for irreducible representations, *Adv. Math.* **94**(1992) 139–179.
- [12] G. Lusztig, On the cleanness of cuspidal character sheaves, *Moscow Math. J.* **12** (2012) 621–631.
- [13] G. Lusztig, Comments on my papers, *ArXiv 1707.09368v6*
- [14] G. Lusztig and N. Spaltenstein, Induced unipotent classes, *J. LMS* **19** (1979) 41–52.
- [15] G. Malle and E. Rotilio, The 2-parameter Green functions for 8-dimensional spin groups, *ArXiv 2003.14231*.
- [16] J. Michel, The development version of the **Chevie** package of **GAP.**, *J. Algebra* **435** (2015), 308–336.
- [17] T. Shoji, Character sheaves and almost characters of reductive groups, II *Adv. Math.* **111** (1995) 314–354
- [18] T. Shoji, On the computation of unipotent characters of finite classical groups. *Appl. Algebra Engrg. Comm. Comput.* **7** (1996) 165–174

(F. Digne) LABORATOIRE AMIÉNOIS DE MATHÉMATIQUE FONDAMENTALE ET APPLIQUÉE, CNRS UMR 7352, UNIVERSITÉ DE PICARDIE-JULES VERNE, 80039 AMIENS CEDEX FRANCE.

Email address: digne@u-picardie.fr

URL: www.lamfa.u-picardie.fr/digne

(J. Michel) INSTITUT MATHÉMATIQUE DE JUSSIEU – PARIS RIVE GAUCHE, CNRS UMR 7586, UNIVERSITÉ DE PARIS, BÂTIMENT SOPHIE GERMAIN, 75013, PARIS FRANCE.

Email address: jean.michel@imj-prg.fr

URL: webusers.imj-prg.fr/~jean.michel