

# Importance of systematic right ventricular assessment in cardiac resynchronization therapy candidates: a machine-learning approach

E Galli, V Le Rolle, O A Smiseth, J Duchenne, J M Aalen, C K Larsen, E Sade, A Hubert, S Anilkumar, M Penicka, et al.

#### ► To cite this version:

E Galli, V Le Rolle, O A Smiseth, J Duchenne, J M Aalen, et al.. Importance of systematic right ventricular assessment in cardiac resynchronization therapy candidates: a machine-learning approach. Journal of The American Society of Echocardiography, 2021, 34 (5), pp.494-502. 10.1016/j.echo.2020.12.025 . hal-03128136

## HAL Id: hal-03128136 https://hal.science/hal-03128136

Submitted on 8 Mar 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Importance of systematic right ventricular assessment in cardiac resynchronization therapy candidates: a machine-learning approach

Short title: machine learning and right ventricular function in CRT

Galli E, MD<sup>1</sup>\*, PhD; Le Rolle V<sup>1</sup>\*, PhD; Smiseth OA<sup>2</sup>, MD, PhD; Duchenne J<sup>3,4</sup>, PhD; Aalen JM<sup>2</sup>, Larsen CK<sup>2</sup>, MD; Sade E<sup>5</sup>, MD, PhD; Hubert A<sup>1</sup>, MD; Anilkumar S<sup>6</sup>, MD; Penicka M<sup>7</sup>, MD, PhD; Cecilia Linde<sup>8</sup>, MD, PhD; Leclercq C<sup>1</sup>, MD, PhD; Hernandez A<sup>1</sup>, PhD; Voigt J-U<sup>3,4</sup>, MD, PhD; Donal E<sup>1</sup>, MD, PhD.

\* These two authors equally contributed to this work.

<sup>1</sup>Univ Rennes, CHU Rennes, Inserm, LTSI – UMR 1099, F-35000 Rennes, France
<sup>2</sup>Institute for Surgical Research and Department of Cardiology, Oslo University Hospital and University of Oslo, Norway
<sup>3</sup>Department of Cardiovascular Disease, KU Leuven, Belgium
<sup>4</sup>Department of Cardiovascular Science, KU Leuven, Belgium
<sup>5</sup>Department of Cardiology, Baskent University Hospital, Ankara, Turkey
<sup>6</sup>Non-Invasive Cardiac Laboratory, Department of Cardiology, Heart Hospital, Hamad Medical Corporation, Doha, Qatar
<sup>7</sup>Cardiovascular Center Aalst, OLV Clinic, Aalst, Belgium
<sup>8</sup>Heart and Vascular Theme, Karolinska University Hospital and Karolinska Institutet, Stockholm,

Sweden

#### **Corresponding Author:**

Dr. Elena Galli & Pr Erwan DONAL

CHU Pontchaillou

2 Rue Henri Le Guilloux, 35000 Rennes - FRANCE

Tel.: +33 2 99 28 78 96

Fax: +33 2 99 28 25 29

E-mail: elena.galli@chu-rennes.fr; erwan.donal@chu-rennes.fr

#### **Declaration of interest:** Nothing to disclose

#### HIGHLIGHTS

- The prediction of the impact of CRT on LV function and outcomes is often difficult
- CRT candidates are a highly heterogeneous population
- Machine learning allows the analysis of a huge amount of clinical and imaging data
- Machine learning can identify clusters of patients with different characteristics and prognosis
- RV-derived features are important for the characterization of CRT-candidates

#### ABSTRACT

#### Background

Despite all having systolic heart failure and broad QRS, patients screened for cardiac resynchronization therapy (CRT) are highly heterogeneous, and it remains extremely challenging to predict the impact of the device on left ventricular (LV) function and outcomes.

#### **Objectives**

We sought to evaluate the relative impact of clinical, electrocardiographic, and echocardiographic data on the left ventricular (LV) remodeling and prognosis of CRT-candidates by the application of machine learning (ML) approaches.

#### Methods

193 patients with systolic heart failure undergoing CRT according to current recommendations were prospectively included in this multicentre study. We used a combination of the Boruta algorithm and random forest methods to identify features predicting both CRT volumetric response and prognosis. The model performance was tested by the area under the receiver operating curve (AUC). We also applied the K-medoid method to identify clusters of phenotypically similar patients.

#### Results

From 28 clinical, electrocardiographic, and echocardiographic-derived variables, 16 features were predictive of CRT response, and 11 features were predictive of prognosis.

Among the predictors of CRT-response, 8 variables (50%) pertained to right ventricular (RV) size or function. Tricuspid annular plane systolic excursion was the main feature associated with prognosis. The selected features were associated with a particularly good prediction of both CRT response (AUC 0.81, 95% CI: 0.74-0.87) and outcomes (AUC 0.84, 95% CI: 0.75-0.93). An unsupervised ML approach allowed the identifications of two phenogroups of patients who differed significantly in clinical variables and parameters of biventricular size, and RV function. The two phenogroups had significant different prognosis (HR 4.70, 95% CI: 2.1-10.0, p<0.0001; log-rank p<0.0001).

#### Conclusions

Machine learning can reliably identify clinical and echocardiographic features associated with CRTresponse and prognosis. The evaluation of both RV-size and function parameters has pivotal importance for the risk stratification of CRT-candidates and should be systematically assessed in patients undergoing CRT.

Keywords: cardiac resynchronization therapy, heart failure, machine learning, right ventricle

#### Abbreviations

- ApR, apical rocking
- AUC, area under the curve
- BNP, brain-type natriuretic peptide
- CRT, cardiac resynchronization therapy
- EDV, end-diastolic volume
- ESV, end-systolic volume
- FAC, fractional area change
- GFR, glomerular filtration rate
- GLS, global longitudinal strain
- HFrEF, heart failure with reduced ejection fraction
- IHD, ischemic heart disease
- LAVi, indexed left atrial volume
- LBBB, left bundle branch block
- LV, left ventricle
- LVEF, left ventricular ejection fraction
- NYHA, New York Heart Association functional class.
- PAPs, estimated systolic pulmonary artery pressure
- RAP, right atrial pressure
- RAV, right atrial volume
- RV, right ventricle
- RVDA, right ventricular diastolic area
- RVSA, right ventricular systolic area
- RVLS, right ventricular free wall longitudinal strain

SF, septal flash

TAPSE, tricuspid annular plane systolic excursion

TRVmax, maximal tricuspid regurgitation velocity

#### **INTRODUCTION**

Cardiac resynchronization therapy (CRT) is an established treatment for patients with heart failure and reduced ejection fraction (HFrEF) who have wide QRS and remain symptomatic despite optimized medical therapy<sup>1</sup>. Nevertheless, nearly 30% of patients undergoing CRT according to recommendations are not-responders to this treatment<sup>2</sup>, and some patients die from heart failure despite the initial improvement in left ventricular ejection fraction (LVEF) and cardiac remodeling<sup>34</sup>. CRT-response and the outcome rely upon several factors which include clinical characteristics, typical ventricular conduction disturbances<sup>5</sup>, and the evaluation of the specific electromechanical substrate responsible for LV discoordination<sup>67</sup>.

The comprehensive integration and interpretation of this huge mass of heterogeneous data is difficult for the human brain and makes CRT an interesting field of application for personalized medicine<sup>8</sup>. Previous studies have shown that the application of machine learning (ML) algorithms to CRT can predict CRT response<sup>910</sup> and prognosis<sup>11</sup>. Some of these studies were focused on the analysis of specific electrocardiographic data such as QRS morphology and duration<sup>11</sup>, on the computational analysis of strain derived-curves<sup>10</sup>, or on the unsupervised examination of strain curve dynamics, clinical and standard echocardiographic data<sup>9</sup>. Our study aimed to apply supervised and nonsupervised machine learning approaches to a comprehensive amount of commonly available clinical, electrocardiographic, and echocardiographic data to 1) evaluate the relative importance of each feature in the prediction of CRT-response and prognosis by a supervised machine learning approach and 2) to identify specific phenogroups of patients which are at increased risk of poor outcomes.

#### **MATERIALS AND METHODS**

#### **Population**

209 patients with systolic heart failure undergoing CRT implantation according to current guidelines<sup>1</sup> at Oslo University Hospital (Norway), Leuven University Hospital (Belgium), Rennes

University Hospital (France), Aalst OLV Hospital (Belgium), and Karolinska University Hospital (Sweden) between August 2015 and November 2017 were prospectively included in this observational, multicentre study. Sixteen patients were excluded from the final analysis because of study withdrawal (n=4), lead extraction due to infective endocarditis (n=1), and lack of fundamental echocardiographic data (n=11). As a result, 193 patients were included in the study. At the time of CRT implantation, all patients were receiving optimized medical therapy. Clinical data including age, sex, and treatments were collected for each patient. The functional status was assessed by the estimation of the New York Heart Association (NYHA) functional class. Ischemic heart disease (IHD) was defined as a history of myocardial infarction and coronary revascularization or angiographic evidence of multiple vessel disease or single-vessel disease with  $\geq$ 75% stenosis of the left main or proximal left anterior descending artery<sup>12</sup>.

All patients gave their written informed consent for study participation.

The study was conducted following the "Good Clinical Practice" guidelines of the Declaration of Helsinki and was approved by the Regional Ethical Committees of every participating centre. The study was registered at clinicaltrials.gov (identifier NCT02525185).

#### *Echocardiography*

All patients underwent standard transthoracic echocardiography using a Vivid E9 and E95 ultrasound system (GE Healthcare, Horten, Norway) equipped with an M5S 3.5-MHz transducer at baseline and 6-month follow-up. Two-dimensional, colour Doppler, pulsed-wave and continuouswave Doppler data were stored on a dedicated workstation for the offline analysis (EchoPAC, GE EchoPAC, GE Healthcare, Horten, Norway). Left atrial, left ventricular (LV) volumes, and left ventricular ejection fraction (LVEF) were measured by the biplane method, as recommended<sup>13</sup>. Right ventricular size and function were measured in the modified four-chamber view as recommended<sup>13</sup>. Tricuspid annular plane systolic excursion (TAPSE) was measured in the apical 4chamber view using M-mode echocardiography. Right atrial volume (RAV) was measured using the disk summation techniques in a dedicated apical four-chamber view<sup>14</sup>. RV fractional area change (FAC) was measured as the percentage difference between RV diastolic and systolic area, divided by RV diastolic area. RV lateral strain was assessed by tracking the RV endocardium as indicated in recommendations<sup>14</sup>. Peak velocity of early (E) and late (A) diastolic filling were derived from transmitral Doppler recordings, and the E/A ratio was calculated. Pulsed-wave TDIderived early diastolic velocity were obtained at the septal and lateral mitral annulus and the mean value (e') was used to estimate the E/e' ratio. In the presence of tricuspid regurgitation, continuous Doppler was used to estimate the maximal tricuspid velocity (TRV<sub>max</sub>). Inferior vena cava diameter and respiratory changes were used to estimate right atrial pressure (RAP)<sup>15</sup>. Systolic pulmonary artery pressure (PAPs) was calculated as  $[4*(TRV_{max})^2+RAP]$  as recommended<sup>15</sup>. Right ventriculoarterial coupling was assessed by the ratio between tricuspid annular plane systolic excursion and PAPs (TAPSE/PAPs)<sup>16</sup>.

#### Assessment of LV dyssynchrony

Septal flash (SF) and apical rocking (ApR) were visually assessed in Leuven by two experienced readers. In case of disagreement, a third reading was performed in Leuven by an independent expert to reach a consensus. Septal flash was defined as pre-ejection septal shortening or rapid leftward septal motion immediately after onset QRS and was assessed visually in apical 2D images or, when in doubt, with longitudinal strain or M-mode in parasternal views<sup>17</sup>. Apical rocking was defined as a transverse rightward motion of the apex immediately after onset QRS, followed by a leftward motion of the apex during ejection<sup>18</sup>. LV mechanical dyssynchrony was defined by the presence of SF and/or ApR.

#### Cardiac resynchronization therapy

CRT delivery followed a standard protocol. The right atrial and ventricular leads were positioned conventionally. The LV lead was inserted in a lateral or postero-lateral vein if possible and coronary venography was used to optimize lead placement. The device was programmed in conventional biventricular pacing and retested before hospital discharge. Positive response to CRT was defined as a decrease in LV end-systolic volume of  $\geq 15\%^2$ . To optimize precision, all volumes were measured independently in three different centres (Rennes, Leuven and Oslo) and a majority decision was used in cases of disagreement on CRT-response.

#### Outcome

The outcome was a composite of heart transplantation, LV assisted device implantation or allcause death during follow-up.

#### Baseline characteristics and machine learning approaches

Twenty-two baseline echocardiographic variables, with <10% missing data were identified. After filtering correlated variables using a cut-off Pearsons' coefficient>0.8, 18 variables, which included right and left ventricular function parameters, were selected. In addition to these echocardiographic data, the following variables were included in the machine learning (ML) algorithm: 1) clinical parameters: age, sex, NYHA class, IHD, ln NT-proBNP and glomerular filtration rate (GFR); 2) ECG-derived parameters, such as QRS duration and presence of typical left bundle branch block<sup>19</sup>; 3) dyssynchrony parameters and 4) CRT-volumetric response. Implementations of data analysis and machine learning methods were performed in R-studio (version 1.2.1335).

Missing data were imputed using the MICE algorithm as described by Van Buuren et Groothuis-Oudshoorn et-al.<sup>20</sup> ("mice" package) which allows imputing mixed datasets with continuous and categorical variables using Fully Conditional Specification (FCS) (Table 1S).

#### Unsupervised machine-learning approach

To identify clusters of phenotypically-similar patients, subjects were clustered using the partitioning around medoids (PAM or k-medoid) algorithm with Gower distance, which allows the analysis of datasets made of mixed-type data with numeric and categorical features ("cluster" package). Figure 1 shows an overview of the input data and the proposed data analysis. The final number of clusters was determined to maximize the silhouette index<sup>21</sup>.

#### Supervised machine-learning approach

A feature selection phase was firstly applied by using the Boruta algorithm<sup>22</sup> ("Boruta" package) to select the best predictors of CRT-response and outcome. According to this method, each original feature is compared to the randomly generated features created by the algorithm. A feature that contributes positively to the predictive model has a performance which is superior to the best random feature, indicated as "shadowMax".

The random forest (RF) method <sup>23</sup> ("randomForest" package) is a popular and versatile supervised ensemble learner method that is based on the training of a set of uncorrelated trees to estimate a target variable for classification or regression purposes. This method was then applied to a reduced feature set to build ensemble classifiers to predict CRT-response and outcome. Receiver Operating Characteristic (ROC) curves and the corresponding area under the curve (AUC) were estimated to evaluate the classification performance of the RF method proposed method.

#### Statistical analysis

The normality of data distribution was assessed using the Shapiro-Wilk test. When the hypothesis of normality was rejected, a Box-Cox transformation was performed. Results were rounded and expressed as mean  $\pm$  SD. The differences between the phenotype groups were tested using a one-way ANOVA analysis. Categorical variables were expressed as percentages. Comparison between groups was performed using the  $\chi^2$  test or Fisher's exact test when appropriate. Statistical significance was considered as a two-sided p-value <0.05. We used Cox regression models to calculate between-group differences in outcomes and the Kaplan–Meier method to calculate survival curves.

We used R statistical software (version 3.3.3) and SPSS (SPSS Version 20.0, IBM, Chicago, IL, USA) for statistical analysis.

#### RESULTS

The baseline characteristics of the population are depicted in Table 1.

The population had a mean age of 67 years. The majority of patients were males (70%), with a high prevalence (67%) of non-ischemic dilated cardiomyopathy. 168 (87%) patients had typical LBBB.

#### **CRT** response

At 6-month follow-up, successful response to CRT was observed in 132 (68%) patients. After the application of the Boruta algorithm, 16 features were significantly associated with CRT-response (Figure 2A). Among all variables, 8 (50%) were obtained through the assessment of RV size or function. SF and ApR were the most important features, followed by IHD, E/e' ratio, RAV, RV diameters and RV systolic area, LV end-diastolic diameter, RV free wall strain, TAPSE/PAPs ratio, QRS width, LV-end diastolic volume, LV end-systolic diameter, TRV<sub>max</sub> and FAC. The model obtained by the application of the RF method to this ensemble of features showed an AUC 0.81 (95% CI: 0.75-0.87) for the prediction of CRT-response (Figure 3A).

#### **Predictors of prognosis**

During a median follow-up of 37 months, the primary endpoint occurred in 29 (15%) patients. There were 11 non-cardiac deaths, 16 cardiac-related deaths, one heart transplantation and one LV assist device implantation.

The main features associated with the outcome are displayed in Figure 2B. The most important feature was TAPSE, followed by LV end-diastolic volume, LV end-systolic diameter, lnNT-proBNP, GFR, NYHA class, RV systolic area, LV-end-diastolic diameter, FAC, and LVEF, and RVLS.

These variables showed high accuracy in the prediction of the primary endpoint, with an overall AUC of 0.84 (95% CI: 0.75-0.93) (Figure 3B)

#### Clustering

Clustering with the K-medoid method allowed the identification of two phenotypically different groups of CRT candidates. Phenogroup 1 included 122 patients, with a mean age of 67 years, and a higher prevalence of women.

The baseline clinical and echocardiographic characteristics were imbalanced between the 2 groups.

Patients in phenogroup 1 had better kidney function, less dilated LV and a less restrictive filling pattern. LVEF was not different between phenogroups, but LV longitudinal function was better in Phenogroup 1. Also, patients in phenogroup 1 had less dilated RV and a significantly better RV function, with less impaired ventriculoarterial coupling. The prevalence of LV dyssynchrony was higher in phenogroup 1. Typical LBBB was more prevalent in phenogroup 1, but there was no significant difference in QRS width between the 2 groups. CRT volumetric response was highly prevalent in phenogroup 1 (Table 1).

Regarding the primary endpoint, phenogroup 2 had a significantly worse prognosis compared to phenogroup 1 (HR 4.70, 95% CI: 2.1-10.0, p<0.0001; log-rank p<0.0001) (Figure 4).

#### DISCUSSION

In this prospective, multicentric study, we have shown the feasibility and validity of the application of both supervised and unsupervised ML approaches to a comprehensive pattern of preimplantation clinical, biochemical, electrocardiographic and echocardiographic data obtained in everyday clinical practice in CRT-candidates. Our analysis allowed: 1) the identification of groups of features which are significantly associated with CRT response and prognosis; 2) the identification of two phenogroups of patients with specific characteristics and differential outcomes. We also demonstrated that 3) RV size and function parameters have pivotal importance in the prediction of CRT response and prognosis.

#### **Prediction of CRT-response**

Current recommendations for CRT implantation rely upon the assessment of the symptomatic status of patients, LVEF and QRS width<sup>1</sup>. Nevertheless, 30-to-40% of patients who receive CRT according to guidelines are not-responders to treatment<sup>2</sup>. Kalsheur et al. have shown that the application of ML to CRT can provide a better classification of patients than the simple assessment of LBBB morphology and QRS duration<sup>11</sup>. Similarly, we were able to demonstrate that the application of the RF method to ECG data and the dynamic analysis of LV strain curves

significantly improves the prediction of CRT-response<sup>10</sup>. The relevance of imaging derived parameters in the selection of CRT candidates remains underestimated. In the current study, the application of supervised ML approaches was able to identify groups of features which are good predictors of CRT-response.

In our study, SF, ApR and IHD, were the most important variables associated with CRT response. These results are in line with an increasing amount of data underscoring the relevance of the visual assessment of LV mechanical discoordination and ischemic cardiomyopathy in determining in CRT response<sup>24,25</sup>.

Noteworthy, QRS width was not the most important feature associated with CRT volumetric response, and its predictive role was improved by several RV-derived parameters such as RV basal and median diameter, right ventricular systolic area, RVLS and RV arterial coupling estimated by the TAPSE/PAPs ratio. These results are particularly relevant because the role of the right heart in CRT response is still debated<sup>26,27,28,29, 30</sup>.

Physiologically, the RV and LV are interdependent, because of the sharing of the interventricular septum, the pericardial space, and myocardial fibres. Several experimental studies have shown that 20-40% of RV systolic pressure and flow results from LV contraction and 4-10% of left ventricular systolic pressure and stroke volume are due to right ventricular contraction. This systolic interaction is decreased by a stiff septum and increased by stiff ventricular free walls<sup>31</sup>. This relationship between the left and the right heart explains why RV dysfunction occurs in nearly 50% of patients with HF and reduced LVEF. On the other hand, RV failure might induce changes in LV geometry which cause an impairment in LV filling and decrease in cardiac output<sup>32</sup>. In the field of CRT, Storsen et al. have shown that LBBB causes an abnormal RV free wall motion which is reversed by CRT only in cases of preserved RV function. This pattern of activation might explain why RV failure is associated with a poor response to CRT<sup>33</sup>.

Some small retrospective studies<sup>27,29</sup> and a post-hoc analysis of the prospective Cardiac Resynchronization Therapy Modular Registry (CRT-MORE)<sup>30</sup> have shown that TAPSE is a

predictor of CRT response. Compared with these previous studies, we performed a comprehensive analysis of RV function and we found that the ability of TAPSE to predict CRT-response was improved by the combination of both LV and RV-derived features. Our comprehensive model was able to predict CRT-response with an AUC of 0.81, which is superior to the value of 0.63 found for TAPSE alone in the CRT-MORE trial<sup>30</sup>. Despite this evidence, a large meta-analysis of different studies was not able to show a significant impact of RV performance on LV remodelling after CRT<sup>26</sup>. These discrepancies might be attributable to the fact that most of the published data have used only one or two parameters to assess RV function. Our study has the advantage of providing a multi-parametric assessment of the RV performance, including RV size, FAC, RV lateral strain and the TAPSE/PAPs ratio, which allow a comprehensive description of RV morphology and performance. The incorporation of these parameters into a model which includes the estimation of LV dyssynchrony, QRS width and LV size, improves the results compared to that achieved by analysis of single parameters and results in the identification of the model that best fits the primary clinical question.

#### Prediction of prognosis

To identify prognostic predictors in patients undergoing CRT, we combined supervised ML, which allowed the identification of variables specifically associated with outcome, and unsupervised ML analysis, which allowed the stratification of our population in two distinct groups of subjects with different characteristics and outcomes.

We found that TAPSE is the most important feature associated with outcome, followed by LV size, NYHA functional class, kidney function, FAC, LVEF and RVLS. The combination of these variables through the application of the RF method enhanced the appraisal of the likelihood of favourable clinical evolution after CRT.

These results are strengthened by the clustering analysis. This agnostic approach allowed the classification of our population in two distinct groups of patients who represent different archetypes

of CRT-candidates. On one side, we have subjects with more advanced heart failure, as witnessed by a pronounced biventricular dilatation, a higher prevalence of RV dysfunction and kidney failure. On the other side, we have patients with less severe disease, higher prevalence of SF and ApR and a better prognosis.

Our findings have several implications: 1) the presence of advanced heart failure features before CRT-implantation can significantly impact prognosis, independently from the result of CRT; 2) the assessment of RV function has pivotal importance in the risk stratification of patients. Previous studies have shown that RV dysfunction assessed by TAPSE is a prognostic determinant in HF patients, regardless of the degree of RV dysfunction<sup>16</sup>. In a post-hoc analysis of the CARE-HF trial, Damy et al. have also demonstrated that baseline TAPSE is a predictor of outcome in patients undergoing CRT, together with baseline NYHA and NTproBNP<sup>28</sup>. Our results reinforce these data and put into perspective the importance of a comprehensive evaluation of HF patients receiving CRT, which should systematically include the evaluation of RV size and function. We need to underscore that the aim of the application of machine learning methods in the clinical field is not to identify the importance of a single parameter -such as TAPSE –, but to handle large datasets with complex interactions between variables to find the algorithm that best suits the clinical problem<sup>34</sup>. Another important finding of our study is that despite the prevalence of CRT-response being higher among patients with a better outcome, volumetric response to CRT was not the main feature associated with prognosis.

This means that some patients who are considered CRT-responders can still experience a poor outcome, probably because of the presence of concomitant factors such as advanced HF, cardiorenal syndrome and RV dysfunction<sup>28</sup>. Whether the severity of HF and/or concomitant RV dysfunction before device implantation should influence the selection of CRT-candidates requires further study and underscores the subtle relationship between outcome and response to treatment<sup>35</sup>. On one hand, it is evident that the dichotomization of CRT-candidates in responders and nonresponders according to volumetric remodelling is simplistic because there is a wide spectrum of response to CRT, which goes from mild LV remodelling to complete functional recovery, and this progressive remodelling can take more than 6 months to settle. On the other hand, some patients might have such an advanced cardiopathy (eg severely dilated heart, biventricular dysfunction and/or cardiorenal syndrome), that the success of CRT might not be sufficient to change the natural history of heart failure disease progression and the outcome<sup>36</sup>. These observations suggest that finding the good therapeutic window for CRT implantation has pivotal importance. By incorporating a large amount of data, ML might be useful for the characterization and identification of patients with different "risk profiles", favouring the generation of pathophysiological hypothesis, and facilitating the tailored management of patients, which represent a significant step forwards personalized medicine.

#### Limitations

This study has several limitations. First, data are obtained from a medium-size population of patients undergoing CRT according to current recommendations. The availability of more data, including electrophysiological data obtained at the moment of CRT implantation and the analysis of a wider population might have strengthened and improved our results. Second, compared to previous studies<sup>2,5,11,28,30</sup>, our population displayed a lower prevalence of ischemic heart disease and a higher prevalence of typical LBBB. This means that our results should be applied with caution to patients with ischemic cardiomyopathy and or non-LBBB QRS morphology, who are known to present a lower rate of CRT-response. Third, we did not perform an external validation of our results, which limit their applicability and should represent an object of further research. Fourth, patients were followed-up for 37 months and a longer follow-up might have been useful to strengthen the value of our model for the prediction of events. Moreover, data on heart failure hospitalizations during follow-up are not available. This is an important adverse event that requires assessment in CRT recipients.

#### **Conclusions**

The application of machine learning methods to a set of common clinical,

electrocardiographic and echocardiographic data emphasize the importance of a multiparametric approach for both the identification of CRT-response and the prediction of prognosis after CRT. Our results underscore the importance of RV function on both CRT response and outcome and the pivotal role of the global assessment of heart function in patients undergoing CRT. Despite these interesting results, additional studies on a broader population will be necessary to fully validate and understand the clinical application of machine learning to CRT. Although there remains much to understand, as demonstrated in our study, it is likely that application of ML-derived algorithms will allow the stratification of CRT-candidates to guide patients towards specific therapeutic approaches and additional focused clinical trials.

#### Acknowledgement

JM Aalen was supported by a grant from the Norwegian Health Association.

CK Larsen was recipient of a clinical research fellowship from the South-Eastern Norway Regional Health Authority.

J Duchenne was supported by a research grant of the University Leuven (OT12/084).

JU Voigt holds a research mandate of the Research Foundation Flanders (FKM1832917N). VLR was supported by the French National Research Agency (ANR) (ANR-16-CE19-0008-01) (project MAESTRo).

#### References

- Brignole M, Auricchio A, Baron-Esquivias G, Bordachar P, Boriani G, et al. 2013 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy: The Task Force on cardiac pacing and resynchronization therapy of the European Society of Cardiology (ESC). Developed in collaboration with the European Heart Rhythm Association (EHRA). Europace. 2013 Aug 1;15(8):1070–118. Doi: 10.1093/europace/eut206.
- Chung ES, Leon AR, Tavazzi L, Sun J-P, Nihoyannopoulos P, Merlino J, et al. Results of the Predictors of Response to CRT (PROSPECT) Trial. Circulation. 2008 May 20;117(20):2608– 16. Doi: 10.1161/CIRCULATIONAHA.107.743120.
- Yang S, Liu Z, Hu Y, Jing R, Gu M, Niu H, et al. A novel risk model for mortality and hospitalization following cardiac resynchronization therapy in patients with non-ischemic cardiomyopathy: the alpha-score. BMC Cardiovasc Disord. 2020 Apr 28;20(1):205. Doi: 10.1186/s12872-020-01460-x.

 Patel D, Trulock KM, Moennich LA, Kiehl EL, Kumar A, Toro S, et al. Predictors of longterm outcomes greater than 10 years after cardiac resynchronization therapy implantation. J Cardiovasc Electrophysiol. 2020 May;31(5):1182-1186. doi: 10.1111/jce.14425. Epub 2020 Mar 15.

 Zareba W, Klein H, Cygankiewicz I, Hall WJ, McNitt S, Brown M, et al. Effectiveness of Cardiac Resynchronization Therapy by QRS Morphology in the Multicenter Automatic Defibrillator Implantation Trial-Cardiac Resynchronization Therapy (MADIT-CRT). Circulation. 2011 Mar 15;123(10):1061–72. Doi:10.1161/CIRCULATIONAHA.110.960898.

- Auricchio A, Prinzen FW. Enhancing Response in the Cardiac Resynchronization Therapy Patient: The 3B Perspective-Bench, Bits, and Bedside. JACC Clin Electrophysiol. 2017 Nov;3(11):1203–19. Doi: 10.1161/CIRCULATIONAHA.110.960898.
- Lumens J, Tayal B, Walmsley J, Delgado-Montero A, Huntjens PR, Schwartzman D, et al. Differentiating Electromechanical From Non-Electrical Substrates of Mechanical Discoordination to Identify Responders to Cardiac Resynchronization Therapy. Circ Cardiovasc Imaging. 2015 Sep;8(9):e003744. Doi: 10.1161/CIRCIMAGING.115.003744.
- Shah SH, Arnett D, Houser SR, Ginsburg GS, MacRae C, Mital S, et al. Opportunities for the Cardiovascular Community in the Precision Medicine Initiative. Circulation. 2016 Jan 12;133(2):226–31. Doi: 10.1161/CIRCULATIONAHA.115.019475.
- Cikes M, Sanchez-Martinez S, Claggett B, Duchateau N, Piella G, Butakoff C, et al. Machine learning-based phenogrouping in heart failure to identify responders to cardiac resynchronization therapy. Eur J Heart Fail. 2019;21(1):74–85. Doi: 10.1002/ejhf.1333.
- Donal E, Hubert A, Le Rolle V, Leclercq C, Martins R, Mabo P, et al. New Multiparametric Analysis of Cardiac Dyssynchrony: Machine Learning and Prediction of Response to CRT. JACC Cardiovasc Imaging. 2019 Sep;12(9):1887–8. Doi: 10.1016/j.jcmg.2019.03.009.
- Kalscheur MM, Kipp RT, Tattersall MC, Mei C, Buhr KA, DeMets DL, et al. Machine Learning Algorithm Predicts Cardiac Resynchronization Therapy Outcomes: Lessons From the COMPANION Trial. Circ Arrhythm Electrophysiol. 2018;11(1):e005499. Doi: 10.1161/CIRCEP.117.005499.
- 12. Felker GM, Shaw LK, O'Connor CM. A standardized definition of ischemic cardiomyopathy for use in clinical research. J Am Coll Cardiol. 2002 Jan 16;39(2):210–8.

- Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, et al. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. J Am Soc Echocardiogr Off Publ Am Soc Echocardiogr. 2015 Jan;28(1):1-39.e14. Doi: 10.1016/j.echo.2014.10.003.
- 14. Badano LP, Kolias TJ, Muraru D, Abraham TP, Aurigemma G, Edvardsen T, et al. Standardization of left atrial, right ventricular, and right atrial deformation imaging using twodimensional speckle tracking echocardiography: a consensus document of the EACVI/ASE/Industry Task Force to standardize deformation imaging. Eur Heart J -Cardiovasc Imaging. 2018 Jun 1;19(6):591–600. Doi: 10.1093/ehjci/jey042.
- 15. Rudski LG, Lai WW, Afilalo J, Hua L, Handschumacher MD, Chandrasekaran K, et al. Guidelines for the echocardiographic assessment of the right heart in adults: a report from the American Society of Echocardiography endorsed by the European Association of Echocardiography, a registered branch of the European Society of Cardiology, and the Canadian Society of Echocardiography. J Am Soc Echocardiogr Off Publ Am Soc Echocardiogr. 2010 Jul;23(7):685–713; quiz 786–8. Doi: 10.1016/j.echo.2010.05.010.
- 16. Ghio S, Guazzi M, Scardovi AB, Klersy C, Clemenza F, Carluccio E, et al. Different correlates but similar prognostic implications for right ventricular dysfunction in heart failure patients with reduced or preserved ejection fraction: RV function in heart failure. Eur J Heart Fail. 2017 Jul;19(7):873–9. Doi: 10.102/ejhf.664.
- Parsai C, Bijnens B, Sutherland GR, Baltabaeva A, Claus P, Marciniak M, et al. Toward understanding response to cardiac resynchronization therapy: left ventricular dyssynchrony is only one of multiple mechanisms. Eur Heart J. 2009 Apr;30(8):940–9. Doi: 10.1093/eurheartj/ehn481.

- Voigt J-U, Schneider T-M, Korder S, Szulik M, Gürel E, Daniel WG, et al. Apical transverse motion as surrogate parameter to determine regional left ventricular function inhomogeneities: a new, integrative approach to left ventricular asynchrony assessment. Eur Heart J. 2009 Apr;30(8):959–68. Doi: 10.1093/eurheartj/ehp062.
- Strauss DG, Selvester RH, Wagner GS. Defining left bundle branch block in the era of cardiac resynchronization therapy. Am J Cardiol. 2011 Mar 15;107(6):927–34. Doi: 10.1016/j.amjcard.2010.11.010.
- Buuren S van, Groothuis-Oudshoorn K. mice : Multivariate Imputation by Chained Equations in *R*. J Stat Softw 2011;45(3). Doi: 10.18637/jss.v045.i03.
- Charrad M, Ghazzali N, Boiteau V, Niknafs A. NbClust : An *R* Package for Determining the Relevant Number of Clusters in a Data Set. J Stat Softw 2014;61(6). Doi: 10.18637/jss.v061.i06.
- Kursa MB, Jankowski A, Rudnicki WR. Boruta A System for Feature Selection. Fundam Informaticae. 2010;101(4):271–85. Doi: 10.3233/FI-2010-288.
- 23. Breiman L. Random forests. Machine learning 2001; 45 (1) 5–32.
- Stankovic I, Prinz C, Ciarka A, Daraban AM, Kotrc M, Aarones M, et al. Relationship of visually assessed apical rocking and septal flash to response and long-term survival following cardiac resynchronization therapy (PREDICT-CRT). Eur Heart J Cardiovasc Imaging. 2016 Mar;17(3):262–9. Doi: 10.1093/ehjci/jev288.
- 25. Delgado V, van Bommel RJ, Bertini M, Borleffs CJW, Marsan NA, Ng ACT, et al. Relative Merits of Left Ventricular Dyssynchrony, Left Ventricular Lead Position, and Myocardial Scar to Predict Long-Term Survival of Ischemic Heart Failure Patients Undergoing Cardiac

Resynchronization Therapy. Circulation. 2011 Jan 4;123(1):70–8. Doi: 10.1161/CIRCULATIONAHA.110.945345.

- 26. Sharma A, Bax JJ, Vallakati A, Goel S, Lavie CJ, Kassotis J, et al. Meta-Analysis of the Relation of Baseline Right Ventricular Function to Response to Cardiac Resynchronization Therapy. Am J Cardiol. 2016 Apr;117(8):1315–21. Doi: 10.1016/j.amjcard.2016.01.029.
- Scuteri L, Rordorf R, Marsan NA, Landolina M, Magrini G, Klersy C, et al. Relevance of echocardiographic evaluation of right ventricular function in patients undergoing cardiac resynchronization therapy. Pacing Clin Electrophysiol PACE. 2009 Aug;32(8):1040–9. Doi: 10.1111/j.1540-8159.2009.02436.x
- Damy T, Ghio S, Rigby AS, Hittinger L, Jacobs S, Leyva F, et al. Interplay between right ventricular function and cardiac resynchronization therapy: an analysis of the CARE-HF trial (Cardiac Resynchronization-Heart Failure). J Am Coll Cardiol. 2013 May 28;61(21):2153–60. Doi: 10.1016/j.jacc.2013.02.049.
- Cappelli F, Cristina Porciani M, Ricceri I, Perrotta L, Ricciardi G, Pieragnoli P, et al. Tricuspid annular plane systolic excursion evaluation improves selection of cardiac resynchronization therapy patients. Clin Cardiol. 2010 Sep;33(9):578–82. Doi: 10.1002/clc.20806.
- Rapacciuolo A, Maffè S, Palmisano P, Ferraro A, Cecchetto A, D'Onofrio A, et al. Prognostic Role of Right Ventricular Function in Patients With Heart Failure Undergoing Cardiac Resynchronization Therapy: Prognostic role of RV function in CRT patients. Clin Cardiol. 2016 Nov;39(11):640–5. Doi: 10.1002/clc.22574.
- Naeije R, Badagliacca R. The overloaded right heart and ventricular interdependence. Cardiovasc Res. 2017 Oct 1;113(12):1474–85. Doi: 10.1093/cvr/cvx160.

- Konstam MA, Kiernan MS, Bernstein D, Bozkurt B, Jacob M, Kapur NK, et al. Evaluation and Management of Right-Sided Heart Failure: A Scientific Statement From the American Heart Association. Circulation 2018 May 15;137(20):e578-e622. Doi: 10.1161/CIR.00000000000560.
- Storsten P, Aalen JM, Boe E, Remme EW, Gjesdal O, Larsen CK, et al. Mechanical Effects on Right Ventricular Function From Left Bundle Branch Block and Cardiac Resynchronization Therapy. JACC Cardiovasc Imaging 2020 Jul;13(7):1475-1484. Doi: 10.1016/j.jcmg.2019.11.016.
- Shah SJ. Precision Medicine for Heart Failure with Preserved Ejection Fraction: An Overview.
   J Cardiovasc Transl Res. 2017 Jun;10(3):233–44. Doi: 10.1007/s12265-017-9756-y.
- Cleland JGF, Tavazzi L, Daubert J-C, Tageldien A, Freemantle N. Cardiac Resynchronization Therapy. J Am Coll Cardiol. 2009 Feb;53(7):608–11. Doi: 10.1016/j.jacc.2008.10.040.
- 36. Cleland JGF, Ghio S. The determinants of clinical outcome and clinical response to CRT are not the same. Heart Fail Rev. 2012 Nov;17(6):755–66. Doi: 10.1007/s10741-011-9268-9.

#### Table 1. Clinical and echocardiographic characteristics of the overall population and

according to phenogroups.

	Overall	Phenogroup 1	Phenogroup 2	
	(n= 193)	(n= 122)	(n=71)	p-value
Age, years	67±11	67 ±11	67±11	0.97
Female sex, n(%)	57(29)	46(37)	11(15)	0.0019
Ischemic heart disease, n(%)	64(33)	23(19)	41(58)	< 0.0001
NYHA class, n(%)				
Ι	14(7)	7(6)	7(10)	
II	107(56)	80(66)	27(38)	
III	70(36)	35(28)	35(49)	
IV	2(1)	0	2(2)	
InNT-proBNP, pg/ml	6.13±1.61	6.07±1.62	6.23±1.62	0.51
eGFR, ml/min	63±23	<u>66±23</u>	57±23	0.01
QRS width, msec	167±21	166±19	169±25	0.80
LBBB, n(%)	169(88)	113(93)	56(79)	0.01
LV-EDD, mm	63±12	62±10	66±13	0.007
LV-ESD, mm	52±13	51±13	55±13	0.024
LV-EDV, mm	211±86	202± 84	226± 88	0.04
LV-ESV, mm	153±73	146±68	165±79	0.084
LVEF, %	29±8	29±7	29±9	0.67
LV-GLS, %	-8.47±3.84	$-9.12 \pm 3.70$	-7.36±3.84	0.002
LAVI, ml/m <sup>2</sup>	47±17	45±16	50±17	0.03
E/Ea	16±9	15±7	19±12	0.02
RVBD, mm	41±8	38±7	45±8	< 0.0001
RVMD, mm	31±8	29±7	35±7	< 0.0001
RV lenght, mm	81±12	78±12	86±12	< 0.0001
Tricuspid annulus, mm	36±7	34±7	39±7	< 0.0001
RAV, ml	63±34	52±29	80±36	< 0.0001
RVDA, cm <sup>2</sup>	22±7	20±5	26±7	< 0.0001
RVSA, cm <sup>2</sup>	13±6	10±4	16±7	< 0.0001
FAC, %	44±12	48±9	38±13	< 0.0001
TAPSE, mm	19±5	10±5	18±6	< 0.0001
RVLS, %	-20.04±6.12	$-20.85 \pm 5.03$	$-18.66 \pm 7.47$	0.030
TRVmax, m/sec	$2.41 \pm 0.63$	$2.30 \pm 0.63$	$2.58 \pm 0.60$	< 0.0001
RAP, mmHg	6.83±3.71	$5.86 \pm 3.24$	8.49±3.91	< 0.0001
PAPs, mmHg	31.60±12.88	$29.00 \pm 11.38$	36.07±14.10	0.0002
TAPSE/PAPs, mm/mmHg	$0.80 \pm 0.86$	$0.92{\pm}1.02$	$0.60 \pm 0.39$	0.0002
Septal Flash, n(%)	130(67)	117 (96)	13 (18)	< 0.0001
Apical Rocking, n(%)	123 (64)	112 (92)	11 (15)	< 0.0001
CRT responder, n(%)	132 (68)	108 (89)	24 (34)	< 0.0001

BNP, brain natriuretic peptide; CRT, cardiac resynchronization therapy; EDD, end-diastolic

diameter; EDV, end-diastolic volume; ESD, end-systolic diameter; ESV, end-systolic volume;

FAC, fractional area change; GFR, glomerular filtration rate; LBBB, left bundle branch block; LV left ventricle; LVEF, left ventricular ejection fraction; NYHA, New York heart association functional class; PAPs, estimated pulmonary artery systolic pressure; RAP, right atrial pressure; RAV, right atrial volume; RV, right ventricle; RVBD, right ventricular basal diameter; RVDA, right ventricular diastolic area; RVLS, RV longitudinal strain; RVMD, right ventricular median diameter; RVSA, right ventricular systolic area; TAPSE, tricuspid annular plane systolic excursion; TRVmax, maximal tricuspid regurgitation velocity

#### Table 1S. Missing data

Age, years	0%
Female sex	0%
<b>IHD, n(%)</b>	0%
NYHA class	0.51 %
lnNT-proBNP, pg/ml	13.98 %
eGFR, ml/min	1.55 %
QRSd, ms	0%
LBBB, n(%)	0.51%
LV-EDD, mm	2.07 %
LV-ESD, mm	2.07 %
LV-EDV, mm	0%
LV-ESV, mm	0%
LV-EF, %	0%
LV-GLS, %	1.03%
LAVI, ml/m <sup>2</sup>	0%
E/Ea	6.21%
RVBD, mm	3.10 %
RVMD, mm	5.18 %
RV lenght, mm	4.66 %
Tricuspid annulus, mm	3.62 %
RAV, ml	2.59 %
RVDA, cm <sup>2</sup>	4.66 %
RVSA, cm <sup>2</sup>	5.18 %
FAC, %	5.18 %
TAPSE, mm	2.59 %
RVLS, %	16.06 %
TRVmax, m/sec	21.24 %
RAP, mmHg	22.27 %
PAPs, mmHg	18.65 %
TAPSE/PAPs, mm/mmHg	20.20 %
Septal Flash, n(%)	0 %
Apical Rocking, n(%)	0 %
CRT responder, n(%)	0 %

#### Figure 1. Framework illustrating the machine learning applied methods

Input data consists of clinical parameters, cardiac biomarkers, electrocardiographic data and echocardiographic variables (left panel).

Unsupervised machine learning (right, upper panel): the k-Medoid clustering method was applied to predefined features to select cluster of similar patients.

Supervised machine learning (right, lower panel): a feature selection phase was firstly applied by using the Boruta algorithm to select the best predictors of CRT-response or outcome. The random forest method is then applied to selected features to ensemble classifiers to predict CRT-response or outcome.

BNP, brain natriuretic peptide; ECG, electrocardiogram; GFR, glomerular filtration rate; IHD, ischemic heart disease; LBBB, left bundle branch block; LA, left atrium; LV, left ventricle; NYHA, New York Heart Association functional class; RV, right ventricle, TRV<sub>max</sub>, maximal tricuspid regurgitation velocity.

### Figure 2. Feature selection for the response to CRT (A) and outcome (B)

The Boruta algorithm was applied to select the best predictors of CRT-response and outcome.

# Figure 3. Receiver Operating Characteristics curve analysis for the prediction of CRT response (A) and outcome (B)

The random forest method was on the most important feature to build ensemble classifiers to predict CRT-response and outcome. Receiver Operating Characteristic (ROC) curves and the corresponding area under the curve (AUC) were estimated to evaluate the classification performance of the RF method proposed method.

#### Figure 4. Kaplan-Meier survival curve showing survival stratified by phenogroups

The final number of cluster was determined to maximize the silhouette index. Two clusters were identified (left panel). The survival stratification of these two phenogroups is depicted in the right panel.

#### Central illustration.

Application of machine learning methods to CRT-candidates allows the identification of phenogroup of patients with significant differently prognosis (upper panel) and the identification of specific features which are predictors of CRT-response and prognosis (lower panel). Parameters derived from the assessment of right ventricular size and function have pivotal importance in the risk stratification of CRT-candidates and should be systematically assessed before CRT-implantation.

CRT, cardiac resynchronization therapy; RV, right ventricle





# Variable importance





# Variable importance





False Positive Percentage

В

Fig 3

А









Supervised Machine Learning

Selection of features Boruta algorithm

Predictors of CRTresponse 16 main features: 11 main features: 50% from RV size/function

Predictors of Prognosis

36% from RV size/function

TAPSE is the most important feature

# **Unsupervised Machine Learning**



# Patients' phenotyping



Predictors of CRT-Predictors of Random Forest prognosis response Features' classification 9.0 AUC 0.81 AUC 0.84 0.4 0.4 0.0 0.2 0.2 0.8 1.0 0.4 0.4



False Positive Percentage