



HAL
open science

***Vibrio aestuarianus* subsp. *cardii* subsp. nov.,
pathogenic to the edible cockles *Cerastoderma edule* in
France, and establishment of *Vibrio aestuarianus* subsp.
aestuarianus subsp. nov. and *Vibrio aestuarianus* subsp.
francensis subsp. nov.**

Céline Garcia, Aurélie Mesnil, Delphine Tourbiez, Mirna Moussa, Christine Dubreuil, Amélie Gonçalves de Sa, Bruno Chollet, Yoann Godfrin, Lionel Degremont, Delphine Serpin, et al.

► **To cite this version:**

Céline Garcia, Aurélie Mesnil, Delphine Tourbiez, Mirna Moussa, Christine Dubreuil, et al.. *Vibrio aestuarianus* subsp. *cardii* subsp. nov., pathogenic to the edible cockles *Cerastoderma edule* in France, and establishment of *Vibrio aestuarianus* subsp. *aestuarianus* subsp. nov. and *Vibrio aestuarianus* subsp. *francensis* subsp. nov.. *International Journal of Systematic and Evolutionary Microbiology*, 2021, 71 (2), pp.004654. 10.1099/ijsem.0.004654 . hal-03128462

HAL Id: hal-03128462

<https://hal.science/hal-03128462>

Submitted on 2 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Description of *Vibrio aestuarianus* subsp. *cardii* subsp. nov. pathogenic to the edible cockles, *Cerastoderma edule* in France and establishment of *Vibrio aestuarianus* subsp. *aestuarianus* subsp. nov. and *Vibrio aestuarianus* subsp. *francensis* subsp. nov.

1.1 Author names

Céline Garcia^{1*#}, Aurélie Mesnil^{1,2#}, Delphine Tourbiez¹, Mirna Moussa¹, Christine Dubreuil¹, Amélie Gonçalves de Sa¹, Bruno Chollet¹, Yoann Godfrin¹, Lionel Degrémont¹, Delphine Serpin¹, Marie-Agnès Travers^{1,2}

1.2 Affiliation

¹ Ifremer, SG2M-LGPMM, Laboratoire de Génétique et Pathologie des Mollusques Marins Avenue de Mus de Loup, 17390 La Tremblade, France

² IHPE, Univ. Montpellier, CNRS, Ifremer, Univ. Perpignan Via Domitia, 34090 Montpellier, France

Participated equally to this work

1.3 Corresponding author

*Corresponding author: Céline Garcia, celine.garcia@ifremer.fr

Tel +33 5 46 76 26 45, Fax +33 5 46 76 26 11

1.4 Keywords

Shellfish; Bivalvia; mortality; *Vibrio aestuarianus*; cockle; bacterial pathogen

1.5 Repositories

The GenBank/EMBL/DDBJ accession numbers for the whole-genome sequences of *Vibrio aestuarianus* subsp. *cardii* subsp. nov. 12/122 3T3 (=LMG 31436=DSM 109723), 15/061 4T2 (=LMG 31438=DSM 109725), 15/064 3T2 (=LMG31439=DSM 109719) and 15/075 3T2 (=LMG 31440=DSM 109720) are JAAKZK000000000, JAAKZL000000000, JAAKZM000000000 and JAAKZN000000000, respectively, under BioProject accession number PRJNA607444. Accession numbers for the 16S, *gyrB* and *ldh* sequences of the same strains are MK307684, MK307691, MK307694, MK307697; MK315009, MK315016, MK315019, MK315022; and MK315026, MK315033, MK315036, MK315039, respectively. Lastly, for these three genes in strains LMG 31437 (=DSM 109724), 12/122 1T3, 12/122 2T3, 12/122 4T3, 12/122 5T3, 15/061 1T1, 15/061 2T2, 15/061 3T1, 15/061 3T2, 15/061 4T4, 15/061 5T2, 15/064 4T2, 15/075 1T2 of *Vibrio aestuarianus* subsp. *cardii* subsp. nov., the accession numbers are MK307681-307683, MK307685-307690, MK307692, MK307693, MK307695, MK307696; MK315006-315008, MK315010-315015, MK315017, MK315018, MK315020, MK315021; and MK315023-315025, MK315027-315032, MK315034, MK315035, MK315037, MK315038, respectively.

For *Vibrio aestuarianus* subsp. *francensis* subsp. nov., accession numbers for strains 01/308, 01/151, 01/032, 02/041 and 07/115 are JABANK000000000, JABANJ000000000, JABANI000000000, JAAZTV000000000 and JABANH000000000, respectively, under BioProject accession number PRJNA623953.

Finally, for *Vibrio aestuarianus* subsp. *aestuarianus* subsp. nov. LMG 7909T, the whole genome accession number is JAAZTU000000000.

ABSTRACT

Cockle mortality events have been reported in northern France since 2012. In the present study, we describe and investigate the implication of a potential bacterial causative agent in cockle mortality. Bacteria isolated from five different cockle mortality events were characterized and studied.

Using phenotypic analysis combined with DNA-DNA hybridization and whole genome sequencing, the isolates were shown to belong to *Vibrio aestuarianus*, a species regularly detected in France during oyster mortality events. Comparison of the strains from cockles with strains from French oysters and the type strain showed that the strains from cockles were genetically different to those from oysters and also different to the *V. aestuarianus* type strain. Moreover, the cockle and oyster strains were classified into two different, but close, groups both separated from the type strain by: (1) analyses of the *ldh* gene sequences; (2) DDH assays between 12/122 3T3 (LMG 31436=DSM 109723), a representative cockle strain, 02/041 (CIP 109791=LMG 24517) representative oyster strain and *V. aestuarianus* type strain LMG 7909T; (3) ANI values calculated on the genomes; and (4) phenotypic traits. Finally, results of MALDI-TOF analyses also revealed specific peaks discriminating the 3 representative strains. The toxicity of representative strains of these cockle isolates was demonstrated by experimental infection of hatchery-produced cockles. The data therefore allow us to propose two novel subspecies of *Vibrio aestuarianus*: *Vibrio aestuarianus* subsp. *cardii* subsp. nov. for cockle strains and *Vibrio aestuarianus* subsp. *francensis* subsp. nov. for the Pacific oyster strains, in addition to an emended description of the species *Vibrio aestuarianus*.

INTRODUCTION

The edible cockle *Cerastoderma edule* is one of the most common bivalve species widely distributed along the European coasts, and is commercially important in the UK, Netherlands, Spain

and France. For many years, mortalities of cockles and dramatic declines in their populations have been reported [1]. Predation, diseases, climatic events, pollution, poor recruitment and over-fishing have all been suggested as driving factors [2]. Cockles are host to a wide variety of viruses, bacteria, fungi and protists. Many of these microorganisms are normally innocuous but can become problematic if prevailing environmental conditions are poor. Some parasites have been isolated during mass mortality events [For review, 1]: Protistan *Marteilia cochillia* in Spain [3], gregarines *Nematopsis scheideri* and *Nematopsis incognito* in Portugal [4, 5], *Amoeba* [6], *Haplosporidia* [7], and a wide range of *Digenea* (trematodes) [8, 9]. Cockles also host emerging parasites, such as *Perkinsus chesapeaki*, newly described in Europe [10]. They can also suffer from a transmissible disease known as disseminated neoplasia [11, 12]. Some studies have suggested that different factors, such as genetic alterations, viruses, retrotransposons and contaminants [12], can act in concert [13]. However, bacteria have rarely been isolated associated with mortalities, apart from mycoplasma-like organisms in Spain [14].

Vibrio are the most abundant cultivable bacteria found in the marine environment. Free, associated with plankton or with marine animals, bacteria belonging to this genus are classified in clades [15-18] clustering more than 100 different species [19]. Some of these species have virulent strains implicated in fish and shellfish diseases either alone, in bacterial consortia [20-22] or in concert with other pathogens [23]. For bivalves, the main pathotypes belong to the *Splendidus* clade (*V. tasmaniensis*, *V. crassostreae*, *V. splendidus*) or the species *V. tapetis*, *V. tubiashii*, *V. bivalvicida*, *V. europaeus* or *V. aestuarianus* [19, 24].

Vibrio aestuarianus was first isolated in the 1980s in the USA, from seawater and several shellfish including oysters, clams and crabs presenting no signs of disease [25]. This species was isolated for the first time in France in 2001, during mortality events of Pacific oyster, *Crassostrea gigas*. Garnier et al. [26] compared the French strains with the American one and found that the French strains had some different biochemical and pathogenic characters. These authors concluded

that the French strains could constitute a new subspecies within *V. aestuarianus* and proposed to name the American strain *V. aestuarianus* subsp. *aestuarianus* and the French strains *V. aestuarianus* subsp. *francensis*, although these two subspecies were never officially validated.

The present study describes the isolation and classification of bacteria from cockle mortality events that occurred in France in 2012 and 2015. Strains were described and sequenced that allowed us to reproduce and describe this new cockle disease. These strains were compared with the French 'oyster' strains [26] and with the type strain of *Vibrio aestuarianus* (LMG 7909T) [25], and led us to propose two novel subspecies of *Vibrio aestuarianus*.

Isolation and Ecology

In 2012, mass mortality events of *Cerastoderma edule* were reported in two wild beds on the English Channel Coast: Baie des Veys and Baie de Somme, the latter of which is the largest cockle bed in France. In 2015, another mortality event occurred in both these locations as well as in Binic, also on the English Channel coast (Figure S1). Fishers reported these mortality events to the competent authority and national network for surveillance and monitoring of mollusc health, REPAMO (REseau de PATHologie des MOllusques), in charge of recording mortality cases and analysing representative samples associated with them. In 2012 and 2015, 15 to 30 adult cockles were collected from the different wild beds. Each collected individual was examined for macroscopic characteristics and processed using either histological or bacteriological analysis in addition to molecular analyses (Table S1). The mortality rate in 2012 was estimated at 80–89% according to site. For each site, the mortality rate was lower in 2015 than in 2012 ranging from 33 to 49% (Table S1). In all cases, cockle mortality was uniform, massive and sudden, concerning mainly adults. No other species were affected by these mortality events except in Baie des Veys where a number of stocks of *Crassostrea gigas* spat and juveniles also suffered mortalities. The cockle mortality occurred during summer, and

lasted from 1 to 2 months, according to fishermen. No specific environmental events were reported during the summers of 2012 or 2015, except a high level of macroalgae at Binic. The sampled individuals were mainly moribund, lying on the bottom and characterized by very slow valve closing valves and limited intrapalleal fluid. No specific macroscopic signs such as pustules, conchyolin deposits or necrosis and degradation of the hinge could be observed.

For the detection and quantification of bacterial pathogens, pieces of gill, muscle and mantle of five moribund cockles per sample were individually crushed in 150 µl sterile seawater, diluted 100-fold and 10,000-fold and spread onto Marine agar (Conda, Spain) incubated for 48–72 h at 20°C. Colonies were enumerated and those bacterial colonies found in abundance were isolated and tested by duplex real time PCR for the detection of *V. aestuarianus* and *V. Splendidus*-related. [27]. Pure cultures of these bacterial strains were conserved frozen at -80°C in Zobell broth with 15% glycerol. These bacteriological analyses revealed that all analysed samples yielded numerous bacterial colonies. The species *V. aestuarianus* was detected in all samples after 48h at 20°C (Table S2). Bacteria of the *Splendidus* clade were also isolated in some samples but these bacteria were not the dominant strains in the Petri dishes. Moreover, *V. aestuarianus* DNA was detected by molecular analyses [28] in all samples, at a generally high detection frequency. DNA of OsHV-1 (Ostreid Herpesvirus type 1) known to contribute to mortality events in *Crassostrea gigas* [23, 29], was not detected in any of the samples (Table S2).

Fifteen to twenty individuals per sample were also processed for histological analyses. After 48 h in Davidson's fixative, tissues were maintained in 70% ethanol until histological preparation by dehydration and embedding in paraffin following standard procedures. Sections of 2–3 µm thickness were stained with haematoxylin and eosin. Histological analyses revealed the presence of other organisms at low infection intensity (a few parasites per individual), despite a high prevalence in some cases such as gregarine spores or rickettsia-like organisms (Table S2).

16S RNA phylogeny

A total of 59 bacterial strains were isolated during periods when the REPAMO surveillance network reported high levels of cockle mortalities. After isolation, the predominant bacteria appeared small (1-2 mm), translucent, regular and white to cream-colored on marine agar after 48 h at 20°C (Figure S2). For genotyping, DNA was extracted by boiling in distilled water [30] and the *ldh* (lactate dehydrogenase) gene was amplified and sequenced according to methods previously described [20], as were the 16S rRNA and *gyrB* genes [17] (Table S4). Sequences were aligned and phylogenetic trees built using Mega X [31]. The trees were drawn using the Neighbor-Joining method with the Kimura two-parameter model [32] and maximum parsimony method [33]. A total of 17 isolates (isolated in 2012 and 2015 from cockle mortality events) were identified as belonging to *Vibrio aestuarianus* by qPCR (Table S3) and gene sequencing (16S rDNA and *gyrB*, Table S4). They showed a high degree of similarity with 16S rDNA reference sequences of *Vibrio aestuarianus* (99% with the type strain LMG 7909T). Phylogenetic analyses based on the *ldh* gene sequence of these strains led to their affiliation to the species *V. aestuarianus*, in a group close but apart from the oyster strains and type strain. Moreover, oyster strains formed an individual branch clearly separated from the type strain of *V. aestuarianus* (Figure 1) [26].

This classification was confirmed by DNA-DNA hybridization (DDH) performed by the BCCM™/LMG laboratory at Gent University. After genomic DNA extraction according to a modified Wilson protocol [34], hybridizations were performed in microplates at 40°C [35] and DNA relatedness values (%) based on means of a minimum of six hybridizations. DDH of two strains isolated from diseased cockles in 2012 and 2015 and one strain isolated from diseased oysters were performed with *Vibrio aestuarianus*, *V. ordalii* and *V. salilacus* reference strains. With more than 70% hybridization to the *V. aestuarianus* type strain, we confirm the affiliation of the cockle and oyster strains to the species *V. aestuarianus*. Indeed, hybridizations were 70, 71 and 79% with the *V.*

aestuarianus type strain for strains 12/122 3T3, 15/061 4T2 and 02/041, respectively, and were 80 and 80% with the *V. aestuarianus* oyster strain for cockle strains 12/122 3T3 and 15/061 4T2, respectively (Table 2). Garnier et al. generally found similar results for hybridizations between the *V. aestuarianus* type strain and different oyster strains: their hybridizations were between 54 and 87% [26].

The 3 representative strains for the 3 subspecies were further studied by whole cell matrix-assisted laser desorption ionization-time of flight mass spectrometry profiling. MALDI-TOF mass spectrum measurements and processing were carried out as described by Mougín et al [36]. The analyses were performed using Flex Analysis (version 3.4) and ClinProTool (version 3.0) softwares (Bruker Daltonics). Figure 2 shows the comparison of the MALDI-TOF spectra. First, all 3 strains were identified with Bruker database as *Vibrio aestuarianus* with correct scores. Moreover, significant differences observed between spectra (12 per strains) with prominent mass ions at 3877.4 and 7321.7, characteristic for *V. aestuarianus francensis* subsp. nov type strain and *V. aestuarianus cardii* subsp. nov type strain, respectively. The dendrogram based on whole-cell MALDI-TOF mass spectra as well as principal component analysis demonstrated a clear separation of the three strains. Comparison in other isolates is now needed to confirm the peak combination allowing discrimination [37].

Genome Features

The existence of three distinct and coherent groups was also supported by results from the comparison of the complete genomes of four cockle isolates, genome of the *V. aestuarianus* type strain, and the five genomes already available for different *V. aestuarianus* oyster strains. Briefly, whole genomes of cockle isolates were sequenced by BIOFIDAL (Biofidal, France) after DNA isolation using NucleoSpin microbial DNA (Macherey-Nagel). DNA Libraries were made from 1 ng of genomic

DNA using a Nextera XT DNA kit (Illumina). Paired-end isolate sequencing (2x300) was performed on a MiSeq system (Illumina). Genomes were assembled *de novo* using Spades (3.11.0 version, [38]), with read error corrections and 20x coverage cut-off. Assembly quality was checked using QUAST [39]. Assemblies were added in MAGE [40], excluding contigs with lengths below 1000 bp. Oyster and cockle *Vibrio aestuarianus* strain draft genomes presented typical features for the species. They ranged between 4.02 and 4.51Mb, with a relatively close number of CDS, genes coding for transfer RNA (between 91 and 102 for cockle isolates and between 84 and 85 for oyster isolates), and G+C contents (42.68 to 42.89% for cockle isolates and 42.52 to 42.83% for oyster isolates) (Table 3). 16S sequences and whole genome comparisons were performed by calculating average nucleotide identity (ANI) between strains (JSpeciesWS, [41]). A distance matrix was calculated inverting ANI results based on whole genomes. Similarity percentage calculated between 16S sequences extracted from whole genomes ranged from 99.09 to 100% (Data not shown). All values were above the 98.7 % cut-off generally accepted for differentiating two species [42]. Moreover, ANI values calculated with whole genomes, indicated an intra-clade similarity of 99.21% to 99.99% (mean value of 99.69%) for *V. aestuarianus* oyster strains, and an intra-clade similarity ranging from 97.88% to 99.99% (mean value of 98.47%) for cockle isolates. The inter-clade similarity ranged from 96.83% and 97.8% depending on the isolates (Table S5) and confirmed that all strains belong to the same species (ANI>95%) but can be classified into three different subspecies.

Comparison in predicted CDS revealed a core genome, shared by all 3 strains, with 2858 CDS and specific genomes containing 611 to 797 specific CDS (figure S3). As oysters and cockles' isolates are mollusc pathogens, predicted virulome was compared in the 3 representative strains. Globally, the 3 strains contain homologous CDS encoding putative virulence effectors, virulence regulators, secretion systems (SST1, SST2, SST4 and SST6) and pili/flagella/chemotaxis-related proteins. But, some predicted virulence factors are missing in the different strains as haemolysin co-regulated protein in LMG7909T, zonular occludens toxin zot and T1SS-2 in 12/122-3T3 and a complete T6SS locus in 02-041. However, their implication in pathogenesis must definitively be demonstrated.

Physiology and Chemotaxonomy

The phenotypes of selected isolates were characterized in different ways. Bacterial growth was observed on Marine Agar (Conda), TCBS medium (AES Chemunex), and CHROMagar™ Vibrio medium (CHROMagar) (Figure S2). The capacity of bacteria to grow in Zobell broth with 0% to 10% NaCl at 20°C, or with 2% NaCl at 4°C, 10°, 37°C and 40°C was determined (Table 1).

Antibiograms were carried out using disc diffusion assays on Mueller–Hinton agar (Oxoid) with commercial impregnated discs (Oxoid): Flumequine 30 µg (UB30), Kanamycin 30 µg (K30), Tetracycline 30 µg (TE30), Sulfamethoxazole/trimethoprim 19:1 25 µg (SXT25), Streptomycin 25 µg (S25), Sulfonamides 300 µg (S3), Erythromycin 30 µg (E30), Penicillin 10 UI (6 µg) (P10), Trimethoprim 5 µg (W5), Chloramphenicol 30 µg (C30), Ampicillin 10 µg (AMP10), Gentamicin 30 µg (CN30) or the Vibriostatic agent, Pteridine 150 µg and 10 µg (O129/150 and O129/10). Inhibition was observed after 24–48 h at 22 °C.

API 20E, API 20NE and API 50CH (Biomérieux) were used in accordance with manufacturer's recommendations. Fresh colonies, obtained from culture bacteria on TSA (Difco™) adjusted to 2% NaCl, were suspended in SASW to obtain about $OD_{600} = 0.1$. After inoculation, the strips were incubated for 48 h at 20°C and results were determined according to the manufacturer's instructions. Finally, FAME (Fatty Acids) determination was performed by the BCCM™/LMG laboratory at Gent University using a standard commercial identification system (MIDI) and gas chromatography.

The 17 cockle strains and two oyster strains were able to grow in Zobell liquid medium at different temperatures from 4°C to 37°C, with a limited growth rate at the extremes. However, no growth was observed at 40°C, with or without 6% salt. Only the *Vibrio aestuarianus* type strain was able to grow in the presence of 6% salt. All the strains, like the type strain, were shown to be sensitive to the tested antibiotics (Table 1). Cockle and oyster strains appeared under blue light on Chromagar medium and, unlike the type strain, had a limited growth capacity on TCBS medium,

where a few yellow colonies appeared (Figure S2). The observations on oyster strains and the type strain were similar to those made by Garnier et al. [26]. Additional biochemical characterization analyses were performed to test this proposition, using API20E and 50CH strips (Table 4) and FAME analyses (Table 5). The strains isolated from cockles and oysters, like the *V. aestuarianus* type strain, could form acids from D-Glucose, Sucrose, D-Trehalose, N-acetyl-D-glucosamine and have Beta-galactosidase activity and indole production. Cockle strains could be distinguished from oyster strains by their utilization of D-mannitol and arginine and fermentation of starch, D-ribose and glycogen. They could be distinguished from the type strain by the non-fermentation of potassium gluconate. Oyster strains cannot use D- mannitol and arginine. However, as fermentation and utilization of substrates appeared variable among the cockle and oyster strains, no character clearly separated the three groups of strains.

Other characters that can also differentiate strains include their fatty acids (Table 5). *V. aestuarianus* strains can be distinguished from *V. ordalii* and *V. furnisii* by their lower level of 3 hydroxy-dodecanoic acid and higher level of iso-C16:0 fatty acid. Between the *V. aestuarianus* strains, differences were observed for the C12:0-3-OH, iso-C14:0 and C14:0, for which cockle strains appeared intermediate and oyster strains had a high level compared with the type strain. However, the most remarkable difference concerned the C18:1 w7c fatty acid, present at a higher level in the type strain compared with the oyster (low level) or cockle strains (intermediate level). In contrast, oyster and cockle strains showed a high level of the C16:0 fatty acid compared to the type strain.

Finally, the pathogenicity of nine cockle isolates (12/122 1T1, 12/122 2 T3, 12/122 3T3, 12/122 4T3, 15/061 1T1, 15/061 4T2, 15/064 3T2, 15/075 1T2, 15/075 3T2) on reared cockles was investigated by experimental injections, which also included oyster strain 02/041, pathogenic for oysters, and *Vibrio splendidus*-related 15/064 1T2, isolated from cockles (Figure 1). Cockles were induced to spawn in September 2015 at the Ifremer hatchery in La Tremblade and larvae were grown

according to standard protocols used for *C. gigas*, except that seawater temperature was maintained at 20°C for cockle larvae [43]. Settlement took place from 11 to 16 days post fertilization. All progenies were maintained under controlled environmental conditions in our facilities using unheated UV treated (40 mj/cm²) seawater in a flow-through system. All animals were fed *ad libitum*. Following an initial screening of 30 of these animals (no detection of *Vibrio aestuarianus* on 159 screened colonies, no detection of *V. aestuarianus* or herpesvirus OsHV-1 DNA and no detection of any other pathogens by histology) and in the absence of mortality over the rearing period, we chose to use this biological material for the controlled infections. Experiments were conducted in May and June 2016 when cockles weighed 4.39 g (s.d. 0.80) and measured 23.28 mm (s.d. 1.52).

Bacteria isolated from diseased cockles as well as controls (*V. splendidus*-related from cockles 15/064 1T2 and *V. aestuarianus* oyster strain 02/041) were grown at 20°C for 20 h in marine broth with constant shaking at 40 rpm (Rotator SB3; Stuart). After a SASW washing, bacterial concentrations were evaluated spectrophotometrically at an optical density (OD) of 600 nm, adjusted to 0.4 and 0.04 and checked by plating. Cockles were anesthetized for 2 to 4 hours in a magnesium chloride solution (MgCl₂, Fluka) at a final concentration of 50 g l⁻¹ (1/4: v/v seawater/freshwater) under aeration. Subsequently, 50 µl of bacterial suspension (10⁷ CFU to 10⁶ per cockle) was injected into the adductor muscle. A group of cockles were injected with SASW to serve as negative controls. After injection, the cockles were transferred to tanks (triplicates of 10 cockles in 2 litres) filled with UV-treated, 1-µm-filtered seawater and maintained under static conditions at 20°C with aeration. Mortality was monitored twice a day and any newly dead cockles were removed from the tanks over a 7-day period. Animals were considered to be dead when the valves did not close following stimulation.

After 24 h, cumulative mortalities ranged from 0 to 83% for cockles injected with *V. aestuarianus* cockle strains. Mortalities continued over 7 days, with a peak during the first two days. For all moribund cockles, the predominant bacterial isolates were obtained from muscle and

haemolymph (205 strains screened from 116 animals, all belonging to *V. aestuarianus* species using the *dnaJ*-qPCR diagnosis assay). We also noticed the high levels of bacteria belonging to the *Splendidus* clade in 31 individuals. No mortality was noted in the control or in cockles injected with *V. splendidus*-related 15/064 1T2 or *V. aestuarianus* oyster strain 02/041.

Garnier et al. [26] also tested the pathogenicity of 11 *Vibrio aestuarianus* oyster strains and *V. aestuarianus* type strain by injection of 10^8 bacteria per oyster. They observed mortality ranging between 2 and 55% depending on the oyster strain, whereas a low mortality (10%) was observed with the *V. aestuarianus* type strain. The same observation was made by Goudenège et al. [20], who classified oyster strains as either highly virulent for oysters (i.e. inducing > 50% mortalities at 10^2 CFU animal⁻¹) or non-virulent (i.e. inducing < 50% mortalities at 10^7 CFU animal⁻¹).

In summary, *Vibrio aestuarianus* isolates obtained from oysters or cockles present phenotypic and genetic traits distinct from the *V. aestuarianus* type strain and these differences support the separation of these oyster and cockle strains into two novel subspecies of *V. aestuarianus* distinct from *V. aestuarianus* LMG 7909T and other *V. aestuarianus* strains. Therefore, we propose that *Vibrio aestuarianus* should be divided into three subspecies, for which the names *Vibrio aestuarianus* subsp. *aestuarianus* subsp. nov., *Vibrio aestuarianus* subsp. *francensis* subsp. nov. and *Vibrio aestuarianus* subsp. *cardii* subsp. nov. are proposed, with strains LMG 7909T, 02/041 and 12/122 3T3 as the type strains, respectively.

Protologue

Emended description of *Vibrio aestuarianus*

The characteristics modified or added to those reported in original descriptions by Tison and Seidler (1983) [25] are as follows:

Vibrio aestuarianus is a gram-negative, oxidase-positive, motile bacterium. Colonial morphology is variable according to the subspecies. Growth occurs on Marine Agar from 4 to 30°C with an optimum between 20 and 25°C and strains do not grow at 40°C or in the absence of NaCl. Most strains utilize N-acetyl-D-glucosamine, Beta-galactosidase and produce indole. Strains ferment the sucrose, D-mannose, D-fructose, D-glucose, D-galactose and D-trehalose. Strains are susceptible to Ampicillin, Erythromycin, Chloramphenicol, Kanamycin and Tetracycline. The major fatty acids are C12:0 (2.2–4.4%), C12:0; 3-OH (1.3–3.4%), C14:0 (4–8.7%), iso-C16:0 (4.1–5.6%), C16:0 (13.4–20.06%) and C18:1 w7c (11.3–22.6%).

The genome size is 3.9–4.5 Mb (Genbank genome accession number JAAZTU000000000), and genomic DNA G+C content is 42.5–42.9%.

The type strain is ATCC 35048 (=DSM 19606, CAIM 592, CCUG 28583, CECT 625, CIP 102971, LMG 7909, NBRC 15629, NCMB 2236). The GenBank/EMBL/DDBJ accession number for the genome of the type strain is JAAZTU000000000.

Description of *Vibrio aestuarianus* subsp. *aestuarianus* subsp. nov.

Vibrio aestuarianus subsp. *aestuarianus* Tison and Seidler [25] (a.es.tu.ri.a'nus. N.L. masc. adj. *aestuarianus*, pertaining to an estuary). The description is essentially the same as that given above for the species *Vibrio aestuarianus*, with the following additions.

Colonies are opaque, 2–3 mm in diameter on marine agar (MA) after 24h at 20°C. It grows on TCBS, with 6% salt and at 37°C. It has gelatinase and arginine dihydrolase activity and utilizes starch, D-ribose, D-maltose, D-cellobiose, glycogen and potassium gluconate. Among the three subspecies, it presents the highest level of C18:1 w7c fatty acid and the lowest level of C12:0; 3-OH and iso-C14:0 fatty acids. The type strain does not induce mortality of Pacific oyster, *Crassostrea gigas*. The genome size is predicted around 4.2 Mb and genomic DNA G+C content is 42.6% (Genbank genome accession number JAAZTU000000000).

The type strain is ATCC 35048 (=DSM 19606, CAIM 592, CCUG 28583, CECT 625, CIP 102971, LMG 7909, NBRC 15629, NCMB 2236).

Description of *Vibrio aestuarianus* subsp. *francensis* subsp. nov.

Vibrio aestuarianus subsp. *francensis* as proposed by Garnier et al. [26] (fran.cen'sis. N.L. masc. adj. *francensis*: of, or belonging to, France)

It displays typical characteristics for the species *V. aestuarianus*. Colonies are white and slightly opaque on Marine Agar and are less than 1 mm in diameter after 24 h at 20°C. No strain can grow in the absence of NaCl or in the presence of 6% NaCl (except for strain 02/041) or on TCBS. Growth occurs from 4 to 30°C. Most strains have no arginine dihydrolase and cannot use D- mannitol. Some strains use citrate. It presents the highest level of C16:0 and C14:0 fatty acids and the lowest level of C18:1 w7c fatty acids compared to the two other subspecies.

The majority of strains, including the type strain 02/041, are pathogenic to the Pacific oyster, *Crassostrea gigas*, but the type strain is not pathogenic to edible cockles, *Cerastoderma edule*. The genome size is estimated around 4.18–4.53 Mb with the draft genomes, and genomic DNA G+C content is about 42.52–42.83 mol%.

The type strain is 02/041 (LMG 24517 = CIP 109791, Genbank genome accession number JAAZTV000000000) and was isolated from a diseased oyster maintained in the Argenton experimental hatchery (Ifremer, France).

Description of *Vibrio aestuarianus* subsp. *cardii* subsp. nov.

Vibrio aestuarianus subsp. *cardii* (car'di.i. N.L. fam. *cardii* referring to *Cardium* – latin: cockle)

It displays typical characteristics for the species *V. aestuarianus*. Colonies are around 1 mm in diameter after 24 h at 20°C. No strain can grow in the absence of NaCl and growth was poor on TCBS. Growth occurs from 4 to 30°C, difficult at 37°C and not at 40°C. Strains can use D- mannitol and do not ferment potassium gluconate. Concerning its fatty acid composition, it presents a high level of

C16:0 and an intermediate level of other fatty acids. Some strains, including the type strain 12/122 3T3, are pathogenic to the edible cockle *Cerastoderma edule* but the type strain is not pathogenic to Pacific oyster *Crassostrea gigas*.

The type strain is 12/122 3T3 (LMG 31436 = DSM 109723, Genbank genome accession number JAAKZK000000000) and was isolated from a diseased cockle sampled from a wild bed (Baie de Somme) in France.

AUTHOR STATEMENTS

1.6 Authors and contributors

DT, CD, AGS, BC, YG, DS, CG, AM and MAT performed the experiments. LD produced and conserved the animals. MAT and CG supervised the project. DT, CD, AM, CG and MAT analysed the data. MAT, AM and CG wrote the manuscript that was revised and approved by all the authors.

1.7 Conflicts of interest

The authors declare that there are no conflicts of interest.

1.8 Funding information

This work was supported by Ifremer; by DGAL (Direction Générale de l'Alimentation) through the REPAMO network and the National Reference Laboratory for Mollusc Diseases, Ifremer, La Tremblade; by DPMA (Direction des Pêches Maritimes et de l'Aquaculture) through MIRAGE project funding and by the European Commission through the European Union Reference Laboratory, Ifremer, La Tremblade. This work was also partly supported by the EU-funded project VIVALDI ('Preventing and Mitigating Farmed Bivalve Diseases') H2020 program, no. 678589.

1.9 Acknowledgements

We wish to thank the coastal correspondents of the REPAMO network (particularly those at Boulogne sur Mer, Port en Bessin and Dinard) for their contribution to collecting and providing

samples and related information. We also thank the hatchery and nursery teams of the Laboratory of Genetics and Pathology at Ifremer La Tremblade and Ifremer Bouin for their assistance with cockle production. We are also grateful to M. Thuillier and M. Le Piouffle from the laboratory LABOCEA, Mme Cauvin and Mme Benoît from the laboratory, LABEO-Manche and M. Treilles from the laboratory QUALYSE for their assistance in MALDI-TOF analyses.

ABBREVIATIONS

ANI: Average Nucleotide Identity

DDH: DNA-DNA hybridization

FAME: Fatty Acid Methyl Esters

MALDI-TOF MS : Matrix-Assisted Laser Desorption Ionization-Time Of Flight Mass Spectrometry

REPAMO: REseau de PATHologie des MOllusques

SASW: Sterilized Artificial SeaWater

TCBS: Thiosulfate Citrate Bile salts Sucrose agar

TSA: Tryptone Soya Agar

CDS: coding DNA sequence

REFERENCES

1. Longshaw M, Malham SK: **A review of the infectious agents, parasites, pathogens and commensals of European cockles (*Cerastoderma edule* and *C. glaucum*)**. *Journal of the Marine Biological Association of the United Kingdom* 2013, **93**(1):227-247. DOI: <https://doi.org/10.1017/S0025315412000537>.
2. Ducrotoy JP, Rybarczyk H, Souprayen J, Bachelet G, Beukema JJ, Desprez M, Dorjes J, Essink K, Guillou J, Michaelis H *et al*: **A comparison of the population dynamics of the cockle (*Cerastoderma edule*, L.) in north-western Europe**. . In: *Proceedings of the Estuarine and Coastal Sciences Association Symposium: 1989*; 1989: 173-184.
3. Carrasco N, Hine PM, Durfort M, Andree KB, Malchus N, Lacuesta B, González M, Roque A, Rodgers C, Furones MD: ***Marteilia cochillia* sp nov., a new *Marteilia* species affecting the**

- edible cockle *Cerastoderma edule* in European waters** *Aquaculture* 2013, **412–413**:223-230. DOI: <https://doi.org/10.1017/S003118201800063X>.
4. Azevedo C, Cachola R: **Fine structure of the apicomplexa oocyst of *Nematopsis* sp. of two marine bivalve mollusks** *Diseases Of Aquatic Organisms* 1992, **14**(1):69-73. DOI: 10.3354/dao014069.
 5. Ordas MC, Figueras A: **Histopathological study of abnormal mortalities of cockle (*Cerastoderma edule*) in Carril (Galicia, NW Spain)**. *Bulletin of the European Association of Fish Pathologists* 2005, **25**(3):124-130.
 6. Azevedo C: **Some observations on an amoeba, destructive agent of the gills of *Cerastoderma edule* (Mollusca, Bivalvia)**. *Diseases of Aquatic Organisms* 1997, **30**(3):223-225. DOI: 10.3354/dao014069.
 7. Azevedo C, Conchas RF, Montes J: **Description of *Haplosporidium edule* n. sp. (Phylum Haplosporidia), a parasite of *Cerastoderma edule* (Mollusca, Bivalvia) with complex spore ornamentation**. *European Journal of Protistology* 2003, **39**(2):161-167. DOI: <https://doi.org/10.1078/0932-4739-00905>.
 8. de Montaudouin X, Thieltges DW, Gam M, Krakau M, Pina S, Bazairi H, Dabouineau L, Russell-Pinto F, Jensen KT: **Digenean trematode species in the cockle *Cerastoderma edule*: identification key and distribution along the north-eastern Atlantic shoreline**. *Journal of the Marine Biological Association of the United Kingdom* 2009, **89**(3):543-556. DOI: 10.1017/s0025315409003130.
 9. Magalhães L, Correia S, de Montaudouin X, Freitas R: **Spatio-temporal variation of trematode parasites community in *Cerastoderma edule* cockles from Ria de Aveiro (Portugal)**. *Environmental Research* 2018, **164**:114-123. DOI: <https://doi.org/10.1016/j.envres.2018.02.018>.
 10. Carrasco N, Rojas M, Aceituno P, Andree KB, Lacuesta B, Furones MD: ***Perkinsus chesapeaki* observed in a new host, the European common edible cockle *Cerastoderma edule*, in the Spanish Mediterranean coast**. *Journal of Invertebrate Pathology* 2014, **117**:56-60. DOI: <https://doi.org/10.1016/j.jip.2014.01.009>.
 11. Metzger MJ, Villalba A, Carballal MJ, Iglesias D, Sherry J, Reinisch C, Muttray AF, Baldwin SA, Goff SP: **Widespread transmission of independent cancer lineages within multiple bivalve species**. *Nature* 2016, **534**(7609):705-709. DOI: 10.1038/nature18599.
 12. Díaz S, Iglesias D, Villalba A, Carballal MJ: **Long-term epidemiological study of disseminated neoplasia of cockles in Galicia (NW Spain): temporal patterns at individual and population levels, influence of environmental and cockle-based factors and lethality**. *Journal of Fish Diseases* 2016, **39**(9):1027-1042. DOI: 10.1111/jfd.12436.
 13. Barber BJ: **Neoplastic diseases of commercially important marine bivalves**. *Aquat Living Resour* 2004, **17**(4):449-466. DOI: <https://doi.org/10.1051/alr:2004052>.
 14. Azevedo C: **Occurrence of an unusual branchial mycoplasma-like infection in cockle *Cerastoderma edule* (Mollusca, Bivalvia)**. *Dis Aquat Org* 1993, **16**:55-59. DOI: 10.3354/dao016055.

15. Sawabe T, Kita-Tsukamoto K, Thompson FL: **Inferring the evolutionary history of vibrios by means of multilocus sequence analysis.** *Journal of Bacteriology* 2007, **189**(21):7932-7936. DOI: 10.1128/JB.00693-07.
16. Sawabe T, Ogura Y, Matsumura Y, Gao F, Amin AKM, Mino S, Nakagawa S, Sawabe T, Kumar R, Fukui Y *et al*: **Updating the Vibrio clades defined by multilocus sequence phylogeny: proposal of eight new clades, and the description of Vibrio tritonius sp. nov.** *Frontiers in Microbiology* 2013, **4**:414. DOI: <https://doi.org/10.3389/fmicb.2013.00414>.
17. Thompson FL, Gevers D, Thompson CC, Dawyndt P, Naser S, Hoste B, Munn CB, Swings J: **Phylogeny and Molecular Identification of Vibrios on the Basis of Multilocus Sequence Analysis.** *Appl Environ Microbiol* 2005, **71**(9):5107-5115. DOI: 10.1128/AEM.71.9.5107-5115.2005.
18. Thompson FL, Iida T, Swings J: **Biodiversity of Vibrios.** *Microbiol Mol Biol Rev* 2004, **68**(3):403-431. DOI: 10.1128/mnbr.68.3.403-431.2004.
19. Romalde JL, Dieguez AL, Lasa A, Balboa S: **New Vibrio species associated to molluscan microbiota: a review.** *Frontiers in Microbiology* 2014, **4**. DOI: 10.3389/fmicb.2013.00413.
20. Goudenege D, Travers MA, Lemire A, Petton B, Haffner P, Labreuche Y, Tourbiez D, Mangenot S, Calteau A, Mazel D *et al*: **A single regulatory gene is sufficient to alter Vibrio aestuarianus pathogenicity in oysters.** *Environ Microbiol* 2015, **17**(11):4189-4199. DOI: 10.1111/1462-2920.12699.
21. Lemire A, Goudenege D, Versigny T, Petton B, Calteau A, Labreuche Y, Le Roux F: **Populations, not clones, are the unit of vibrio pathogenesis in naturally infected oysters.** *ISME J* 2015, **9**(7):1523-1531. DOI: 10.1038/ismej.2014.233.
22. Lasa A, di Cesare A, Tassistro G, Borello A, Gualdi S, Furones D, Carrasco N, Cheslett D, Brechon A, Paillard C *et al*: **Dynamics of the Pacific oyster pathobiota during mortality episodes in Europe assessed by 16S rRNA gene profiling and a new target enrichment next-generation sequencing strategy.** *Environ Microbiol* 2019, **21**(12):4548-4562. DOI: 10.1111/1462-2920.14750.
23. de Lorgeril J, Lucasson A, Petton B, Toulza E, Montagnani C, Clerissi C, Vidal-Dupiol J, Chaparro C, Galinier R, Escoubas JM *et al*: **Immune-suppression by OsHV-1 viral infection causes fatal bacteraemia in Pacific oysters.** *Nat Commun* 2018, **9**(1):4215. DOI: 10.1038/s41467-018-06659-3.
24. Travers MA, Miller KB, Roque A, Friedman CS: **Bacterial diseases in marine bivalves.** *Journal of Invertebrate Pathology* 2015, **131**:11-31. DOI: 10.1016/j.jip.2015.07.010.
25. Tison DL, Seidler RJ: **Vibrio-Aestuarianus - a New Species from Estuarine Waters and Shellfish.** *International Journal of Systematic Bacteriology* 1983, **33**(4):699-702. DOI: <https://doi.org/10.1099/00207713-33-4-699>.
26. Garnier M, Labreuche Y, Nicolas J-L: **Molecular and phenotypic characterization of Vibrio aestuarianus subsp francensis subsp nov., a pathogen of the oyster Crassostrea gigas.** *Systematic and Applied Microbiology* 2008, **31**(5):358-365. DOI: <https://doi.org/10.1016/j.syapm.2008.06.003>.

27. Saulnier D, De Decker S, Tourbiez D, Travers MA: **Development of a duplex Taqman real-time PCR assay for rapid identification of *Vibrio splendidus*-related and *V. aestuarianus* strains from bacterial cultures.** *J Microbiol Methods* 2017, **140**:67-69. DOI: 10.1016/j.mimet.2017.07.002.
28. Saulnier D, De Decker S, Haffner P: **Real-time PCR assay for rapid detection and quantification of *Vibrio aestuarianus* in oyster and seawater: A useful tool for epidemiologic studies.** *Journal of Microbiological Methods* 2009, **77**(2):191-197. DOI: 10.1016/j.mimet.2009.01.021.
29. Garcia C, Thebault A, Degremont L, Arzul I, Miossec L, Robert M, Chollet B, Francois C, Joly J-P, Ferrand S *et al*: **Ostreid herpesvirus 1 detection and relationship with *Crassostrea gigas* spat mortality in France between 1998 and 2006.** *Veterinary Research* 2011, **42**. DOI: 10.1186/1297-9716-42-73.
30. Saulnier D, De Decker S, Haffner P, Cobret L, Robert M, Garcia C: **A large-scale epidemiological study to identify bacteria pathogenic to Pacific oyster *Crassostrea gigas* and correlation between virulence and metalloprotease-like activity.** *Microbial Ecology* 2010, **59**(4):787-798. DOI: <https://doi.org/10.1007/s00248-009-9620-y>.
31. Kumar S, Stecher G, Li M, Knyaz C, Tamura K: **MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms.** *Mol Biol Evol* 2018, **35**(6):1547-1549. DOI: 10.1093/molbev/msy096.
32. Saitou N, Nei M: **The Neighbor-Joining Method: a new method for reconstructing phylogenetic trees.** *Molecular Biology and Evolution* 1987, **4**(4):406-425. DOI: 10.1093/oxfordjournals.molbev.a040454.
33. Eck RV, Dayhoff MO: **Atlas of protein sequence and structure.** Maryland: Silver Springs; 1966.
34. Wilson K: **Preparation of genomic DNA from bacteria.** In: *Current protocols in molecular biology*. Edited by Ausubel FM BR, Kingston RE, Seidman JG, Smith JA, Struhl K. New York: Greene Publishing and Wiley Interscience; 1987.
35. Ezaki T, Hashimoto Y, Yabuubhi E: **Fluorometric Deoxyribonucleic Acid-Deoxyribonucleic Acid Hybridization in Microdilution Wells as an Alternative to Membrane Filter Hybridization in which Radioisotopes Are Used To Determine Genetic Relatedness among Bacterial Strains.** *International Journal of Systematic and Evolutionary Microbiology* 1989, **39**(3):224-229. DOI: <https://doi.org/10.1099/00207713-39-3-224>.
36. Mougin J, Flahaut C, Roquigny R, Bonnin-Jusserand M, Grard T, Le Bris C: **Rapid Identification of *Vibrio* Species of the Harveyi Clade Using MALDI-TOF MS Profiling With Main Spectral Profile Database Implemented With an In-House Database: Luvibase.** *Frontiers in Microbiology* 2020, **11**:2494.
37. Rahmani A, Vercauteren M, Vranckx K, Boyen F, Bidault A, Pichereau V, Decostere A, Paillard C, Chiers K: **MALDI-TOF MS as a promising tool to assess potential virulence of *Vibrio tapetis* isolates.** *Aquaculture* 2021, **530**:735729. <https://doi.org/10.1016/j.aquaculture.2020.735729>.
38. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD *et al*: **SPAdes: a new genome assembly algorithm and its applications**

- to single-cell sequencing.** . *Journal of computational biology* 2012, **19**(5):455-477. DOI: <https://doi.org/10.1089/cmb.2012.0021>.
39. Gurevich A, Saveliev V, Vyahhi N, Tesler G: **QUAST: quality assessment tool for genome assemblies.** *Bioinformatics*, 2013, **29**(8):1072-1075. DOI: 10.1093/bioinformatics/btt086.
40. Vallenet D, Calteau A, Dubois M, Amours P, Bazin A, Beuvin M, Burlot L, Bussell X, Fouteau S, Gautreau G *et al*: **MicroScope: an integrated platform for the annotation and exploration of microbial gene functions through genomic, pangenomic and metabolic comparative analysis.** *Nucleic Acids Research* 2020, **48**(D1):579-589. DOI: <https://doi.org/10.1093/nar/gkz926>.
41. Richter M, Rosselló-Móra R, Glöckner FO, Peplies J: **JSpeciesWS: a web server for prokaryotic species circumscription based on pairwise genome comparison.** . *Bioinformatics* 2015, pii: **btv681**. DOI: <https://doi.org/10.1093/bioinformatics/btv681>.
42. Kim M, Oh H-S, Park S-C, Chun J: **Towards a taxonomic coherence between average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of prokaryotes.** *International Journal of Systematic and Evolutionary Microbiology* 2014, **64**(2):346-351. DOI: <https://doi.org/10.1099/ijs.0.059774-0>.
43. Degremont L, Bedier E, Soletchnik P, Ropert M, Huvet A, Moal J, Samain JF, Boudry P: **Relative importance of family, site, and field placement timing on survival, growth, and yield of hatchery-produced Pacific oyster spat (*Crassostrea gigas*).** *Aquaculture* 2005, **249**(1-4):213-229. DOI: <https://doi.org/10.1016/j.aquaculture.2005.03.046>.

FIGURES AND TABLES

Figure 1. Phylogenetic position of *Vibrio aestuarianus* subsp. *cardii* subsp. nov. and *V. aestuarianus* subsp. *francensis* subsp. nov. strains and virulence of *V. aestuarianus* subsp. *cardii* subsp. nov. Molecular phylogenetic analysis by a Maximum Likelihood method based on the Kimura 2-parameter model conducted in MEGA X. The analysis involved 42 nucleotide sequences with a total of 570 positions. Proposed type strains are indicated in bold.

Virulence was tested by intramuscular injection of hatchery-produced cockles, using two bacterial doses (high: OD 600 nm = 0.4 intermediate OD 600 nm = 0.04). Strains causing more than 70% mortality are indicated by a red oval.

Figure 2. MALDI TOF analyses. Spectra of three representative strains were assigned and compared by principal component analysis: LMG7909T for *V. aestuarianus aestuarianus* subsp. nov, 02-041 for *V. aestuarianus francensis* subsp. nov and 12-122 3T3 for *V. aestuarianus cardii* subsp. nov

(A) Bacterial strains identification by MALDI-TOF analysis using Bruker database. Presence/absence of discriminatory peaks revealed by PCA is also indicated.

(B) Twelve spectra were compared by PCA clustering with MBT compass

(C) MALDI-TOF spectra overview. Major discriminatory peaks are indicated

Table 1. Key phenotypic characteristics (antibiotic sensitivity and growth capacity) of *Vibrio aestuarianus* strains isolated from cockles and oysters compared with the *V. aestuarianus* type strain.

Strains: 1, 17 strains from cockles; 2, *V. aestuarianus* type strain LMG 7909T; 3, oyster strain 02/041; 4, oyster strain 02/093; 5, Results of Garnier et al. for 11 oyster strains [26]

Flumequine 30 µg (UB30), Kanamycin 30 µg (K30), Tetracycline 30 µg (TE30), Sulfamethoxazole / trimethoprim 19:1 25 µg (SXT25), Streptomycin 25 µg (S25), Sulfonamides 300 µg (S3), Erythromycin 30 µg (E30), Penicillin 10 UI (6 µg) (P10), Trimethoprim 5 µg (W5), Chloramphenicol 30 µg (C30), Ampicillin 10 µg (AMP10), Gentamicin 30 µg (CN30) or the vibriostatic agent, Pteridine 150 µg and 10 µg (O129/150 and O129/10).

nd: no data

	1	2	3	4	5
AMP10, C30, E30, K30, TE30	+	+	+	+	+
P10, UB30, SXT25, S25, S3, W5, CN30, O129/150, O129/10	+	+	+	+	nd
Growth at 4°C and 20°C	+	+	+	+	+
Growth at 35 and 37°C	+	+	+	+	-
Growth at 40°C or in absence of salt	-	-	-	-	-
Growth in 6% NaCl	-	+	-	-	-
Growth on TCBS	-	+	-	-	-

1 **Table 2.** DDH values measured by comparing *Vibrio aestuarianus* strains from cockles and oysters with *V. aestuarianus* type strain (LMG 7909T) and other
 2 close species

3 Taxa. 1, *V. aestuarianus* subsp. *cardii* subsp. nov. 12/122 3T3; 2, *V. aestuarianus* subsp. *cardii* subsp. nov. 15/061 4T2; 3, *V. aestuarianus* subsp. *francensis*
 4 subsp. nov. 02/041; 4, *V. aestuarianus* subsp. *aestuarianus* subsp. nov. LMG 7909T; 5, *Vibrio furnisii* LMG 7910T; 6, *Vibrio ordalii* LMG 13544T; 7, *Vibrio*
 5 *salilacus* JCM 19265T.

6

	1	2	3	4	5	6	7
<i>V. aestuarianus</i> subsp. <i>cardii</i> subsp. nov. 12/122 3T3	100						
<i>V. aestuarianus</i> subsp. <i>cardii</i> subsp. nov. 15/061 4T2	84 (7)	100					
<i>V. aestuarianus</i> subsp. <i>francensis</i> subsp. nov. 02/041	80 (18)	80 (11)	100				
<i>V. aestuarianus</i> subsp. <i>aestuarianus</i> subsp. nov. LMG 7909T	70 (2)	71 (5)	79 (22)	100			
<i>V. furnisii</i> LMG 7910T	14 (3)	18 (4)	16 (5)	17 (20)	100		
<i>V. ordalii</i> LMG 13544T	14 (12)	16 (5)	16 (5)	9 (16)	8 (19)	100	
<i>V. salilacus</i> JCM 19265T	15 (4)	16 (4)	12 (2)	n.d.	n.d.	n.d.	100

7

8 **Table 3.** Genome characteristics for *Vibrio aestuarianus* subsp. *cardii* subsp. nov., *V. aestuarianus* subsp. *francensis* subsp. nov and *V. aestuarianus* subsp.
9 *aestuarianus* subsp. nov. type strain (LMG 7909T).

Strain	Source	Genome size (bp)	No. of CDS	No. of tRNA	G+C content (%)	N ₅₀ (bp)	L ₅₀	CheckM Completnes	CheckM contamin	Accession no.	No. of reads
12/122_3T3	<i>Cerastoderma edule</i>	4,350,835	4,449	97	42.73	50,782	25	100%	0.91%	JAAKZN000000000	694,942
15/061_1T1	<i>Cerastoderma edule</i>	4,099,086	4,205	102	42.89	43,962	29	100%	0.91%	JAAKZK000000000	843,350
15/064_3T2	<i>Cerastoderma edule</i>	4,024,371	4,101	91	42.79	44,282	29	99.93%	0.83%	JAAKZM000000000	703,751
15/075_3T2	<i>Cerastoderma edule</i>	4,394,098	4,614	98	42.68	55,001	26	99.90%	1.10%	JAAKZL000000000	1,091,426
LMG 7909T	Type strain	4,229,739	4,020	103	42.67	96,002	16	100%	0.09%	JAAZTU000000000	897,933
01/032	<i>Crassostrea gigas</i>	4,204,160	4,188	84	42.83	-	-	99.72	0.12%	JABANI000000000	-
01/151	<i>Crassostrea gigas</i>	4,368,412	4,350	84	42.73	-	-	100%	0.12%	JABANJ000000000	-
01/308	<i>Crassostrea gigas</i>	4,510,343	4,534	85	42.52	-	-	100%	0.16%	JABANK000000000	-
02/041	<i>Crassostrea gigas</i>	4,196,441	4,063	84	42.83%	-	-	99.72%	0.12%	JAAZTV000000000	-

07/115	<i>Crassostrea gigas</i>	4,245,988	4,243	84	42.79	-	-	99.72%	0.12%	JABANH000000000	-
--------	------------------------------	-----------	-------	----	-------	---	---	--------	-------	-----------------	---

11 **Table 4.** Key phenotypic characteristics (API) for *Vibrio aestuarianus* strains isolated from cockles
 12 and oysters from *Vibrio aestuarianus* type strain.

13 Strains: 1, 17 strains from cockles; 2, *V. aestuarianus* LMG 7909T; 3, Oyster strain 02/041; 4, Oyster
 14 strain 02/093; 5, Results of Garnier et al. for 11 oyster strains [26].

15 +, positive for >90% of the strains; (+), positive for 75-89%; -, negative for <10%; (-), negative for 25–
 16 11%; v, variable for 26–74%.

	1	2	3	4	5
Fermentation of D-glucose	+	+	-	-	(+)
Fermentation of N-acetyl-D-glucosamine	(+)	+	+	+	(+)
Fermentation of sucrose and D-trehalose	+	+	+	+	+
Fermentation of potassium gluconate	-	+	+	-	v
Fermentation of D-galactose	+	+	-	-	+
Fermentation of D-maltose	(+)	+	-	+	v
Fermentation of D-cellobiose	(+)	+	-	-	v
Fermentation of D-fructose and D-mannose	+	+	+	-	+
Fermentation of starch, D-ribose and glycogen	(+)	+	-	-	v
Arginine dihydrolase	+	+	+	+	(-)
Beta-galactosidase	+	+	+	+	+
Tryptophan deaminase	-	-	-	-	v
Gelatinase	v	+	-	+	v
Indole production	+	+	-	+	+
Utilisation of D-mannitol	v	-	-	-	-
Utilisation of citrate	(-)	-	-	-	v
LDC, ODC, VP, H ₂ S, URE, INO, SOR, RHA, MEL, ARA, GLY, ERY, DXYL, LXYL, ADO, MDX, SBE, DUL, MDM, MDG, AMY, ARB, ESC, SAL, LAC, INU, MLZ, RAF, XLT, GEN, TUR, LYX, TAG, DFUC, LFUC, DARL, LARL, 2KG, 5KG	-	-	-	-	-

18 **Table 5.** Fatty acid compositions (as percentage of the total fatty acids present) in *Vibrio*
 19 *aestuarianus* strains from cockles and oysters, *V. aestuarianus* type strain and related species.

20 Taxa. 1, *V. aestuarianus* subsp. *cardii* subsp. nov. 12/122 3T3; 2, *V. aestuarianus* subsp. *francensis*
 21 subsp. nov. 02/041T; 3, *V. aestuarianus* subsp. *aestuarianus* subsp. nov. LMG 7909T; 4, *V. furnisii*
 22 LMG 7910T; 5, *V. ordalii* LMG 13544T.

23 nd = no data

24

	1	2	3	4	5
C12:0	4.4 (0.4)	4	3.2	6.7	-
C12:0; 3-OH	2.1 (0.4)	3.4	1.3	6.1	4.5
iso-C14:0	1.2 (0.1)	2.1	TR	-	-
C14:0	4.8 (1.7)	8.7	4	4.5	5.9
iso-C16:0	4.1 (1.4)	5.4	5.6	TR	-
C16:0	18.6 (1.5)	16.3	13.4	12	19.2
C18:1 w7c	14.5 (2.7)	11.3	22.6	17.1	23.4
summed features 2	2.9 (1.6)	2.3	1.1	6.7	1.5
summed features 3	41.6 (1.8)	34.6	41.8	37.5	39.8

25


Virulence by injection

Intermediate dose

High dose

Vibrio aestuarianus subspecies and Host isolation source

77 12/122 1T3
 12/122 2T3
 12/122 1T1
 12/122 4T3
12/122 3T3
 57 64 12/122 5T3
 15/075 3T2
 15/064 3T2
 15/064 4T2
 15/064 1T2
 15/061 5T2
 15/061 3T2
 15/061 3T1
 15/061 2T2
 15/061 1T1
 15/075 1T2
 83 15/061 4T2
 75 15/061 4T4



V. aestuarianus subsp. *cardii* subsp. nov.



V. aestuarianus subsp. *aestuarianus* subsp. nov.

V. aestuarianus LMG 7909T

12/016 1T1
02/041
 08/042 1T1
 08/055 3T1
 08/110 4T5
 08/132 3T1
 08/162 1T1
 09/006 1T1
 70 10/027 1T1
 11/009 1T1
 60 11/010 2T1
 11/010 3T1
 11/025/5b
 11/026/3a
 11/035 1T1
 11/051 11MT1
 97 08/056 1T1
 08/081 5T1
 08/090 3T1
 08/093 3T1
 10/075 3T2
 11/142 10T1

V. aestuarianus subsp. *francensis* subsp. nov.



Vibrio anguillarum strain 4299



0.020

A

	Identification (score)	m/z					
		3660.7	3674.3	3877.4	7321.7	7349.8	7440.1
<i>V. aestuarianus aestuarianus</i> subsp. nov (strain LMG 7909T)	<i>V. aestuarianus</i> (2.520)						+
<i>V. aestuarianus francensis</i> subsp. nov (strain 02-041)	<i>V. aestuarianus</i> (2.450)			++			+
<i>V. aestuarianus cardii</i> subsp. nov (strain 12-122 3T3)	<i>V. aestuarianus</i> (2.360)	+	+		++	+	

