

Supplementary Information

**A journey in lanthanide coordination chemistry: from evaporable
dimers to magnetic materials and luminescent devices**

Kevin Bernot*, Carole Daiguebonne, Guillaume Calvez, Yan Suffren and Olivier Guillou

Table S1. Main geometric parameters of the dimers

Dimer	Dimensionality	Site symmetry and CShM	Dy-Dy distance (Å)	Remark	Ref
DyPyNO	0D	D_{4d} , 0.757	4.07(2)	Double-bridged dimer	¹
Dy₂PyNO₃	0D	Dy ₁ , C _{4v} , 0.927 Dy ₂ , C _{4v} , 0.605	3.86(1)	Triple-bridge dimer	²
[(Dy₂PyNO₃)(biPyNO)]_∞	3D	Dy ₁ , C _{4v} , 0.828 Dy ₂ , C _{4v} , 0.682	3.94(1) 3.93(2)	3D assembly of triple-bridged dimers	²
DyttaPyNO	0D	D_{2d} , 0.403	4.15 (6)	Double-bridged dimer	³
161DyttaPyNO	0D	Idem	Idem-	Isostructural to DyttaPyNO	³
164DyttaPyNO	0D	Idem	Idem	Isostructural to DyttaPyNO	³
YDyttaPyNO	0D	Idem	Idem	Isostructural to DyttaPyNO	³
DyPyNOS1	0D	D_{4d} , 0.618	4.02(8)	Double-bridged dimer	⁴
DyPyNOS2	0D	D_{4d} , 0.345	4.05(2)	Double-bridged dimer	⁴
DyPyNO-NO₂	0D	D_{4d} , 0.551	4.07(4)	Double-bridged dimer	⁵
DyPyNO-NO₂-film	Thick film	-	-	330 nm thick film on Teflon substrate	⁵

Table S2. Main magnetic figures of merit of the dimers

Dimer	g_z	Dy-Dy interaction (cm ⁻¹)*	H _{dc} (Oe)	τ_0 (s)	U _{eff} (K)	C (s ⁻¹ .K ⁻¹), n, and τ_{tun} (s) in $\tau^{-1} = CT^n + \tau_{\text{tun}}^{-1}$	α_{2K}	τ_{2K} (s)	Relaxing fraction (1 - χ_s/χ_T) _{2K}	Ref
DyPyNO	18.69 exp 19.53 theo	J _{exp} = -2.46 (0.03) J _{theo} = J _{dip} + J _{exch} J _{theo} = -2.44 - 0.25 = -2.69 J _{crossing} = -3.36 cm ⁻¹	0	1.87 (2.2) × 10 ⁻¹²	192 (1)	C = 0.037(6), n = 4.79(4), $\tau_{\text{tun}} = 0.55$ (1)	0.05	0.38	0.99	¹
Dy₂PyNO₃	n.a.	n.a.	0 1900	n.a. 3.18 (0.22) × 10 ⁻⁶	n.a. 24.4 (1)	n.a. n.a.	0.22 0.45	1.95 × 10 ⁻⁴ 0.01	0.57 0.94	²
[(Dy₂PyNO₃)(biPyNO)]_∞	n.a.	n.a.	1900	1.25 10 ⁻⁵	10.3	n.a.	0.51	1.11 10 ⁻³	0.85	²
DyttaPyNO	19.59	J _{exp} = 2.92 J _{theo} = J _{dip} + J _{exch} = -2.36 - 0.50 = -2.86	0	n.a.	n.a.	C = 2.25(12) × 10 ⁻³ n = 5.42(19) $\tau_{\text{tun}} = 2.2$ (1)	0.21	1.13	0.99	³
¹⁶¹DyttaPyNO	**	**	0	n.a.	n.a.	C = 0.00199(78) n = 5.38(17) $\tau_{\text{tun}} = 1.8$ (1)	0.20	1.42	0.98	³
¹⁶⁴DyttaPyNO	**	**	0	n.a.	n.a.	C = 0.00219(11) n = 5.34(18) $\tau_{\text{tun}} = 3.0$ (1)	0.20	2.04	0.99	³
YDyttaPyNO	n.a.	n.a.	0	1.81 (2.8) 10 ⁻⁷	117(7)	C = 0.149(1) n = 3.27 (1) $\tau_{\text{tun}} = 0.08$ (0.05)	0.32	0.09	0.97	³
DyPyNOS1	19.50	J _{theo} = J _{dip} + J _{exch} = -2.525 - 0.125 = -2.65	0	n.a.	n.a.	C = 0.200(249) n = 4.05(54) $\tau_{\text{tun}} = 0.08$ (1)	0.24	0.061	0.91	⁴
DyPyNOS2	19.52	J _{theo} = J _{dip} + J _{exch} = -2.49 - 0.5 = -2.99	0	n.a.	n.a.	C = 0.279(148) n = 4.12(29) $\tau_{\text{tun}} = 0.07$ (1)	0.28	0.047	0.97	⁴

DyPyNO₂	19.56	J _{exp} = 2.9	0	3.1 (1.5) 10 ⁻¹⁰	187 (3)	C=0.037(3) n=3.93(5)	0.42	1.54	0.96	5
DyPyNO₂-film	-	-		6.6 (2.2) 10 ⁻¹¹	166 (4)	C=0.104(6) n=4.25(4)**	0.24 **	0.128**	0.95	5

Fitted values may differ from the originally reported ones because up-to-date homogenous fitting procedure is applied.

*In the S=1/2 ground state approximation. ** isotopic substitution does not influence g values and magnetic coupling. *** Values at 4K.

- (1) Yi, X.; Bernot, K.; Pointillart, F.; Poneti, G.; Calvez, G.; Daiguebonne, C.; Guillou, O.; Sessoli, R. A Luminescent and Sublimable Dy-III-Based Single-Molecule Magnet. *Chem.-Eur. J.* **2012**, *18*, 11379-11387.
- (2) Yi, X.; Calvez, G.; Daiguebonne, C.; Guillou, O.; Bernot, K. Rational Organization of Lanthanide-Based SMM Dimers into Three-Dimensional Networks. *Inorg. Chem.* **2015**, *54*, 5213-5219.
- (3) Huang, G.; Yi, X.; Jung, J.; Guillou, O.; Cador, O.; Pointillart, F.; Le Guennic, B.; Bernot, K. Optimization of magnetic relaxation and isotopic enrichment in dimeric Dy^{III} single-molecule magnets. *Eur. J. Inorg. Chem.* **2018**, 326-332.
- (4) Yi, X.; Pointillart, F.; Le Guennic, B.; Jung, J.; Daiguebonne, C.; Calvez, G.; Guillou, O.; Bernot, K. Rational engineering of dimeric Dy-based Single-Molecule Magnets for surface grafting. *Polyhedron* **2019**, *164*, 41-47.
- (5) Cimatti, I.; Yi, X.; Sessoli, R.; Puget, M.; Guennic, B. L.; Jung, J.; Guizouarn, T.; Magnani, A.; Bernot, K.; Mannini, M. Chemical tailoring of Single Molecule Magnet behavior in films of Dy(III) dimers. *Appl. Surf. Sci.* **2018**, *432*, 7-14.