

EDGE: two routes to dark matter core formation in ultra-faint dwarfs

Matthew D.A. Orkney, Justin I. Read, Martin P. Rey, Imran Nasim, Andrew Pontzen, Oscar Agertz, Stacy Y. Kim, Maxime Delorme, Walter Dehnen

► To cite this version:

Matthew D.A. Orkney, Justin I. Read, Martin P. Rey, Imran Nasim, Andrew Pontzen, et al.. EDGE: two routes to dark matter core formation in ultra-faint dwarfs. Monthly Notices of the Royal Astronomical Society, 2021, 504 (3), pp.3509-3522. 10.1093/mnras/stab1066 . hal-03129407

HAL Id: hal-03129407 https://hal.science/hal-03129407

Submitted on 5 May 2023 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EDGE: two routes to dark matter core formation in ultra-faint dwarfs

Matthew D. A. Orkney[®],¹* Justin I. Read[®],¹ Martin P. Rey[®],² Imran Nasim,¹ Andrew Pontzen,³ Oscar Agertz[®],² Stacy Y. Kim,¹ Maxime Delorme⁴ and Walter Dehnen^{®5,6,7}

¹Department of Physics, University of Surrey, Guildford GU2 7XH, UK

²Lund Observatory, Department of Astronomy and Theoretical Physics, Lund University, Box 43, SE-221 00 Lund, Sweden

³Department of Physics and Astronomy, University College London, London WC1E 6BT, UK

⁴Département d'Astrophysique/AIM, CEA/IRFU, CNRS/INSU, Université Paris-Saclay, F-91191 Gif-Sur-Yvette, France

⁵Astronomisches Recheninstitut, Zentrum für Astronomie der Universität Heidelberg, Mönchhofstraße 12-14, D-69120 Heidelberg, Germany

⁶Universitäts-Sternwarte München, Scheinerstraße 1, D-81679 München, Germany

⁷School for Physics and Astronomy, University of Leicester, University Road, Leicester LE1 7RH, UK

Accepted 2021 April 13. Received 2021 April 2; in original form 2020 December 21

ABSTRACT

In the standard Lambda cold dark matter paradigm, pure dark matter simulations predict dwarf galaxies should inhabit dark matter haloes with a centrally diverging density 'cusp'. This is in conflict with observations that typically favour a constant density 'core'. We investigate this 'cusp-core problem' in 'ultra-faint' dwarf galaxies simulated as part of the 'Engineering Dwarfs at Galaxy formation's Edge' project. We find, similarly to previous work, that gravitational potential fluctuations within the central region of the simulated dwarfs kinematically heat the dark matter particles, lowering the dwarfs' central dark matter density. However, these fluctuations are not exclusively caused by gas inflow/outflow, but also by impulsive heating from minor mergers. We use the genetic modification approach on one of our dwarf's initial conditions to show how a delayed assembly history leads to more late minor mergers and, correspondingly, more dark matter heating. This provides a mechanism by which even ultra-faint dwarfs ($M_* < 10^5 \, M_{\odot}$), in which star formation was fully quenched at high redshift, can have their central dark matter density lowered over time. In contrast, we find that late major mergers can regenerate a central dark matter cusp, if the merging galaxy had sufficiently little star formation. The combination of these effects leads us to predict significant stochasticity in the central dark matter density slopes of the smallest dwarfs, driven by their unique star formation and mass assembly histories.

Key words: methods: numerical – galaxies: dwarf – galaxies: evolution – galaxies: formation – galaxies: haloes – dark matter.

1 INTRODUCTION

The Lambda cold dark matter (ACDM) paradigm presents us with a Universe that has an energy budget dominated by dark matter and dark energy, and in which galaxies are assembled through successive hierarchical mergers (White & Rees 1978). It has proven to be extremely successful in predicting the formation of cosmic structure on large scales (Clowe et al. 2006; Springel, Frenk & White 2006; Tegmark et al. 2006; Dawson et al. 2013; Oka et al. 2014; Planck Collaboration XVI 2014; Wang et al. 2016). However, disagreements between theory and observation endure at (sub-)galactic scales that have become collectively known as 'small-scale puzzles' (e.g. Bullock & Boylan-Kolchin 2017).

The oldest, and perhaps most challenging, of the small-scale puzzles is the 'cusp-core (CC) problem' (e.g. Flores & Primack 1994; Moore 1994; Read et al. 2017). Pure dark matter structure formation simulations in Λ CDM predict a self-similar radial dark matter density profile – the NFW profile (Navarro, Frenk & White 1997). This scales as $\rho \propto r^{-1}$ within a scale radius r_s , referred to as a 'cusp' due to its divergence towards the origin. By contrast, observations of rotation curves in dwarf galaxies appear to

* E-mail: m.d.a.orkney@gmail.com

favour instead a constant inner dark matter density, referred to as a 'core' (e.g. Carignan & Freeman 1988; Flores & Primack 1994; Moore 1994; McGaugh, Rubin & de Blok 2001; Read et al. 2017).

Many solutions to the CC have been proposed in the literature to date. First, observations could have been misinterpreted due to incorrect modelling assumptions. Typical assumptions include spherical symmetry, circular orbits and dynamical pseudo-equilibrium, all of which can be reasonably questioned (e.g. Kuzio de Naray & Kaufmann 2011; Oman et al. 2016; Read et al. 2016b). Secondly, the assumed underlying dark matter model could be incorrect. Alternatives such as warm dark matter (e.g. Hogan & Dalcanton 2000; Avila-Reese et al. 2001; Bode, Ostriker & Turok 2001), selfinteracting dark matter (e.g. Spergel & Steinhardt 2000; Tulin & Yu 2018) or ultra-light dark matter (e.g. Schive, Chiueh & Broadhurst 2014; Ferreira 2020) all predict a lower dark matter density at the centres of dwarf galaxies whilst retaining the predictions of Λ CDM on larger scales. However, in recent years a third class of solution has been gaining traction.

The CC problem originates from a comparison of pure dark matter simulations – that do not model stars or gas (hereafter baryons) – with observations. This opens up the possibility that purely gravitational interactions between dark matter particles and baryons could act to push dark matter out from the centres of dwarf

doi:10.1093/mnras/stab1066

galaxies, transforming a cusp into a core. Three main mechanisms have been proposed to date:

(i) Dynamical friction from infalling dense clumps (e.g. El-Zant, Shlosman & Hoffman 2001; Mo & Mao 2004; Romano-Díaz et al. 2009; Goerdt et al. 2010; Cole, Dehnen & Wilkinson 2011; Nipoti & Binney 2015; Del Popolo & Pace 2016). These clumps impart energy and angular momentum to the dark matter halo, causing it to expand.

(ii) Dynamical friction from a central stellar or gaseous bar that acts similarly to infalling clumps, kinematically 'heating' the background dark matter halo (e.g. Weinberg & Katz 2007).

(iii) A fluctuating gravitational potential driven by gas inflow/outflow due to cooling, stellar winds, and supernovae. This causes dark matter particle orbits to slowly migrate outwards (e.g. Navarro, Eke & Frenk 1996; Read & Gilmore 2005; Mashchenko, Wadsley & Couchman 2008; Pontzen & Governato 2012).

All three mechanisms owe, ultimately, to a time-varying gravitational potential. This allows dark matter particles to exchange orbital energy both with one another and with the stars and gas in the galaxy. What differs is only the physical mechanism that drives the time-dependent gravitational field. In principle, all three mechanisms can act in tandem as galaxies form and evolve.

Despite this diversity of mechanisms in the literature, to date high-resolution galaxy formation simulations have typically favoured mechanism (iii) at the scale of dwarf galaxies (Pontzen & Governato 2012, 2014; Teyssier et al. 2013; Di Cintio et al. 2014; Oñorbe et al. 2015; Dutton et al. 2016). Once gas is allowed to cool (T < 10^4 K) and reach high density ($\rho > 10$ atoms cm⁻³; e.g. Pontzen & Governato 2012; Dutton et al. 2016), these simulations find that gas flows drive repeated fluctuations in the central galaxy mass of amplitude 10-20 per cent over a period less than the local dynamical time. Such fluctuations gradually lower the inner dark matter density on the scale of the stellar half-mass radius, $R_{1/2}$, transforming a dark matter cusp to a core (e.g. Chan et al. 2015; Read, Agertz & Collins 2016a). There is mounting observational evidence that this process occurs in real dwarf galaxies (e.g. Kauffmann 2014; El-Badry et al. 2016; Sparre et al. 2017; Read, Walker & Steger 2019; Hirtenstein et al. 2019; but see also Bose et al. 2019; Oman et al. 2019; Genina et al. 2020).

While dark matter heating may solve the cusp-core problem in isolated gas-rich dwarfs, a new puzzle has recently presented itself: there is a growing body of evidence for small dark matter cores even within 'ultra-faint' dwarf galaxies, typically defined to have stellar masses $M^* < 10^5 \,\mathrm{M_{\odot}}$ (e.g. Amorisco 2017; Contenta et al. 2018; Sanders, Evans & Dehnen 2018; Malhan, Valluri & Freese 2020; Simon et al. 2021). Several papers have suggested that galaxies with so few stars have insufficient energy from stellar feedback to carve out a dark matter core of size 0.5-1 kpc (e.g. Peñarrubia et al. 2012; Munshi et al. 2013; Oñorbe et al. 2015; Tollet et al. 2016). However, dark matter cores form on the scale of the half-mass radius, $R_{1/2}$ (Read et al. 2016a), which can be as small as 30-300 pc in ultrafaint dwarfs (e.g. Simon 2019). Such small cores form much more rapidly and require significantly less energy, raising the possibility that dark matter core formation could proceed 'all the way down' to even the smallest dwarfs (e.g. Read et al. 2016a; Contenta et al. 2018). Furthermore, such small cores remain dynamically important by construction since they exist precisely where the stars and gas do i.e. precisely where we can hope to measure the inner dark matter potential.

In this paper, we use a suite of high-resolution cosmological zoom simulations from the Engineering Dwarfs at Galaxy formation's Edge (EDGE) project (Rey et al. 2019; Agertz et al. 2020; Pontzen et al.

2020; Rey et al. 2020) to explore whether dark matter core formation can proceed even in the very smallest dwarf galaxies. Our simulations model galaxies over the mass range $M_{200c} \sim 1-4 \times 10^9 \,\mathrm{M_{\odot}}$, consistent with ultra-faint dwarfs, and reach a spatial resolution of $\sim 3 \,\mathrm{pc}$, sufficient to resolve even very small dark matter cores.

This paper is organized as follows. In Section 2.1, we describe the EDGE simulations and our numerical methods. In Section 3, we present a first visual impression of the EDGE simulation suite. In Section 4.1, we investigate how gas flows driven by bursty star formation drive early-time core formation in our simulations. In Section 4.2, we show that minor mergers can also drive the formation of cores in EDGE and we validate the robustness of our results using the *N*-body code GRIFFIN. In Section 4.3, we show how late major mergers can reintroduce a dark matter cusp. In Section 5, we discuss the implications of our results for dark matter cusps and cores in the smallest dwarf galaxies. Finally, in Section 6 we present our conclusions.

2 METHOD

2.1 Simulations

The EDGE project is described in detail in Agertz et al. (2020). Here, we briefly summarize the key points. We start with a 512³ resolution cosmological dark matter simulation of a 50 Mpc void region (Fig. 1). All simulations assume cosmological parameters $\Omega_{\rm m} = 0.309$, $\Omega_{\Lambda} = 0.691$, $\Omega_{\rm b} = 0.045$, and $H_0 = 67.77$ km s⁻¹ Mpc⁻¹, in line with data from the *Planck* satellite (Planck Collaboration XVI 2014).

We draw a selection of target haloes from the void volume, chosen from a range in halo mass of $10^9 < M/M_{\odot} < 5 \times 10^9$. These target haloes are resimulated following the zoom simulation technique (Katz & White 1993; Oñorbe et al. 2014) with the adaptive mesh refinement (AMR) code RAMSES (Teyssier 2002). This grants us a highly resolved target galaxy within its lower resolution wider cosmological context. The velocity in our initial conditions is then adjusted to match the velocity of the target halo, which reduces the impact of numerical diffusion effects (see Pontzen et al. 2020).

Key details of our RAMSES simulations are presented in Table 1. Each simulation is run over the redshift range $99 \ge z \ge 0$, with a minimum of 100 outputs spaced linearly with the scale factor, *a*. The contamination fraction, defined as the fraction of lower resolution dark matter particles within the virial radius, is never greater than 2×10^{-5} in any of our simulations. This is relevant for the impact of numerical relaxation, which we discuss further in Appendix A. Our simulations are run at a resolution in which the dark matter particle mass approaches 100 M_{\odot} in the high-resolution Lagrangian region of the target galaxy, with a spatial resolution ~ 3 pc in the most resolved zoom regions. At this resolution, the momentum injection of individual supernova into the interstellar medium can be accurately resolved (Kimm et al. 2015), avoiding the need for delayed cooling, inflated supernovae energies, or sub-grid wind models (e.g. Read et al. 2016a; Agertz et al. 2020).

Star formation is described with a Schmidt law (Schmidt 1959; Kennicutt 1998) in gas cells that meet certain density and temperature requirements:

$$\dot{\rho_*} = \epsilon_{\rm ff} \frac{\rho_{\rm g}}{t_{\rm ff}} \text{ for } \rho_{\rm g} > \rho_{\star} \text{ and } T_{\rm g} < T_{\star}, \tag{1}$$

where $\rho_{\star} = 300 \, m_{\text{proton}} \, \text{cm}^{-3}$ and $T_{\star} = 100 \, \text{K}$. Here, $\dot{\rho_{\star}}$ is the star formation rate density in a gas cell, ρ_{g} is the density per gas cell, $t_{\text{ff}} = \sqrt{3\pi/32G\rho_{\text{g}}}$ is the local free-fall time of the gas, and ϵ_{ff} is the star



Figure 1. The location of the EDGE dark matter haloes selected for higher resolution resimulation from a lower resolution void at z = 0. Each panel shows a surface density plot of a cube. The zoomed panels for each halo are taken from the 'DMO' simulations out to r_{200c} (see Table 1), and show the corresponding M_{200c} mass, as marked. For Halo600, there is a partial-zoom to help illustrate how small the selected haloes are in comparison to the total box size (50 Mpc).

Table 1. Details of all simulations. The simulation labels denote different reference haloes selected from the initial void simulation. From left to right the columns are the simulation label, the physics scheme employed, the mass resolution, the halo mass (M_{200c}) at z = 0, total stellar mass within the virial radius (r_{200c}) at z = 0, the virial radius at z = 0, the projected half-light radius at z = 0, and the V-band magnitude at z = 0. The simulations are ordered by the M_{200c} mass of the full physics simulations.

Name	Physics	Resolution $[m_{\text{DM}}, m_{\text{gas}}, m_*]/M_{\odot}$	$M_{200c}~(\mathrm{M}_{\odot})$	r _{200c} (kpc)	$M_{*}~({ m M}_{\odot})$	<i>R</i> _{half} (pc) (projected)	M_V (mag)
Halo1445 DMO	DM-only	[139, -, -]	1.39×10^{9}	24.92	_	_	_
Halo1459 DMO	DM-only	[139, -, -]	1.44×10^{9}	25.20	_	_	-
Halo600 DMO	DM-only	[139, -, -]	3.42×10^{9}	33.62	_	_	-
Halo605 DMO	DM-only	[139, -, -]	3.33×10^{9}	33.31	_	_	_
Halo624 DMO	DM-only	[139, -, -]	3.58×10^9	34.13	-	-	-
Halo1459 DMO GM:Later	DM-only	[139, -, -]	1.47×10^{9}	25.34	_	_	_
Halo1459 DMO GM:Latest	DM-only	[139, -, -]	1.47×10^9	23.34	-	-	-
Halo1445	Agertz $+ 2020$	[117, 18, 300]	1.32×10^{9}	23.10	1.35×10^{5}	100.80	-6.93
Halo1459	Agertz $+ 2020$	[117, 18, 300]	1.43×10^{9}	23.75	3.77×10^{5}	98.80	-8.03
Halo600	Agertz $+ 2020$	[117, 18, 300]	2.65×10^{9}	31.17	9.84×10^{5}	109.65	-9.19
Halo605	Agertz $+ 2020$	[117, 18, 300]	3.20×10^{9}	31.08	1.93×10^{6}	101.83	-9.84
Halo624	Agertz $+$ 2020	[117, 18, 300]	3.23×10^9	29.18	1.08×10^{6}	107.04	-9.44
Halo1459 GM:Later	Agertz $+ 2020$	[117, 18, 300]	1.43×10^{9}	23.73	1.11×10^{5}	168.16	-6.70
Halo1459 GM:Latest	Agertz + 2020	[117, 18, 300]	1.38×10^9	23.45	8.65×10^4	303.80	-6.44

formation efficiency per free-fall time which is set to 10 per cent in line with arguments from Grisdale et al. (2019). Each stellar particle is initialized at 300 M_{\odot} and is representative of a single-age stellar population described by a Chabrier initial mass function (Chabrier 2003). Stellar feedback from both Type II and Ia supernovae are included, and stellar winds from massive and asymptotic giant branch stars (see Agertz et al. 2013; Agertz & Kravtsov 2015; Agertz et al. 2020 for details).

The epoch of reionization is modelled as a time-dependent uniform UV background around z = 8.5, as in the public release of RAMSES (Haardt & Madau 1996). The exact implementation is discussed further in Rey et al. (2020), and is consistent with a late reionization expected for a cosmic void (e.g. Keating et al. 2020).

In addition, to further test our numerical results, we use the GRIFFIN code (Dehnen 2014) to run a controlled investigation into the effects of the merger history on the dark matter halo of one of our RAMSES

simulations. The motivation for these additional simulations will be made clear in Section 4.2.1. GRIFFIN is a high performance *N*-body integrator that exploits the fast multipole method (FMM), and is an ideal tool to compare against RAMSES because it is based on a fundamentally different numerical integration scheme that has comparable force accuracy to direct summation codes (e.g. Dehnen 2014; Gualandris et al. 2017; Nasim et al. 2020; Nasim et al. 2021). These simulations are run at both the equivalent and $10 \times$ better mass resolution compared to our RAMSES runs, with the former being used as a convergence study. We employ a force softening length of 10 pc in all of these simulations. Additional tests with a softening length of 5 pc showed no measurable change in the results.

2.2 Halo finding

We use the HOP halo finder (Eisenstein & Hut 1998) to identify all distinct bound structures in each simulation output. HOP does not identify haloes within haloes (subhaloes), and so, where necessary, our analysis is supplemented with the AHF (Amiga Halo Finder) (Knollmann & Knebe 2009). Merger trees and halo properties are calculated using PYNBODY (Pontzen et al. 2013) and TANGOS (Pontzen & Tremmel 2018), respectively. We locate the centre of each bound structure using the shrinking spheres method of Power et al. (2003), performed exclusively on the dark matter component, as implemented in PYNBODY. Results from our dark matter only (DMO) simulations are in all cases corrected for the universal baryon fraction.

Throughout this paper, the virial radius, r_{200c} , is defined as the spherical region that is at least 200 times the critical mass density of the universe at that redshift. The halo mass, M_{200c} , is then the total mass of all matter contained within that radius.

3 AN OVERVIEW OF THE SIMULATIONS

Several dwarfs presented in this paper have been discussed already in previous EDGE collaboration papers,¹ however run at a lower 'fiducial' resolution ($m_{\rm DM} = 1112 \, {\rm M}_{\odot}$). Here, since we are interested in resolving potentially very small dark matter cores, the same dwarfs are resimulated at what we called 'hires' resolution in Agertz et al. (2020) ($m_{\rm DM} = 117 \, {\rm M}_{\odot}$). We show convergence tests between these two resolutions in Appendix B, demonstrating that our results presented here do not depend on resolution.

Fig. 1 shows a surface density plot for the dark matter of the total void region from which our initial conditions were selected. The locations for each of our haloes in Table 1 are indicated with zoomed surface density plots, with images taken from the 'DMO' simulations. This highlights that our haloes are chosen from a particularly underdense region, without any major cosmic structure in the near vicinity.

Fig. 2 shows a visual representation of each high-resolution baryonic simulation at z = 0. The left-most panels show the centred dark matter surface density out to r_{200c} , which is indicated by a dashed circle. The middle panels show the central gas density averaged in a 0.2 kpc thick slice through the *z*-axis. The plot is zoomed into the inner five half-light radii, where the 3D half-light radius is marked with a solid circle. Each halo is oriented side-on on the angular momentum vector of the central cold gas (<10⁴ K), where available, which represents the central gas disc if it is present. The right-most panels are a PYNBODY rendering of the halo stars.

¹Halo1459, Halo1459 GM:Later, and Halo1459 GM:Latest appear in Rey et al. (2019). Halo600, Halo605, and Halo624 appear in Rey et al. (2020).

In Fig. 3, we present merger trees for each of our haloes using the 'DMO' simulations. For simplicity, only major mergers on to the main progenitor (coloured line) are shown. The line thickness is representative of each halo mass. The general form of these merger trees are identical for different resolutions and physics, with the one exception that the final merger in Halo624 DMO occurs just after z =0 in Halo624. The present-day main progenitor halo is not necessarily the most massive halo at all times (for instance, Halo600 DMO at z = 6). This is because the main progenitor is defined as the most massive halo at the time of each merger.

4 RESULTS

4.1 Core formation from gas flows

In Fig. 4, we show the evolution of the central dark matter density for all of our EDGE simulations. In the upper panels, the inner 3D dark matter density is plotted at 40 pc. This approaches the inner limit of the region that we consider numerically resolved (see Appendix A). The opaque lines are the results for the baryonic simulations, whereas faint lines are the results for the pure dark matter (DMO) simulations. The corresponding star formation rates of both the main progenitor and all progenitors for the baryonic simulation are plotted in the middle panels, averaged over 100 Myr bins. Included in the lower panels are the 3D density profiles for both baryonic and DMO simulations at z = 0.

From Fig. 4, we see that the inner dark matter density of the baryonic simulations – that include star formation, gas cooling and stellar feedback – is in all cases lower than the DMO simulations after \sim 1 Gyr. This disparity is driven by the early star formation period seen in all our dwarfs. The reduction in inner density occurs over this star-forming period, as expected from the gas-flow mechanism. Star formation reignites at later times in some of our dwarfs, a result that is explored in detail for our fiducial resolution simulations in Rey et al. (2020). The intensity of this late rejuvenation is insufficient to drive a further reduction of the inner density in Halo624 and Halo605, but is great enough in Halo600 to manifest as a sudden dip in the inner density. However, the late rejuvenation in Halo600 lacks the continuous bursts of star formation necessary to grow a large core (Read & Gilmore 2005; Pontzen & Governato 2012).

The lower panels of Fig. 4 show at which radius the density profile slopes in the baryonic simulations depart from the DMO simulations. In all cases, the bulk of the profile flattening occurs within the 3D half-light radius (black dashed line) at approximately 100 pc. This is also consistent with prior work on the gas flow mechanism (see e.g. Chan et al. 2015; Read et al. 2016a; and Section 1).

In Fig. 5, we show the gas-to-dark matter central density ratio $(\rho_{\text{gas}}/\rho_{\text{DM}})$ for two representative dwarf galaxies. This is computed at time intervals of $\Delta a = 0.01$ (the cadence of our simulation outputs), and so the true peak $\rho_{\text{gas}}/\rho_{\text{DM}}$ may be higher. The upper panel shows Halo1445, which is quenched permanently by z = 4 due to the effects of reionization. There are large-scale fluctuations in the density ratio before quenching, corresponding to gas outflow/inflow



Figure 2. A visual representation of the high-resolution baryonic simulations in Table 1. *Left-hand panels*: The dark matter surface density out to r_{200c} (dashed circle). *Middle panels*: The gas density averaged along a 0.2 kpc slice through *z*. The 3D half-light radius is indicated by the solid circle. *Right-hand panels*: The halo stars are rendered in PYNBODY with the *i*, *v*, and *u* bands over the range $23 \le \text{mag arcsec}^{-2} \le 28$, shown at the same scale as the gas density panels. All images are oriented side-on along the angular momentum vector of the cool gas (<10⁴ K) within 1 kpc of the halo centre. The physical size of each frame is indicated by a scale bar in the top left corner.



Figure 3. The accretion history of each halo as taken from the 'DMO' simulations, where <1/30 mass ratio mergers and sub-mergers are excluded for clarity. The central coloured branch is the main progenitor halo and the line thickness represents the M_{200c} mass of each dark matter halo.



Figure 4. Upper panels: The evolution of the 3D dark matter density at 40 pc in the main progenitor halo for the high-resolution dwarf galaxies. The opaque lines show simulations run with baryonic physics, whereas the faint lines show pure dark matter (DMO) simulations using the same initial conditions. Our results are qualitatively similar when selecting the density at different inner radii. *Middle panels:* The star formation rate of the baryonic simulations averaged over bins of 100 Myr. The opaque bars show stars formed within r_{200c} of the main progenitor, whereas fainter bars include stars that are brought in with mergers. *Lower panels:* A comparison of the 3D dark matter density profiles between the baryonic and DMO simulations at z = 0. The black dashed lines mark the 3D half-light radii in each case.

triggered by bursty star formation. These fluctuations cease after quenching, and the gas gradually photoevaporates. The lower panel shows Halo624, which is a more massive dwarf that is able to rejuvenate at late times. As in Halo1445, there are large-scale density ratio fluctuations at early times that diminish after the initial quenching. However, as the halo grows more massive it is able to increase its central gas density (see Rey et al. 2020). Star formation then rejuvenates after 10 Gyr, but the intensity of this late star formation is not sufficient to drive large enough fluctuations in $\rho_{\rm gas}/\rho_{\rm DM}$ to further heat the central dark matter.

4.2 Core formation from minor mergers

Our results so far are in line with previous studies in the literature that suggest that forming kpc-scale dark matter cores becomes inefficient

in ultra-faint dwarfs (Peñarrubia et al. 2012; Garrison-Kimmel et al. 2013; Di Cintio et al. 2014; Madau, Shen & Governato 2014; Maxwell, Wadsley & Couchman 2015; Tollet et al. 2016; Read et al. 2016a). However, the puzzle of apparent dark matter cores in at least some ultra-faints remains (Amorisco 2017; Contenta et al. 2018), see Section 1.

Perhaps the most compelling case for a small dark core in an ultra-faint dwarf comes from the survival of a lone and extended star cluster, offset from the centre of the Eridanus II galaxy (Amorisco 2017; Contenta et al. 2018; Simon et al. 2021). Eridanus II is substantially more extended than any of our reference EDGE dwarfs, with $R_{half} = 299 \pm 12$ pc. Using the genetic modification approach (Roth, Pontzen & Peiris 2016; Rey & Pontzen 2018; Stopyra et al. 2021), we can create alternative mass accretion histories for a galaxy. Rey et al. (2019) modified one of the fiducial resolution EDGE



Figure 5. Evolution of the inner gas-to-dark matter density ratio at 40 pc (ρ_{gas}/ρ_{DM}) for two representative EDGE dwarfs. Prior to quenching by reionization (black dashed line), there are large fluctuations in ρ_{gas}/ρ_{DM} caused by repeated cycles of gas cooling, star formation, and stellar feedback. These fluctuations excite 'dark matter heating' that lowers the central dark matter density (Fig. 4). Halo1445 experiences no further star formation or dark matter heating after it quenches. By contrast, Halo624 grows in mass, accumulating cold gas, and rejuvenating its star formation after ~10 Gyr. Subsequent fluctuations are orders of magnitude smaller than at early times, and so there is no late time dark matter heating.

dwarfs, Halo1459, such that it assembled later. They found that this leads to a larger, lower surface brightness, and lower metallicity dwarf – more similar to Eridanus II. Furthermore, Boldrini, Mohayaee & Silk (2020a) have recently shown that eccentric minor mergers can themselves fluctuate the inner gravitational potential, flattening a dark matter cusp. As such, in this section, we resimulate the genetically modified later forming dwarfs from Rey et al. (2019) at higher resolution to study how late formation impacts the central dark matter density.

We run two modifications of Halo1459 which we call Halo1459 GM:Later and Halo1459GM:Latest (Table 1). The assembly histories of these modifications were already shown at our fiducial resolution in fig. 1 of Rey et al. (2019); they are indistinguishable from the higher resolution trajectories which we show in Fig. 6. The modified haloes are approximately $3 \times (2 \times)$ less massive at z = 8.5 for Halo1459 GM:Latest), but grow to the same mass within a 4 per cent margin by z = 0.

In the upper panel of Fig. 7, we show the evolution of the central density in Halo1459 GM:Later. Now, the inner dark matter density continues to fall long after star formation has ceased. This occurs in both the simulation with baryonic physics and in the DMO simulation. The baryonic simulation is also extremely gas deficient



Figure 6. The halo mass growth history of Halo1459 and two modified variants which we call Halo1459 GM:Later and Halo1459 GM:Latest. Included are grey bands indicating the 68 and 95 per cent scatter for the mass growth histories of haloes throughout our lower resolution void simulation. The bands are truncated at $\sim 3 \times 10^7$ due to the resolution limit of the void simulation. This shows that our modified haloes are within the expected scatter of assembly histories in Λ CDM.

after quenching, with $M_{\rm gas}/M_{\rm DM}(< r_{\rm half-light}) < 10^{-5}$, so any gas flows driven by residual feedback from old stars (as in Rey et al. 2020) have a negligible impact on the overall mass distribution. This reduction in the central density is not seen to the same extent in Halo1459 GM:Latest. We will discuss the reasons for this in Section 4.3.

The lower panel of Fig. 7 shows the radial dark matter density profiles of Halo1459 GM:Later (darker) and Halo1459 DMO GM:Later (lighter) at z = 4 (dashed) and z = 0 (solid). A schematic shows the expected density slopes for a cusp and a core, indicating that whilst the central density slope has declined by z = 0, it still has a slope of $\rho \propto r^{-0.5}$.

It is important to rule out the possibility that any apparent dark matter heating is caused by numerical relaxation. In Appendix A, we calculate the 'relaxation radius', r_{relax} , for our EDGE simulations. This is the radius inside which the numerical relaxation time is equal to the simulation run time and so numerical relaxation will become important. From this calculation, we conclude that our simulations should still be well resolved above $r_{relax} > 25$ pc at z = 0, yet the density clearly evolves on scales larger than this at all times. Therefore, there must be some other mechanism by which the central dark matter density is lowering. It is well established that dynamical friction heating from dense clumps can contribute to core formation (e.g. El-Zant et al. 2001; Mo & Mao 2004; Romano-Díaz et al. 2009; Goerdt et al. 2010; Cole et al. 2011; Nipoti & Binney 2015; Del Popolo & Pace 2016; and Section 1). In Halo1495 GM:Later, the only dense clumps available to drive such a process at late times are merging dark matter subhaloes.

To investigate the veracity of the above late time dark matter heating, and to explore whether merging dark matter subhaloes are indeed the culprit, in the next section we resimulate the sequence of mergers



Figure 7. Upper panel: The evolution of the 3D dark matter density at 40 pc, but for a modified simulation that has a delayed formation history (Halo1459 GM:Later). A black dashed line marks the approximate time at which the galaxy is permanently quenched. *Middle panel:* The star formation rate of the baryonic simulation, averaged over bins of 100 Myr. Opaque bars are stars formed within r_{200c} of the main progenitor, whereas faint bars include stars that are brought in with mergers. *Lower panel:* A comparison of the 3D dark matter density profiles between the baryonic and DMO simulations at both z = 4 (after quenching) and z = 0. A black dashed line marks the 3D half-light radius at z = 0.

that form Halo1459 GM:Later using the non-cosmological *N*-body FMM code GRIFFIN (Dehnen 2014), as described in Section 2.1. This allows us to isolate the heating effects of minor mergers in a non-cosmological setting, and to verify that numerical effects unique to RAMSES are not responsible for the heating.

4.2.1 Testing minor merger induced dark matter core formation with the GRIFFIN code

In this section, we use the GRIFFIN *N*-body code to reproduce the assembly history of Halo1459 GM:Later in a controlled manner. For this, we simulate a series of halo mergers based upon profile fits to the reference RAMSES simulation. We use the coreNFW profile as in Read et al. (2016a), which is a NFW profile (Navarro et al. 1997) adapted to include a parameter *n* that controls the flatness of the central density slope.

We first fit the spherically symmetric dark matter density profile of the main progenitor halo at z = 4. By this time, the central dark matter halo is well established and star formation has permanently quenched. The total stellar mass is low ($\sim 10^5 M_{\odot}$) and the gas content is negligible, so the system can be safely resimulated using exclusively its dark matter component.

Mergers are defined based on every halo identified with HOP at z = 4, on the condition that they contain at least 800 dark matter particles and are destined to merge with the main progenitor. Each merging halo is tracked until the snapshot prior to infall (defined as the point where the merger crosses over r_{200c} of the main progenitor), by which time many of them have coalesced. Despite the large number of individual haloes at z = 4, there are a manageable 38 distinct subhaloes at the time of merging.

For each of these mergers, we perform a spherically symmetric coreNFW profile fit and generate initial conditions using AGAMA (Vasiliev 2018). We use a multipole potential approximation with one hundred grid nodes, an isotropic velocity distribution function, and model the density out to $3 \times r_{200c}$ for each coreNFW profile fit (which is necessary to ensure there is sufficient dynamical friction between interacting haloes at large radii; e.g. Read et al. 2008). A further improvement could include fitting the halo triaxiality, but should not be necessarily for resolving the leading order effects of halo mergers. The resulting merger masses range from $5 \times 10^5 \leq M_{merger}/M_{\odot} \leq 10^8$, with the majority falling around the $2 \times 10^6 M_{\odot}$ mark.

The initial conditions of the main progenitor are integrated forward in time with the GRIFFIN code, as described in Section 2.1. As the simulation reaches the time of each merger infall, the corresponding initial conditions are inserted at the same phase-space location as in the reference RAMSES simulation. In this way, the merger history of the RAMSES simulation from $4 \le z \le 0$ is recreated.

The above method has a few caveats. First, the number density of mergers in a Λ CDM cosmology increases with decreasing halo mass (Stewart et al. 2009; Rodriguez-Gomez et al. 2015). Therefore, a significant amount of mass accretion is neglected by only considering mergers above a certain mass threshold. This yields a final z = 0 mass roughly $1.5 \times$ less that of the original RAMSES simulation, although much of this missing mass is located in the halo outskirts (~80 per cent of the missing mass is exterior to 5 kpc). We also see that the orbits of merging haloes begin to diverge from the reference simulation after two pericentre passages, and it is already established that reproducing exact orbital behaviour of mergers is challenging (e.g. Lux, Read & Lake 2010). Lastly, any mergers already within the

Table 2. The GRIFFIN simulations used to investigate the late-time density reduction in Halo1459 GM:Later. From left to right, the columns give the simulation names, whether mergers were included or the main progenitor was isolated, the dark matter particle mass resolution, and the force softening length.

Name	Mergers	Resolution $(m_{\rm DM}/{\rm M}_{\odot})$	Force softening (pc)
Isolated low	×	117	10
Mergers low	1	117	10
Isolated	×	11.7	10
Mergers	\checkmark	11.7	10



Figure 8. The evolution of the 3D dark matter density at 40 pc in the GRIFFIN simulations. We start the *x*-axis at redshift z = 4 to mimic the age of the reference RAMSES simulation. The grey line shows a control GRIFFIN simulation run to z = 0 without any mergers. The black line shows the same simulation including 38 mergers. The hatched grey areas mark time intervals over which the output cadence is increased by an order of magnitude to resolve high-frequency effects. Notice that the dark matter density is lowered substantially in the simulation with mergers as compared to the control simulation. Notice, further, the three dark matter density spikes marked in red. These correspond to close subhalo passages, suggesting that the 'dark matter heating' effect seen here is driven by repeated gravitational shocks from merging subhaloes (see Fig. 10).

virial radius of the main progenitor by z = 4 are ignored, meaning that the onset of dynamical heating due to mergers may be delayed as compared to the reference simulation.

A control simulation without mergers was also run to distinguish any physical reduction in density from numerical relaxation. Our GRIFFIN simulations are summarized in Table 2.

The evolution of the central density for our isolated control (grey) and full assembly (black) GRIFFIN simulations is shown in Fig. 8. At 40 pc, there is still a small numerical heating effect in the isolated simulation (grey line). The inner density falls by $\approx 2.6 \times 10^7 \, M_\odot \, \rm kpc^{-3}$ over 12 Gyr. However, this contrasts with a much more substantial drop in the central density of the simulation with mergers of $\approx 9.1 \times 10^7 \, M_\odot \, \rm kpc^{-3}$. The small heating present in the isolated simulation is not immediately obvious in any of our non-GM RAMSES simulations, and the differences in numerical setup

and lack of cosmological growth may be contributing to this. We do not run any RAMSES simulations without cosmological accretion, so a strict comparison is difficult.

Finally, notice the three prominent 'spikes' in the inner dark matter density at \sim 4.9, \sim 6.5, and \sim 7.5 Gyr, as marked on Fig. 8 in red. These times correspond to close subhalo passages, suggesting that the 'dark matter heating' effect is driven by repeated gravitational shocks from merging subhaloes. We discuss this further in Section 5.

4.3 Cusp replenishment

Along with mechanisms that flatten the central dark matter density, there are mechanisms that can rebuild it. Laporte & Penarrubia (2015) investigate a scenario where dense mergers can reintroduce dynamically CDM into a cored parent halo, thereby rebuilding the central density cusp. These events require that the merger is able to fall into the centre of the parent halo intact, which would demand the merging structures are resistant to the tidal disruption of the parent galaxy.

In Fig. 9, we show an example of such a merger in Halo1459 GM:Latest. The left-hand panel shows the central dark matter densities of the merging system at several key times. The merging halo is shown in red at z = 0.79, by which time it has permanently crossed over the r_{200c} radius of the parent halo. The parent halo is shown at the same time in black, and has already begun to depart from a primordial density cusp (black dotted line), primarily due to the action of minor mergers as in Section 4.2. By z = 0.51, the dark grey line shows that the central density of the parent halo has continued to decline to its lowest point. Finally, the light grey line shows the parent and merging halo combined at z = 0.47, with the central density returning to a steep primordial cusp (which is coincidentally well described by the cuspy profile fit made at z = 0.79).

The right three panels show the progression of the merger event in surface density plots, with the accreted material highlighted in red. The outer regions of the merging halo have been stripped away between z = 0.79 and z = 0.51, but the central density is retained. The final panel conveys how the increase in central density at z =0.47 is correlated with the central deposition of the merger material.

5 DISCUSSION

5.1 Two mechanisms for fluctuating the gravitational potential

We have established that gas flows driven by star formation are able to drive sufficiently large potential fluctuations to erode central density cusps in our EDGE simulations (Section 4.1). However, this is seen only at early times when star formation rates exceed $1 \times 10^{-4} \, M_{\odot} \, yr^{-1}$ and fluctuate on a time-scale of order the local dynamical time. Despite the late-time rejuvenation of star formation in several of our simulations, this second phase of star formation is not sufficient to drive significant gas flows and the central dark matter density is unaffected.

However, bursty star formation is not the only means by which the central gravitational force can be varied. We have also shown that passing subhaloes act to fluctuate the central density driving dark matter heating (Section 4.2). In Fig. 10, we show an example orbit of one merging subhalo taken from our GRIFFIN simulation. Notice that this merger repeatedly punctures the inner region (0.1 kpc) of its host galaxy on a time-scale shorter than the local dynamical time (red). The first close passage of this subhalo corresponds to the dark matter density spike at 6.5 Gyr shown in Fig. 8. Shortly before this passage at ~ 6 Gyr, the mass enclosed within 0.1 kpc of the host



Figure 9. This plot illustrates how the reintroduction of denser dark matter material to the Halo1459 GM:Latest simulation erases the effect of earlier dark matter heating. *Left-most panel*: The central 3D dark matter density profiles of a cuspy subhalo (red line) and of the parent halo evolution (black, dark grey and light grey lines). The density profile of the host at z = 0.51 is centred on only the host particles to avoid any bias due to the merger. Removing the merger particles from the density profile calculation does not qualitatively change the results. A black dashed line represents a NFW profile fit to the parent halo at z = 0.79 (Navarro et al. 1997). *Right-hand panels*: The evolution of the merger system shown in three panels, where the parent halo is in grey-scale and the merging halo is red. The merging halo has permanently fallen into the central 100 pc of the parent halo by z = 0.47. In all cases, the merger system has been oriented such that the centres of both haloes are in the *xy*-plane.



Figure 10. The orbit of one example merger from our 'Mergers' GRIFFIN simulation. The orbit between each simulation output has been reconstructed with a two-body integration in AGAMA, assuming a spherically symmetric background potential and using a multipole fit to each simulation snapshot. Whilst these orbit reconstructions are imperfect due to perturbations from other subhaloes and triaxiality, they provide a reasonable estimate for our purposes. A red bar shows the width of one dynamical time $t_{dyn} = 2\pi \sqrt{3/4\pi G\rho(r)}$ over a range of orbital radii, and a black star marks the time at which the merger dissolves. The approximate inner region is indicated at 0.1 kpc with a horizontal dashed line.

halo is $7.92\times10^5\,M_\odot$. In comparison, the merging subhalo contains $6.54\times10^5\,M_\odot$ within 0.1 kpc of its centre, which is among the most massive of the mergers we model. This suggests the heating effect is greatest when the interacting masses are similar, as expected for impulsive heating (Hills 1980). The other density spikes in that figure correspond to close passages from different merging subhaloes. Taken together, this indicates that the late time dark matter heating is being driven by tidal shocks from the merging subhaloes on their host.

However, in the DMO simulations the inner dark matter density, while lower, remains cuspy (Fig. 7, bottom panel). By contrast, in the baryonic simulation – and in our GRIFFIN replica of this simulation – these minor mergers flatten the cusp. This occurs because in these simulations, the cusp is already weakened at early times by dark matter heating due to star formation (Fig. 7, upper panel). Previous literature has suggested that minor mergers can 'heat up' a host dark matter profile. Naab, Johansson & Ostriker (2009) propose a mechanism by which the central concentrations of massive

The above late time dark matter heating due to minor mergers

occurs in both our baryonic and DMO RAMSES simulations (Fig. 7).

propose a mechanism by which the central concentrations of massive elliptical galaxies are reduced through repeated minor mergers, with Bédorf & Portegies Zwart (2013) finding the same for massive compact galaxies. Similarly, Boldrini et al. (2020a) are able to enact a cusp-core transition in a model of M31 via the accretion of both minor and major mergers on highly eccentric orbits. Towards the scale of dwarf galaxies, Macciò et al. (2017) show that minor mergers can boost star formation, exciting dark matter heating via the usual gas-flow mechanism. This effect cannot be important for ultra-faint dwarfs, since they do not show evidence for late-time star formation. However, Boldrini, Mohayaee & Silk (2020b) find that minor mergers can directly heat the central dark matter density in a model of the Fornax dwarf via dynamical friction. Finally, Leung et al. (2020) propose that mergers could expand the orbits of globular clusters in the Fornax dwarf spheroidal galaxy, solving a long-standing puzzle as to why they have not sunk to the centre of Fornax via dynamical friction. This same mechanism would also expand the orbits of the dark matter particles.

6 CONCLUSIONS

We have presented a suite of cosmological zoom simulations of the ultra-faint dwarf galaxies performed with the AMR code RAMSES as part of the EDGE project. These simulations have a spatial and mass resolution of 3 pc and 120 M_{\odot} , respectively, sufficient to resolve the formation of very small dark matter cores.

Our key result is that we uncover two distinct pathways to dark matter core formation at sub-kpc scales in the $10^9 < M_{200c}/M_{\odot} < 5 \times 10^9$ halo mass regime. These are able to drive reductions in the central (40 pc) dark matter density of up to approximately a factor of

However, even after quenching by reionization, we found that a second mechanism can cause dark matter cores to continue to grow: impulsive heating from minor mergers. To demonstrate this, we 'genetically modified' the initial conditions for one dwarf such that it assembled later from many minor mergers. We found that, in this case, the dwarf's inner dark matter density continued to drop long after star formation ceased. We tested the veracity of this result by running an independent 'replica' simulation using the GRIFFIN *N*-body code, finding excellent agreement between the GRIFFIN and RAMSES calculations.

While all of our dwarfs experienced some dark matter heating prior to reionization, we showed that dense major mergers can replenish kinematically CDM, thereby reintroducing a density cusp at late times. This demonstrates that the central density of the smallest dwarf galaxies at z = 0 is sensitive to both their mass assembly histories and their star formation histories. This will drive stochasticity in the central dark matter densities of isolated ultra-faint dwarfs. We will study this in more detail in future work.

Finally, none of our simulated EDGE dwarfs experienced sufficient dark matter heating to produce a flat density core. In a companion paper, we will consider whether the dark matter heating we do find in EDGE is sufficient to explain the survival and properties of the lone star cluster in Eridanus II.

ACKNOWLEDGEMENTS

The author acknowledges the UK Research and Innovation (UKRI) Science and Technology Facilities Council (STFC) for support (grant ST/R505134/1). This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement no. 818085 GMGalaxies. AP was further supported by the Royal Society. OA and MPR acknowledge support from the Knut and Alice Wallenberg Foundation and the Swedish Research Council (grants 2014-5791 and 2019-04659).

MD acknowledges support by ERC-Syg 810218 WHOLE SUN. This work was performed using the Distributed Research utilising Advanced Computing (DiRAC) Data Intensive service at Leicester, operated by the University of Leicester IT Services, which forms part of the STFC DiRAC HPC Facility (www.dirac.ac.uk). The equipment was funded by Business, Energy & Industrial Strategy (BEIS) capital funding via STFC capital grants ST/K000373/1 and ST/R002363/1 and STFC DiRAC Operations grant ST/R001014/1. DiRAC is part of the National e-Infrastructure. We thank the anonymous referee for their review and comments.

DATA AVAILABILITY

Data available upon request.

REFERENCES

Agertz O., Kravtsov A. V., 2015, ApJ, 804, 18 Agertz O., Kravtsov A. V., Leitner S. N., Gnedin N. Y., 2013, ApJ, 770, 25 Agertz O. et al., 2020, MNRAS, 491, 1656

- Amorisco N. C., 2017, ApJ, 844, 64
- Avila-Reese V., Colín P., Valenzuela O., D'Onghia E., Firmani C., 2001, ApJ, 559, 516
- Bédorf J., Portegies Zwart S., 2013, MNRAS, 431, 767
- Binney J., Tremaine S., 1987, Galactic Dynamics. Princeton University Press, Princeton N.J.
- Bode P., Ostriker J. P., Turok N., 2001, ApJ, 556, 93
- Boldrini P., Mohayaee R., Silk J., 2020a, preprint (arXiv:2002.12192)
- Boldrini P., Mohayaee R., Silk J., 2020b, MNRAS, 492, 3169
- Bose S. et al., 2019, MNRAS, 486, 4790
- Bullock J. S., Boylan-Kolchin M., 2017, ARA&A, 55, 343
- Carignan C., Freeman K. C., 1988, ApJ, 332, L33
- Chabrier G., 2003, PASP, 115, 763
- Chan T. K., Kereš D., Oñorbe J., Hopkins P. F., Muratov A. L., Faucher-Giguère C. A., Quataert E., 2015, MNRAS, 454, 2981
- Clowe D., Bradač M., Gonzalez A. H., Markevitch M., Randall S. W., Jones C., Zaritsky D., 2006, ApJ, 648, L109
- Cole D. R., Dehnen W., Wilkinson M. I., 2011, MNRAS, 416, 1118
- Contenta F. et al., 2018, MNRAS, 476, 3124
- Dawson K. S. et al., 2013, AJ, 145, 10
- Dehnen W., 2014, Comput. Astrophys. Cosmol., 1, 1
- Dehnen W., Read J. I., 2011, Eur. Phys. J. Plus, 126, 55
- Del Popolo A., Pace F., 2016, Ap&SS, 361, 162
- Di Cintio A., Brook C. B., Dutton A. A., Macciò A. V., Stinson G. S., Knebe A., 2014, MNRAS, 441, 2986
- Diemand J., Moore B., Stadel J., Kazantzidis S., 2004a, MNRAS, 348, 977
- Diemand J., Moore B., Stadel J., 2004b, MNRAS, 353, 624
- Dutton A. A. et al., 2016, MNRAS, 461, 2658
- Eisenstein D. J., Hut P., 1998, ApJ, 498, 137
- El-Badry K., Wetzel A., Geha M., Hopkins P. F., Kereš D., Chan T. K., Faucher-Giguère C.-A., 2016, ApJ, 820, 131
- El-Zant A., Shlosman I., Hoffman Y., 2001, ApJ, 560, 636
- Ferreira E. G. M., 2020, preprint (arXiv:2005.03254)
- Flores R. A., Primack J. R., 1994, ApJ, 427, L1
- Garrison-Kimmel S., Rocha M., Boylan-Kolchin M., Bullock J. S., Lally J., 2013, MNRAS, 433, 3539
- Genina A., Read J. I., Fattahi A., Frenk C. S., 2020, preprint (arXiv:2011.09482)
- Goerdt T., Moore B., Read J. I., Stadel J., 2010, ApJ, 725, 1707
- Grisdale K., Agertz O., Renaud F., Romeo A. B., Devriendt J., Slyz A., 2019, MNRAS, 486, 5482
- Gualandris A., Read J. I., Dehnen W., Bortolas E., 2017, MNRAS, 464, 2301
- Haardt F., Madau P., 1996, ApJ, 461, 20
- Hills J. G., 1980, ApJ, 235, 986
- Hirtenstein J. et al., 2019, ApJ, 880, 54
- Hogan C. J., Dalcanton J. J., 2000, Phys. Rev. D, 62, 063511
- Katz N., White S. D. M., 1993, ApJ, 412, 455
- Kauffmann G., 2014, MNRAS, 441, 2717
- Keating L. C., Weinberger L. H., Kulkarni G., Haehnelt M. G., Chardin J., Aubert D., 2020, MNRAS, 491, 1736
- Kennicutt R. C., Jr, 1998, ApJ, 498, 541
- Kimm T., Cen R., Devriendt J., Dubois Y., Slyz A., 2015, MNRAS, 451, 2900
- Knollmann S. R., Knebe A., 2009, ApJS, 182, 608
- Kuzio de Naray R., Kaufmann T., 2011, MNRAS, 414, 3617
- Laporte C. F. P., Penarrubia J., 2015, MNRAS, 449, L90
- Leung G. Y. C., Leaman R., van de Ven G., Battaglia G., 2020, MNRAS, 493, 320
- Lux H., Read J. I., Lake G., 2010, MNRAS, 406, 2312
- Macciò A. V., Frings J., Buck T., Penzo C., Dutton A. A., Blank M., Obreja A., 2017, MNRAS, 472, 2356
- McGaugh S. S., Rubin V. C., de Blok W. J. G., 2001, AJ, 122, 2381
- Madau P., Shen S., Governato F., 2014, ApJ, 789, L17
- Malhan K., Valluri M., Freese K., 2020, MNRAS, 501, 179
- Mashchenko S., Wadsley J., Couchman H. M. P., 2008, Science, 319, 174
- Maxwell A. J., Wadsley J., Couchman H. M. P., 2015, ApJ, 806, 229
- Mo H. J., Mao S., 2004, MNRAS, 353, 829
- Moore B., 1994, Nature, 370, 629

- Munshi F. et al., 2013, ApJ, 766, 56
- Naab T., Johansson P. H., Ostriker J. P., 2009, ApJ, 699, L178
- Nasim I., Gualandris A., Read J., Dehnen W., Delorme M., Antonini F., 2020, MNRAS, 497, 739
- Nasim I. T., Gualandris A., Read J. I., Antonini F., Dehnen W., Delorme M., 2021, MNRAS, 502, 4794
- Navarro J. F., Eke V. R., Frenk C. S., 1996, MNRAS, 283, L72
- Navarro J. F., Frenk C. S., White S. D. M., 1997, ApJ, 490, 493
- Nipoti C., Binney J., 2015, MNRAS, 446, 1820
- Oka A., Saito S., Nishimichi T., Taruya A., Yamamoto K., 2014, MNRAS, 439, 2515
- Oman K. A., Navarro J. F., Sales L. V., Fattahi A., Frenk C. S., Sawala T., Schaller M., White S. D. M., 2016, MNRAS, 460, 3610
- Oman K. A., Marasco A., Navarro J. F., Frenk C. S., Schaye J., Benítez-Llambay A. R., 2019, MNRAS, 482, 821
- Oñorbe J., Garrison-Kimmel S., Maller A. H., Bullock J. S., Rocha M., Hahn O., 2014, MNRAS, 437, 1894
- Oñorbe J., Boylan-Kolchin M., Bullock J. S., Hopkins P. F., Kereš D., Faucher-Giguère C.-A., Quataert E., Murray N., 2015, MNRAS, 454, 2092
- Peñarrubia J., Pontzen A., Walker M. G., Koposov S. E., 2012, ApJ, 759, L42
- Planck Collaboration XVI, 2014, A&A, 571, A16
- Pontzen A., Governato F., 2012, MNRAS, 421, 3464
- Pontzen A., Governato F., 2014, Nature, 506, 171
- Pontzen A., Tremmel M., 2018, ApJS, 237, 23
- Pontzen A., Roškar R., Stinson G. S., Woods R., Reed D. M., Coles J., Quinn T. R., 2013, Astrophysics Source Code Library, record ascl:1305
- Pontzen A., Rey M. P., Cadiou C., Agertz O., Teyssier R., Read J., Orkney M. D. A., 2020, MNRAS, 501, 1755
- Power C., Navarro J. F., Jenkins A., Frenk C. S., White S. D. M., Springel V., Stadel J., Quinn T., 2003, MNRAS, 338, 14
- Read J. I., Gilmore G., 2005, MNRAS, 356, 107
- Read J. I., Lake G., Agertz O., Debattista V. P., 2008, MNRAS, 389, 1041
- Read J. I., Agertz O., Collins M. L. M., 2016a, MNRAS, 459, 2573
- Read J. I., Iorio G., Agertz O., Fraternali F., 2016b, MNRAS, 462, 3628
- Read J. I., Iorio G., Agertz O., Fraternali F., 2017, MNRAS, 467, 2019
- Read J. I., Walker M. G., Steger P., 2019, MNRAS, 484, 1401
- Rey M. P., Pontzen A., 2018, MNRAS, 474, 45
- Rey M. P., Pontzen A., Agertz O., Orkney M. D. A., Read J. I., Saintonge A., Pedersen C., 2019, ApJ, 886, L3
- Rey M. P., Pontzen A., Agertz O., Orkney M. D. A., Read J. I., Rosdahl J., 2020, MNRAS, 497, 1508
- Rodriguez-Gomez V. et al., 2015, MNRAS, 449, 49
- Romano-Díaz E., Shlosman I., Heller C., Hoffman Y., 2009, ApJ, 702, 1250
- Roth N., Pontzen A., Peiris H. V., 2016, MNRAS, 455, 974
- Sanders J. L., Evans N. W., Dehnen W., 2018, MNRAS, 478, 3879
- Schive H.-Y., Chiueh T., Broadhurst T., 2014, Nat. Phys., 10, 496
- Schmidt M., 1959, ApJ, 129, 243
- Simon J. D., 2019, ARA&A, 57, 375
- Simon J. D. et al., 2021, ApJ, 908, 18
- Sparre M., Hayward C. C., Feldmann R., Faucher-Giguère C.-A., Muratov A. L., Kereš D., Hopkins P. F., 2017, MNRAS, 466, 88
- Spergel D. N., Steinhardt P. J., 2000, Phys. Rev. Lett., 84, 3760
- Springel V., Frenk C. S., White S. D. M., 2006, Nature, 440, 1137
- Stewart K. R., Bullock J. S., Barton E. J., Wechsler R. H., 2009, ApJ, 702, 1005
- Stopyra S., Pontzen A., Peiris H., Roth N., Rey M. P., 2021, ApJS, 252, 28
- Tegmark M. et al., 2006, Phys. Rev. D, 74, 123507
- Teyssier R., 2002, A&A, 385, 337
- Teyssier R., Pontzen A., Dubois Y., Read J. I., 2013, MNRAS, 429, 3068
- Tollet E. et al., 2016, MNRAS, 456, 3542
- Tulin S., Yu H.-B., 2018, Phys. Rep., 730, 1
- Vasiliev E., 2018, AGAMA: Action-Based Galaxy Modeling Framework, Astrophysics Source Code Library (record ascl:1805.008)
- Wang W., White S. D. M., Mandelbaum R., Henriques B., Anderson M. E., Han J., 2016, MNRAS, 456, 2301
- Weinberg M. D., Katz N., 2007, MNRAS, 375, 460
- White S. D. M., Rees M. J., 1978, MNRAS, 183, 341

APPENDIX A: DERIVING THE RELAXATION RADIUS FOR OUR EDGE SIMULATIONS

The dark matter 'particles' in collisionless cosmological simulations like RAMSES are really 'superparticles' that represent unresolved patches of the dark matter fluid. This approximation leads to overly large two-body relaxation that causes an artificial reduction in the inner dark matter density of a halo over time (e.g. Power et al. 2003; Diemand et al. 2004a; Diemand, Moore & Stadel 2004b; Dehnen & Read 2011).

The two-body relaxation time-scale is given by Binney & Tremaine (1987):

$$t_{\rm relax} = \frac{N t_{\rm orb}}{16\pi \log \Lambda},\tag{A1}$$

where *N* is the number of particles within some radius *r*, t_{orb} is the orbital time of the system given by $t_{orb} = 2\pi \sqrt{R^3/GM}$, and Λ is the 'Coulomb logarithm' given by $\Lambda = b_{max}/b_{min}$. The maximum impact parameter b_{max} and the minimum impact parameter b_{min} define the largest and smallest scales at which particles are expected to interact with each other. The relaxation time gives the time taken for the particle velocities to change by 90°, and can be considered as the time taken for a system to lose dynamical memory of its initial conditions. A large relaxation time is preferred because this implies two-body relaxation has a minimal influence on the particle kinematics. For standard *N*-body methods, the simplest way to increase the relaxation time is by increasing the number of dark matter particles, *N*, that sample the system (e.g. Dehnen & Read 2011).

There is some debate in the literature over the best choices for the impact parameters b_{max} and b_{min} . Here, we define b_{max} to be the total virial size of the system r_{200} , and b_{min} to be the side length of the highest resolution grid cell in RAMSES. This is the approximate distance above which Newtonian gravity is recovered. Although multiple resolutions of grid cell are used throughout our simulated haloes, the central regions that are of interest here are predominantly at the highest resolution.

We now perform a brief resolution study in order to determine at what radius numerical relaxation effects become important. In Fig. A1, we plot dark matter density profiles from a DMO simulation at three different resolutions. These three resolutions each exhibit a different amount of central density flattening due to numerical relaxation, with the flattening becoming stronger with increasing particle mass and with time. At z = 0, all three resolutions are converged beyond ≈ 300 pc, but begin to deviate from the expected NFW form within some critical radius.

We define the 'relaxation radius', r_{relax} , to be the radius at which the relaxation time (equation A1) for the enclosed dark matter particles is equal to the simulation age for any particular simulation output:

$$\left(\frac{r_{\rm relax}}{\rm kpc}\right) = \eta \left(\frac{t_{\rm sim}}{\rm Gyr}\right)^{\alpha} \left(\frac{M(< r_{\rm relax})}{\rm M_{\odot}}\right)^{-\frac{1}{3}} \left(\frac{\langle m \rangle}{\rm M_{\odot}}\right)^{\frac{2}{3}},\tag{A2}$$

where we have substituted t_{relax} for the total run time of the simulation, $\langle m \rangle$ is the mean particle mass and η and α are fitting parameters. From the data in Fig. A1, we find $\alpha = 1/3$ and $\eta = 64G\log \Lambda/100^2$. The mass within the relaxation radius, $M(r_{relax})$, is calculated directly from our simulation data and so equation (A2) can be solved numerically to find r_{relax} . This is then used to predict the resolution limit for our EDGE simulations in this paper. The relaxation radii, calculated in this way, are marked in Fig. A1 by the vertical lines. Notice that the dark matter density profiles are shallower leftwards of r_{relax} in the lower resolution simulations as compared to the higher resolution simulations.



Figure A1. The evolution of the 3D density profiles for Halo1459 DMO at three dark matter particle resolutions, as stated in the legend. The two lowest particle masses (green and cyan lines) correspond to the high resolution and fiducial resolutions, respectively (see Table 1). The fainter lines indicate the radii within which at least one relaxation time (estimated using equation A1) has passed, with the transition marked using an additional vertical line for clarity. In some cases, the density profiles are not plotted at smaller radii because there are insufficient particles.

APPENDIX B: SIMULATION CONVERGENCE

The dark matter particle mass resolution used in this paper is an order of magnitude smaller than used in the fiducial EDGE simulations (Rey et al. 2019; Agertz et al. 2020; Pontzen et al. 2020; Rey et al. 2020). In this appendix, we perform a convergence study to determine how our results are impacted by dark matter resolution. As previously in this paper, we distinguish between the lower and higher resolution simulations by appending the labels 'fiducial' and 'hires', respectively, to their name.

In Fig. B1, we plot the total stellar mass within r_{200c} versus M_{200c} as a function of time for the fiducial (faint dashed) and hires (opaque solid) EDGE simulations. We find good convergence in shape of the general evolution of each simulated dwarf. The higher resolution simulations form systematically more stars (as first noted in Agertz et al. 2020). However, the final stellar masses typically agree within ~30 per cent and at worst differ by a factor of ~2 (for Halo600). This is within the expected uncertainties due to modelling galaxy formation (Agertz et al. 2020).

In Fig. B2, we compare the 3D dark matter density profiles for our main simulation suite at two redshifts, z = 4 (top panels) and z = 0 (bottom panels). The DMO simulations are marked by the faint lines, the baryonic simulations by the opaque lines. The fiducial and hires simulations are marked by dashed and solid lines, respectively.

Overall, there is good convergence between the fiducial and hires simulations. The DMO simulations are all converged for $r > r_{relax}$ (vertical lines), and there is good agreement also for the baryonic simulations Halo1445, Halo1459, and Halo624, at both z = 4 and z = 0. However, two of our EDGE simulations show poor convergence. Halo600 fiducial substantially rejuvenates its star formation after z = 1, forming 17 per cent of its final stellar mass after this time. This extended period of star formation is replaced by a singular burst in Halo600 hires. As a result, Halo600 fiducial undergoes significantly more dark matter heating that causes its final inner density profile to be substantially lower than Halo600 hires. Similarly, Halo605 fiducial forms a larger dark matter core than Halo605 hires at early times. However, this disparity is largely resolved before z = 0 due to a late cuspy merger in Halo605 fiducial (see Section 4.3 for a discussion of this cusp reintroduction mechanism).



Figure B1. The stellar mass–halo mass relation through time for our main simulation suite, where we compare our fiducial resolution simulations (faint dashed lines) to our high-resolution simulations (solid lines). The black circles on the high-resolution lines mark intervals of 1 Gyr in time. The grey diagonal lines show constant ratios of M_*/M_{200c} in powers of 10, as marked.

The above results highlight an important point. It is often stated in the literature that whether a galaxy will be cusped or cored is determined by the stellar mass to halo mass ratio, M_*/M_{200c} (e.g. Peñarrubia et al. 2012; Di Cintio et al. 2014; Read et al. 2019). This is true to leading order. However, it also matter *how* those stars form, as illustrated by Halo600 fiducial versus Halo600 hires. The latter actually forms more stars, but because at late time they form in a single burst, this leads to less dark matter heating. And, at least at the very edge of galaxy formation, it also matters what the merger history is. Halo605 fiducial goes on to partially lose its dark matter core due to a late cuspy merger.



Figure B2. The 3D dark matter density profiles of our main simulation suite, where we compare our DMO simulations (faint lines) and baryonic simulations (opaque lines) at fiducial (dashed lines) and high (solid lines) resolution. The upper panels show this comparison at z = 4, by which time all haloes are quenched due to reionization (some will later reignite their star formation). The lower panels show this comparison at z = 0. The numerical relaxation radius, as defined in Appendix A, is indicated by a short vertical line for the baryonic simulations. This limit is similar for the corresponding DMO simulations.

This paper has been typeset from a $T_EX/I \ge T_EX$ file prepared by the author.