HAL
open science

A convex function satisfying the Lojasiewicz inequality but failing the gradient conjecture both at zero and infinity

Aris Daniilidis, Mounir Haddou, Olivier Ley

To cite this version:

Aris Daniilidis, Mounir Haddou, Olivier Ley. A convex function satisfying the Lojasiewicz inequality but failing the gradient conjecture both at zero and infinity. 2021. hal-03136550v1

HAL Id: hal-03136550
https://hal.science/hal-03136550v1

Preprint submitted on 9 Feb 2021 (v1), last revised 31 Aug 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A convex function satisfying the Łojasiewicz inequality but failing the gradient conjecture both at zero and infinity.

Aris Daniilidis, Mounir Haddou, Olivier Ley

Abstract. We construct an example of a smooth convex function on the plane with a strict minimum at zero, which is real analytic except at zero, for which Thom's gradient conjecture fails both at zero and infinity. More precisely, the gradient orbits of the function spiral around zero and at infinity. Besides, the function satisfies the Łojasiewicz gradient inequality at zero.

Key words. Gradient conjecture, gradient conjecture at infinity, Kurdyka-Łojasiewicz inequality, convex function, convergence of secants.

AMS Subject Classification *Primary* 37C10 ; *Secondary* 34A26, 34C08, 52A41.

1 Introduction

Answering a question of Whitney, Łojasiewicz [19] showed that every analytic variety $f^{-1}(0)$, where $f : \mathcal{U} \subset \mathbb{R}^N \to \mathbb{R}$ is real-analytic ($\mathcal{U} \neq \emptyset$, open), is a deformation retract of its open neighborhood. The deformation was given by the flow of the Euclidean gradient ∇f. The main argument of Łojasiewicz was based on a famous lemma, nowadays known as the Łojasiewicz (gradient) inequality, which asserts that for some $\vartheta \in (0,1)$ and $c > 0$ we have

$$\|\nabla f(x)\| \geq c|f(x) - f(a)|^{\vartheta} \tag{1.1}$$

for all x sufficiently close to $a \in f^{-1}(0)$. The above inequality ensures that every bounded gradient orbit $t \mapsto \gamma(t)$ (*i.e.*, $\dot{\gamma} = \nabla f(\gamma)$) has finite length and therefore converges to a singular point γ_∞ with $\nabla f(\gamma_\infty) = 0$.

Some years later, Thom conjectured that in this case, up to a change of coordinates that identifies γ_∞ to 0, the spherical part of the orbit also converges. In other words, the limit of secants

$$\lim_{t \to +\infty} \frac{\gamma(t) - \gamma_\infty}{||\gamma(t) - \gamma_\infty||} \quad \text{exists.} \tag{1.2}$$

For decades, this has been known as the (Thom) gradient conjecture, see [1, 29]. (For the more general problem of non-oscillation of trajectories, we refer to [4, 12, 24].) The gradient conjecture makes sense for any gradient dynamics for which bounded orbits converge. Partial results revealed that (1.2) should hold in the real-analytic case, see [13, 18, 27], fact that was eventually published in full generality by Kurdyka, Mostowski and Parusiński [16] in 2000. The proof was based on (1.1) together with concrete analytic estimations.

Łojasiewicz showed that the gradient inequality (1.1) remains valid also for $\mathcal{C}^1$ semialgebraic (respectively, globally subabalytic) functions, see [20]. In 1998, Kurdyka generalized (1.1) for $\mathcal{C}^1$ functions that are *definable* in some *o-minimal structure*, an axiomatic definition due to van den Dries [30, 31] which encompasses semialgebraic and globally subanalytic functions, but also larger classes that include the exponential function [23]. More precisely, Kurdyka showed that

for every definable function f and critical value r_∞ (which is necessarily isolated) there exists $\delta > 0$ and a continuous function $\Psi : [r_\infty, r_\infty + \delta) \to \mathbb{R}$ which is $\mathcal{C}^1$ on $(r_\infty, r_\infty + \delta)$ with $\Psi' > 0$ such that

$$||\nabla(\Psi \circ f)(x)|| \geq 1 \tag{1.3}$$

for all $x \in \mathbb{R}^N$ such that $r_\infty < f(x) < r_\infty + \delta$. In addition, Kurdyka's proof showed that the function Ψ can be taken in the same o-minimal structure as f. Consequently, if f is semialgebraic or globally subanalytic, then so is Ψ and thanks to Puiseux's theorem we may take $\Psi(r) = r^{1-\vartheta}$, for $\vartheta \in (0, 1)$. It is then straightforward to see that (1.3) actually yields (1.1) for $c = (1 - \vartheta)^{-1}$.

We refer to (1.3) as the Kurdyka-Łojasiewicz (in short, KŁ) inequality and we call KŁ-function any function with (upper) isolated critical values that satisfies the KŁ-inequality around any of them. Similarly to the gradient inequality (1.1), bounded gradient orbits of a KŁ-function have finite length. There are well-known examples of $\mathcal{C}^\infty$ functions in $\mathbb{R}^2$ with isolated critical values that are not KŁ-functions (they have bounded gradient orbits which fail to converge), see [10, 25]. Bounded gradient orbits of convex functions have finite length [7, 22] and therefore converge, but there are also examples of $\mathcal{C}^2$-smooth convex functions failing KŁ-property, see [3, §4.3] or [2, §5.1]. In [3] we characterized the class of KŁ-functions (among the ones with upper isolated critical values) and gave criteria for a convex function to be KŁ.

In [17], Kurdyka and Parusinski used KŁ-inequality together with a quasiconvex cell decomposition of o-minimal sets and concrete estimates to show that the gradient conjecture holds for $\mathcal{C}^1$ o-minimal functions provided either $N = 2$ (planar case) or the structure is *polynomially bounded* (in particular if f is semialgebraic or globally subanalytic). On the other hand, mere convexity is not sufficient to guarantee (1.2): there exist examples of convex functions whose orbits either spiral [8, §7.2] or oscillate between two secants [2].

In [11], Grandjean considered the behavior of the secants at infinity: he showed that if f is a $\mathcal{C}^1$ semialgebraic function and $t \mapsto \gamma(t)$ is a gradient orbit satisfying $||\gamma(t)|| \to \infty$, as $t \to +\infty$, then the limit of secants at infinity

$$\lim_{t \to +\infty} \frac{\gamma(t)}{||\gamma(t)||} \quad \text{exists} \qquad \text{(gradient conjecture at infinity).} \tag{1.4}$$

The proof is based on a Łojasiewicz type gradient inequality at infinity previously obtained by the author together with D'Acunto in [6].

The behavior of secants at infinity has recently become relevant in Machine Learning. If a deep network model is unbiased and homogeneous (max-pooling, ReLu, linear and convolutional layers), then minimizing the cross-entropy or other classification losses forces the parameters of the model to diverge in norm to infinity [21]. In this setting, convergence of the secants at infinity is important. In [14] the authors manage to establish that for a certain type of prediction functions (L-homogeneous and definable in the log-exp structure) (1.4) holds. For the time being, no further results have been reported.

In a nutshell, proving the gradient conjecture (respectively, the gradient conjecture at infinity) seems to require at least the KŁ-inequality (1.3) together with other properties of o-minimal functions, but it is still unknown if these conjectures are true for general o-minimal functions.

In this work we present an example of a smooth convex function in $\mathbb{R}^2$, which is real-analytic outside zero (its unique critical point), it satisfies the Łojasiewicz inequality (1.1) and fails the gradient conjecture both at zero and at infinity. In particular, all gradient orbits spiral both

at zero and at infinity, underlying in this way the two failures of o-minimality of the function, despite the fact that the function is convex and satisfies the Łojasiewicz gradient inequality.

Theorem 1.1 (main result)**.** *For every $k \in \mathbb{N}$, there exists a $\mathcal{C}^k$-convex function $f : \mathbb{R}^2 \to \mathbb{R}$ with a unique minimum at $\mathcal{O} := (0,0)$ such that:*

- *f is real analytic on $\mathbb{R}^2 \setminus \{\mathcal{O}\}$;*
- *f satisfies the Łojasiewicz inequality at $\mathcal{O}$ and*
- *every gradient orbit $\gamma : (-\infty, T) \to \mathbb{R}^2$ of f spirals infinitely many times both when $t \to T$ and $t \to -\infty$.*

Throughout the manuscript, by gradient orbits (or gradient trajectories) we refer to maximal solutions of the ordinary differential equation:

$$\gamma'(t) = \nabla f(\gamma(t)).$$

In our example, the function f will be convex, with unique critical point (global minimizer) at $\mathcal{O}$, where we tacitly assume that $\gamma(0) \neq \mathcal{O}$ (avoiding stationary orbits).

Let us briefly describe our strategy for the construction of this example: in Section 2 we prescribe a family of convex sets, all being delimited by ellipses, centered at the origin, and obtained via rotations and size adjustments of a basic ellipse $E(0)$. This is done in a way that convex foliation is obtained, which can be represented by some (quasiconvex) function.

In Section 3, we further calibrate the parameters so that we can apply a criterium due to de Finetti [9] and Crouzeix [5] that guarantees that the aforementioned quasiconvex function is in fact convex. The construction yields that the function is real-analytic on $\mathbb{R}^2 \setminus \mathcal{O}$, which of course cannot be further improved to real analycity on the whole space, due to the proof of Thom's gradient conjecture [16]. Instead, we are able to show that the function can be taken $\mathcal{C}^k$-smooth at $\mathcal{O}$ for arbitrary large $k \in \mathbb{N}$. Still our construction fails to ensure $\mathcal{C}^\infty$. Finally, applying a result of [3] which gives conditions for a convex function to satisfy (1.3), we show that our function satisfies KŁ-inequality and in fact even (1.1) (the Łojasiewicz inequality).

Gradient orbits are perpendicular to the foliation and explicit calculations, conducted in Section 4, show that the orbits turn around both at the origin and at infinity, which disproves the conjecture. An additional difficulty to establish spirality is that the evolution of the spherical part of the orbit (the rotation angle $\alpha(t)$ of $\gamma(t)$ in polar coordinates) is not monotone in time, so that the decrease rate is established in average, see Figure 3 and Figure 4. For a study of monotonic spiraling of orbits of general analytic vector fields in dimensions 2 and 3, we refer to [28].

2 Construction of a convex real analytic foliation in $\mathbb{R}^2 \setminus \{\mathcal{O}\}$.

Let us first consider two smooth increasing functions $a, b : \mathbb{R} \to (0, +\infty)$ for which we assume:

$$\begin{cases} \lim\limits_{t\to+\infty} a(t) = \lim\limits_{t\to+\infty} b(t) = +\infty \\ \lim\limits_{t\to-\infty} a(t) = \lim\limits_{t\to-\infty} b(t) = 0 \quad \text{and} \\ \qquad a(t) \geq b(t), \quad \text{for all } t \in \mathbb{R}. \end{cases} \tag{2.1}$$

The exact definition of the functions $a(t)$ and $b(t)$ will be given in Lemma 3.1 (Section 3). We also consider the rotation matrix by an angle t denoted by:

$$R(t) = \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix} \tag{2.2}$$

For $t \in \mathbb{R}$ and $\theta \in \mathbb{T} := \mathbb{R}/2\pi\mathbb{Z}$ we set

$$m(t,\theta) := (x(t,\theta),\, y(t,\theta)) = (a(t)\cos\theta,\, b(t)\sin\theta),$$

and

$$M(t,\theta) := R(t)\, m(t,\theta) \;=\; (X(t,\theta), Y(t,\theta)). \tag{2.3}$$

Therefore

$$\begin{cases} X(t,\theta) = x(t,\theta)\cos t - y(t,\theta)\sin t \;=\; a(t)\cos t\cos\theta - b(t)\sin t\sin\theta \\ Y(t,\theta) = \; x(t,\theta)\sin t + y(t,\theta)\cos t \;=\; a(t)\sin t\cos\theta + b(t)\cos t\sin\theta\,. \end{cases} \tag{2.4}$$

The subset

$$\mathcal{E}(t) := \{M(t,\theta) : \theta \in \mathbb{T}\} \tag{2.5}$$

is an ellipse with major axis of length $a(t)$ and minor axis of length $b(t)$ (see Figure 1 for illustration). Notice that $\mathcal{E}(t)$ is the rotation by angle t of the ellipse

$$E(t) := \big\{m(t,\theta) :\, \theta \in \mathbb{T}\big\} \;=\; \left\{ (x,y) \in \mathbb{R}^2 :\; \frac{x^2}{a^2(t)} + \frac{y^2}{b^2(t)} = 1 \right\}.$$

Under an additional condition on the functions a, b, the family of ellipses $\{\mathcal{E}(t)\}_{t\in\mathbb{R}}$ defined in (2.5) is disjoint with union equal to $\mathbb{R}^2 \setminus \{\mathcal{O}\}$. More precisely, denoting by a', b' the derivatives of the functions a, b respectively, we have the following result:

Lemma 2.1 (Convex foliation by ellipses)**.** *Let $a, b : \mathbb{R} \to (0, +\infty)$ satisfy* (2.1) *and assume*

$$4\, a(t)\, b(t)\, a'(t)\, b'(t) \;>\; (a(t)^2 - b(t)^2)^2\,, \qquad \textit{for all } \; t \in \mathbb{R}. \tag{2.6}$$

Then $(\mathcal{E}(t))_{t\in\mathbb{R}}$ defines an analytic convex foliation of $\mathbb{R}^2 \setminus \{\mathcal{O}\}$.

Proof. The proof is divided in three steps:

Step 1. The map $M : \mathbb{R} \times \mathbb{T} \to \mathbb{R}^2 \setminus \{\mathcal{O}\}$ is a local analytic diffeomorphism.

Indeed, let us first notice that the map M, defined by (2.3)–(2.4), is real-analytic as composition of analytic functions. Therefore, if we show that the Jacobian matrix $\mathcal{J}M = \begin{pmatrix} \frac{\partial X}{\partial t} & \frac{\partial X}{\partial \theta} \\ \frac{\partial Y}{\partial t} & \frac{\partial Y}{\partial \theta} \end{pmatrix}$ is invertible at each point $(t,\theta) \in \mathbb{R} \times \mathbb{T}$, the assertion follows from the local analytic inverse function theorem [15, Theorem 2.5.1]. To this end, we shall prove that

$$\det(\mathcal{J}M) = \frac{\partial X}{\partial t}\frac{\partial Y}{\partial \theta} - \frac{\partial Y}{\partial t}\frac{\partial X}{\partial \theta} = \Big\langle \frac{\partial M}{\partial t}, n \Big\rangle > 0, \tag{2.7}$$



Figure 1: The ellipse $\mathcal{E}(t)$ and the map $(t,\theta) \mapsto M(t,\theta)$

where $n(t,\theta) = -R(\frac{\pi}{2})\,\frac{\partial M}{\partial \theta} = (\frac{\partial Y}{\partial \theta}, -\frac{\partial X}{\partial \theta})$ is the outer unit normal to the convex set $\operatorname{conv}\mathcal{E}(t)$ (convex envelope of $\mathcal{E}(t)$) at $M(t,\theta)$. Recalling that $M(t,\theta) = R(t)\,m(t,\theta)$ (see (2.3)) and that the rotation matrix (2.2) satisfies

$$R'(t) = R(t+\frac{\pi}{2}), \quad R(t)^{-1} = R(t)^T = R(-t) \quad \text{and} \quad R(t)\,R(s) = R(t+s),$$

we deduce

$$\begin{aligned}
\langle \frac{\partial M}{\partial t}, n\rangle &= \langle \frac{\partial}{\partial t}(R(t)m), -R(\frac{\pi}{2})\,\frac{\partial}{\partial \theta}(R(t)m)\,\rangle \\
&= \langle R'(t)m + R(t)\frac{\partial m}{\partial t},\, -R(\frac{\pi}{2})\,R(t)\,\frac{\partial m}{\partial \theta}\rangle \\
&= \langle R(t+\frac{\pi}{2})\,m + R(t)\frac{\partial m}{\partial t},\, R(t-\frac{\pi}{2})\,\frac{\partial m}{\partial \theta}\rangle \\
&= \langle R(t-\frac{\pi}{2})^T R(t+\frac{\pi}{2})m,\, \frac{\partial m}{\partial \theta}\rangle + \langle R(t-\frac{\pi}{2})^T R(t)\frac{\partial m}{\partial t},\, \frac{\partial m}{\partial \theta}\rangle \\
&= -\langle m, \frac{\partial m}{\partial \theta}\rangle + \langle R(\frac{\pi}{2})\frac{\partial m}{\partial t},\, \frac{\partial m}{\partial \theta}\rangle.
\end{aligned}$$

Plugging

$$\frac{\partial m}{\partial \theta} = (-a\sin\theta, b\cos\theta) \qquad \text{and} \qquad \frac{\partial m}{\partial t} = (a'\cos\theta, b'\sin\theta)$$

into the above equality, we end up with the expression:

$$\det(\mathcal{J}M) = \langle \frac{\partial M}{\partial t}, n\rangle = a'b\cos^2\theta + ab'\sin^2\theta + (a^2-b^2)\cos\theta\sin\theta. \tag{2.8}$$

This is a quadratic expression with respect to $\cos\theta$ and $\sin\theta$, which is positive for all $\theta \in \mathbb{T}$ if and only if the discriminant $(a^2-b^2)^2 - 4aa'bb'$ is negative. The result follows in view of (2.6).

Step 2. The map $M : \mathbb{R}\times\mathbb{T} \to \mathbb{R}^2 \setminus \{\mathcal{O}\}$ is injective.

Fix $t \in \mathbb{R}$. From (2.7)–(2.8), using compactness of $\mathcal{E}(t)$ and smoothness of M, we deduce the existence of $\delta_t, \rho_t > 0$ such that, for all $s \in [t, t+\delta_t]$, $\theta \in \mathbb{T}$,

$$\Big\langle \frac{\partial M}{\partial t}(s,\theta), n(t,\theta) \Big\rangle \geq \rho_t > 0,$$

which yields

$$\big\langle M(s,\theta) - M(t,\theta), n(t,\theta)\big\rangle \;\geq\; \rho_t(s-t) \;>\; 0, \quad \text{for } t < s \leq t+\delta_t \text{ and } \theta \in \mathbb{T}.$$

It follows that $\operatorname{conv}\mathcal{E}(t) \subset \operatorname{int}\operatorname{conv}\mathcal{E}(s)$ for all $s > t$. Therefore, the family $(\operatorname{conv}\mathcal{E}(t))_{t\in\mathbb{R}}$ is nested and the map M is injective.

Step 3. The map $M : \mathbb{R}\times\mathbb{T} \to \mathbb{R}^2 \setminus \{\mathcal{O}\}$ is surjective.

Fix $(x,y) \in \mathbb{R}^2 \setminus \{\mathcal{O}\}$ and set, for $t \in \mathbb{R}$ and $D(t) = \begin{pmatrix} a(t) & 0 \\ 0 & b(t) \end{pmatrix}$,

$$\rho(t) \;:=\; ||D(t)^{-1}R(t)^{-1}(x,y)||^2 = \frac{1}{a^2(t)}(x\cos t + y\sin t)^2 + \frac{1}{b^2(t)}(-x\sin t + y\cos t)^2.$$

We claim that ρ is a smooth decreasing function with $\lim\limits_{-\infty}\rho = +\infty$ and $\lim\limits_{+\infty}\rho = 0$.

Indeed, since $(x,y) \neq (0,0)$, we get $R(t)^{-1}(x,y) \neq (0,0)$ and either $x\cos t + y\sin t \neq 0$ or $-x\sin t + y\cos t \neq 0$. Recalling that $a(t), b(t) \to 0$ as $t \to -\infty$, we deduce $\lim\limits_{-\infty}\rho = +\infty$. We also observe that $\lim\limits_{+\infty}\rho = 0$ is a direct consequence of the fact $a(t), b(t) \to +\infty$ as $t \to +\infty$.

It remains to prove that ρ' is negative. To this end, set $q(t) := x\cos t + y\sin t$ and notice that $\rho = a^{-2}q^2 + b^{-2}(q')^2$. Using that $q'' = -q$, we infer

$$\begin{aligned} \rho'(t) &= -2a'a^{-3}q^2 + 2a^{-2}q'q - 2b'b^{-3}(q')^2 + 2b^{-2}q''q' \\ &= -2a^{-2}b^{-2}\left(a'a^{-1}b^2q^2 + (a^2-b^2)qq' + b'b^{-1}a^2(q')^2\right). \end{aligned}$$

The quadratic expression $a'a^{-1}b^2q^2 + (a^2-b^2)qq' + b'b^{-1}a^2(q')^2$ with respect to q and q' is positive if and only if its discriminant is negative, which is equivalent, once again, to assume (2.6). Thus ρ is strictly decreasing and the claim follows.

Using the claim, we infer that there exists a unique $\bar{t} \in \mathbb{R}$ such that

$$\rho(\bar{t}) = ||D(\bar{t})^{-1}R(\bar{t})^{-1}(x,y)||^2 = 1.$$

Therefore, there exists a unique $\bar{\theta} \in \mathbb{T}$ such that $D(\bar{t})^{-1}R(\bar{t})^{-1}(x,y) = (\cos\bar{\theta}, \sin\bar{\theta})$. It follows that $M(\bar{t},\bar{\theta}) = (x,y)$, which proves that M is onto. □

A typical instance where Lemma 2.1 applies is to take $a = \mu b$ for some constant $\mu > 1$. Then for $b(t) = e^{\nu t}$ with $\nu > \frac{\mu^2-1}{2\mu}$, it is straightforward to check that a, b satisfy (2.1) and (2.6). Figure 2 represents the explicit choice $\mu = 2$ and $\nu = 1$ leading to $a(t) = 2e^t$ and $b(t) = e^t$.



Figure 2: The convex foliation $(\mathcal{E}(t))_{t\in\mathbb{R}}$ for $a(t) = 2b(t) = 2e^t$.

3 Defining the convex function and regularity properties

In this section we shall show that for a more precise choice of the functions $a(t), b(t)$ we can construct a convex function whose level sets are exactly the foliation $\{\mathcal{E}(t)\}_{t\in\mathbb{R}}$. Moreover, we shall show that this convex function is smooth, real-analytic on $\mathbb{R}^2 \setminus \{\mathcal{O}\}$ and satisfies (1.1).

Concretely, let us denote by $\varphi : \mathbb{R} \to \mathbb{R}$ a smooth strictly increasing function (the concrete definition will be given in (3.2), see Lemma 3.1) and let us set for all $M \in \mathbb{R}^2$

$$f(M) = \begin{cases} 0\,, & \text{if } M = (0,0), \\ \varphi(t), & \text{if } M \in \mathcal{E}(t), \end{cases} \tag{3.1}$$

where $\mathcal{E}(t)$ is the ellipse given in (2.5). We shall now show that we can adjust the parameters and choose φ in a way that (3.1) gives a well-defined convex function.

Lemma 3.1 (Construction of the convex function)**.** *Setting for $t \in \mathbb{R}$*

$$\begin{aligned} a(t) = \sqrt{2}\,\exp(t), \quad b(t) = \exp(t) \quad &\textit{in } (2.4), \\ \varphi(t) = \exp(t/\tau), \quad \tau \in (0, \tfrac{1}{10}), \quad &\textit{in } (3.1), \end{aligned} \tag{3.2}$$

the function f defined by (3.1) *is convex, with level sets the ellipses $\mathcal{E}(t)$ and* $\operatorname{argmin} f = \{\mathcal{O}\}$.

Proof. Since the functions a, b satisfy (2.1) and (2.6), we deduce by Lemma 2.1 that $\operatorname{conv}(\mathcal{E}(t))_{t\in\mathbb{R}}$ is a convex foliation. In particular, the function f is well defined from (3.1) with sublevel sets

$$[f \le \lambda] := \{M \in \mathbb{R}^2 : \, f(M) \le \lambda\} = \operatorname{conv}\left[\mathcal{E}(\varphi^{-1}(\lambda))\right]$$

compact and convex. Therefore f is a coercive, quasiconvex function.

We shall now use a result due to de Finetti and Crouzeix [5,9] which asserts that the quasiconvex function f is convex if and only if

$$\lambda \mapsto \sigma_{[f\leq\lambda]}(p) \text{ is concave for every } p \in \mathbb{R}^2,$$

where $\sigma_A(p) = \max_{M\in A} \langle p, M\rangle$ is the support function to the subset A. Without loss of generality, we may restrict to unit vectorss $p \in \mathbb{R}^2$, which results in assuming that $p = (\cos\alpha, \sin\alpha)$, for some $\alpha \in \mathbb{T}$. Therefore, we are led to prove that the function

$$\begin{aligned} G_\alpha(\lambda) &:= \sup\Big\{\Big\langle (x,y), (\cos\alpha, \sin\alpha)\Big\rangle : \, f(x,y) \leq \lambda\Big\} \\ &= \sup\Big\{\Big\langle M(t,\theta), (\cos\alpha, \sin\alpha)\Big\rangle : \, f(M(t,\theta)) = \varphi(t) \leq \lambda\Big\} \\ &= \max\Big\{\Big\langle M(t,\theta), (\cos\alpha, \sin\alpha)\Big\rangle : \theta \in \mathbb{T}, t = t(\lambda) = \varphi^{-1}(\lambda)\Big\} \end{aligned}$$

is concave. To this, end, after straightforward calculations we obtain

$$\begin{aligned} \Big\langle M(t,\theta), (\cos\alpha, \sin\alpha)\Big\rangle &= \Big\langle R(t)\, m(t,\theta),\ (\cos\alpha, \sin\alpha)\Big\rangle \\ &= \Big\langle (a(t)\cos\theta,\ b(t)\sin\theta),\ R(-t)\,(\cos\alpha, \sin\alpha)\Big\rangle \\ &= \Big\langle (\cos\theta, \sin\theta),\ (a(t)\cos(\alpha - t), b(t)\sin(\alpha - t))\ \Big\rangle \end{aligned}$$

whence we deduce

$$G_\alpha(\lambda) \,=\, \Big\| \, a(t(\lambda))\cos(\alpha - t(\lambda)),\ b(t(\lambda))\sin(\alpha - t(\lambda)) \, \Big\| \,=\, \sqrt{g_\alpha(\lambda)} \tag{3.3}$$

with

$$g_\alpha(\lambda) = a(t(\lambda))^2 \cos^2(t(\lambda) - \alpha) \,+\, b(t(\lambda))^2 \sin^2(t(\lambda) - \alpha). \tag{3.4}$$

Calculating the second derivative of G_α in (3.3) yields

$$G''_\alpha = \frac{2g''_\alpha g_\alpha - (g'_\alpha)^2}{4g_\alpha^{3/2}}.$$

Therefore, the functions $\{G_\alpha\}_{\alpha\in\mathbb{T}}$ are concave provided we establish:

$$2g''_\alpha g_\alpha - (g'_\alpha)^2 \leq 0, \quad \text{for all } \alpha \in \mathbb{T}\,. \tag{3.5}$$

At this step, we replace in (3.4) the choice for a, b and φ given in (3.2):

$$a(t) = \sqrt{2}\, e^t\,, \quad b(t) = e^t \quad \text{and } \lambda = \varphi(t) = e^{t/\tau}, \quad \text{for all } t \in \mathbb{R},$$

and we seek for the values of $\tau > 0$ that ensure inequality (3.5). In particular,

$$t := t(\lambda) = \tau \log\lambda, \quad \text{whence } t'(\lambda) = \frac{\tau}{\lambda} \text{ and } t''(\lambda) = -\frac{\tau}{\lambda^2} < 0.$$

After tedious computations, we get

$$g_\alpha = e^{2t}\left(\cos^2(t-\alpha)+1\right), \qquad g'_\alpha = 2\,e^{2t}\,t'\left(\cos^2(t-\alpha)+1-\cos(t-\alpha)\sin(t-\alpha)\right)$$

and

$$g''_\alpha = 2e^{2t}\Big((t')^2\big(3-4\cos(t-\alpha)\sin(t-\alpha)\big)+t''\left(\cos^2(t-\alpha)+1-\cos(t-\alpha)\sin(t-\alpha)\right)\Big).$$

Hence

$$\begin{aligned}
&2g''_\alpha g_\alpha-(g'_\alpha)^2 = \\
&= \quad 4e^{4t}(t')^2\Big\{\big(\cos^2(t-\alpha)+1\big)\big(3-4\cos(t-\alpha)\sin(t-\alpha)\big)-\big(\cos^2(t-\alpha)+1-\cos(t-\alpha)\sin(t-\alpha)\big)^2 \\
&\qquad\qquad + 4\,e^{4t}\,t''\big(\cos^2(t-\alpha)+1\big)\big(\cos^2(t-\alpha)+1-\cos(t-\alpha)\sin(t-\alpha)\big)\Big\} \\
&\le \quad 4e^{4t}\Big(5(t')^2+\frac{1}{2}t''\Big) \quad \le \quad \frac{2\tau(10\tau-1)e^{4t}}{\lambda^2},
\end{aligned}$$

which is negative provided we choose $\tau < 1/10$. □

We fix $M:\mathbb{R}\times\mathbb{T}\mapsto\mathbb{R}^2\backslash\{\mathcal{O}\}$ under the choice made in Lemma 3.1, that is,

$$M(t,\theta)=(X(t,\theta),Y(t,\theta))=e^t\left(\sqrt{2}\cos t\cos\theta-\sin t\sin\theta,\sqrt{2}\sin t\cos\theta+\cos t\sin\theta\right). \tag{3.6}$$

Setting

$$\begin{cases} \tilde{f}:\mathbb{R}\times\mathbb{T}\mapsto\mathbb{R} \\ \tilde{f}(t,\theta)=\varphi(t)=\exp(t/\tau) \end{cases} \tag{3.7}$$

we observe that the convex function f defined in (3.1) satisfies:

$$f(x,y)=\begin{cases} (\tilde{f}\circ M^{-1})(x,y), & \text{if } (x,y)\neq\mathcal{O}, \\ 0\,, & \text{if } (x,y)=\mathcal{O}. \end{cases} \tag{3.8}$$

With the next couple of lemmas we show that the function f, apart from being convex, enjoys several other good properties.

Lemma 3.2 (Properties of the convex function)**.** *Let $f:\mathbb{R}^2\mapsto[0,+\infty)$ be the convex function defined by* (3.6)–(3.8) *for $0<\tau<1/10$. Then*

(i). *f is strictly positive on $\mathbb{R}^2\setminus\{\mathcal{O}\}$ with $f(\mathcal{O})=0$.*

(ii). *For all $(x,y)\in\mathbb{R}^2$, it holds*

$$\left(1/\sqrt{2}\right)^{1/\tau}\left\|(x,y)\right\|^{1/\tau} \le f(x,y)\le \left\|(x,y)\right\|^{1/\tau}. \tag{3.9}$$

In particular, f is coercive.

(iii). *f is real analytic on $\mathbb{R}^2\setminus\{\mathcal{O}\}$ and $f\in\mathcal{C}^1(\mathbb{R}^2)$.*

(iv). *f satisfies the Łojasiewicz inequality* (1.1) *with* $\vartheta = 1-\tau$, $c = \tau/\sqrt{2}$, $a \equiv \mathcal{O}$ *and* $f(\mathcal{O}) = 0$, *that is*

$$\|\nabla f(x,y)\| \;\geq\; \left(\frac{\tau}{\sqrt{2}}\right) f(x,y)^{1-\tau}, \qquad \textit{for all } (x,y) \in \mathbb{R}^2. \tag{3.10}$$

Proof. (i). It is straightforward from the definition of f in (3.1) and the choice of φ.

(ii). From Lemma 2.1, for every $(x,y) \in \mathbb{R}^2 \setminus \{\mathcal{O}\}$, there exists a unique $t \in \mathbb{R}$ such that $(x,y) \in \mathcal{E}(t)$ and we have

$$\frac{x^2+y^2}{a^2(t)} \;\leq\; \frac{1}{a^2(t)}(x\cos t + y\sin t)^2 + \frac{1}{b^2(t)}(-x\sin t + y\cos t)^2 \;=\; 1 \;\leq\; \frac{x^2+y^2}{b^2(t)},$$

whence

$$e^t = b(t) \leq \|(x,y)\| \leq a(t) = \sqrt{2}e^t.$$

We deduce easily that

$$2^{-1/(2\tau)}\,\|(x,y)\|^{1/\tau} \;\leq\; f(x,y) = \varphi(t) = e^{t/\tau} \;\leq\; \|(x,y)\|^{1/\tau}.$$

(iii). It follows from (3.1) that $f = \varphi \circ p_1 \circ M^{-1}$ on $\mathbb{R}^2 \setminus \{\mathcal{O}\}$, where $p_1 : \mathbb{R}\times\mathbb{T} \mapsto \mathbb{R}$ with $p_1(t,\theta) = t$. By Lemma 2.1, the map $M : \mathbb{R}\times\mathbb{T} \mapsto \mathbb{R}^2 \setminus \{\mathcal{O}\}$ given in (3.6) is a real analytic diffeomorphism. Since p_1 and φ are analytic, the first part of the assertion follows. In particular, the function f is $\mathcal{C}^\infty$-smooth on $\mathbb{R}^2 \setminus \{\mathcal{O}\}$.
Since $1/\tau > 1$, the function $(x,y) \mapsto \|(x,y)\|^{1/\tau}$ is $\mathcal{C}^1$ over $\mathbb{R}^2$ and (3.9) yields that f is differentiable at $\mathcal{O}$ with $\nabla f(\mathcal{O}) = 0$. Therefore f is differentiable everywhere in $\mathbb{R}^2$ and, since it is convex, it is $\mathcal{C}^1$ (see for instance, [26, p. 20]).

(iv) Since $S := \operatorname{argmin} f = \{\mathcal{O}\}$, we have $\operatorname{dist}_S(M) = \|M\|$ for all $M = (x,y) \in \mathbb{R}^2$. Therefore, the first inequality in (3.9) can be written

$$f(M) \;\geq\; \mathbf{m}(\operatorname{dist}_S(M)) \quad \text{for all } M \in \mathbb{R}^2,$$

where $\mathbf{m}(r) = 2^{-1/(2\tau)}\, r^{1/\tau}$. Since

$$\frac{\mathbf{m}^{-1}(s)}{s} = \sqrt{2}\, s^{\tau-1} \in L^1_{\mathrm{loc}}((0,+\infty)),$$

we deduce from [3, Theorem 30] that the KŁ-inequality

$$\|\nabla(\psi \circ f)(M)\| \geq 1,$$

holds for all $M \in [f > 0] := \mathbb{R}^2 \setminus \{\mathcal{O}\}$, where

$$\psi(s) = \int_0^s \frac{\mathbf{m}^{-1}(\sigma)}{\sigma}\, d\sigma \;=\; \frac{\sqrt{2}}{\tau}\, s^\tau.$$

A straightforward calculation shows that (3.10) holds. □

Lemma 3.3 ($\mathcal{C}^k$-smoothness of the convex function)**.** *Let f be the convex function defined by (3.7)–(3.8) for $0 < \tau < 1/10$. Let $k \in \mathbb{N}$ be the biggest integer such that $k < \frac{1}{\tau}$. Then $f \in \mathcal{C}^k(\mathbb{R}^2)$ and $f \notin \mathcal{C}^{k+1}(\mathbb{R}^2)$.*

Proof. Recalling that f is real analytic in $\mathbb{R}^2 \setminus \{\mathcal{O}\}$ with $f(\mathcal{O}) = 0$ and $\nabla f(\mathcal{O}) = 0$, in order to prove that f is $\mathcal{C}^k$, it is sufficient to show that all the partial derivatives

$$\frac{\partial^{l_1+l_2} f}{\partial x^{l_1} \partial y^{l_2}}, \quad l_1 + l_2 \leq k, \tag{3.11}$$

which exist in $\mathbb{R}^2 \setminus \{\mathcal{O}\}$, converge to 0 at $\mathcal{O}$. To this end, it is more convenient to start by computating the partial derivatives of $\tilde{f}$ defined in (3.7). We have

$$\tilde{f}(t,\theta) := f(M(t,\theta)) = e^{t/\tau} = f(x,y) \quad \text{for } (x,y) = M(t,\theta) = (X(t,\theta), Y(t,\theta)),$$

and by differentiation, we obtain

$$\begin{pmatrix} \frac{\partial \tilde{f}}{\partial t} \\ \frac{\partial \tilde{f}}{\partial \theta} \end{pmatrix} = \begin{pmatrix} \frac{1}{\tau} e^{t/\tau} \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{\partial X}{\partial t} & \frac{\partial Y}{\partial t} \\ \frac{\partial X}{\partial \theta} & \frac{\partial Y}{\partial \theta} \end{pmatrix} \begin{pmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \end{pmatrix} \tag{3.12}$$

We can compute explicitely the partial derivatives of X and Y, see (3.6), to obtain

$$\frac{\partial X}{\partial t}, \frac{\partial Y}{\partial t}, \frac{\partial X}{\partial \theta}, \frac{\partial Y}{\partial \theta} = e^t P(t,\theta),$$

where $P(t,\theta)$ denotes generically a smooth periodic (hence bounded) function with respect to t and θ. More generally, in what follows, $P_{n,m}(t,\theta)$ (respectively $B_{n,m}(t,\theta)$) denotes a $n \times m$ matrix, the coefficients of which are smooth and periodic with respect to t and θ (respectively bounded in $(-\infty, 1] \times \mathbb{R}$). It follows that

$$\begin{pmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \end{pmatrix} = \frac{1}{\frac{\partial X}{\partial t}\frac{\partial Y}{\partial \theta} - \frac{\partial Y}{\partial t}\frac{\partial X}{\partial \theta}} \begin{pmatrix} \frac{\partial Y}{\partial \theta} & -\frac{\partial Y}{\partial t} \\ -\frac{\partial X}{\partial \theta} & \frac{\partial X}{\partial t} \end{pmatrix} \begin{pmatrix} \frac{1}{\tau} e^{t/\tau} \\ 0 \end{pmatrix}$$

Since

$$0 < e^{2t}(\sqrt{2} - \frac{1}{2}) \leq \frac{\partial X}{\partial t}\frac{\partial Y}{\partial \theta} - \frac{\partial Y}{\partial t}\frac{\partial X}{\partial \theta} = e^{2t}(\sqrt{2} + \cos\theta \sin\theta) \leq e^{2t}(\sqrt{2} + \frac{1}{2}),$$

we obtain

$$\begin{pmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \end{pmatrix} = e^{(\frac{1}{\tau}-1)t} P_{2,1}(t,\theta), \tag{3.13}$$

from which we infer that $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \to 0$ as $(x,y) \to \mathcal{O}$ or equivalently as $t \to -\infty$, since $\frac{1}{\tau} > 1$. We then recover the fact that f is $\mathcal{C}^1$, with $\nabla f(\mathcal{O}) = (0,0)$.

To prove that f is $\mathcal{C}^2$ (when $\frac{1}{\tau} > 2$), we differentiate again (3.12) to obtain

$$\begin{pmatrix} \frac{\partial^2 \tilde{f}}{\partial t^2} \\ \frac{\partial^2 \tilde{f}}{\partial t \partial \theta} \\ \frac{\partial^2 \tilde{f}}{\partial \theta^2} \end{pmatrix} = \begin{pmatrix} \frac{1}{\tau^2} e^{t/\tau} \\ 0 \\ 0 \end{pmatrix} = e^{2t} P_{3,3}(t,\theta) \begin{pmatrix} \frac{\partial^2 f}{\partial x^2} \\ \frac{\partial^2 f}{\partial x \partial y} \\ \frac{\partial^2 f}{\partial y^2} \end{pmatrix} + e^t P_{3,2}(t,\theta) \begin{pmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \end{pmatrix}, \tag{3.14}$$

where the coefficients of $e^{2t}P_{3,3}(t,\theta)$ are of the form

$$Z_1 Z_2, \quad \text{with } Z_1, Z_2 \in \mathcal{D}_1 := \left\{ \frac{\partial X}{\partial t}, \frac{\partial Y}{\partial t}, \frac{\partial X}{\partial \theta}, \frac{\partial Y}{\partial \theta} \right\}$$

and the coefficients of $e^{t}P_{3,2}(t,\theta)$ are second derivatives of X, Y. The matrix $P_{3,3}(t,\theta)$ is invertible since $(t,\theta) \in \mathbb{R} \times \mathbb{T} \mapsto M(t,\theta) := (x,y) \in \mathbb{R}^2 \setminus \{\mathcal{O}\}$ is an analytic diffeomorphism. Finally, we get

$$\begin{pmatrix} \frac{\partial^2 f}{\partial x^2} \\ \frac{\partial^2 f}{\partial x \partial y} \\ \frac{\partial^2 f}{\partial y^2} \end{pmatrix} = e^{(\frac{1}{\tau}-2)t} P_{3,1}(t,\theta) + e^{(\frac{1}{\tau}-1)t} B_{3,1}(t,\theta),$$

which proves that the second derivatives of f converge to 0 as $(x,y) \to \mathcal{O}$ if $\frac{1}{\tau} > 2$. Therefore f is $\mathcal{C}^2$ with $\nabla^2 f(\mathcal{O}) = 0_{2\times 2}$.

Continuing along the same lines, when differentiating l times, the invertible matrix in front of the l-th order derivatives of f has coefficients of the form $Z_1 Z_2 \cdots Z_l$ with $Z_1, \cdots, Z_l \in \mathcal{D}_1$ and, after tedious computations, we obtain

$$\begin{pmatrix} \frac{\partial^l f}{\partial x^l} \\ \vdots \\ \frac{\partial^l f}{\partial x^{l-i} \partial y^i} \\ \vdots \\ \frac{\partial^l f}{\partial y^l} \end{pmatrix} = e^{(\frac{1}{\tau}-l)t} P_{l+1,1}(t,\theta) + e^{(\frac{1}{\tau}-(l-1))t} B_{l+1,1}(t,\theta), \tag{3.15}$$

which converges to 0 as $(x,y) \to \mathcal{O}$ as long as $\frac{1}{\tau} > l$. Therefore f is $\mathcal{C}^l$ and all the l-th order derivatives of f are zero at $\mathcal{O}$ and we conclude that $f \in \mathcal{C}^k(\mathbb{R}^2)$, where k is the biggest integer such that $k < \frac{1}{\tau}$.

Let us now assume, towards a contradiction, that f is $\mathcal{C}^{k+1}$. Then we can write a Taylor expansion of f up to the order $k+1$ at $\mathcal{O}$. Since $\nabla^l f(\mathcal{O}) = 0$ for $l \leq k$, we obtain that

$$f(x,y) = O(||(x,y)||^{k+1}) \quad \text{in a neighborhood of } \mathcal{O}, \tag{3.16}$$

where $O(r^{k+1})/r^{k+1}$ is bounded near 0. If $\frac{1}{\tau} \notin \mathbb{N}$, then $k+1 > \frac{1}{\tau}$, and we obtain a straightforward contradiction with the first inequality in (3.9). If now $k+1 = \frac{1}{\tau} \in \mathbb{N}$, then (3.16) is not anymore contradictory with (3.9). But writing (3.15) with $l = k+1$, we get

$$\begin{pmatrix} \frac{\partial^{k+1} f}{\partial x^{k+1}} \\ \vdots \\ \frac{\partial^{k+1} f}{\partial y^{k+1}} \end{pmatrix} = P_{k+2,1}(t,\theta) + e^{t} B_{k+2,1}(t,\theta).$$

The second term above converges to zero as $t \to -\infty$, or equivalently as $(x,y) \to \mathcal{O}$, but $P_{k+2,1}(t,\theta)$ is a periodic nonconstant matrix with respect to t and θ so cannot converge as $t \to -\infty$, contradicting our assumption. This ends the proof. $\square$

4 Oscillating gradient trajectories

Suppose that f is the convex function defined in the previous section, see (3.1)–(3.2), and consider the ordinary differential equation for the gradient orbits:

$$\begin{cases} \gamma'(t) = \nabla(f(\gamma(t))), & t \in \mathbb{R}, \\ \gamma(0) = \gamma_0 \in \mathbb{R}^2 \setminus \{\mathcal{O}\}. \end{cases} \tag{4.1}$$

Since f is convex analytic in $\mathbb{R}^2 \setminus \{\mathcal{O}\}$ and coercive with a unique minimum at $\mathcal{O}$, there exists a unique maximal solution γ in $(-\infty, T)$, $T \leq +\infty$, with

$$\lim_{t \to T} \|\gamma(t)\| = +\infty \quad \text{and} \quad \lim_{t \to -\infty} \gamma(t) = \mathcal{O}.$$

In fact, finding gradient orbits is a geometric problem. We seek the unique curve γ passing through γ_0, which is orthogonal to the level sets of f. It is convenient to parametrize γ as

$$\gamma(s) = M(t(s), \theta(s)) = (X(t(s), \theta(s)), Y(t(s), \theta(s))), \ s \in \mathbb{R} \tag{4.2}$$

using the notations (2.3)–(2.4). Under this parametrization $\gamma(s) \in \mathcal{E}(t(s))$, for every $s \in \mathbb{R}$ and $\gamma'(s)$ is a normal vector at $\gamma(s)$ to the (convex) sublevel set $[f \leq f(\gamma(s))] = \operatorname{conv} \mathcal{E}(t(s))$. Therefore:

$$\gamma'(s) \perp \partial_\theta M(t(s), \theta(s)), \quad \text{for all } s \in \mathbb{R}. \tag{4.3}$$

We define the rotation angle $s \mapsto \alpha(s)$ as the angle between the x-axis and the secant $\frac{\gamma(s)}{\|\gamma(s)\|}$ (spherical part of the orbit) varying in a continuous way. Therefore

$$\begin{cases} \cos \alpha(s) = \frac{X(t,\theta)}{\sqrt{X(t,\theta)^2 + Y(t,\theta)^2}}, \\ \sin \alpha(s) = \frac{Y(t,\theta)}{\sqrt{X(t,\theta)^2 + Y(t,\theta)^2}}. \end{cases}$$

In particular, according to the notation used in (2.3)–(2.5), if $\phi(s)$ is the angle in polar coordinates of the point $m(t, \theta)$, then we have (see Figure 1):

$$\alpha(s) = t(s) + \phi(s), \quad \text{for all } s \in \mathbb{R}.$$

Lemma 4.1 (Spiraling around the origin)**.** *Let f be the convex function defined in* (3.1) *under the assumption* (3.2) *and let $s \mapsto \gamma(s)$ be a maximal orbit of the convex foliation $(\mathcal{E}(t))_{t \in \mathbb{R}}$. Then the rotation angle $s \mapsto \alpha(s)$ satisfies*

$$\lim_{s \to \pm\infty} \alpha(s) = \pm\infty. \tag{4.4}$$

See Figure 3 for a generic numerical simulation of the maximal orbit of the function f associated with the convex foliation of Figure 2.



Figure 3: Gradient orbit $\gamma(s)$ with initial point $\gamma(0) = (2, 0)$, then zoom and extra-zoom.

Proof. We use the parametrization given by (4.2). Since

$$\lim_{s\to+\infty} \|\gamma(s)\| = +\infty \quad \text{and} \quad \lim_{t\to-\infty} \gamma(s) = \mathcal{O},$$

we can assume that the function $s \mapsto t(s)$ satisfies

$$t'(s) > 0 \quad \text{and} \quad \lim_{s\to\pm\infty} t(s) = \pm\infty. \tag{4.5}$$

The goal is to compute $\alpha(s)$ using the orthogonality condition (4.3), which is equivalent to

$$\big\langle\, \gamma'(s),\, \partial_\theta M(t(s), \theta(s))\,\big\rangle = 0, \quad \text{for all } s \in \mathbb{R}. \tag{4.6}$$

Using the notations of Section 2, we have

$$\gamma'(s) = \frac{d}{ds} M(t(s), \theta(s)) = t'\partial_t(Rm) + \theta'\partial_\theta(Rm) = t'(R'm + R\partial_t m) + \theta' R\partial_\theta m$$

and $\partial_\theta M = \partial_\theta(Rm) = R\partial_\theta m$. It follows

$$\begin{aligned}
\big\langle \gamma'(s), \partial_\theta M\big\rangle &= t'\big\langle R'm, R\partial_\theta m\big\rangle + t'\big\langle R\partial_t m, R\partial_\theta m\big\rangle + \theta'\big\langle R\partial_\theta m, R\partial_\theta m\big\rangle \\
&= t'\big\langle R(\frac{\pi}{2})m, \partial_\theta m\big\rangle + t'\big\langle \partial_t m, \partial_\theta m\big\rangle + \theta'\|\partial_\theta m\|^2 \\
&= t'\left(ab + (bb' - aa')\cos\theta\sin\theta\right) + \theta'\left(a^2\sin^2\theta + b^2\cos^2\theta\right).
\end{aligned}$$

By (4.3), we have $\big\langle \gamma'(s), \partial_\theta M\big\rangle = 0$ and after substitution $a(t) = \sqrt{2}e^t$ and $b(t) = e^t$ we get

$$t'e^{2t}(\sqrt{2} - \cos\theta\sin\theta) + \theta' e^{2t}(1 + \sin^2\theta) = 0$$

whence we deduce the following relation between $t(s)$ and $\theta(s)$:

$$t'(s) = -\frac{1 + \sin^2\theta(s)}{\sqrt{2} - \cos\theta(s)\sin\theta(s)}\theta'(s). \tag{4.7}$$

Since for every $\theta \in \mathbb{R}$ we have

$$0 < \frac{1}{\sqrt{2} + \frac{1}{2}} \le \frac{1 + \sin^2\theta}{\sqrt{2} - \cos\theta\sin\theta} \le \frac{2}{\sqrt{2} - \frac{1}{2}},$$

we get

$$-\frac{1}{\sqrt{2}+\frac{1}{2}}\,\theta'(s) \;\le\; t'(s) \;\le\; -\frac{2}{\sqrt{2}-\frac{1}{2}}\,\theta'(s).$$

Therefore, from (4.5) we deduce

$$\theta'(s) < 0, \quad \theta(s) \underset{s\to-\infty}{\to} +\infty, \quad \theta(s) \underset{s\to+\infty}{\to} -\infty. \tag{4.8}$$

Next, we establish the relation between $\theta(s)$ and $\phi(s)$, see Figure 1. We have

$$\cos\phi = \frac{a\cos\theta}{\sqrt{a^2\cos^2\theta + b^2\sin^2\theta}} = \frac{\sqrt{2}\cos\theta}{\sqrt{2\cos^2\theta+\sin^2\theta}},$$
$$\sin\phi = \frac{b\sin\theta}{\sqrt{a^2\cos^2\theta + b^2\sin^2\theta}} = \frac{\sin\theta}{\sqrt{2\cos^2\theta+\sin^2\theta}}.$$

Differentiating $\cos\phi$ and plugging the result in the second expression, we end up with

$$\phi' = \frac{\sqrt{2}}{1+\cos^2\theta}\theta'. \tag{4.9}$$

Assembling (4.7) and (4.9), we obtain

$$\alpha' = t' + \phi' = \left(\frac{\sqrt{2}}{1+\cos^2\theta} - \frac{1+\sin^2\theta}{\sqrt{2}-\cos\theta\sin\theta}\right)\theta' =: h(\theta)\theta', \tag{4.10}$$

The function h is analytic and 2π-periodic, see Figure 4. We can expand it in Fourier series and integrate (4.10) to obtain

$$\alpha(s) = \frac{a_0}{2}\theta(s) + O(1), \tag{4.11}$$

where $O(1)$ is a bounded function and

$$a_0 = \frac{1}{\pi}\int_0^{2\pi} h(\theta)d\theta \simeq -0.84 < 0.$$

We finally conclude from (4.11) and (4.5) that (4.4) holds. □

5 Proof of Theorem 1.1

Consider the convex foliation by ellipses $\{\mathcal{E}(t)\}_{t\in\mathbb{R}}$ given by Lemma 2.1. Let $k \ge 1$ be any integer and f be the convex function defined by Lemma 3.1 for $0 < \tau < \min\{1/10, 1/k\}$. Then, by Lemma 3.2, the function f is coercive, has its unique minimum at the origin $\mathcal{O}$, is real analytic in $\mathbb{R}^2 \setminus \{\mathcal{O}\}$ and satisfies the Łojasiewicz inequality (1.1). Further, Lemma 3.3, ensures that f is $\mathcal{C}^k$-smooth. Finally, Lemma 4.1 asserts that all nontrivial gradient orbits spiral infinitely many times both near the origin (bounded part) and at infinity. □

Acknowledgement. This work was partially supported by the Centre Henri Lebesgue ANR-11-LABX-0020-01 and the grants CMM AFB170001, ECOS-Sud/ANID C18E04 and FONDECYT 1211217. Major part of this work has been done during a research visit of the first author to INSA Rennes. This author is indebted to his hosts for hospitality.



Figure 4: Plot of $h(\theta) = \dfrac{\sqrt{2}}{1+\cos^2\theta} - \dfrac{1+\sin^2\theta}{\sqrt{2}-\cos\theta\sin\theta}$.

References

[1] V. I. Arnold. Some open problems in the theory of singularities. In *Singularities, Part 1 (Arcata, Calif., 1981)*, volume 40 of *Proc. Sympos. Pure Math.*, pages 57–69. Amer. Math. Soc., Providence, R.I., 1983. Translated from the Russian.

[2] J. Bolte and E. Pauwels. Curiosities and counterexamples in smooth convex optimization. *TSE Working Paper, n. 20-1080*, 2020.

[3] J. Bolte, A. Daniilidis, O. Ley, and L. Mazet. Characterizations of Lojasiewicz inequalities: subgradient flows, talweg, convexity. *Trans. Amer. Math. Soc.*, 362(6):3319–3363, 2010.

[4] F. Cano, R. Moussu, F. Sanz. Nonoscillating projections for trajectories of vector fields. *J. Dyn. Control Syst.* 13(2):173–176, 2007.

[5] J.-P. Crouzeix. Conditions for convexity of quasiconvex functions. *Math. Oper. Res.*, 5(1):120–125, 1980.

[6] D. D'Acunto and V. Grandjean. On gradient at infinity of semialgebraic functions. *Ann. Polon. Math.*, 87:39–49, 2005.

[7] A. Daniilidis, G. David, E. Durand-Cartagena, and A. Lemenant. Rectifiability of self-contracted curves in the Euclidean space and applications. *J. Geom. Anal.*, 25(2):1211–1239, 2015.

[8] A. Daniilidis, O. Ley, and S. Sabourau. Asymptotic behaviour of self-contracted planar curves and gradient orbits of convex functions. *J. Math. Pures Appl. (9)*, 94(2):183–199, 2010.

[9] B. de Finetti. Sulle stratificazioni convesse. *Ann. Mat. Pura Appl. (4)*, 30:173–183, 1949.

[10] M. Fokin. Limit sets of trajectories of dynamical systems of gradient type (Russian). *Mat. Sb. (N.S.)*, 606(4):502–514, 1981.

[11] V. Grandjean. On the limit set at infinity of a gradient trajectory of a semialgebraic function. *J. Differential Equations*, 233(1):22–41, 2007.

[12] V. Grandjean, F. Sanz. On restricted analytic gradients on analytic isolated surface singularities. *J. Differential Equations*, 255(7):1684–1708, 2013.

[13] F. Ichikawa. Thom's conjecture on singularities of gradient vector fields. *Kodai Math. J.*, 15(1):134–140, 1992.

[14] Z. Ji and M. Telgarsky. Directional convergence and alignment in deep learning. *Preprint*, 2020.

[15] S. G. Krantz and H. R. Parks. *A primer of real analytic functions*. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser Boston, Inc., Boston, MA, second edition, 2002.

[16] K. Kurdyka, T. Mostowski, and A. Parusiński. Proof of the gradient conjecture of R. Thom. *Ann. of Math. (2)*, 152(3):763–792, 2000.

[17] K. Kurdyka and A. Parusiński. Quasi-convex decomposition in o-minimal structures. Application to the gradient conjecture. In *Singularity theory and its applications*, volume 43 of *Adv. Stud. Pure Math.*, pages 137–177. Math. Soc. Japan, Tokyo, 2006.

[18] H. X. Lin. Sur la structure des champs de gradients de fonctions analytiques réelles. *PhD Thesis, Université Paris VII*, 1992.

[19] S. Łojasiewicz. Une propriété topologique des sous-ensembles analytiques réels. In *Les Équations aux Dérivées Partielles (Paris, 1962)*, pages 87–89. Éditions du Centre National de la Recherche Scientifique, Paris, 1963.

[20] S. Łojasiewicz. Sur les trajectoires du gradient d'une fonction analytique. In *Geometry seminars, 1982–1983 (Bologna, 1982/1983)*, pages 115–117. Univ. Stud. Bologna, Bologna, 1984.

[21] K. Lyu and J. Li. Gradient descent maximizes the margin of homogeneous neural networks. *Preprint*, 2019.

[22] P. Manselli and C. Pucci. Maximum length of steepest descent curves for quasi-convex functions. *Geom. Dedicata*, 38(2):211–227, 1991.

[23] C. Miller. Exponentiation is hard to avoid. *Proc. Amer. Math. Soc.*, 122(1):257–259, 1994.

[24] R. Moussu. Sur la dynamique des gradients. Existence de variétés invariantes (French). *Math. Ann.* 307(3): 445–460, 1997.

[25] J. Palis, Jr. and W. de Melo. *Geometric theory of dynamical systems.* Springer-Verlag, New York, 1982. An introduction, Translated from the Portuguese by A. K. Manning.

[26] R. R. Phelps. *Convex functions, monotone operators and differentiability*, volume 1364 of *Lecture Notes in Mathematics.* Springer-Verlag, Berlin, 1989.

[27] F. Sanz. Non-oscillating solutions of analytic gradient vector fields. *Ann. Inst. Fourier (Grenoble)*, 48(4):1045–1067, 1998.

[28] F. Sanz. Balanced coordinates for spiraling dynamics. *Qual. Theory Dyn. Syst.*, 3(1):181–226, 2002.

[29] R. Thom. Problèmes rencontrés dans mon parcours mathématique: un bilan. *Inst. Hautes Études Sci. Publ. Math.*, (70):199–214 (1990), 1989.

[30] L. van den Dries. *Tame topology and o-minimal structures*, volume 248 of *London Mathematical Society Lecture Note Series.* Cambridge University Press, Cambridge, 1998.

[31] L. van den Dries and Chris Miller. Geometric categories and o-minimal structures. *Duke Math. J.*, 84(2):497–540, 1996.

Aris Daniilidis

DIM–CMM, CNRS IRL 2807
Beauchef 851, FCFM, Universidad de Chile

E-mail: `arisd@dim.uchile.cl`
`http://www.dim.uchile.cl/~arisd/`

Research supported by the grants:

CMM AFB170001, ECOS-ANID C18E04, Fondecyt 1211217 (Chile),
PGC2018-097960-B-C22 (Spain and EU).

Mounir Haddou, Olivier Ley

Univ Rennes, INSA, CNRS, IRMAR - UMR 6625, F-35000 Rennes, France

E-mail: {`mounir.haddou, olivier.ley`}`@insa-rennes.fr`
`http://`{`haddou, ley`}`.perso.math.cnrs.fr/`

Research supported by the Centre Henri Lebesgue ANR-11-LABX-0020-01.