



HAL
open science

An example of symmetry space group for a quasicrystal

A. Elkharrat, C. Frougny, J.-P. Gazeau, Jean-Louis Verger-Gaugry

► **To cite this version:**

A. Elkharrat, C. Frougny, J.-P. Gazeau, Jean-Louis Verger-Gaugry. An example of symmetry space group for a quasicrystal. *Ferroelectrics*, 2004, 305 (1), pp.9-13. 10.1080/00150190490462243 . hal-03136727

HAL Id: hal-03136727

<https://hal.science/hal-03136727>

Submitted on 13 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An Example of Symmetry Space-Group for a Quasicrystal

*A. Elkharrat[‡], C. Frougny[§], J.-P. Gazeau[‡],
and J.-L. Verger-Gaugry[†].*

[‡] *Laboratoire de Physique Théorique de la Matière Condensée, Boîte 7020, Université Paris 7-Denis Diderot, 75251 Paris Cedex 05, France*

[§] *Laboratoire d'Informatique Algorithmique : Fondements et Applications, UMR 7089 CNRS, Boîte 7014, Université Paris 7-Denis Diderot, 75251 Paris Cedex 05, France*

[†] *Institut Fourier, Université Joseph Fourier Grenoble, CNRS URA 188, BP 74- Domaine Universitaire, 38402- Saint Martin d'Hères, France.*

Abstract.

We present the construction of a symmetry plane-group for a quasiperiodic point-set in the plane named τ -lattice, τ being the golden-ratio. The τ -lattice generalizes the notion of periodic lattice to quasiperiodicity. The algebraic framework is issued from the counting system of τ -integers. The set of τ -integers can be equipped with an abelian group structure and an internal multiplicative law. These arithmetic structures lead to a freely generated symmetry plane-group for the τ -lattice, based on repetitions of discrete " τ -rotations" and " τ -translations" in the plane. Hence the τ -lattice, endowed with these adapted rotations and translations, can be viewed as a lattice with "rotational" symmetries.

1. Introduction

Historically, periodic tilings and lattices merge from Crystallography, and are associated with crystals. A crystallographic group of \mathbb{R}^d , is a discrete group of isometries whose maximal translation subgroup is of rank d . A crystal, as it is commonly used in Physics, is the orbit under the action of a crystallographic group of a finite number of points of \mathbb{R}^d . Let us recall the example of the square lattice $\Lambda = \mathbb{Z} + \mathbb{Z}e^{i\frac{\pi}{2}}$. The symmetry space-group G of Λ is the semi-direct product of the group of translations of Λ by its group of rotations $G = \Lambda \times \{1, -1, e^{i\frac{\pi}{2}}, e^{-i\frac{\pi}{2}}\}$, its composition law being defined by $(\lambda, R)(\lambda', R') = (\lambda + R\lambda', RR')$, with $\lambda, \lambda' \in \Lambda$ and $R, R' \in \{1, -1, e^{i\frac{\pi}{2}}, e^{-i\frac{\pi}{2}}\}$.

For quasicrystals, as a consequence of their aperiodicity, we generally do not have such an algebraic frame. For quasicrystals determined by some Pisot number $\beta > 1$, we can introduce an underlying structure, generically called *beta-lattice* [3]. Experimentally observed quasicrystals are related to well known Pisot numbers: $\tau = \frac{1+\sqrt{5}}{2}$ with symmetry of order 5 or 10, $\delta = 1 + \sqrt{2}$ with symmetry of order 8, and $\theta = 2 + \sqrt{3}$ with symmetry of order 12. Beta-lattices are Meyer sets, which have the property to be self-similar, with β as self-similarity factor [5]. Beta-lattices are eligible “counting systems” for quasicrystals, that is an algebraic framework in which one could study the properties of quasiperiodic point-sets and tilings.

The objective of the present work consists in generalizing the above context of crystals to beta-lattices, the τ -lattice, $\Gamma(\tau)$, being an illustration (Fig. 1). The algebraic framework




FIGURE 1 Two tilings of the τ -lattice $\Gamma(\tau)$.

is issued from the counting system called τ -integers and denoted by \mathbb{Z}_τ . We show that this τ -lattice is a “crystal” for a “plane-group” that we will explicitly determine. First, we recall the internal laws, called τ -addition and quasi-multiplication, on the set \mathbb{Z}_τ , then we establish the algebraic operations of τ -translations and τ -rotations, leaving the τ -lattice invariant, and finally we construct a plane-group for the τ -lattice which possesses the same algebraic structure as a plane-group in the classical lattice case.

2. The Set of τ -Integers \mathbb{Z}_τ

Any real number $x > 0$ can be developed in series of powers of τ , $x = \sum_{i \leq k} x_i \tau^i$, such that the sequence of digits $(x_i)_{i \leq k}$, called the τ -expansion, is computed through the *greedy algorithm* (see [4] and references therein). The set of positive τ -integers is the set of positive real numbers x who's τ -expansions are such that $(x_i)_{0 \leq i \leq k}$, and $x = \sum_{0 \leq i \leq k} x_i \tau^i$. For example $\tau^2 + 1$ and τ^3 are τ -integers, but $1/\tau$ and 2 are not (Table 1). The set of τ -integers \mathbb{Z}_τ is self-similar and symmetrical with respect to the origin

$$\tau \mathbb{Z}_\tau \subset \mathbb{Z}_\tau, \quad \mathbb{Z}_\tau = -\mathbb{Z}_\tau. \quad (1)$$

The set of τ -integers is associated to a tiling of the real line, with two tiles L and S , the lengths of the tiles being $\ell(L) = 1$, and $\ell(S) = 1/\tau$.

The set of τ -integers verifies the following arithmetic property. Let b_m and b_n be the m th and n th τ -integers respectively. Then $b_m + b_n = b_{m+n} + \{0, \pm 1/\tau\}$, and $b_m \times b_n = b_{mn - \rho_S(m)\rho_S(n)} + \{0, \pm 1/\tau\}$. The function $\rho_S(j)$ counts the number of small tiles between 0 and b_j in the tiling associated to \mathbb{Z}_τ . For instance, $\rho_S(2) = 1$, since there is one long tile and one small tile between 0 and $\tau = b_2$. From these relations, we introduce the following

TABLE 1 Some τ -expansions

	...	τ^3	τ^2	τ	1	$\frac{1}{\tau}$	$\frac{1}{\tau^2}$...
τ				1	0	·	0	0
$\tau^2 + 1$			1	0	1	·	0	0
$\tau^3 = \tau^2 + \tau$		1	0	0	0	·	0	0
2				1	0	·	0	1

internal laws [2]

$$b_m \oplus b_n = b_{m+n}, \quad (2)$$

$$b_m \otimes b_n = b_{mn - \rho_S(m)\rho_S(n)}, \quad (3)$$

respectively called here τ -addition and quasi-multiplication. The τ -addition is consistent with addition on \mathbb{R} , and the quasi-multiplication is consistent with multiplication on \mathbb{R} , with regards to the additive and multiplicative properties of τ -integers stated above. Note that $(\mathbb{Z}_\tau, \oplus)$ is an abelian group, but since the quasi-multiplication is neither associative, nor is it distributive with respect to the τ -addition, $(\mathbb{Z}_\tau, \oplus, \otimes)$ is not a ring.

3. Tau-Lattices and Plane-Groups

We now focus on the τ -lattices in the plane

$$\Gamma(\tau) = \mathbb{Z}_\tau + \mathbb{Z}_\tau \zeta, \quad \zeta = e^{i\frac{\pi}{5}}. \quad (4)$$

Note that the latter is not rotationally invariant. The material of this section can be found [1].

A point group for the τ -lattice in the plane—Since the τ -lattice $\Gamma(\tau)$ is not rotationally invariant, we shall enforce invariance by changing the additive and multiplicative laws by τ -additive and quasi-multiplicative laws. Let η_q and ν_q the real numbers such that $\zeta^q = \eta_q + \nu_q \zeta$.

Definition 1. *The following 10 operators r_q , $q = 0, 1, \dots, 9$, are defined on $\Gamma(\tau)$ by:*

$$r_q \odot (b_m + b_n \zeta) = \eta_q b_m \oplus (-\nu_q b_n) + ((\eta_q + \tau \nu_q) b_n \oplus \nu_q b_m) \zeta. \quad (5)$$

Introducing the identity action Id , the space inversion σ in the complex plane, $\sigma \odot z = -z$, and the associative composition rule, $(r r') \odot z = r \odot (r' \odot z)$, we trivially check that $r_0 = Id$ and $r_{q+n/2} = \sigma r_q = r_q \sigma$ for $q = 0, 1, \dots, 4$.

Now the computing of composition of any two of such τ -rotations r_q yields the following important result.

Proposition 1. *Let \mathfrak{R}_n denote the semi-group freely generated by all r_q , $q \in \{0, 1, \dots, n-1\}$. Among all such extended rotations, only $r_0, r_1, r_{n/2-1}, r_{n/2+1}, r_{n-1}, \sigma$ have their inverse in \mathfrak{R}_n .*

As an immediate consequence, we can assert the following on the existence of an extended symmetry group for $\Gamma(\tau)$.

Corollary 1. *The group \mathcal{R}_{10} freely generated by the four-element set $\{r_0, \sigma, r_1, r_{n/2-1}\}$ is a symmetry group for the τ -lattice $\Gamma(\tau)$. It will be called in the present context the point symmetry group of $\Gamma(\tau)$.*

A plane-group for the τ -lattice—We now introduce into the present formalism the following τ -translations acting on $\Gamma(\tau)$.

Definition 2. Let $z_0 = b_{m_0} + b_{n_0}\zeta$ be an element of $\Gamma(\tau)$. There corresponds to it the following internal action $t_{z_0} : \Gamma(\tau) \mapsto \Gamma(\tau)$

$$t_{z_0}(z) = z \oplus z_0 \stackrel{\text{def}}{=} b_m \oplus b_{m_0} + (b_n \oplus b_{n_0})\zeta. \quad (6)$$

The set of such τ -translations forms an abelian group isomorphic to $\Gamma(\tau)$, considered itself as a group for the law \oplus . For this reason it will be also denoted by $\Gamma(\tau)$.

A simple computation leads to the following result.

Proposition 2. The group \mathcal{S}_{10} , freely generated by the five-element set $\{r_0, \sigma, r_1, r_4, t_1\}$ is a symmetry group for the τ -lattice $\Gamma(\tau)$. This group is the semi-direct product of $\Gamma(\tau)$ and \mathcal{R}_{10}

$$\mathcal{S}_{10} = \Gamma(\tau) \times \mathcal{R}_{10} = \{(b, R) \in \Gamma(\tau) \times \mathcal{R}_{10}\} \quad (7)$$

with the composition rule

$$(b, R)(b', R') = (b \oplus R \odot b', RR'). \quad (8)$$

In the present context, \mathcal{S}_n will be called the space symmetry group of $\Gamma(\tau)$. The action of an element of \mathcal{S}_{10} on $\Gamma(\tau)$ is thus defined as

$$(b, R) \cdot z = b \oplus R \odot z = t_b(R \odot z) \in \Gamma(\tau). \quad (9)$$

Cut-and-project sets under group actions—Now that we have constructed a space-group for $\Gamma(\tau)$ we can observe the action on this group on cut-and-project sets. Figure 2 displays a cut-and-project set of decagonal window, and its transforms under the action of t_1 and of t_2r_1 .

Tiles under group action—We also would like to illustrate the action of \mathcal{S}_{10} by showing how a tile of $\Gamma(\tau)$ is transformed under the action of an element of \mathcal{S}_{10} . Let $z = b_m + b_n\zeta \in \Gamma(\tau)$.




FIGURE 2 Embedding of a decagonal cut-and-project set in $\Gamma(\tau)$ (left), and of its transforms under t_1 (center) and t_2r_1 (right).




FIGURE 3 Rotation operator r_1 applied to elementary tiles of the τ -lattice. Note how the tiles are deformed in order for the vertices to remain in $\Gamma(\tau)$.

An elementary quadrilateral tile on z is $\mathbf{T}(z) = \{z, z \oplus 1, z \oplus \zeta, z \oplus (1 + \zeta)\}$. Let us apply the rotation operator r_1 on $\mathbf{T}(z)$ and around one of the edges of $\mathbf{T}(z)$. For instance, a rotation around z is defined as

$$t_z(r_1 \circ t_{-z}(\mathbf{T}(z))) = \{z, z \oplus \zeta, z \oplus (-1 + \tau\zeta), z \oplus (-1 + \tau^2\zeta)\}. \quad (10)$$

Examples of such rotation operations are displayed on Fig. 3. This operation not only rotates, but distorts the tiles.

References

1. A. Elkharrat, Ch. Frougny, J. P. Gazeau, and J. L. Verger-Gaugry, to appear in *Theor. Comp. Sci.*, special issue on Tilings.
2. Ch. Frougny, J. P. Gazeau, and R. Krejcar, *Theor. Comp. Sci.* **303**, vol. 2-3, pp. 491–516.
3. J. P. Gazeau and R. Krejcar, in *From Quasicrystals to More Complex Systems* (F. Axel, F. Denoyer, and J. -P. Gazeau, eds.), Springer & Les éditions de physique (2000).
4. M. Lothaire, *Algebraic Combinatorics on Words*, Cambridge University Press (2002).
5. Y. Meyer, *Algebraic Numbers and Harmonic Analysis*, North-Holland (1972).