



HAL
open science

Symmetry groups for beta-lattices

Avi Elkharrat, Christiane Frougny, Jean-Pierre Gazeau, Jean-Louis
Verger-Gaugry

► **To cite this version:**

Avi Elkharrat, Christiane Frougny, Jean-Pierre Gazeau, Jean-Louis Verger-Gaugry. Symmetry groups for beta-lattices. *Theoretical Computer Science*, 2004, 319 (1-3), pp.281 - 305. 10.1016/j.tcs.2004.02.013 . hal-03136734

HAL Id: hal-03136734

<https://hal.science/hal-03136734>

Submitted on 12 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Symmetry groups for beta-lattices

*A. Elkharrat[‡], C. Frougny[§], J.-P. Gazeau[‡],
and J.-L. Verger-Gaugry[†].*

[‡] *Laboratoire de Physique Théorique de la Matière Condensée, Boîte 7020, Université Paris 7-Denis Diderot, 75251 Paris Cedex 05, France*

[§] *Laboratoire d'Informatique Algorithmique : Fondements et Applications, UMR 7089 CNRS, Boîte 7014, Université Paris 7-Denis Diderot, 75251 Paris Cedex 05, France*

[†] *Institut Fourier, Université Joseph Fourier Grenoble, CNRS URA 188, BP 74- Domaine Universitaire, 38402- Saint Martin d'Hères, France.*

Abstract. We present a construction of symmetry plane-groups for quasiperiodic point-sets named beta-lattices. The framework is issued from beta-integers counting systems. Beta-lattices are vector superpositions of beta-integers. When $\beta > 1$ is a quadratic Pisot-Vijayaraghavan algebraic unit, the set of beta-integers can be equipped with an abelian group structure and an internal multiplicative law. When $\beta = (1 + \sqrt{5})/2, 1 + \sqrt{2}$ and $2 + \sqrt{3}$, we show that these arithmetic and algebraic structures lead to freely generated symmetry plane-groups for beta-lattices. These plane-groups are based on repetitions of discrete adapted rotations and translations we shall refer to as “beta-rotations and “beta-translations. Hence beta-lattices, endowed with beta-rotations and beta-translations, can be viewed like lattices. The quasiperiodic function $\rho_S(n)$, defined on the set of beta-integers as counting the number of small tiles between the origin and the n th beta-integer, plays a central part in these new group structures. In particular, this function behaves asymptotically like a linear function. As an interesting consequence, beta-lattices and their symmetries behave asymptotically like lattices and lattice symmetries, respectively.

1. Introduction

Underlying the notion of a *tiling* there is the notion of a *point-set*. In this paper, we assume point-sets to be *Delaunay sets* [16,17]. There exist infinitely many possibilities to build a tiling from a Delaunay set, and conversely, there are infinitely many ways to build a Delaunay set from its associated tiling. A possible method is to consider the set of vertices of a tiling as an associated Delaunay set [12], which is the correspondence we will assume in the following. We will indifferently mention a tiling or its associated Delaunay set, displaying or not the edges in the figures.

In general, there does not exist a symmetry group for a tiling nor for its associated Delaunay set, except for periodic tilings and *lattices*. Historically, the latter merge from crystallography, and are associated with crystals. Note that in 1991, after the discovery of modulated phases and of quasicrystals, crystallography have been divided in two categories: periodic crystallography, and aperiodic crystallography [10]. Let us sketch the general algebraic frame of periodic crystallography.

Definition 1. A crystallographic group in \mathbb{R}^d , or a space-group in \mathbb{R}^d , is a discrete group of isometries whose maximal translation subgroup is of rank d , hence isomorphic to \mathbb{Z}^d .

Definition 2. A periodic crystal is the orbit under the action of a crystallographic group of a finite number of points of \mathbb{R}^d .

We can illustrate these definitions with the square lattice $A = \mathbb{Z} + \mathbb{Z}e^{i\pi/2}$, which is a classical lattice case. This set presents a 4-fold rotational symmetry. The symmetry space-group G associated with A is the semi-direct product of the translation-group of A by its rotation-group

$$G = A \rtimes \{1, -1, e^{i\pi/2}, e^{-i\pi/2}\}$$

its internal law being

$$(\lambda, R)(\lambda', R') = (\lambda + R\lambda', RR')$$

with $\lambda, \lambda' \in A$ and $R, R' \in \{1, -1, e^{i\pi/2}, e^{-i\pi/2}\}$.

In the context of the 18th problem of Hilbert, Bieberbach has shown that the number of isomorphism classes (equivalently of conjugation classes) of crystallographic groups is finite for all d [25]. Therefore, the number of crystallographic groups leaving invariant a fixed crystal of \mathbb{R}^d is finite.

For quasicrystals, as a consequence of aperiodicity, we do not have such a convenient algebraic structure of symmetry space-groups, as in the periodic case. For quasicrystals determined by some quadratic Pisot–Vijayaraghavan (PV) unit, generically denoted by $\beta > 1$, we can introduce an underlying structure, the so-called *beta-lattice* [1]. Experimentally observed quasicrystals are related to well-known PV numbers [11], namely for $\beta = \tau = \frac{1+\sqrt{5}}{2}$, $\beta = \delta = 1 + \sqrt{2}$, and $\beta = \theta = 2 + \sqrt{3}$.

Beta-lattices are based on beta-integers. When β is a PV number, the set of beta-integers, denoted by \mathbb{Z}_β , is a self-similar Meyer set, with self-similarity factor β . Recall that a Delaunay set is a Meyer set $A \in \mathbb{R}^d$ if $A - A \subset A + F$, where F is a finite set. We generically define a beta-lattice $\Gamma = \Gamma(\beta) \in \mathbb{R}^d$ by

$$\Gamma = \sum_{i=1}^d \mathbb{Z}_\beta \mathbf{e}_i$$

with (\mathbf{e}_i) a base of \mathbb{R}^d . Therefore, Γ is a self-similar Meyer set with self-similarity factor β . With this respect, beta-lattices are eligible frames in which one could think of the properties of quasiperiodic point-sets and tilings, thus generalizing the notion of lattice in periodic cases.

The aim of the present work is to extend the algebraic frame of periodic crystals to beta-lattices: we construct a space-group matching Definition 1 such that the beta-lattice is the orbit under the action of this space-group of a finite set of points of \mathbb{R}^2 , as in Definition 2. In other words, we show that a beta-lattice is at least a “crystal” for a “space-group” that we determine explicitly.

We consider the cases in which β is one of the “quasicrystallographic” numbers mentioned above. Since we restrict ourselves to the case $d = 2$, we rather talk of “plane-groups”. We proceed by first recalling the internal additive and multiplicative laws on the set of beta-integers $\mathbb{Z}_\beta \subset \mathbb{R}$, which “almost” endow this set with a structure of ordered ring (order induced by that of \mathbb{R}) [6], then by establishing a set of algebraic operations, acting on the given beta-lattice by leaving it invariant. We report on the algebraic constructions of such extended plane-groups, leaving aside the delicate questions of compatible metrics and of the number (finite or infinite) of possible “space-groups” leaving invariant a given beta-lattice. However we show that the internal transformations defined on beta-lattices are *compatible* with Euclidean transformations. Compatibility property is given by the following definition.

Definition 3. Let \top be an internal law defined on \mathbb{R}^d , and let $A \subset \mathbb{R}^d$ be a set. We say that an internal operation $\hat{\top}$ defined on A is \top -compatible with the operation \top if for all $\lambda, \lambda' \in A$, $\lambda \top \lambda' \in A$ implies $\lambda \hat{\top} \lambda' = \lambda \top \lambda'$.

The article is organized as follows. In Section 2, we recall some definitions on Delaunay sets, Meyer sets, and on cyclotomic PV numbers. In Section 3, we recall results on the arithmetics and the internal laws on \mathbb{Z}_β . Most of this material can be found in [6], and is essential for the understanding of the present article. In Section 4, we give the definition of beta-lattices in the plane, together with their rotational and translational properties. A general form for beta-lattices is $\Gamma_1(\beta) = \mathbb{Z}_\beta + \mathbb{Z}_\beta e^{i2\pi/N}$ for β a cyclotomic PV unit of symmetry N . Fig. 1 is a possible tiling of such a beta-lattice with $\beta = \tau$, the golden mean, namely a τ -lattice. Section 5 is the central part of the article, with its main result: the construction of the plane-groups associated with the beta-lattices. We use the internal additive and multiplicative laws on \mathbb{Z}_β to define a symmetry point-group for $\Gamma_1(\beta)$ in Theorem 1, and the free symmetry plane-group of $\Gamma_1(\beta)$ in Theorem 2. Then we illustrate the action of the symmetry plane-group of




Fig. 1. A tiling of the τ -lattice $F_1(\tau)$.

$F_1(\tau)$ on the tiles of a τ -lattice. Section 6 is dedicated to the asymptotic properties of beta-lattices. The striking feature which is shown there is that asymptotically the set of beta-integers behaves like a ring, but with a contraction factor. We touch here the fundamental question of whether a beta-lattice can be considered as a module over an ordered ring. If it were the case, the present construction would enter into the realm of the Artin–Schreier theory ([13, Chapter 6]). Eventually, we make explicit the rotation actions for the quasicrystallographic numbers τ , δ and θ in the appendix.

2. Preliminaries

2.1. Delaunay sets and Meyer sets

Delaunay sets were introduced as a mathematical idealization of a solid-state structure, see [12]. A set $A \subset \mathbb{R}^d$ is said to be *uniformly discrete* if there exists $r > 0$ such that $\|x - y\| \geq r$, for all $x, y \in A$. We can equivalently say that every closed ball of radius r contains at most a point of A . A set A is said to be *relatively dense* if there exists $R > 0$ such that for all $y \in \mathbb{R}^d$, there exists $x \in A$ such that $\|x - y\| < R$. We can equivalently say that every open ball of radius R contains at least a point of A . If both

conditions are satisfied, A is said to be a *Delaunay set*. The possible range of ratios R/r is studied in [21] as a function of d . The action of the group of rigid motions (or Euclidean displacements) of \mathbb{R}^d on the set of uniformly discrete sets and Delaunay sets can be found in [22].

The first models of quasicrystal were introduced by Meyer [16–18], and they are now known as *Meyer sets*. A set $A \subset \mathbb{R}^d$ is said to be a *Meyer set* if it is a Delaunay set and if there exists a finite set F such that

$$A - A \subset A + F.$$

This is equivalent to $A - A$ being a Delaunay set. A review on Meyer sets can be found in [19,20].

2.2. Crystals and Bravais lattices

Bravais lattices are used as mathematical models for crystals. A Bravais lattice is an infinite discrete point-set such that the neighborhoods of a point are the same whichever point of the set is considered. Geometrically, a Bravais lattice is characterized by all Euclidean transformations (translations and possibly rotations) that transform the lattice into itself. The condition $2 \cos 2\pi/N \in \mathbb{Z}$ characterizes Bravais lattices which are left invariant under rotation of $2\pi/N$, N -fold Bravais lattices, in \mathbb{R}^2 (and in \mathbb{R}^3). Let us put $\zeta = e^{i2\pi/N}$, $\zeta^N = 1$. If we consider the \mathbb{Z} -module in the plane

$$\mathbb{Z}[\zeta] = \mathbb{Z} + \mathbb{Z}\zeta + \mathbb{Z}\zeta^2 + \cdots + \mathbb{Z}\zeta^{N-1} = \mathbb{Z} \left[2 \cos \frac{2\pi}{N} \right] + \mathbb{Z} \left[2 \cos \frac{2\pi}{N} \right] \zeta,$$

we get the cyclotomic ring of order N . This N -fold structure is generically dense in \mathbb{C} , except precisely for the crystallographic cases. We indeed check that $\mathbb{Z}[\zeta] = \mathbb{Z}$ for $N = 1$ or 2 , $\mathbb{Z}[\zeta] = \mathbb{Z} + \mathbb{Z}i$ for $N = 4$ (square lattice), and $\mathbb{Z}[\zeta] = \mathbb{Z} + \mathbb{Z}e^{i\pi/3}$ for the triangular and hexagonal cases $N = 3$ and 6 . Note that a Bravais lattice is a Meyer set such that $F = \{0\}$.

2.3. Non-crystallographic cases

For a general N , the number $2 \cos 2\pi/N$ is an algebraic integer of degree $m = \varphi(N)/2 \leq \lfloor (N-1)/2 \rfloor$ where φ is the Euler function and $\lfloor y \rfloor$ denotes the integer part of a real number y . We shall now recall some definitions on numbers.

A *Pisot–Vijayaraghavan number*, or PV number in short, is an algebraic integer $\beta > 1$ such that all its Galois conjugates (i.e. other roots of the involved algebraic equation) have their moduli strictly smaller than 1. A *cyclotomic PV number* with symmetry of order N is a PV number β such that

$$\mathbb{Z} \left[2 \cos \frac{2\pi}{N} \right] = \mathbb{Z}[\beta]. \tag{1}$$

Then $\mathbb{Z}[\zeta] = \mathbb{Z}[\beta] + \mathbb{Z}[\beta]\zeta$, with $\zeta = e^{i2\pi/N}$, is a ring invariant under rotation of order N (see [1]). This ring is the natural framework for two-dimensional structures having

β as scaling factor, and $2\pi/N$ rotational symmetry. In this paper, we will focus on quadratic PV units. They are of two kinds. The first kind is such that β is solution of

$$X^2 = aX + 1, \quad a \geq 1$$

and its conjugate is $\beta' = -1/\beta$. The second kind is such that β is solution of

$$X^2 = aX - 1, \quad a \geq 3$$

and its conjugate is $\beta' = 1/\beta$. Let us give some examples of those numbers, together with their respective Galois conjugates, related to non-crystallographic cyclotomic structures in the plane, and minimal polynomials, the following notations being used throughout the article:

$$N = 5, \quad \beta = \tau = \frac{1 + \sqrt{5}}{2} = 1 + 2 \cos \frac{2\pi}{5},$$

$$\tau' = -\frac{1}{\tau} = 1 - \tau, \quad X^2 - X - 1 \quad (\text{pentagonal case}),$$

$$N = 10, \quad \beta = \tau = \frac{1 + \sqrt{5}}{2} = 2 \cos \frac{2\pi}{10},$$

$$\tau' = -\frac{1}{\tau} = 1 - \tau, \quad X^2 - X - 1 \quad (\text{decagonal case}),$$

$$N = 8, \quad \beta = \delta = 1 + \sqrt{2} = 1 + 2 \cos \frac{2\pi}{8},$$

$$\delta' = -\frac{1}{\delta} = 2 - \delta, \quad X^2 - 2X - 1 \quad (\text{octogonal case}),$$

$$N = 12, \quad \beta = \theta = 2 + \sqrt{3} = 2 + 2 \cos \frac{2\pi}{12},$$

$$\theta' = \frac{1}{\theta} = 4 - \theta, \quad X^2 - 4X + 1 \quad (\text{dodecagonal case}).$$

Note that in the case $N = 7$, we have $\beta = 1 + 2 \cos 2\pi/7$ which is solution of the cubic equation $X^3 - 2X^2 - X + 1 = 0$. At this point, we should be aware that finding a PV number such that the cyclotomic condition (1) is fulfilled for $N \geq 16$ is an open problem!

3. Additive and multiplicative properties of beta-integers

3.1. Beta-expansions

When a number $\beta > 1$ appears as a kind of fundamental invariant in a given structure, it is tempting to introduce into the procedure of understanding the latter a *counting system* based precisely on this β . Let us explain here what we mean by counting system.

Among all *beta-representations* of a real number $x \geq 0$, i.e. infinite sequences $(x_i)_{i \leq k}$, such that $x = \sum_{i \leq k} x_i \beta^i$ for a certain integer k , there exists a particular one, called the *beta-expansion*, which is obtained through the “greedy algorithm” (see [23,24]). Recall that $\lfloor y \rfloor$ is the integer part of the real number y , and denote by $\{y\}$ the fractional part of y . There exists $k \in \mathbb{Z}$ such that $\beta^k \leq x < \beta^{k+1}$. Let $x_k = \lfloor x/\beta^k \rfloor$ and $r_k = \{x/\beta^k\}$. For $i < k$, put $x_i = \lfloor \beta r_{i+1} \rfloor$, and $r_i = \{\beta r_{i+1}\}$. Then we get the expansion $x = x_k \beta^k + x_{k-1} \beta^{k-1} + \dots$. If $x < 1$ then $k < 0$, and we put $x_0 = x_{-1} = \dots = x_{k+1} = 0$. The beta-expansion of x is denoted by

$$\langle x \rangle_\beta = x_k x_{k-1} \dots x_1 x_0 \cdot x_{-1} x_{-2} \dots$$

The digits x_i obtained by this algorithm are integers from the set $A = \{0, \dots, \lceil \beta \rceil - 1\}$, called the *canonical alphabet*, where $\lceil \beta \rceil$ denotes the smallest integer larger than β . If an expansion ends in infinitely many zeros, it is said to be *finite*, and the ending zeros are omitted. For instance, if $\beta = \tau \approx 1.618 \dots$, then $x_i \in \{0, 1\}$. The τ -expansion of, say, $4 = \tau^2 + 1 + 1/\tau^2$ is $\langle 4 \rangle_\tau = 101.01$. There is a representation which plays an important role in the theory. The *beta-expansion* of 1, denoted by $d_\beta(1)$, is computed by the following process [24]. Let the *beta-transformation* be defined on $[0, 1]$ by $T_\beta(x) = \beta x \bmod 1$. Then $d_\beta(1) = (t_i)_{i \geq 1}$, where $t_i = \lfloor \beta T_\beta^{i-1}(1) \rfloor$. Bertrand has proved that if β is a PV number, then $d_\beta(1)$ is eventually periodic [2]. For instance, $d_\tau(1) = 11$, $d_\delta(1) = 21$, and $d_\theta(1) = 322 \dots = 3(2)^\omega$, where $(\cdot)^\omega$ means that the digit between parenthesis is repeated an infinite number of times. A number β such that $d_\beta(1)$ is eventually periodic is traditionally called a *beta-number*. Since these numbers were introduced by Parry [23], we propose to call them *Parry numbers*. When $d_\beta(1)$ is finite, β is said to be a *simple Parry number*.

3.2. The set of beta-integers

We now come to the notion of *beta-integer*. The set of beta-integers is the set of real numbers whose beta-expansions are polynomial,

$$\begin{aligned} \mathbb{Z}_\beta &= \{x \in \mathbb{R} \mid \langle |x| \rangle_\beta = x_k \dots x_0\} \\ &= \mathbb{Z}_\beta^+ \cup (-\mathbb{Z}_\beta^+), \end{aligned}$$

where \mathbb{Z}_β^+ is the set of non-negative beta-integers. The set \mathbb{Z}_β is self-similar and symmetrical with respect to the origin

$$\beta \mathbb{Z}_\beta \subset \mathbb{Z}_\beta, \quad \mathbb{Z}_\beta = -\mathbb{Z}_\beta.$$

It has been shown in [3] that if β is a PV number then \mathbb{Z}_β is a Meyer set. This means that there exists a finite set F such that $\mathbb{Z}_\beta - \mathbb{Z}_\beta \subset \mathbb{Z}_\beta + F$. This beta-dependent set F has to be characterized in order to see to what extent beta-integers differ from ordinary integers with respect to additive and multiplicative structures. This problem is solved in [3,4,6] for all quadratic PV units and for a few higher-degree cases (see also [27]). We now restrict the presentation to quadratic PV units. There are two cases to consider.

Case 1 (β is solution of $X^2 = aX + 1$, $a \geq 1$): The Galois conjugate is $\beta' = -1/\beta$. The canonical alphabet is equal to $A = \{0, \dots, a\}$, the beta-expansion of 1 is finite, equal to $d_\beta(1) = a1$, and every positive number of $\mathbb{Z}[\beta]$ has a finite beta-expansion [7]. Denote $\mathbb{A} = \{L, S\}$. Define the substitution σ_β by

$$\sigma_\beta : \begin{cases} L \mapsto L^a S, \\ S \mapsto L. \end{cases}$$

The fixed point of the substitution, denoted by $\sigma_\beta^\infty(L)$, is associated with a tiling of the positive real line, made with the two tiles L and S , where the lengths of the tiles are $\ell(L) = 1$, $\ell(S) = T_\beta(1) = \beta - a = 1/\beta$, see [26,5]. The nodes of this tiling are the positive beta-integers.

Case 2 (β is solution of $X^2 = aX - 1$, $a \geq 3$): The Galois conjugate is $\beta' = 1/\beta$. The canonical alphabet is equal to $A = \{0, \dots, a-1\}$, the beta-expansion of 1 is eventually periodic, equal to $d_\beta(1) = (a-1)(a-2)^\omega$, and every positive number of $\mathbb{Z}[\beta]$ has an eventually periodic beta-expansion, which is finite for numbers from $\mathbb{N}[\beta]$, [7]. The substitution σ_β is defined on $\mathbb{A} = \{L, S\}$ by

$$\sigma_\beta : \begin{cases} L \mapsto L^{a-1} S, \\ S \mapsto L^{a-2} S. \end{cases}$$

As in Case 1, the fixed point of the substitution is denoted by $\sigma_\beta^\infty(L)$, and is associated with a tiling of the positive real line, made with the two tiles L and S . The lengths of the tiles are $\ell(L) = 1$, $\ell(S) = T_\beta(1) = \beta - (a-1) = 1 - 1/\beta$ [26,5]. The nodes of this tiling are the positive beta-integers.

In both cases, we shall denote by $|\sigma_\beta^q(L)|$ the number of letters in the word generated by $\sigma_\beta^q(L)$, and by $|\sigma_\beta^q(L)|_L$, respectively $|\sigma_\beta^q(L)|_S$, the number of letters L , respectively S , in the later word.

3.3. Beta-integers arithmetics

Since \mathbb{Z}_β is a Meyer set symmetrical with respect to the origin, we have $\mathbb{Z}_\beta - \mathbb{Z}_\beta = \mathbb{Z}_\beta + \mathbb{Z}_\beta \subset \mathbb{Z}_\beta + F$. Hence, the set \mathbb{Z}_β can be qualified as ‘‘quasi-additive’’. It can also be qualified as ‘‘quasi-multiplicative’’. Accordingly, addition and multiplication of beta-integers are characterized below.

- In Case 1, we have

$$\mathbb{Z}_\beta + \mathbb{Z}_\beta \subset \mathbb{Z}_\beta + \left\{ 0, \pm \left(1 - \frac{1}{\beta} \right) \right\} \subset \mathbb{Z}_\beta / \beta^2, \quad (2)$$

$$\mathbb{Z}_\beta \times \mathbb{Z}_\beta \subset \mathbb{Z}_\beta + \left\{ 0, \pm \frac{1}{\beta}, \dots, \pm \frac{a}{\beta} \right\} \subset \mathbb{Z}_\beta / \beta^2. \quad (3)$$

For instance, for $\beta = \tau$, $1 + 1 = 2 = \tau + (1 - 1/\tau)$, and $(\tau^2 + 1)(\tau^2 + 1) = \tau^5 + \tau^2 - 1/\tau$.

- In Case 2, we have

$$\mathbb{Z}_\beta + \mathbb{Z}_\beta \subset \mathbb{Z}_\beta + \left\{0, \pm \frac{1}{\beta}\right\} \equiv \tilde{\mathbb{Z}}_\beta, \quad (4)$$

$$\begin{aligned} \mathbb{Z}_\beta^+ + \mathbb{Z}_\beta^+ &\subset \mathbb{Z}_\beta^+/\beta, \\ \mathbb{Z}_\beta \times \mathbb{Z}_\beta &\subset \mathbb{Z}_\beta + \left\{0, \pm \frac{1}{\beta}, \dots, \pm \frac{a-1}{\beta}\right\} \subset \mathbb{Z}_\beta/\beta. \end{aligned} \quad (5)$$

For instance, for $\beta = \theta$, $2 + 2 = \theta + 1/\theta = 2 \times 2$.

The set $\tilde{\mathbb{Z}}_\beta$, introduced in Eq. (4), is called the set of decorated beta-integers. This set plays an important role in the theory of algebraic model sets, and is to be mentioned in the two-dimensional case (Fig. 4).

3.4. Beta-integers as an additive group

Let b_m and b_n be the m th and n th beta-integer.

Definition 4. We define the *beta-addition* as the following internal additive law on the set of beta-integers

$$b_m \oplus b_n = b_{m+n}.$$

The beta-substraction is defined by

$$b_m \ominus b_n = b_{m-n} = b_m \oplus (-b_n).$$

The set of beta-integers endowed with the beta-addition has an abelian group structure [4,6]. Actually, we can endow any countable strictly increasing sequence $\mathcal{S} = (s_n)_{n \in \mathbb{Z}}$ of real numbers, $s_0 = 0$, with such an internal additive law by simple isomorphic transport of the additive group structure of the integers, the additive law of \mathcal{S} being defined by

$$s_m \oplus s_n \stackrel{\text{def}}{=} s_{m+n}.$$

Recall that the internal additive law \oplus defined on \mathcal{S} , is said to be *compatible* with addition of real numbers if for all $(m, n) \in \mathbb{Z}^2$, $s_m + s_n \in \mathcal{S}$ implies $s_m + s_n = s_m \oplus s_n$, and obviously, for an arbitrary sequence \mathcal{S} , the law \oplus is *not* compatible with the addition of real numbers. Yet this property holds true for \mathbb{Z}_β !

Lemma 1. *Beta-addition is compatible with addition if β is a quadratic PV unit.*

Proof. It has been proven in [4,6] that beta-addition has the following *minimal distortion property* with respect to addition: for all $(b_m, b_n) \in \mathbb{Z}_\beta^2$ with β a quadratic PV unit,

$$b_m + b_n - (b_m \oplus b_n) \in \begin{cases} \{0, \pm(1 - \frac{1}{\beta})\} & \text{in Case 1,} \\ \{0, \pm 1/\beta\} & \text{in Case 2.} \end{cases} \quad (6)$$

Suppose that for a given couple of integers (m, n) , there exist q such that $b_m + b_n = b_q$. Then $b_q - (b_m \oplus b_n)$ verifies (6), and this implies $b_q - (b_m \oplus b_n) = 0$. Indeed the distances between two non-equal beta-integers are larger than or equal to $\ell(S) = 1/\beta$ in Case 1, and $\ell(S) = 1 - 1/\beta$ in Case 2. So we have $b_q = b_m \oplus b_n = b_{m+n}$, which gives $q = m + n$. \square

For instance, if $\beta = \tau$, then $1 \oplus 1 = \tau$ and $2 - \tau = 1 - 1/\tau$, and if $\beta = \theta$, then $2 \oplus 2 = \theta$ and $4 - \theta = 1/\theta$.

3.5. Internal multiplicative law for beta-integers

We could attempt to play the same game with multiplication by defining

$$b_m \times \times b_n \stackrel{\text{def}}{=} b_{mn}$$

for all $(b_m, b_n) \in \mathbb{Z}_\beta^2$. However, we reject this definition of an internal multiplicative law since it is not compatible with multiplication in \mathbb{R} . For instance, for $\beta = \tau$, $b_2 \times b_2 = \tau \times \tau = \tau^2 = b_3 \neq b_4$.

Definition 5. We define the *quasi-multiplication* as the following internal multiplicative law on the set of beta-integers:

$$b_m \otimes b_n = \begin{cases} b_{(mn - \rho_S(m)\rho_S(n))} & \text{in Case 1,} \\ b_{(mn - \rho_S(m)\rho_S(n))} & \text{in Case 2,} \end{cases} \quad (7)$$

where, for $n \geq 0$, $\rho_S(n)$ denotes the number of tiles S between $b_0 = 0$ and b_n [6]. For instance, for τ , $\rho_S(5) = 2$ while for θ , $\rho_S(5) = 1$. Geometrically, for $n \geq 0$, the n th beta-integer is the right vertex of the n th tile of the tiling associated with \mathbb{Z}_β , which can be expressed as $b_n = n + (-1 + \ell(S))\rho_S(S)$ and from which we derive the following:

$$\begin{aligned} \rho_S(n) &= \frac{1}{1 - 1/\beta} (n - b_n) \quad \text{Case 1,} \\ \rho_S(n) &= \beta(n - b_n) \quad \text{Case 2.} \end{aligned}$$

For $n < 0$, $\rho_S(n) = -\rho_S(-n)$.

Lemma 2. *Quasi-multiplication is compatible with multiplication of real numbers if β is a quadratic PV unit.*

Proof. Quasi-multiplication has minimal distortion property with respect to multiplication [4,6]: for all $(b_m, b_n) \in \mathbb{Z}_\beta^2$ with β quadratic PV unit,

$$b_m b_n - (b_m \otimes b_n) \in \begin{cases} \{(0, \pm 1, \dots, \pm a)(1 - \frac{1}{\beta})\} & \text{Case 1,} \\ \{(0, 1, \dots, a - 1) \frac{\text{sgn}(b_m b_n)}{\beta}\} & \text{Case 2.} \end{cases} \quad (8)$$

Suppose that for a given couple of integers (m, n) , there exist a q such that $b_m b_n = b_q$. Then $b_q - (b_m \otimes b_n)$ verifies (8), and this implies $b_q - (b_m \otimes b_n) = 0$. Indeed the distances

between two non-equal beta-integers are larger than or equal to $\ell(S) = 1/\beta$ in Case 1, and $\ell(S) = 1 - 1/\beta$ in Case 2. We then have $b_q = b_m \otimes b_n = b_{mn - a\rho_S(m)\rho_S(n)}$ in Case 1 and $b_q = b_m \otimes b_n = b_{mn - \rho_S(m)\rho_S(n)}$ in Case 2, which gives $q = mn - a\rho_S(m)\rho_S(n)$ in Case 1, and $q = mn - \rho_S(m)\rho_S(n)$ in Case 2. \square

An interesting outcome of this multiplicative structure is the following explicit result concerning self-similarity properties of the set of beta-integers.

Let $U = (u_q)_{q \in \mathbb{N}}$ be the linear recurrent sequence of integers associated with β . In Case 1, the u_q are defined by $u_{q+2} = au_{q+1} + u_q$ with $u_0 = 1, u_1 = a + 1$. In Case 2, the u_q are defined by $u_{q+2} = au_{q+1} - u_n$ with $u_0 = 1, u_1 = a$. The recurrence is possibly extended to negative indices.

Proposition 1. *Let β be a quadratic PV unit, and \mathbb{Z}_β the corresponding set of beta-integers. Then for $q \in \mathbb{N}$ and $b_n \in \mathbb{Z}_\beta$ we have the self-similarity formulas:*

$$\begin{aligned} \beta^q b_n &= b_{u_q} b_n = b_{u_q} \otimes b_n = b_{u_q n - a\rho_S(u_q)\rho_S(n)} = b_{u_q n - (u_q - u_{q-1})\rho_S(n)} && \text{(in Case 1),} \\ \beta^q b_n &= b_{u_q} b_n = b_{u_q} \otimes b_n = b_{u_q n - \rho_S(u_q)\rho_S(n)} = b_{u_q n - (u_q - 1)\rho_S(n)} && \text{(in Case 2).} \end{aligned}$$

The proof is a direct consequence of the definition of the quasi-multiplication and of the following lemma giving some of the properties of the counting function ρ_S .

Lemma 3. *The values assumed by the counting function $\rho_S(n)$ when $n = u_q \in U$ are*

$$\begin{aligned} \rho_S(u_q) &= \frac{u_q - u_{q-1}}{a} && \text{(in Case 1),} \\ \rho_S(u_q) &= u_{q-1} && \text{(in Case 2).} \end{aligned}$$

Proof. *Case 1:* Let $w_q = \rho_S(u_q)$. By construction, $u_q = |\sigma_\beta^q(L)|$ and $w_q = |\sigma_\beta^q(L)|_S$. Therefore, the sequence (w_q) satisfies the same linear recurrence as (u_q) , that is $w_q = aw_{q-1} + w_{q-2}$, with $w_0 = 0, w_1 = 1$. Thus $w_2 = aw_1 + w_0 = (u_2 - u_1)/a = a$ and $w_3 = aw_2 + w_1 = (u_3 - u_2)/a = a^2 + 1$. The recurrence is proved through $w_{q+1} = aw_q + w_{q-1} = a(u_q - u_{q-1})/a + (u_{q-1} - u_{q-2})/a = (u_{q+1} - u_q)/a$.

Case 2: Let $w_q = \rho_S(u_q)$. We have $w_q = aw_{q-1} - w_{q-2}$, with $w_0 = 0$ and $w_1 = 1$. Then $w_2 = aw_1 - w_0 = u_1 = a$ and $w_3 = aw_2 - w_1 = u_2 = a^2 - 1$. The recurrence is proved through $w_{q+1} = aw_q - w_{q-1} = a(u_{q+1} - u_q) - u_{q-1} = u_q$. \square

It should be noticed that quasi-multiplication does not define a group for not being associative and is not distributive with respect to beta-addition. So it seems hopeless to obtain a ring structure, like we have with integers, with such an internal multiplicative law. Note that beta-addition and quasi-multiplication are related to some operations in numeration systems studied in [9,14,15]. Nevertheless, the set of beta-integers recovers a ring structure asymptotically, as shall be explained in Section 6.

4. Beta-lattices in the plane

4.1. General considerations

We have seen that the condition $2 \cos(2\pi/N) \in \mathbb{Z}$, i.e. $N = 1, 2, 3, 4$ and 6 , characterizes N -fold Bravais lattices in \mathbb{R}^2 (and in \mathbb{R}^3). We would like to generalize this notion when N is quasicrystallographic i.e. $N = 5, 10, 8$ and 12 , respectively, associated with one of the cyclotomic Pisot units $\tau = 2 \cos(2\pi/10)$, $\delta = 1 + 2 \cos(2\pi/8)$ and $\theta = 2 + 2 \cos(2\pi/12)$. As a consequence of the results presented above, if (\mathbf{e}_i) is a basis of \mathbb{R}^d , then the point set

$$\Gamma = \sum_{i=1}^d \mathbb{Z}_\beta \mathbf{e}_i$$

is a Meyer set and a lattice for the law \oplus . Moreover $\mathbb{Z}_\beta \otimes \Gamma \subset \Gamma$. We shall adopt the generic name of *beta-lattice* for such a Γ . Examples of beta-lattices in the plane are point-sets of the form

$$\Gamma_q(\beta) = \mathbb{Z}_\beta + \mathbb{Z}_\beta \zeta^q,$$

with $\zeta = e^{i2\pi/N}$, for $1 \leq q \leq N - 1$. Note that the latter are not rotationally invariant. Examples of rotationally invariant point-sets based on beta-integers are

$$A_q \stackrel{\text{def}}{=} \bigcup_{j=0}^{N-1} \Gamma_q \zeta^j, \quad 1 \leq q \leq N - 1$$

and

$$\mathbb{Z}_\beta[\zeta] \stackrel{\text{def}}{=} \sum_{j=0}^{N-1} \mathbb{Z}_\beta \zeta^j.$$

These sets A_q and $\mathbb{Z}_\beta[\zeta]$ are Meyer sets.

Let us now focus on the simplest case, namely $N = 5$ or 10 . It is more convenient to introduce the root of unity $\zeta = e^{i\pi/5}$, since $\tau = 2 \cos \pi/5 = \zeta + \zeta^c$, where ζ^c is the complex conjugate of ζ . We obtain the set

$$\mathbb{Z}_\tau[\zeta] \equiv \mathbb{Z}_\tau + \mathbb{Z}_\tau \zeta + \mathbb{Z}_\tau \zeta^2 + \mathbb{Z}_\tau \zeta^3 + \mathbb{Z}_\tau \zeta^4.$$

Consider now the following τ -lattices in the plane:

$$\Gamma_q = \mathbb{Z}_\tau + \mathbb{Z}_\tau \zeta^q, \quad q = 1, 2, 3, \text{ or } 4.$$

The following inclusions were proven in [3]:

$$\Gamma_q \subset \mathbb{Z}_\tau[\zeta] \subset \frac{\Gamma_q}{\tau^4}.$$

It has been shown that a large class of aperiodic sets can be embedded in beta-lattices such as $\Gamma_q(\beta)$ (see [3]).




Fig. 2. The τ -lattice $I_1(\tau)$ with points (left), and its trivial tiling made by joining points along the horizontal axis, and along the direction defined by ζ .




Fig. 3. The δ -lattice $I_1(\delta)$ with points (left), and its trivial tiling obtained by joining points along the horizontal axis, and along the direction defined by ζ .

On Figs. 2–4, we displayed the τ -lattice $I_1(\tau)$, the δ -lattice $I_1(\delta)$ and the decorated θ -lattice, $\tilde{I}_1(\theta) = \tilde{\mathbb{Z}}_\beta + \tilde{\mathbb{Z}}_\beta \zeta$, respectively, both as point-sets, and as tilings.

4.2. Rotational properties of the beta-lattices $I_1(\beta)$

Although beta-lattices are not rotationally invariant, we can nevertheless study the action of rotations on them. In this section, and throughout the rest of the article, we focus on $I_1(\beta)$. For $\beta = \tau$ and δ , any beta-lattice $I_q(\beta)$ is a subset of the properly scaled beta-lattice $I_1(\beta)$. Therefore, the rotational properties of $I_q(\beta)$ can always be




Fig. 4. The decorated θ -lattice $\tilde{\Gamma}_1(\theta)$ with points (left), and its trivial tiling obtained by joining points along the horizontal axis, and along the direction defined by ζ .

reexpressed in terms of the rotational properties of $\Gamma_1(\beta)$. Note that since θ is a quadratic PV unit of the second kind, the statement is slightly different, since the θ -lattices $\Gamma_q(\theta)$ are not subsets of the properly scaled $\Gamma_1(\theta)$, for $q \neq 1$, but of its decorated version $\tilde{\Gamma}_1(\theta)$.

We introduce the algebraic integer associated with ζ , $\chi = \zeta + \bar{\zeta} = 2 \cos(2\pi/N)$, which entails $\zeta^2 = -1 + \chi\zeta$, and

$$\zeta^q = \eta_q + v_q \zeta, \quad q \in \{0, 1, \dots, N-1\}. \quad (9)$$

A rotation by $q2\pi/N$ on an arbitrary element $b_m + b_n \zeta$ of $\Gamma_1(\beta)$ then gives

$$\zeta^q (b_m + b_n \zeta) = (\eta_q b_m - v_q b_n) + (v_q b_m + (\eta_q + v_q \chi) b_n) \zeta. \quad (10)$$

This is not an element of $\Gamma_1(\beta)$ in general, but belongs to a deflated version of $\Gamma_1(\beta)$ by a certain factor. If we consider the values of the pairs (η_q, v_q) and of $\eta_q + v_q \chi$, when β assumes the specific values τ and δ , we can determine this deflation factor. When $\beta = \theta$, $\zeta^q (b_m + b_n \zeta)$ belongs to the twice decorated θ -lattice $\tilde{\tilde{\Gamma}}_1(\theta)$, as will be shown explicitly.

- When $\beta = \tau$, the results are given for $\zeta = e^{i2\pi/10}$, $\chi = \tau$.

$$\begin{array}{rcccccc} q & = & 0 & 1 & 2 & 3 & 4 \\ (\eta_q, v_q) & = & (1, 0) & (0, 1) & (-1, \tau) & (-\tau, \tau) & (-\tau, 1) \\ \eta_q + v_q \chi & = & 1 & \tau & \tau & 1 & 0, \end{array}$$

together with $(\eta_{q+5}, v_{q+5}) = (-\eta_q, -v_q)$. Hence

$$\begin{aligned} \zeta^q \Gamma_1(\tau) &\subset \Gamma_1(\tau) + \left(\left\{ 0, \pm \left(1 - \frac{1}{\tau} \right) \right\} + \left\{ 0, \pm \left(1 - \frac{1}{\tau} \right) \right\} \zeta \right) \\ &\subset \frac{\Gamma_1(\tau)}{\tau^2}. \end{aligned}$$

Note that since $\chi = \tau$, $\Gamma_1(\tau)$ is endowed with specific properties which are not encountered in other cases, namely when $\beta = \delta$, and $\beta = \theta$. These properties are given by the following lemma.

Lemma 4. For $\zeta = e^{i\pi/5}$, all elements of the cyclic group $\{\zeta^q, q \in \{0, 1, 2, \dots, 9\}\}$ are elements of the τ -lattice $\Gamma_1(\tau)$.

Proof. The demonstration is trivial from the values assumed by η_q and v_q in the case of τ ,

$$\zeta^q = \eta_q + v_q \zeta, \quad \text{with } \eta_q, v_q \in \{0, \pm 1, \pm \tau\}. \quad \square$$

Also note that from the self-similarity property of \mathbb{Z}_τ we have $\eta_q b_n \in \mathbb{Z}_\tau$, $v_q b_n \in \mathbb{Z}_\tau$ and $(\eta_q + \tau v_q) b_n \in \mathbb{Z}_\tau$, for all q and n .

- When $\beta = \delta$, $\zeta = e^{i2\pi/8}$ and $\chi = \delta - 1$.

$$\begin{array}{rcccc} q & = & 0 & 1 & 2 & 3 \\ (\eta_q, v_q) & = & (1, 0) & (0, 1) & (-1, \delta - 1) & (-\delta + 1, 1) \\ \eta_q + v_q \chi & = & 1 & \delta - 1 & 1 & 0, \end{array}$$

together with $(\eta_{q+4}, v_{q+4}) = (-\eta_q, -v_q)$. Hence

$$\begin{aligned} \zeta^q \Gamma_1(\delta) &\subset \Gamma_1(\delta) + \left\{ 0, \pm \left(1 - \frac{1}{\delta} \right), \pm 2 \left(1 - \frac{1}{\delta} \right) \right\} \\ &\quad + \left\{ 0, \pm \left(1 - \frac{1}{\delta} \right), \pm 2 \left(1 - \frac{1}{\delta} \right) \right\} \zeta \\ &\subset \frac{\Gamma_1(\delta)}{\delta^3}. \end{aligned}$$

Note that $\delta - 1 = \sqrt{2}$ is not a δ -integer. Its δ -expansion is $\langle \delta - 1 \rangle_\delta = 1 \cdot 1$. It turns out that only ζ , ζ^5 , ζ^4 and 1 are in $\Gamma_1(\delta)$.

- When $\beta = \theta$, $\zeta = e^{i2\pi/12}$ and $\chi = \theta - 2$.

$$\begin{array}{rcccccc} q & = & 0 & 1 & 2 & 3 & 4 & 5 \\ (\eta_q, v_q) & = & (1, 0) & (0, 1) & (-1, \theta - 2) & (-\theta + 2, 2) & (-2, \theta - 2) & (-\theta + 2, 1) \\ \eta_q + v_q \chi & = & 1 & \theta - 2 & 2 & \theta - 2 & 1 & 0, \end{array}$$

together with $(\eta_{q+6}, v_{q+6}) = (-\eta_q, -v_q)$. Note that $\theta - 2 = \sqrt{3}$ is not a θ -integer. Moreover, the θ -expansion of $\theta - 2$ is infinite: $\langle \theta - 2 \rangle_\theta = 1 \cdot (2)^\omega$. Then, only ζ , ζ^6 , ζ^7 and 1 are in $\Gamma_1(\theta)$. Let us introduce the decorated θ -lattice $\tilde{\Gamma}_1(\theta)$, as we have done in the one-dimensional case (Eq. (4)),

$$\Gamma_1(\theta) \subset \tilde{\Gamma}_1(\theta) = \tilde{\mathbb{Z}}_\theta + \tilde{\mathbb{Z}}_\theta \zeta.$$

Since $\theta - 2 = 2 - 1/\theta$, then all ζ^q are in $\tilde{\Gamma}_1(\theta)$, and

$$\begin{aligned} \zeta^q \Gamma_1(\theta) &\subset \Gamma_1(\theta) + \left(\left\{ 0, \pm \frac{1}{\theta}, \pm \frac{2}{\theta} \right\} + \left\{ 0, \pm \frac{1}{\theta}, \pm \frac{2}{\theta} \right\} \zeta \right) \\ &\subset \tilde{\Gamma}_1(\theta) \equiv \tilde{\mathbb{Z}}_\theta + \tilde{\mathbb{Z}}_\theta \zeta, \end{aligned} \quad (11)$$

where $\tilde{\mathbb{Z}}_\theta = \mathbb{Z}_\theta + \{0, \pm 1/\theta, \pm 2/\theta\}$.

4.3. Translational properties

They are deduced from Eqs. (2) and (4). In Case 1,

$$\Gamma_q(\beta) + \Gamma_q(\beta) \subset \Gamma_q(\beta)/\beta^2$$

and in Case 2,

$$\Gamma_q(\beta) + \Gamma_q(\beta) \subset \tilde{\Gamma}_q(\beta).$$

5. A plane-group for beta-lattices

Since beta-lattices of the type $\Gamma_q(\beta)$ are not rotationally and translationally invariant, we shall enforce invariance by replacing the usual additive and multiplicative laws by the beta-addition and the quasi-multiplication.

5.1. A point-group for beta-lattices in the plane

Explicit calculations of internal rotation actions on $\Gamma_1(\beta)$, referred to as beta-rotations, are given in the appendix. Note that since the quasi-multiplication is not distributive with respect to beta-addition, we find several candidates for internal rotational operators on $\Gamma_1(\beta)$. The choice for the beta-rotations presented in the following proposition is driven by compatibility property. Other internal rotational operator *are not* compatible with Euclidean rotations!

We formally imitate the expressions of successive rotations given by Eq. (10), by replacing in the equations, + and - by \oplus and \ominus , and \times by \otimes , when necessary. Proposition 2 below defines the beta-rotations on $\Gamma_1(\beta)$.

Proposition 2.

- When $\beta = \tau$, with the notations of (9), the following 10 operators r_q , $q = 0, 1, \dots, 9$, leave $\Gamma_1(\tau)$ invariant:

$$r_q \odot (b_m + b_n \zeta) = \eta_q b_m \ominus \nu_q b_n + (\nu_q b_m \oplus (\eta_q + \tau \nu_q) b_n) \zeta.$$

- When $\beta = \delta$, the following operators leave $\Gamma_1(\delta)$ invariant:

$$r_1 \odot (b_m + b_n \zeta) = -b_n + (b_m \oplus \delta b_n \ominus b_n) \zeta = -b_n + b_{m+2n-2\rho_S(n)} \zeta,$$

$$\begin{aligned}
r_2 \odot (b_m + b_n \zeta) &= -(b_m \oplus \delta b_n \oplus b_n) + (\delta b_m \oplus b_m \oplus b_n) \zeta \\
&= -b_{m+2n-2\rho_S(n)} + b_{2m+n-2\rho_S(m)} \zeta, \\
r_3 \odot (b_m + b_n \zeta) &= -(\delta b_m \oplus b_m \oplus b_n) + b_m \zeta = -b_{2m+n-2\rho_S(m)} + b_m \zeta.
\end{aligned}$$

• When $\beta = \theta$, the following operators leave $\Gamma_1(\theta)$ invariant:

$$\begin{aligned}
r_1 \odot (b_m + b_n \zeta) &= -b_n + (b_m \oplus \theta b_n \oplus 2b_n) \zeta = -b_n + b_{m+2n-\rho_S(n)} \zeta, \\
r_2 \odot (b_m + b_n \zeta) &= -(b_m \oplus \theta b_n \oplus 2b_n) + (\theta b_m \oplus 2b_m \oplus 2b_n) \zeta \\
&= -b_{m+2n-\rho_S(n)} + b_{2m+2n-\rho_S(m)} \zeta, \\
r_3 \odot (b_m + b_n \zeta) &= -(\theta b_m \oplus 2b_m \oplus 2b_n) + (2b_m \oplus \theta b_n \oplus 2b_n) \zeta \\
&= -b_{2m+2n-\rho_S(m)} + b_{2m+2n-\rho_S(n)} \zeta, \\
r_4 \odot (b_m + b_n \zeta) &= -(2b_m \oplus \theta b_n \oplus 2b_n) + (\theta b_m \oplus 2b_m \oplus b_n) \zeta \\
&= -b_{2m+2n-\rho_S(m)} + b_{2m+n-\rho_S(n)} \zeta, \\
r_5 \odot (b_m + b_n \zeta) &= -(\theta b_m \oplus 2b_m \oplus b_n) + b_m \zeta = -b_{n+2m-\rho_S(m)} + b_m \zeta.
\end{aligned}$$

For $\beta = \tau, \delta$ or θ , let the composition rule of these operators on $\Gamma_1(\beta)$ be defined by

$$(rr') \odot z = r \odot (r' \odot z)$$

and denote by Id the identity and by ι the space inversion

$$\iota \odot z = -z.$$

Then, the composition rule $(r, r') \rightarrow rr'$ is associative and the following identities hold: $r_0 = Id$ and $r_{q+N/2} = \iota r_q$ for $q = 0, 1, \dots, (N/2) - 1$, where N is the symmetry order of β .

Lemma 5. Beta-rotations defined in Proposition 2 on $\Gamma_1(\beta)$ are compatible with rotations when β assumes one of the specified values τ, δ and θ .

Proof. We deduce from Eqs. (6) and (8) that beta-rotations have minimal distortion property with respect to ordinary rotations: let $z_{m,n} = b_m + b_n \zeta \in \Gamma_1(\beta)$, then

- $\beta = \tau$, $\zeta^q z_{m,n} - r_q \odot z_{m,n} \in \{0, \pm(1 - \frac{1}{\tau})\} + \{0, \pm(1 - \frac{1}{\tau})\} \zeta$,
- $\beta = \delta$, $\zeta^q z_{m,n} - r_q \odot z_{m,n} \in \{0, \pm(1 - \frac{1}{\delta}), \pm 2(1 - \frac{1}{\delta})\} + \{0, \pm(1 - \frac{1}{\delta}), \pm 2(1 - \frac{1}{\delta})\} \zeta$,
- $\beta = \theta$, $\zeta^q z_{m,n} - r_q \odot z_{m,n} \in \{0, \pm \frac{1}{\theta}, \pm \frac{2}{\theta}\} + \{0, \pm \frac{1}{\theta}, \pm \frac{2}{\theta}\} \zeta$.

Proposition 2 shows that beta-rotations can be decomposed in terms of beta-additions and quasi-multiplications. Compatibility of beta-rotation with euclidian rotation is thus a consequence of +-compatibility of beta-addition and \times -compatibility of quasi-multiplication. \square

Computing the composition of any two of such beta-rotations r_q yields the following important result.

Proposition 3. For $\beta = \tau, \delta$ and θ and for $N = 10, 8$ and 12 , respectively, let $\mathfrak{R}_N = \mathfrak{R}_N(\beta)$ denote the semi-group freely generated by all $r_q, q \in \{0, 1, \dots, N-1\}$. Among all beta-rotations, only $r_0, r_1, r_{N/2-1}, r_{N/2+1}, r_{N-1}$, and ι have their inverse in \mathfrak{R}_N .

Proof. The following identities are straightforwardly checked:

$$\begin{aligned} r_1 r_{N/2-1} &= r_{N/2-1} r_1 = r_{N/2+1} r_{N-1} = r_{N-1} r_{N/2-1} = \iota, \\ r_1 r_{N-1} &= r_{N-1} r_1 = r_{N/2-1} r_{N/2+1} = r_{N/2+1} r_{N/2-1} = r_0. \end{aligned}$$

A case study of all possible combinations of r_q shows that no other such operators are invertible. \square

An immediate consequence is the existence of a symmetry group for $\Gamma_1(\beta)$, i.e. a group of planar transformations leaving $\Gamma_1(\beta)$ invariant.

Theorem 1. For $\beta = \tau, \delta$ and θ , the group $\mathcal{R}_N = \mathcal{R}_N(\beta)$, freely generated by the four element set $\{r_0, \iota, r_1, r_{N/2-1}\}$, is a symmetry group for the beta-lattice $\Gamma_1(\beta)$. It is called the symmetry point-group of $\Gamma_1(\beta)$.

Proof. An easy computation shows that the elements of \mathcal{R}_N are invertible. Associativity of the law of internal composition of elements of \mathcal{R}_N is a consequence of Proposition 2. \square

5.2. A plane-group for beta-lattices $\Gamma_1(\beta)$

We now introduce into the present formalism the beta-translations acting on $\Gamma_1(\beta)$.

Proposition 4. Let $z_0 = b_{m_0} + b_{n_0} \zeta$ be an element of the beta-lattice $\Gamma_1(\beta)$. There corresponds to it the internal action $t_{z_0} : \Gamma_1(\beta) \mapsto \Gamma_1(\beta)$

$$t_{z_0}(z) = z \oplus z_0 \stackrel{\text{def}}{=} b_m \oplus b_{m_0} + (b_n \oplus b_{n_0}) \zeta = b_{m+m_0} + b_{n+n_0} \zeta.$$

The set of beta-translations forms an abelian group isomorphic to the beta-lattice $\Gamma_1(\beta)$ considered itself as a group for the law \oplus . For this reason it will be also denoted by $\Gamma_1(\beta)$.

Proof. The beta-translation is a simple two-dimensional generalization of the one-dimensional beta-addition. \square

As a direct generalization of one-dimensional beta-addition, it is obvious that beta-translation has minimal distortion property with respect to translation, and is compatible with it. Using Proposition 4, we come to the main result of this article.

Theorem 2. For $\beta = \tau, \delta$ and θ , and for $N = 10, 8$ and 12 , respectively, the group $\mathcal{S}_N = \mathcal{S}_N(\beta)$ freely generated by the five-element set $\{r_0, \iota, r_1, r_{(N/2)-1}, t_1\}$ is a symmetry group for the beta-lattice $\Gamma_1(\beta)$. This group is the semi-direct product of $\Gamma_1(\beta)$ and \mathcal{R}_N

$$\mathcal{S}_n = \Gamma_1(\beta) \rtimes \mathcal{R}_N$$

with the composition rule

$$(b, R)(b', R') = (b \oplus R \odot b', RR').$$

In the present context, \mathcal{S}_N is called the symmetry plane-group of $\Gamma_1(\beta)$.

The action of an element of \mathcal{S}_N on $\Gamma_1(\beta)$ is thus defined as

$$(b, R) \cdot z = b \oplus R \odot z = t_b(R \odot z) \in \Gamma_1(\beta).$$

Proof. An easy computation shows that the elements of \mathcal{S}_N are invertible. Associativity of the law of internal composition of elements of \mathcal{B}_N is a consequence of Proposition 2 and of Theorem 1. \square

5.3. Tile transformations using internal operations on $\Gamma_1(\tau)$

We would like to illustrate the action of \mathcal{S}_n on $\Gamma_1(\beta)$, in the case of τ , by showing how a tile of $\Gamma_1(\tau)$ is transformed under the action of an element of \mathcal{S}_{10} .

Let $z = b_m + b_n \zeta \in \Gamma_1(\tau)$. An elementary quadrilateral tile on z is the following:

$$\mathbf{T}(z) = \{z, z \oplus 1, z \oplus \zeta, z \oplus (1 + \zeta)\}.$$

From the definition of $\Gamma_1(\tau)$, we trivially see that there exist four kinds of elementary tiles, which we shall denote by LL , LS , SL and SS , as a reference to the length of their edges (see Fig. 5).

In case of a translation operation by z_0 , t_{z_0} , the elementary quadrilateral tile $\mathbf{T}(z)$ is transformed into another elementary quadrilateral tile, whether of the same kind or of another kind, according to

$$t_{z_0}(\mathbf{T}(z)) = \mathbf{T}(z \oplus z_0) = \{z_0 \oplus z, z_0 \oplus z \oplus 1, z_0 \oplus z \oplus \zeta, z_0 \oplus z \oplus (1 + \zeta)\}.$$

Another interesting transformation arises when one applies the beta-rotation operator r_1 on $\mathbf{T}(z)$ and around one of the vertex of $\mathbf{T}(z)$. For instance, the beta-rotation around z is given by

$$t_z(r_1 \odot t_{-z}(\mathbf{T}(z))) = \{z, z \oplus \zeta, z \oplus (-1 + \tau\zeta), z \oplus (-1 + \tau^2\zeta)\}.$$

Examples of such rotation operations are displayed on Fig. 6. This operation not only rotates, but distorts the tiles, in general. Therefore, the beta-rotated tile is not elementary anymore.



Fig. 5. Elementary quadrilateral tiles for the τ -lattice $\Gamma_1(\tau)$. From left to right: LL , LS , SL , SS . See also Fig. 2.




Fig. 6. Rotation operator r_1 applied to elementary tiles of the τ -lattice $I_1(\tau)$, $\mathbf{T}(0)$, $\mathbf{T}(1)$ (up), $\mathbf{T}(\zeta)$ and $\mathbf{T}(1 + \zeta)$ (down). Note how the tiles are deformed, by this operation, in order for the vertices to remain in $I_1(\tau)$. The arrows indicate the vertices of the new tile in which are mapped the vertices of the original tile.

6. Asymptotic properties

An interesting feature of beta-lattices is that they behave like lattices asymptotically.

Lemma 6. *The asymptotic behavior of the counting function ρ_S is given by*

$$\begin{aligned} \rho_S(n) &\underset{|n| \rightarrow \infty}{\approx} \left(1 - \frac{1}{\beta}\right) \frac{n}{a} & (\text{Case 1}), \\ \rho_S(n) &\underset{|n| \rightarrow \infty}{\approx} \frac{n}{\beta} & (\text{Case 2}). \end{aligned}$$

Proof. *Case 1:* The proof is based on the development of integers in the linear system $U = (u_q)_{q \in \mathbb{N}}$. We have $n = \sum_{i=0}^k u_i d_i$. Then $\rho_S(n) = \sum_{i=0}^k \rho_S(u_i) d_i = \sum_{i=0}^k (u_i/a)(1 - u_{i-1}/u_i) d_i$. When $n \rightarrow \infty$ we know that $u_{i-1}/u_i \rightarrow 1/\beta$ and $\rho_S(n) \approx (1/a)(1 - 1/\beta) \sum_{i=0}^k u_i d_i = (n/a)(1 - 1/\beta)$, as $n \rightarrow \infty$.

Case 2: As in the first case, the proof is based on the development of integers in (u_q) : $n = \sum_{i=0}^k u_i d_i$, $\rho_S(n) = \sum_{i=0}^k \rho_S(u_i) d_i = \sum_{i=0}^k u_{i-1} d_i = \sum_{i=0}^k (u_i/a)(1 + u_{i-2}/u_i) d_i$. When $n \rightarrow \infty$ we know that $u_{i-2}/u_i = (u_{i-2}/u_{i-1})(u_{i-1}/u_i) = 1/\beta^2 = (a/\beta) - 1$. Therefore $\rho_S(n) \approx (1/\beta) \sum_{i=0}^k u_i d_i = n/\beta$, as $n \rightarrow \infty$. \square

Lemma 6 tells us what is the asymptotic behavior of beta-integers for large n , and of the multiplication \otimes for large m and n . From Eq. (7) and Lemma 6 is deduced the following result.

Proposition 5. *Let β be a quadratic PV unit number. Then the following asymptotic behaviour of beta-integers holds true*

$$\begin{aligned} b_n &\underset{|n| \rightarrow \infty}{\approx} \gamma n, \\ b_m \otimes b_n &\underset{|m|, |n| \rightarrow \infty}{\approx} \gamma^2 mn, \end{aligned}$$

where

$$\gamma = \begin{cases} 1 - \frac{1}{a}(1 - \frac{1}{\beta})^2 = \frac{(a+2)\beta - a^2 - a - 2}{a} & (\text{Case 1}), \\ 1 - \frac{1}{\beta^2} = a(\beta - a) + 2 & (\text{Case 2}). \end{cases}$$

Proof. *Case 1:* Any beta-integer b_n can be written $b_n = n - \rho_S(n)(1 - 1/\beta)$. When n becomes large, we can replace $\rho_S(n)$ by its asymptotic value. We then have $b_n \approx n(1 - (1/a)(1 - 1/\beta)^2) = \gamma n$.

Case 2: In the same fashion, we have $b_n = n - \rho_S(n)1/\beta$, and by replacing $\rho_S(n)$ by its asymptotic value for large n we obtain $b_n = n(1 - 1/\beta^2) = \gamma n$.

The second part of the proposition is a direct consequence of the first part. \square

We then *almost* recover the definition of multiplication we were thinking about at the beginning of Section 3.5, left alone that in both cases we have a contraction of the resulting index by a factor $\gamma < 1$. We should notice that the multiplication \otimes is *asymptotically* associative and distributive with respect to the addition \oplus . In this sense, we can say that \mathbb{Z}_β is asymptotically a ring

$$\begin{aligned} b_m \otimes (b_n \oplus b_p) - (b_m \otimes b_n) \oplus (b_m \otimes b_p) &\underset{|m|, |n|, |p| \rightarrow \infty}{\approx} 0, \\ b_m \otimes (b_n \otimes b_p) - (b_m \otimes b_n) \otimes b_p &\underset{|m|, |n|, |p| \rightarrow \infty}{\approx} 0. \end{aligned}$$

Note that m , n and p must be such that $m \pm n$ and $m \pm p$ are large numbers, otherwise the above equations are not true.

Consequently, we compute the asymptotic behavior of rotational internal laws of beta-lattices, as defined in Section 5.1 in the studied cases.

- When $\beta = \tau$, we have for invertible operators

$$\begin{aligned} r_1 \odot (b_m + b_n \zeta) &\underset{|m|, |n| \rightarrow \infty}{\approx} \gamma(-n + (m + \tau n)\zeta), \\ r_4 \odot (b_m + b_n \zeta) &\underset{|m|, |n| \rightarrow \infty}{\approx} \gamma(-\tau m - n - m\zeta). \end{aligned}$$

- When $\beta = \delta$, we have for invertible operators

$$\begin{aligned} r_1 \odot (b_m + b_n \zeta) &\underset{|m|, |n| \rightarrow \infty}{\approx} \gamma(-n + (m + (\delta - 1)n)\zeta), \\ r_3 \odot (b_m + b_n \zeta) &\underset{|m|, |n| \rightarrow \infty}{\approx} \gamma(-(\delta - 1)m - n + m\zeta). \end{aligned}$$

- When $\beta = \theta$, we have for invertible operators

$$\begin{aligned} r_1 \odot (b_m + b_n \zeta) &\underset{|m|, |n| \rightarrow \infty}{\approx} \gamma(-n + (m + (\theta - 2)n)\zeta), \\ r_5 \odot (b_m + b_n \zeta) &\underset{|m|, |n| \rightarrow \infty}{\approx} \gamma(-(\theta - 2)m - n + m\zeta). \end{aligned}$$

At this point one should be aware that these asymptotic beta-rotations are equivalent to rotations for large $|m|$ and $|n|$, and an easy computation shows that for $z_{m,n} \in I_1(\beta)$

$$\begin{aligned} \zeta z_{m,n} - r_1 \odot z_{m,n} &\underset{|m|, |n| \rightarrow \infty}{\approx} 0, \\ \zeta^{N/2-1} z_{m,n} - r_{N/2-1} \odot z_{m,n} &\underset{|m|, |n| \rightarrow \infty}{\approx} 0 \end{aligned}$$

with $N = 10, 8$ and 12 .

7. Conclusion

The main result of this article is the construction of a symmetry plane-group for beta-lattices for three quadratic PV units. Though preliminary, this study shows the richness of the beta-lattices as far as all the operations of the plane-group can be made arithmetically explicit. Many questions seem to be open, such as the number of possible plane-groups leaving a beta-lattice invariant. Another important issue is to determine whether there is or not a metric left invariant under the action of such groups. It has been shown that a large class of point sets, such as model sets, can be embedded in beta-lattices [8]. A question related to distortion of distances is the action of beta-rotations and beta-translations over a point set embedded in a beta-lattice and over the tiling associated to this point set. The point group $\mathcal{R}_N(\beta)$ also deserves to be carefully studied. The link between beta-lattices and the class of finitely generated modules over ordered rings would deserve to be handled nicely in the framework of the Artin-Schreier theory. The case of PV of higher degree remains open. The present contribution shows the potentiality offered by a class of beta-lattices to provide structure models of more general quasiperiodic crystals, and possibly to predict new crystals.

Appendix A. Explicit internal beta-rotations actions on beta-lattices

In this section, we make the beta-rotation explicit for the quasicrystallographic numbers τ , δ , and θ , and for all the corresponding q , the remaining beta-rotation being deduced from them by combining with space inversion. We give the resulting integer indexes in terms of m, n , and the counting function ρ_S as all involved relations have been introduced in Eqs. (6) and (7).

A.1. Case of the τ -lattice $\Gamma_1(\tau)$

$$\begin{aligned}
r_1 \odot (b_m + b_n \zeta) &= b_{-n} + b_{m+2n-\rho_S(n)} \zeta, \\
r_2 \odot (b_m + b_n \zeta) &= b_{-m-2n+\rho_S(n)} + b_{2(m+n)-\rho_S(m)-\rho_S(n)} \zeta, \\
r_3 \odot (b_m + b_n \zeta) &= b_{-2(m+n)+\rho_S(m)+\rho_S(n)} + b_{2m+n-\rho_S(m)} \zeta, \\
r_4 \odot (b_m + b_n \zeta) &= b_{-2m-n+\rho_S(m)} + b_m \zeta.
\end{aligned}$$

A.2. Case of the δ -lattice $\Gamma_1(\delta)$

For the δ -rotations we would like to proceed to the formal imitation of Eq. (10) as in the case of τ . The case of δ however is slightly more complicated since $\eta_q b_n$ and $v_q b_n$ are not in \mathbb{Z}_δ . When we compute the rotation of an arbitrary element of $\Gamma_1(\delta)$, we need to determine the value of $(\delta - 1)b_n$, which is of course not a δ -integer in the general case. Recall that \otimes is not distributive with respect to \oplus . Therefore, we have to replace $(\delta - 1)b_n$ either by $(\delta \oplus 1) \otimes b_n = b_{2n}$ or by $\delta b_n \oplus b_n = b_{2n-\rho_S(n)}$ (recall that from self-similarity of δ -integers we have $\delta \otimes b_n = \delta b_n$). We then have to make a choice about which operation to choose to build the point-group of $\Gamma_1(\delta)$. We chose to replace $(\delta - 1)b_n$ by $\delta b_n \oplus b_n$ in Section 5, since this case satisfies the compatibility property. Other operations may be interesting. For example, the other internal rotation laws do not satisfy the compatibility property and do not have the same asymptotic behavior.

$$\begin{aligned}
r_1 \odot (b_m + b_n \zeta) &= \begin{cases} -b_n + b_{m+2n-\rho_S(n)} \zeta, \\ -b_n + b_{m+2n} \zeta, \end{cases} \\
r_2 \odot (b_m + b_n \zeta) &= \begin{cases} -b_{m+2n-\rho_S(n)} + b_{2m+n-\rho_S(m)} \zeta, \\ -b_{m+2n} + b_{2m+n-\rho_S(m)} \zeta, \\ -b_{m+2n-\rho_S(n)} + b_{2m+n} \zeta, \\ -b_{m+2n} + b_{2m+n} \zeta, \end{cases} \\
r_3 \odot (b_m + b_n \zeta) &= \begin{cases} -b_{2m+n-\rho_S(m)} + b_m \zeta, \\ -b_{2m+n} + b_m \zeta. \end{cases}
\end{aligned}$$

A.3. Case of the θ -lattice $\Gamma_1(\theta)$

As in the case of the δ -lattice, we have to decide which operation to use to build the point-group of $\Gamma_1(\theta)$ because of the factor $(\theta - 2)b_n$, introduced in the computation of rotations of $\Gamma_1(\theta)$. Once again, we have replaced $(\theta - 2)b_n$ by $\theta b_n \oplus 2 \otimes b_n = b_{2n-\rho_S(n)}$ in Section 5.2. We give now all possibilities.

$$r_1 \odot (b_m + b_n \zeta) = \begin{cases} -b_n + b_{m+2n-\rho_S(n)} \zeta, \\ -b_n + b_{m+2n} \zeta, \end{cases}$$

$$r_2 \odot (b_m + b_n \zeta) = \begin{cases} -b_{m+2n-\rho_S(n)} + b_{2m+2n-\rho_S(m)} \zeta, \\ -b_{m+2n} + b_{2m+2n-\rho_S(m)} \zeta, \\ -b_{m+2n-\rho_S(n)} + b_{2m+2n} \zeta, \\ -b_{m+2n} + b_{2m+2n} \zeta, \end{cases}$$

$$r_3 \odot (b_m + b_n \zeta) = \begin{cases} -b_{2m+2n-\rho_S(m)} + b_{2m+2n-\rho_S(n)} \zeta, \\ -b_{2m+n} + b_{2m+2n-\rho_S(m)} \zeta, \\ -b_{2m+2n-\rho_S(m)} + b_{2m+2n} \zeta, \\ -b_{2m+2n} + b_{2m+2n} \zeta, \end{cases}$$

$$r_4 \odot (b_m + b_n \zeta) = \begin{cases} b_{2m+2n-\rho_S(n)} + b_{n+2m-\rho_S(m)} \zeta, \\ b_{2m+2n} + b_{n+2m-\rho_S(m)} \zeta, \\ b_{2m+2n-\rho_S(n)} + b_{n+2m} \zeta, \\ b_{2m+2n} + b_{n+2m} \zeta, \end{cases}$$

$$r_5 \odot (b_m + b_n \zeta) = \begin{cases} -b_{2m+n-\rho_S(m)} + b_m \zeta, \\ -b_{2m+n} + b_m \zeta. \end{cases}$$

References

- [1] D. Barache, B. Champagne, J.P. Gazeau, Pisot-Cyclotomic quasilattices and their symmetry semi-groups, in: J. Patera (Ed.), *Quasicrystals and Discrete Geometry*, Fields Institute Monograph Series, Vol. 10, American Mathematical Society, Providence, RI, 1998.
- [2] A. Bertrand, Développements en base de Pisot et répartition modulo 1, *C. R. Acad. Sci., Paris* 285 (1977) 419–421.
- [3] Č. Burdík, Ch. Frougny, J.P. Gazeau, R. Krejcar, Beta-integers as natural counting systems for quasicrystals, *J. Phys. A: Math. Gen.* 31 (1998) 6449–6472.
- [4] Č. Burdík, Ch. Frougny, J.P. Gazeau, R. Krejcar, Beta-integers as a group, in: *Dynamical Systems: From Crystal to Chaos*, World Scientific, Singapore, 2000, pp. 125–136.
- [5] S. Fabre, Substitutions et β -systèmes de numération, *Theoret. Comput. Sci.* 137 (1995) 219–236.
- [6] Ch. Frougny, J.P. Gazeau, R. Krejcar, Additive and multiplicative properties of point-sets based on beta-integers, *Theoret. Comput. Sci.* 303 (2003) 491–516.
- [7] Ch. Frougny, B. Solomyak, Finite beta-expansions, *Ergodic Theory Dynamical Systems* 12 (1992) 713–723.
- [8] J.P. Gazeau, R. Krejcar, in: F. Axel, F. Denoyer, J.-P. Gazeau (Eds.), *From Quasicrystals to More Complex Systems*, Les éditions de physique, Springer, Berlin, 2000.
- [9] P.J. Grabner, A. Pethő, R.F. Tichy, G.J. Wöginger, Associativity of recurrence multiplication, *Appl. Math. Lett.* 7 (1994) 85–90.
- [10] International Union of Crystallography, Report of the Executive Committee for 1991, *Acta Crystallogr. A* 48 (1992) 922–946.
- [11] C. Janot, *Quasicrystals, A Primer*, Oxford University Press, Oxford, 1996.
- [12] J. Lagarias, Geometric models for quasicrystals I. Delone sets of finite type, *Discrete Comput. Geom.* 21 (1999) 161–191.
- [13] T.Y. Lam, *A First Course in Noncommutative Rings*, Springer, Berlin, 1991.
- [14] A. Messaoudi, Généralisation de la multiplication de Fibonacci, *Math. Slovaca* 50 (2000) 135–148.
- [15] A. Messaoudi, Tribonacci multiplication, *Appl. Math. Lett.* 15 (2002) 981–985.
- [16] Y. Meyer, Nombres de Pisot, nombres de Salem et analyse harmonique, in: *Lecture Notes in Mathematics*, Vol. 117, Springer, Berlin, 1970.
- [17] Y. Meyer, *Algebraic Numbers and Harmonic Analysis*, North-Holland, Amsterdam, 1972.
- [18] Y. Meyer, Quasicrystals, diophantine approximation and algebraic numbers, in: F. Axel, D. Gratias (Eds.), *Beyond Quasicrystals*, Les éditions de physique, Springer, Berlin, 1995.

- [19] R. Moody, Meyer sets and their duals, in: R. Moody (Ed.), *The Mathematics of Long-Range Aperiodic Order*, Kluwer, Dordrecht, 1997.
- [20] R. Moody, Model sets: a survey, in: F. Axel, F. Denoyer, J.-P. Gazeau (Eds.), *From Quasicrystals to More Complex Systems*, Les éditions de physique, Springer, Berlin, 2000.
- [21] G. Muraz, J.-L. Verger-Gaugry, On lower bounds of the density of packings of equal spheres of \mathbb{R}^n , preprint Institut Fourier, Vol. 580, 2003.
- [22] G. Muraz, J.-L. Verger-Gaugry, On a metric on the set of uniformly discrete sets of \mathbb{R}^n invariant by the group of rigid motions and densest packings of equal spheres, preprint Institut Fourier, Vol. 622, 2003.
- [23] W. Parry, On the β -expansions of real numbers, *Acta Math. Acad. Sci. Hungar.* 11 (1960) 401–416.
- [24] A. Rényi, Representations for real numbers and their ergodic properties, *Acta Math. Acad. Sci. Hungar.* 8 (1957) 477–493.
- [25] R.L.E. Schwarzenberger, *N-dimensional Crystallography*, Pitman Publishing Limited, London, 1980.
- [26] W.P. Thurston, *Groups, tilings, and finite state automata*, Geometry supercomputer Project Research Report GCG1, University of Minnesota, 1989.
- [27] J.-L. Verger-Gaugry, J.P. Gazeau, Geometric study of the beta-integers for a Perron number and mathematical quasicrystals, *J. Th. Nombres Bordeaux* 16, fasc. 1 (2004).