



HAL
open science

Quantum and Semiquantum Pseudometrics and applications

François Golse, Thierry Paul

► **To cite this version:**

François Golse, Thierry Paul. Quantum and Semiquantum Pseudometrics and applications. Journal of Functional Analysis, In press. hal-03136855

HAL Id: hal-03136855

<https://hal.science/hal-03136855v1>

Submitted on 9 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

QUANTUM AND SEMIQUANTUM PSEUDOMETRICS AND APPLICATIONS

FRANÇOIS GOLSE AND THIERRY PAUL

ABSTRACT. We establish a Kantorovich duality for the pseudometric \mathcal{E}_\hbar introduced in [F. Golse, T. Paul, Arch. Rational Mech. Anal. **223** (2017), 57–94], obtained from the usual Monge-Kantorovich distance $\text{dist}_{\text{MK},2}$ between classical densities by quantization of one side of the two densities involved. We show several type of inequalities comparing $\text{dist}_{\text{MK},2}$, \mathcal{E}_\hbar and MK_\hbar , a full quantum analogue of $\text{dist}_{\text{MK},2}$ introduced in [F. Golse, C. Mouhot, T. Paul, Commun. Math. Phys. **343** (2016), 165–205], including an up to \hbar triangle inequality for MK_\hbar . Finally, we show that, when nice optimal Kantorovich potentials exist for \mathcal{E}_\hbar , optimal couplings induce classical/quantum optimal transports and the potentials are linked by a semiquantum Legendre type transform.

CONTENTS

1. Introduction and statement of some main results	1
2. Preliminaries	3
2.1. Monotone Convergence	3
2.2. Finite Energy Condition	5
2.3. Energy and Partial Trace	7
3. Couplings	8
4. Triangle Inequalities	10
5. Applications	18
6. Kantorovich duality for \mathcal{E}_\hbar	19
7. Applications of duality for \mathcal{E}_\hbar I: inequalities between MK_\hbar , \mathcal{E}_\hbar and $\text{dist}_{\text{MK},2}$.	28
8. Applications of duality for \mathcal{E}_\hbar II: “triangle” inequalities	28
9. Applications of duality for \mathcal{E}_\hbar III: Classical/quantum optimal transport and semiquantum Legendre transform	30
9.1. A classical/quantum optimal transport	30
9.2. A semiquantum Legendre transform	32
References	33

1. INTRODUCTION AND STATEMENT OF SOME MAIN RESULTS

The Monge-Kantorovich distance, also called Wasserstein distance, of exponent two on the phase-space $T^*\mathbf{R}^d \sim \mathbf{R}^{2d}$ is defined, for two probability measures by

$$(1) \quad \text{dist}_{\text{MK},2}(\mu, \nu)^2 = \inf_{\pi \in \pi[\mu, \nu]} \int_{\mathbf{R}^{2d} \times \mathbf{R}^{2d}} ((q - q')^2 + (p - p')^2) \pi(dq dp, dq' dp')$$

Date: February 9, 2021.

where $\pi[\mu, \nu]$ is the set of couplings π of μ, ν , i.e. the set of probability measures π on $\mathbf{R}^{2d} \times \mathbf{R}^{2d}$ such that for all test functions $a, b \in C_c(\mathbf{R}^{2d})$ we have that

$$\int_{\mathbf{R}^{2d} \times \mathbf{R}^{2d}} (a(q, p) + b(q', p')) \pi(dq dp, dq' dp') = \int_{\mathbf{R}^{2d}} (a(q, p) \mu(dq dp) + b(q', p') \nu(dq' dp')).$$

Among the many properties of $\text{dist}_{\text{MK},2}$, let us mention the Kantorovich duality which stipulates that

$$(2) \quad \text{dist}_{\text{MK},2}(\mu, \nu)^2 = \max_{\substack{a, b \in C_b(\mathbf{R}^d) \\ a(q, p) + b(q', p') \leq (q - q')^2 + (p - p')^2}} \int_{\mathbf{R}^{2d}} (a(q, p) \mu(dq dp) + b(q, p) \nu(dq dp)),$$

and the Knott-Smith-Brenier Theorem which says that, under certain conditions on μ, ν , any coupling π_{op} satisfying

$$(3) \quad \text{dist}_{\text{MK},2}(\mu, \nu)^2 = \int_{\mathbf{R}^{2d} \times \mathbf{R}^{2d}} ((q - q')^2 + (p - p')^2) \pi_{op}(dq dp, dq' dp')$$

is supported in the graph of the convex function $\frac{1}{2}(q^2 + p^2) - a_{op}(q, p)$ where a_{op} is an optimal function such that a_{op}, b_{op} provide the max in (2) for some b_{op} . Finally, $\frac{1}{2}(q^2 + p^2) - a_{op}(q, p)$ and $\frac{1}{2}(q^2 + p^2) - b_{op}(q, p)$ are proven to be the Legendre transform of each other.

A quantum version of $\text{dist}_{\text{MK},2}$ was proposed in [6] following the general rules of quantization consisting in replacing

- probability measures μ, ν on phase-space $T^*\mathbf{R}^d$ by quantum states R, S , i.e. density operators, i.e. positive trace one operators on $L^2(\mathbf{R}^d)$
- $\int_{T^*\mathbf{R}^d}$ by $\text{trace}_{L^2(\mathbf{R}^d)}$
- couplings of μ, ν by density operators Π on $L^2(\mathbf{R}^d) \otimes L^2(\mathbf{R}^d)$ such that, for any bounded operators A, B , $\text{trace}_{L^2(\mathbf{R}^d) \otimes L^2(\mathbf{R}^d)} (A \otimes I) \Pi = \text{trace}_{L^2(\mathbf{R}^d)} A S$ and $\text{trace}_{L^2(\mathbf{R}^d) \otimes L^2(\mathbf{R}^d)} (I \otimes B) \Pi = \text{trace}_{L^2(\mathbf{R}^d)} B R$.
- the cost function $(q - q')^2 + (p - p')^2$ by its Weyl pseudodifferential quantization $C = (x - x')^2 + (-i\hbar \nabla_x + i\hbar \nabla_{x'})^2$ on $L^2(\mathbf{R}^d \times \mathbf{R}^d)$.

These considerations lead to the definition, for two density operators R, S on $L^2(\mathbf{R}^d)$,

$$(4) \quad MK_{\hbar}(R, S)^2 = \inf_{\Pi \text{ coupling } R \text{ and } S} \text{trace}_{L^2(\mathbf{R}^d) \otimes L^2(\mathbf{R}^d)} C \Pi.$$

The pseudometric MK_{\hbar} has been extensively studied in [6], with applications to the study of the quantum mean-field limit uniformly in \hbar , used in [2] for quantum optimal transport considerations and applied in [3] for the quantum bipartite matching problem. In particular, a Kantorovich duality was proven for MK_{\hbar} in [2] expressed as the following identity

$$(5) \quad MK_{\hbar}(R, S)^2 = \sup_{\substack{A=A^*, B=B^* \in \mathcal{L}(L^2(\mathbf{R}^d)) \\ A \otimes I + I \otimes B \leq C}} \text{trace}(AR + BS)$$

and the supremum was proven to be attained for two operators \bar{A}, \bar{B} defined respectively on two Gelfand triplets surrounding $L^2(\mathbf{R}^d)$ (see [2])

Though MK_{\hbar} is symmetric in its argument, it is not a distance as one can easily show ([6]) that $MK_{\hbar}^2 \geq 2d\hbar$. Nevertheless, one of the main result of this article

will be to prove the following (approximate) triangle inequality, valid for density operators R, S, T (see Theorem 8.1 (iii) below)

$$(6) \quad MK_{\hbar}(R, T) \leq MK_{\hbar}(R, S) + MK_{\hbar}(S, T) + d\hbar.$$

Actually, (6) is proved by using a kind of ‘‘semiquantum’’ generalisation of $\text{dist}_{\text{MK},2}$, defined in [7] and constructed by, roughly speaking, applying the quantization rule aforementioned to only one on the two parts involved in $\text{dist}_{\text{MK},2}(\mu, \nu)$:

for f probability density on \mathbf{R}^{2d} and R density operator on $L^2(\mathbf{R}^d)$ we define

$$(7) \quad \mathcal{E}_{\hbar}(f, R)^2 = \sup_{\substack{\Pi(q,p) \\ \text{density operators} \\ \text{such that} \\ \text{trace} \Pi(q,p) = f(q,p) \\ \text{and } \int_{\mathbf{R}^{2d}} \Pi(q,p) = R}} \int_{\mathbf{R}^{2d}} \text{trace}_{L^2(\mathbf{R}^d, dx)} ((q-x)^2 + (p+i\hbar\nabla_x)^2) \Pi(q,p) dq dp.$$

The pseudometric \mathcal{E}_{\hbar} has been used in [7] in order to derive several results concerning the quantum, uniform in \hbar , mean-field derivation and in [7, 8] for semiclassical propagation estimates involving low regularity of the potential and the initial data (in particular with respect to the dimension, i.e. also to the number of particles present in the quantum evolution).

In the present paper, we prove a Kantorovich duality for \mathcal{E}_{\hbar} (Section 6, Theorem 6.1), namely

$$(8) \quad \mathcal{E}_{\hbar}(f, R)^2 = \sup_{\substack{a \in C_b(\mathbf{R}^{2d}), B \in \mathcal{L}(L^2(\mathbf{R}^d)) \\ a(q,p) + B \leq (q-x)^2 + (p+i\hbar\nabla_x)^2}} \int_{\mathbf{R}^{2d}} a(q,p) f(q,p) dq dp + \text{trace}_{L^2(\mathbf{R}^d)} BR,$$

and then apply this duality to derive inequalities, such as (6), involving MK_{\hbar} , \mathcal{E}_{\hbar} and $\text{dist}_{\text{MK},2}$, Theorems 7.1 and 8.1.

In the last section of the paper, Section 9, we investigate the semiquantum analogue of the Knott-Smith-Brenier Theorem and a semiquantum analogue of the Legendre transform: if

$$\mathcal{E}_{\hbar}(f, R)^2 = \int_{\mathbf{R}^{2d}} a_{op}(q,p) f(q,p) dq dp + \text{trace}_{L^2(\mathbf{R}^d)} B_{op} R,$$

then $a(q,p) := \frac{1}{2}(p^2 + q^2 - a_{op}(q,p))$ is the semiquantum-Legendre transform of $B := \frac{1}{2}(-\nabla_x^2 + x^2 - B_{op})$, in the sense that

$$a(q,p) = \sup_{\phi \in \text{Dom}(B)} (q \cdot \langle \phi | x | \phi \rangle + p \cdot \langle \phi | -i\hbar\nabla_x | \phi \rangle - \langle \phi | B | \phi \rangle).$$

2. PRELIMINARIES

We have gathered together in this section some functional analytic remarks used repeatedly in the sequel.

2.1. Monotone Convergence. We recall the analogue of the Beppo Levi monotone convergence theorem for operators in the form convenient for our purpose.

Let \mathcal{H} be a separable Hilbert space and $0 \leq T = T^* \in \mathcal{L}(\mathfrak{H})$. For each complete orthonormal system $(e_j)_{j \geq 1}$ of \mathcal{H} , set

$$\text{trace}_{\mathcal{H}}(T) = \|T\|_1 := \sum_{j \geq 1} \langle e_j | T | e_j \rangle \in [0, +\infty].$$

See Theorem 2.14 in [13]; in particular the expression on the last right hand side of these equalities is independent of the complete orthonormal system $(e_j)_{j \geq 1}$. Then

$$T \in \mathcal{L}^1(\mathfrak{H}) \iff \|T\|_1 < \infty.$$

Lemma 2.1 (Monotone convergence). *Consider a sequence $T_n = T_n^* \in \mathcal{L}^1(\mathcal{H})$ such that*

$$0 \leq T_1 \leq T_2 \leq \dots \leq T_n \leq \dots, \quad \text{and } \sup_{n \geq 1} \langle x|T_n|x \rangle < \infty \text{ for all } x \in \mathcal{H}.$$

Then

- (a) there exists $T = T^* \in \mathcal{L}(\mathcal{H})$ such that $T_n \rightarrow T$ weakly as $n \rightarrow \infty$, and
- (b) $\text{trace}_{\mathcal{H}}(T_n) \rightarrow \text{trace}_{\mathcal{H}}(T)$ as $n \rightarrow \infty$.

Proof. Since the sequence $\langle x|T_n|x \rangle \in [0, +\infty)$ is nondecreasing for each $x \in \mathcal{H}$,

$$\langle x|T_n|x \rangle \rightarrow \sup_{n \geq 1} \langle x|T_n|x \rangle =: q(x) \in [0, +\infty) \quad \text{for all } x \in \mathcal{H}$$

as $n \rightarrow \infty$. Hence

$$\langle x|T_n|y \rangle = \langle y|T_n|x \rangle \rightarrow \frac{1}{4}(q(x+y) - q(x-y) + iq(x-iy) - iq(x+iy)) =: b(x, y) \in \mathbf{C}$$

as $n \rightarrow +\infty$. By construction, b is a nonnegative sesquilinear form on \mathcal{H} .

Consider, for each $k \geq 0$,

$$F_k := \{x \in \mathcal{H} \text{ s.t. } \langle x|T_n|x \rangle \leq k \text{ for each } n \geq 1\}.$$

The set F_k is closed for each $k \geq 0$, being the intersection of the closed sets defined by the inequality $\langle x|T_n|x \rangle \leq k$ as $n \geq 1$. Since the sequence $\langle x|T_n|x \rangle$ is bounded for each $x \in \mathcal{H}$,

$$\bigcup_{k \geq 0} F_k = \mathcal{H}.$$

Applying Baire's theorem shows that there exists $N \geq 0$ such that $F_N \overset{\circ}{\neq} \emptyset$. In other words, there exists $r > 0$ and $x_0 \in \mathcal{H}$ such that

$$|x - x_0| \leq r \implies |\langle x|T_n|x \rangle| \leq N \text{ for all } n \geq 1.$$

By linearity and positivity of T_n , this implies

$$|\langle z|T_n|z \rangle| \leq \frac{2}{r}(M + N)\|z\|^2 \text{ for all } n \geq 1, \quad \text{with } M := \sup_{n \geq 1} \langle x_0|T_n|x_0 \rangle.$$

In particular

$$\sup_{|z| \leq 1} q(z) \leq \frac{2}{r}(M + N), \quad \text{so that } |b(x, y)| \leq \frac{2}{r}(M + N)\|x\|_{\mathcal{H}}\|y\|_{\mathcal{H}}$$

for each $x, y \in \mathcal{H}$ by the Cauchy-Schwarz inequality. By the Riesz representation theorem, there exists $T \in \mathcal{L}(\mathcal{H})$ such that

$$T = T^* \geq 0, \quad \text{and } b(x, y) = \langle x|T|y \rangle.$$

This proves (a). Observe that $T \geq T_n$ for each $n \geq 1$, so that

$$\sup_{n \geq 1} \text{trace}_{\mathcal{H}}(T_n) \leq \text{trace}_{\mathcal{H}}(T).$$

In particular

$$\sup_{n \geq 1} \text{trace}_{\mathcal{H}}(T_n) = +\infty \implies \text{trace}_{\mathcal{H}}(T) = +\infty.$$

Since the sequence $\text{trace}_{\mathcal{H}}(T_n)$ is nondecreasing,

$$\text{trace}_{\mathcal{H}}(T_n) \rightarrow \sup_{n \geq 1} \text{trace}_{\mathcal{H}}(T_n) \quad \text{as } n \rightarrow \infty.$$

By the noncommutative variant of Fatou's lemma (Theorem 2.7 (d) in [13]),

$$\sup_{n \geq 1} \text{trace}_{\mathcal{H}}(T_n) < \infty \implies T \in \mathcal{L}^1(\mathcal{H}) \text{ and } \text{trace}_{\mathcal{H}}(T) \leq \sup_{n \geq 1} \text{trace}_{\mathcal{H}}(T_n).$$

Since the opposite inequality is already known to hold, this proves (b). \square

Here is a convenient variant of this lemma.

Corollary 2.2. *Consider a sequence $T_n = T_n^* \in \mathcal{L}^1(\mathcal{H})$ such that*

$$0 \leq T_1 \leq T_2 \leq \dots \leq T_n \leq \dots, \quad \text{and } \sup_{n \geq 1} \text{trace}_{\mathcal{H}}(T_n) < \infty.$$

Then there exists $T \in \mathcal{L}^1(\mathfrak{H})$ such that $T_n \rightarrow T$ weakly as $n \rightarrow \infty$, and

$$T = T^* \geq 0, \quad \text{and } \text{trace}_{\mathcal{H}}(T) = \lim_{n \rightarrow \infty} \text{trace}_{\mathcal{H}}(T_n).$$

Proof. Since any $x \in \mathcal{H} \setminus \{0\}$ can be normalized and completed into a complete orthonormal system of \mathcal{H} , one has

$$\sup_{n \geq 1} \langle x|T|x \rangle \leq \|x\|_{\mathcal{H}}^2 \sup_{n \geq 1} \text{trace}_{\mathcal{H}}(T_n) < \infty.$$

One concludes by applying Lemma 2.1 (a) and (b). \square

2.2. Finite Energy Condition. In the sequel, we shall repeatedly encounter the following typical situation. Let $A = A^* \geq 0$ be an unbounded self-adjoint operator on \mathcal{H} with domain $\text{Dom}(A)$, and let E be its spectral decomposition.

Let $T \in \mathcal{L}^1(\mathfrak{H})$ satisfy $T = T^* \geq 0$, and let $(e_j)_{j \geq 1}$ be a complete orthonormal system of eigenvectors of T with $Te_j = \tau_j e_j$ and $\tau_j \in [0, +\infty)$ for each $j \geq 1$.

Lemma 2.3. *Assume that*

$$(9) \quad \sum_{j \geq 1} \tau_j \int_0^\infty \lambda \langle e_j | E(d\lambda) | e_j \rangle < \infty.$$

Then

$$T^{1/2} A T^{1/2} := \sum_{j, k \geq 1} \tau_j^{1/2} \tau_k^{1/2} \left(\int_0^\infty \lambda \langle e_j | E(d\lambda) | e_k \rangle \right) |e_j\rangle \langle e_k|$$

satisfies

$$0 \leq T^{1/2} A T^{1/2} = (T^{1/2} A T^{1/2})^* \in \mathcal{L}^1(\mathcal{H})$$

and

$$\text{trace}_{\mathcal{H}}(T^{1/2} A T^{1/2}) = \sum_{j \geq 1} \tau_j \int_0^\infty \lambda \langle e_j | E(d\lambda) | e_j \rangle.$$

Proof. For each Borel $\omega \subset \mathbf{R}$ and each $x, y \in \mathcal{H}$, one has

$$|\langle x | E(\omega) | y \rangle| = |\langle E(\omega)x | E(\omega)y \rangle| \leq \|E(\omega)x\| \|E(\omega)y\| = \langle x | E(\omega) | x \rangle^{1/2} \langle y | E(\omega) | y \rangle^{1/2}$$

since $E(\omega)$ is a self-adjoint projection. In particular, for each $\alpha > 0$, one has

$$2|\langle x | E(\omega) | y \rangle| \leq \alpha \langle x | E(\omega) | x \rangle + \frac{1}{\alpha} \langle y | E(\omega) | y \rangle.$$

Hence

$$a_{jk} := \int_0^\infty \lambda \langle e_j | E(d\lambda) | e_k \rangle \in \mathbf{C}$$

and satisfies

$$2|a_{jk}|^2 \leq \alpha a_{jj} + \frac{1}{\alpha} a_{kk}$$

for all $\alpha > 0$, so that

$$|a_{jk}|^2 \leq a_{jj} a_{kk}$$

for all $j, k \geq 1$.

Since $(\tau_j a_{jj})_{j \geq 1} \in \ell^1(\mathbf{N}^*)$ by (9) and since

$$\langle e_j | T^{1/2} A T^{1/2} | e_k \rangle = \tau_j^{1/2} \tau_k^{1/2} a_{jk} = \overline{\langle e_k | T^{1/2} A T^{1/2} | e_j \rangle},$$

one concludes that $T^{1/2} A T^{1/2} = (T^{1/2} A T^{1/2})^* \in \mathcal{L}^2(\mathcal{H})$. Moreover, for each $x \in \mathcal{H}$

$$\begin{aligned} \langle x | T^{1/2} A T^{1/2} | x \rangle &= \sum_{j, k \geq 1} \tau_j^{1/2} \tau_k^{1/2} \overline{\langle e_j | x \rangle} \langle e_k | x \rangle \int_0^\infty \lambda \langle e_j | E(d\lambda) | e_k \rangle \\ &\geq \int_0^\infty \lambda \left\langle \sum_{j \geq 1} \tau_j^{1/2} \langle e_j | x \rangle e_j | E(d\lambda) \right\rangle \left\langle \sum_{j \geq 1} \tau_j^{1/2} \langle e_j | x \rangle e_j \right\rangle \\ &= \int_0^\infty \lambda \langle T^{1/2} x | E(d\lambda) | T^{1/2} x \rangle \geq 0, \end{aligned}$$

so that $T^{1/2} A T^{1/2} \geq 0$.

Finally

$$\begin{aligned} \sum_{l \geq 1} \langle e_l | T^{1/2} A T^{1/2} | e_l \rangle &= \sum_{l \geq 1} \sum_{j, k \geq 1} \tau_j^{1/2} \tau_k^{1/2} \left(\int_0^\infty \lambda \langle e_j | E(d\lambda) | e_k \rangle \right) \langle e_l | e_j \rangle \langle e_k | e_l \rangle \\ &= \sum_{l \geq 1} \sum_{j, k \geq 1} \tau_j^{1/2} \tau_k^{1/2} \left(\int_0^\infty \lambda \langle e_j | E(d\lambda) | e_k \rangle \right) \delta_{lj} \delta_{lk} = \sum_{l \geq 1} \tau_l \int_0^\infty \lambda \langle e_l | E(d\lambda) | e_l \rangle < \infty \end{aligned}$$

so that

$$\|T^{1/2} A T^{1/2}\|_1 = \text{trace}_{\mathcal{H}}(T^{1/2} A T^{1/2}) = \sum_{l \geq 1} \tau_l \int_0^\infty \lambda \langle e_l | E(d\lambda) | e_l \rangle < \infty$$

and in particular $T^{1/2} A T^{1/2} \in \mathcal{L}^1(\mathcal{H})$. \square

Corollary 2.4. *Let $T \in \mathcal{L}(\mathfrak{H})$ satisfy $T = T^* \geq 0$ and (9). Let $\Phi_n : \mathbf{R}_+ \rightarrow \mathbf{R}_+$ be a sequence of continuous, bounded and nondecreasing functions such that*

$$0 \leq \Phi_1(r) \leq \Phi_2(r) \leq \dots \leq \Phi_n(r) \rightarrow r \quad \text{as } n \rightarrow \infty.$$

Set

$$\Phi_n(A) := \int_0^\infty \Phi_n(\lambda) E(d\lambda) \in \mathcal{L}(\mathcal{H}).$$

Then $\Phi_n(A) = \Phi_n(A)^* \geq 0$ for each $n \geq 1$ and, for each $T \in \mathcal{L}^1(\mathcal{H})$ such that $T = T^* \geq 0$, the sequence $T^{1/2} \Phi_n(A) T^{1/2}$ converges weakly to $T^{1/2} A T^{1/2}$ as $n \rightarrow \infty$. Moreover

$$\text{trace}_{\mathcal{H}}(T \Phi_n(A)) \rightarrow \text{trace}_{\mathcal{H}}(T^{1/2} A T^{1/2}) \quad \text{as } n \rightarrow \infty.$$

Proof. Since E is a resolution of the identity on $[0, +\infty)$, and since Φ_n is continuous, bounded and with values in $[0, +\infty)$, the operators $\Phi_n(A)$ satisfy

$$0 \leq \Phi_n(A) = \Phi_n(A)^* \leq \left(\sup_{z \geq 0} \Phi_n(z) \right) I_{\mathcal{H}}$$

and

$$0 \leq \Phi_1(A) \leq \Phi_2(A) \leq \dots \leq \Phi_n(A) \leq \dots$$

Set $R_n := T^{1/2} \Phi_n(A) T^{1/2}$; by definition $0 \leq R_n = R_n^* \in \mathcal{L}(\mathfrak{H})$ and one has

$$0 \leq R_1 \leq R_2 \leq \dots \leq R_n \leq \dots$$

together with

$$\text{trace}_{\mathcal{H}}(R_n) = \sum_{j \geq 1} \tau_j \int_0^\infty \Phi_n(\lambda) \langle e_j | E(d\lambda) | e_j \rangle \leq \sum_{j \geq 1} \tau_j \int_0^\infty \lambda \langle e_j | E(d\lambda) | e_j \rangle < \infty$$

by (9). Applying Corollary 2.2 shows that R_n converges weakly to some $R \in \mathcal{L}^1(\mathcal{H})$ such that $R = R^* \geq 0$. Finally

$$T^{1/2} A T^{1/2} - R_n = \sum_{j, k \geq 1} \tau_j^{1/2} \tau_k^{1/2} \left(\int_0^\infty (\lambda - \Phi_n(\lambda)) \langle e_j | E(d\lambda) | e_k \rangle \right) | e_j \rangle \langle e_k |$$

so that

$$\begin{aligned} \langle x | T^{1/2} A T^{1/2} - R_n | x \rangle &= \int_0^\infty (\lambda - \Phi_n(\lambda)) \left\langle \sum_{j \geq 1} \tau_j^{1/2} \langle e_j | x \rangle e_j | E(d\lambda) | \sum_{k \geq 1} \tau_k^{1/2} \langle e_k | x \rangle e_k \right\rangle \\ &= \int_0^\infty (\lambda - \Phi_n(\lambda)) \langle T^{1/2} x | E(d\lambda) | T^{1/2} x \rangle \geq 0. \end{aligned}$$

Hence

$$0 \leq T^{1/2} A T^{1/2} - R_n = (T^{1/2} A T^{1/2} - R_n)^* \in \mathcal{L}^1(\mathcal{H})$$

so that

$$\begin{aligned} \|T^{1/2} A T^{1/2} - R_n\|_1 &= \text{trace}_{\mathcal{H}}(T^{1/2} A T^{1/2} - R_n) \\ &= \sum_{j \geq 1} \tau_j \int_0^\infty (\lambda - \Phi_n(\lambda)) \langle e_j | E(d\lambda) | e_j \rangle \rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$ by monotone convergence. Hence $R_n \rightarrow T^{1/2} A T^{1/2}$ in $\mathcal{L}^1(\mathcal{H})$ and one has in particular

$$\text{trace}_{\mathcal{H}}(T \Phi_n(A)) = \text{trace}_{\mathcal{H}}(T^{1/2} \Phi_n(A) T^{1/2}) \rightarrow \text{trace}_{\mathcal{H}}(T^{1/2} A T^{1/2}).$$

□

2.3. Energy and Partial Trace. Let \mathcal{H}_1 and \mathcal{H}_2 be two separable Hilbert spaces. Let $A = A^* \geq 0$ be an unbounded self-adjoint operator on \mathcal{H}_1 with domain $\text{Dom}(A)$, and let E be its spectral decomposition. Let $S \in \mathcal{L}^1(\mathcal{H}_1)$ satisfy $S = S^* \geq 0$, and let $(e_j)_{j \geq 1}$ be a complete orthonormal system of \mathcal{H}_1 of eigenvectors of S , with eigenvalues $(\sigma_j)_{j \geq 1}$ such that $S e_j = \sigma_j e_j$ for each $j \geq 1$. Assume that

$$\sum_{j \geq 1} \sigma_j \int_0^{+\infty} \lambda \langle e_j | E(d\lambda) | e_j \rangle < \infty.$$

Lemma 2.5. *Let $T \in \mathcal{L}^1(\mathcal{H}_1 \otimes \mathcal{H}_2)$ satisfy the partial trace condition*

$$\text{trace}(T | \mathcal{H}_2) = S.$$

Then $T^{1/2}(A \otimes I_{\mathcal{H}_2})T^{1/2} \in \mathcal{L}^1(\mathcal{H}_1 \otimes \mathcal{H}_2)$ and

$$\text{trace}_{\mathcal{H}_1 \otimes \mathcal{H}_2}(T^{1/2}(A \otimes I)T^{1/2}) = \text{trace}_{\mathcal{H}_1}(S^{1/2} A S^{1/2}).$$

Proof. For all $n \geq 1$, set $A_n = \Phi_n(A) \in \mathcal{L}(\mathfrak{H})$, with

$$\Phi_n(r) := \frac{r}{1 + \frac{1}{n}r}, \quad \text{for all } r \geq 0.$$

By construction, one has

$$A_n = A_n^* \geq 0 \quad \text{and} \quad A_1 \leq A_2 \leq \dots \leq A_n \leq \dots$$

Hence $T^{1/2}(A_n \otimes I_{\mathcal{H}_2})T^{1/2} = (T^{1/2}(A_n \otimes I_{\mathcal{H}_2})T^{1/2})^* \geq 0$ for all $n \geq 1$, and

$$T^{1/2}(A_1 \otimes I_{\mathcal{H}_2})T^{1/2} \leq T^{1/2}(A_2 \otimes I_{\mathcal{H}_2})T^{1/2} \leq \dots \leq T^{1/2}(A_n \otimes I_{\mathcal{H}_2})T^{1/2} \leq \dots$$

and since

$$\text{trace}_{\mathcal{H}_1 \otimes \mathcal{H}_2}(T(A_n \otimes I_{\mathcal{H}_2})) = \text{trace}_{\mathcal{H}_1}(SA_n) \rightarrow \text{trace}_{\mathcal{H}_1}(S^{1/2}AS^{1/2})$$

as $n \rightarrow \infty$ by the partial trace condition and Corollary 2.4, we conclude from Corollary 2.2 that

$$T^{1/2}(A \otimes I_{\mathcal{H}_2})T^{1/2} = (T^{1/2}(A \otimes I_{\mathcal{H}_2})T^{1/2})^* \geq 0$$

and that

$$\text{trace}_{\mathcal{H}_1 \otimes \mathcal{H}_2}(T^{1/2}(A \otimes I)T^{1/2}) = \text{trace}_{\mathcal{H}_1}(S^{1/2}AS^{1/2}).$$

□

3. COUPLINGS

Let $\mathfrak{H} := L^2(\mathbf{R}^d)$. An operator $R \in \mathcal{L}(\mathfrak{H})$ is a density operator if

$$R = R^* \geq 0 \quad \text{and} \quad \text{trace}(R) = 1.$$

We denote by $\mathcal{D}(\mathfrak{H})$ the set of density operators on \mathfrak{H} , and define

$$\mathcal{D}_2(\mathfrak{H}) := \{R \in \mathcal{D}(\mathfrak{H}) \text{ s.t. } \text{trace}(R^{1/2}(|y|^2 - \Delta_y)R^{1/2}) < \infty\}.$$

The set of Borel probability measures on $\mathbf{R}^d \times \mathbf{R}^d$ is denoted by $\mathcal{P}(\mathbf{R}^d \times \mathbf{R}^d)$. We denote by $\mathcal{P}_2(\mathbf{R}^d \times \mathbf{R}^d)$ the set of Borel probability measures μ on $\mathbf{R}^d \times \mathbf{R}^d$ such that

$$\iint_{\mathbf{R}^d \times \mathbf{R}^d} (|x|^2 + |\xi|^2) \mu(dx d\xi) < \infty.$$

The set of Borel probability measures on $\mathbf{R}^d \times \mathbf{R}^d$ which are absolutely continuous with respect to the Lebesgue measure on $\mathbf{R}^d \times \mathbf{R}^d$ is denoted $\mathcal{P}^{ac}(\mathbf{R}^d \times \mathbf{R}^d) \subset \mathcal{P}(\mathbf{R}^d \times \mathbf{R}^d)$. We set $\mathcal{P}_2^{ac}(\mathbf{R}^d \times \mathbf{R}^d) = \mathcal{P}^{ac}(\mathbf{R}^d \times \mathbf{R}^d) \cap \mathcal{P}_2(\mathbf{R}^d \times \mathbf{R}^d)$, and we identify elements of $\mathcal{P}^{ac}(\mathbf{R}^d \times \mathbf{R}^d)$ with their densities with respect to the Lebesgue measure.

Let $R_1, R_2 \in \mathcal{D}(\mathfrak{H})$; a coupling of R_1 and R_2 is an element $\mathcal{R} \in \mathcal{D}(\mathfrak{H} \otimes \mathfrak{H})$ such that

$$\text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((A \otimes I + I \otimes B)\mathcal{R}) = \text{trace}_{\mathfrak{H}}(R_1 A) + \text{trace}_{\mathfrak{H}}(R_2 B).$$

The set of couplings of R_1 and R_2 will be denoted by $\mathcal{C}(R_1, R_2)$. Obviously the tensor product $R_1 \otimes R_2 \in \mathcal{C}(R_1, R_2)$, so that $\mathcal{C}(R_1, R_2) \neq \emptyset$.

Let f be a probability density on $\mathbf{R}^d \times \mathbf{R}^d$, and let $R \in \mathcal{D}(\mathfrak{H})$. A coupling of f and R is an ultraweakly measurable operator-valued function $(x, \xi) \mapsto Q(x, \xi)$ defined a.e. on $\mathbf{R}^d \times \mathbf{R}^d$ with values in $\mathcal{L}(\mathfrak{H})$ such that

$$Q(x, \xi) = Q(x, \xi)^* \geq 0, \quad \iint_{\mathbf{R}^d \times \mathbf{R}^d} Q(x, \xi) dx d\xi = R$$

$$\text{and } \text{trace}_{\mathfrak{H}}(Q(x, \xi)) = f(x, \xi) \text{ for a.e. } (x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d.$$

The set of couplings of f and R will also be denoted by $\mathcal{C}(f, R)$. Since the map $(x, \xi) \mapsto f(x, \xi)R$ (henceforth denoted $f \otimes R$) obviously belongs to $\mathcal{C}(f, R)$, one has $\mathcal{C}(f, R) \neq \emptyset$.

In general, one does not know much about the general structure of couplings between two density operators. However, the case where one of the density operators is a rank 1 projection is particularly simple.

Lemma 3.1. *Let $P = P^* \in \mathcal{L}(\mathfrak{H})$ be a rank 1 projection. Then*
 (i) *for each probability density f on $\mathbf{R}^d \times \mathbf{R}^d$, one has $\mathcal{C}(f, P) = \{f \otimes P\}$;*
 (ii) *for each $R \in \mathcal{D}(\mathfrak{H})$, one has $\mathcal{C}(P, R) = \{P \otimes R\}$ and $\mathcal{C}(R, P) = \{R \otimes P\}$.*

This is in complete analogy with the following elementary observation: if $\mu \in \mathcal{P}(\mathbf{R}^d)$ and $y_0 \in \mathbf{R}^d$, the only coupling of μ and δ_{y_0} is $\mu \otimes \delta_{y_0}$. In other words, self-adjoint rank-1 projections are the quantum analogue of points in this picture.

Proof. Let $Q \in \mathcal{C}(f, P)$; one has

$$\begin{aligned} & \iint_{\mathbf{R}^d \times \mathbf{R}^d} \text{trace}_{\mathfrak{H}}((I - P)Q(x, \xi)(I - P)) dx d\xi \\ &= \text{trace}_{\mathfrak{H}} \left((I - P) \iint_{\mathbf{R}^d \times \mathbf{R}^d} Q(x, \xi) dx d\xi (I - P) \right) \\ &= \text{trace}_{\mathfrak{H}}((I - P)P(I - P)) = 0. \end{aligned}$$

Since $(I - P)Q(x, \xi)(I - P) \geq 0$ for a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$, this implies that

$$(I - P)Q(x, \xi)(I - P) = 0 \quad \text{for a.e. } (x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d.$$

Since $Q(x, \xi) = Q(x, \xi)^* \geq 0$ for a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$, we deduce from the Cauchy-Schwarz inequality that, for all $\phi, \psi \in \mathfrak{H}$

$$\begin{aligned} & |\langle P\phi | Q(x, \xi) | (I - P)\psi \rangle|^2 = |\langle (I - P)\psi | Q(x, \xi) | P\phi \rangle|^2 \\ & \leq \langle P\phi | Q(x, \xi) | P\phi \rangle^{1/2} \langle (I - P)\psi | Q(x, \xi) | (I - P)\psi \rangle^{1/2} = 0. \end{aligned}$$

Hence $(I - P)Q(x, \xi)P = PQ(x, \xi)(I - P) = 0$ for a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$, so that

$$Q(x, \xi) = PQ(x, \xi)P \quad \text{for a.e. } (x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d.$$

Writing P as $P = |u\rangle\langle u|$ where $u \in \mathfrak{H}$ is a unit vector, we conclude that

$$Q(x, \xi) = \langle u | Q(x, \xi) | u \rangle P \quad \text{for a.e. } (x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d.$$

Finally

$$\text{trace}(Q(x, \xi)) = f(x, \xi) = \langle u | Q(x, \xi) | u \rangle \quad \text{for a.e. } (x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d.$$

This concludes the proof of (i).

As for (ii), let $Q \in \mathcal{C}(R, P)$. Then

$$\text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((I \otimes (I - P))Q(I \otimes (I - P))) = \text{trace}_{\mathfrak{H}}(I - P)P(I - P) = 0.$$

Hence

$$(I \otimes (I - P))Q(I \otimes (I - P)) = 0.$$

Since $Q = Q^* \geq 0$, the Cauchy-Schwarz inequality implies that, for all $\phi, \phi', \psi, \psi' \in \mathfrak{H}$

$$\begin{aligned} & |\langle \phi \otimes \psi | (I \otimes P)Q(I \otimes (I - P)) | \phi' \otimes \psi' \rangle| = |\langle \phi' \otimes \psi' | (I \otimes (I - P))Q(I \otimes P) | \phi \otimes \psi \rangle| \\ & \leq \langle \phi \otimes \psi | (I \otimes P)Q(I \otimes P) | \phi \otimes \psi \rangle^{1/2} \langle \phi' \otimes \psi' | (I \otimes (I - P))Q(I \otimes (I - P)) | \phi' \otimes \psi' \rangle^{1/2} \end{aligned}$$

so that

$$(I \otimes P)Q(I \otimes (I - P)) = (I \otimes (I - P))Q(I \otimes P) = 0.$$

Hence

$$Q = (I \otimes P)Q(I \otimes P).$$

Writing $P = |u\rangle\langle u|$ with $u \in \mathfrak{H}$ and $|u| = 1$ as above, we conclude that

$$\langle \phi \otimes \psi | Q | \phi' \otimes \psi' \rangle = \langle \phi \otimes u | Q | \phi' \otimes u \rangle \langle u | \psi \rangle \langle u | \psi' \rangle.$$

This shows that $\mathcal{Q} = L \otimes |u\rangle\langle u| = L \otimes P$, where $L = L^*$ is the element of $\mathcal{L}(\mathfrak{H})$ such that

$$\langle \phi | L | \phi' \rangle = \langle \phi \otimes u | \mathcal{Q} | \phi' \otimes u \rangle$$

for each $\phi, \phi' \in \mathfrak{H}$. (Observe indeed that $(\phi, \phi') \mapsto \langle \phi \otimes u | \mathcal{Q} | \phi' \otimes u \rangle$ is a continuous, symmetric bilinear functional on \mathfrak{H} , and is therefore represented by a unique self-adjoint element of $\mathcal{L}(\mathfrak{H})$.) We conclude by observing that

$$\text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((A \otimes I)\mathcal{Q}) = \text{trace}_{\mathfrak{H}}(AR) = \text{trace}_{\mathfrak{H}}(LR)$$

for each finite rank operator $A \in \mathcal{L}(\mathfrak{H})$, and this implies that $\mathcal{Q} = R \otimes P$.

The case of $\mathcal{Q}' \in \mathcal{C}(P, R)$ is handled similarly. \square

Next we explain how to “disintegrate” a coupling with respect to one of its marginals when this marginal is a probability density.

Lemma 3.2. *Let $f \in \mathcal{P}^{ac}(\mathbf{R}^d \times \mathbf{R}^d)$, let $R \in \mathcal{D}(\mathfrak{H})$ and let $Q \in \mathcal{C}(f, R)$. There exists a $\sigma(\mathcal{L}^1(\mathfrak{H}), \mathcal{L}(\mathfrak{H}))$ weakly measurable function $(x, \xi) \mapsto Q_f(x, \xi)$ defined a.e. on $\mathbf{R}^d \times \mathbf{R}^d$ with values in $\mathcal{L}^1(\mathfrak{H})$ such that*

$$Q_f(x, \xi) = Q_f^*(x, \xi) \geq 0, \quad \text{trace}(Q_f(x, \xi)) = 1, \quad \text{and } Q(x, \xi) = f(x, \xi)Q_f(x, \xi)$$

for a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$.

Proof. Let f_1 be a Borel measurable function defined on $\mathbf{R}^d \times \mathbf{R}^d$ and such that $f(x, \xi) = f_1(x, \xi)$ for a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$. Let \mathcal{N} be the Borel measurable set defined as follows: $\mathcal{N} := \{(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d \text{ s.t. } f(x, \xi) = 0\}$, and let $u \in \mathfrak{H}$ satisfy $|u| = 1$. Consider the function

$$(x, \xi) \mapsto Q_f(x, \xi) := \frac{Q(x, \xi) + \mathbf{1}_{\mathcal{N}}(x, \xi)|u\rangle\langle u|}{f_1(x, \xi) + \mathbf{1}_{\mathcal{N}}(x, \xi)} \in \mathcal{L}(\mathfrak{H})$$

defined a.e. on $\mathbf{R}^d \times \mathbf{R}^d$. The function $f_1 + \mathbf{1}_{\mathcal{N}} > 0$ is Borel measurable on $\mathbf{R}^d \times \mathbf{R}^d$ while $(x, \xi) \mapsto \langle \phi | Q(x, \xi) | \psi \rangle$ is measurable and defined a.e. on $\mathbf{R}^d \times \mathbf{R}^d$ for each $\phi, \psi \in \mathfrak{H}$. Set $\mathcal{A} : \mathcal{L}(\mathfrak{H}) \times (0, +\infty) \ni (T, \lambda) \mapsto \lambda^{-1}T \in \mathcal{L}(\mathfrak{H})$; since \mathcal{A} is continuous, the function $Q_f := \mathcal{A}(Q + \mathbf{1}_{\mathcal{N}} \otimes |u\rangle\langle u|, f_1 + \mathbf{1}_{\mathcal{N}})$ is weakly measurable on $\mathbf{R}^d \times \mathbf{R}^d$. Since $f_1 + \mathbf{1}_{\mathcal{N}} > 0$, and since $Q(x, \xi) = Q^*(x, \xi) \geq 0$, one has $(Q(x, \xi) + \mathbf{1}_{\mathcal{N}} \otimes |u\rangle\langle u|)^* = Q(x, \xi) + \mathbf{1}_{\mathcal{N}} \otimes |u\rangle\langle u| \geq 0$ for a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$. On the other hand, for a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$, one has $\text{trace}(Q(x, \xi) + \mathbf{1}_{\mathcal{N}} \otimes |u\rangle\langle u|) = f(x, \xi) + \mathbf{1}_{\mathcal{N}}(x, \xi)$, so that $\text{trace}(Q_f(x, \xi)) = 1$. Finally

$$f(x, \xi)Q_f(x, \xi) = \frac{f(x, \xi)Q(x, \xi)}{f_1(x, \xi) + \mathbf{1}_{\mathcal{N}}(x, \xi)} = Q(x, \xi) \quad \text{for a.e. } (x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d,$$

since $f = f_1$ a.e. on $\mathbf{R}^d \times \mathbf{R}^d$ and $\mathbf{1}_{\mathcal{N}}(x, \xi) = 0$ for a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$ such that $f(x, \xi) > 0$. Since Q_f satisfies $\text{trace}(Q_f(x, \xi)) = 1$ for a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$ and is weakly measurable on $\mathbf{R}^d \times \mathbf{R}^d$, it is $\sigma(\mathcal{L}^1(\mathfrak{H}), \mathcal{L}(\mathfrak{H}))$ weakly measurable. \square

4. TRIANGLE INEQUALITIES

The following “pseudo metrics” have been defined in [6] and in [7] respectively.

Definition 4.1. *For all $R, S \in \mathcal{D}_2(\mathfrak{H})$ and all $f \in \mathcal{P}_2^{ac}(\mathbf{R}^d \times \mathbf{R}^d)$, we set*

$$MK_h(R, S) := \inf_{A \in \mathcal{C}(R, S)} \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}(A^{1/2} C A^{1/2})^{1/2}$$

where

$$C := C(x, y, \hbar D_x, \hbar D_y) = |x - y|^2 + |\hbar D_x - \hbar D_y|^2.$$

Similarly, we set

$$\mathcal{E}_h(f, R) := \inf_{a \in \mathcal{C}(f, R)} \left(\iint_{\mathbf{R}^d \times \mathbf{R}^d} \text{trace}_{\mathfrak{H}}(a(x, \xi)^{1/2} c(x, \xi) a(x, \xi)^{1/2}) dx d\xi \right)^{1/2}$$

where

$$c(x, \xi) := c(x, \xi, y, \hbar D_y) = |x - y|^2 + |\xi - \hbar D_y|^2.$$

The above ‘‘pseudometrics’’ satisfy the following inequalities.

Theorem 4.2. *Let $f, g \in \mathcal{P}_2^{ac}(\mathbf{R}^d \times \mathbf{R}^d)$, and let $R_1, R_2, R_3 \in \mathcal{D}_2(\mathfrak{H})$. The following inequalities hold true:*

- (a) $\mathcal{E}_h(f, R_1) \leq \text{dist}_{\text{MK}, 2}(f, g) + \mathcal{E}_h(g, R_1)$;
- (b) $MK_h(R_1, R_3) \leq \mathcal{E}_h(f, R_1) + \mathcal{E}_h(f, R_3)$;
- (c) if $\text{rank}(R_2) = 1$, then

$$MK_h(R_1, R_3) \leq MK_h(R_1, R_2) + MK_h(R_2, R_3),$$

- (d) if $\text{rank}(R_2) = 1$, then

$$\text{dist}_{\text{MK}, 2}(f, g) \leq \mathcal{E}_h(f, R_2) + \mathcal{E}_h(g, R_2),$$

- (e) if $\text{rank}(R_2) = 1$, then

$$\mathcal{E}_h(f, R_3) \leq \mathcal{E}_h(f, R_2) + MK_h(R_2, R_3).$$

The proofs of all these triangle inequalities make use of some inequalities between the (classical and/or quantum) transportation cost operators. We begin with an elementary, but useful lemma, which can be viewed as the Peter-Paul inequality for operators.

Lemma 4.3. *Let T, S be unbounded self-adjoint operators on $\mathfrak{H} = L^2(\mathbf{R}^n)$, with domains $\text{Dom}(T)$ and $\text{Dom}(S)$ respectively such that $\text{Dom}(T) \cap \text{Dom}(S)$ is dense in \mathfrak{H} . Then, for all $\alpha > 0$, one has*

$$\langle v | TS + ST | v \rangle \leq \alpha \langle v | T^2 | v \rangle + \frac{1}{\alpha} \langle v | S^2 | v \rangle, \quad \text{for all } v \in \text{Dom}(T) \cap \text{Dom}(S).$$

Proof. Indeed, for each $\alpha > 0$ and each $v \in \text{Dom}(T) \cap \text{Dom}(S)$, one has

$$\begin{aligned} & \alpha \langle v | T^2 | v \rangle + \frac{1}{\alpha} \langle v | S^2 | v \rangle - \langle v | TS + ST | v \rangle \\ &= |\sqrt{\alpha} T v|^2 + \left| \frac{1}{\sqrt{\alpha}} S v \right|^2 - \langle \sqrt{\alpha} T v | \frac{1}{\sqrt{\alpha}} S v \rangle - \langle \frac{1}{\sqrt{\alpha}} S v | \sqrt{\alpha} T v \rangle \\ &= \left| \sqrt{\alpha} T v - \frac{1}{\sqrt{\alpha}} S v \right|^2 \geq 0. \end{aligned}$$

□

Lemma 4.4. *For each $x, \xi, y, \eta, z \in \mathbf{R}^d$ and each $\alpha > 0$, one has*

$$\begin{aligned} c(x, \xi; z, \hbar D_z) &\leq (1 + \alpha)(|x - y|^2 + |\xi - \eta|^2) + (1 + \frac{1}{\alpha})c(y, \eta; z, \hbar D_z), \\ C(x, z, \hbar D_x, \hbar D_z) &\leq (1 + \alpha)c(y, \eta; x, \hbar D_x) + (1 + \frac{1}{\alpha})c(y, \eta; z, \hbar D_z), \\ C(x, z, \hbar D_x, \hbar D_z) &\leq (1 + \alpha)C(x, y; \hbar D_x, \hbar D_y) + (1 + \frac{1}{\alpha})C(y, z; \hbar D_y, \hbar D_z), \\ |x - z|^2 + |\xi - \zeta|^2 &\leq (1 + \alpha)c(x, \xi; y, \hbar D_y) + (1 + \alpha)c(z, \zeta; y, \hbar D_y), \\ c(x, \xi; z, \hbar D_z) &\leq (1 + \alpha)c(x, \xi; y, \hbar D_y) + (1 + \frac{1}{\alpha})C(y, z; \hbar D_y, \hbar D_z). \end{aligned}$$

All these inequalities are of the form $A \leq B$ where A and B are unbounded self-adjoint operators on $L^2(\mathbf{R}^n)$ for some $n \geq 1$, with

$$\mathcal{W} := \{\psi \in H^1(\mathbf{R}^n) \text{ s.t. } |x|\psi \in \mathcal{H}\} \subset \text{Dom}_f(A) \cap \text{Dom}_f(B),$$

denoting by $\text{Dom}_f(A)$ (resp. $\text{Dom}_f(B)$) the form-domain of A (resp. of B) — see §VIII.6 in [9] on pp. 276–277. The inequality $A \leq B$ means that the bilinear form associated to $B - A$ is nonnegative, i.e. that

$$\langle w|A|w \rangle \leq \langle w|B|w \rangle, \quad \text{for all } w \in \mathcal{W}.$$

Proof. All these inequalities are proved in the same way. Let us prove for instance the third inequality:

$$\begin{aligned} C(x, z, \hbar D_x, \hbar D_z) &= |x - y + y - z|^2 + |\hbar D_x - \hbar D_y + \hbar D_y - \hbar D_z|^2 \\ &= C(x, y; \hbar D_x, \hbar D_y) + C(y, z; \hbar D_y, \hbar D_z) \\ &\quad + 2(x - y) \cdot (y - z) + 2(\hbar D_x - \hbar D_y) \cdot (\hbar D_y - \hbar D_z). \end{aligned}$$

Observe indeed that the multiplication operators by $(x - y)$ and by $(y - z)$ commute; likewise $(\hbar D_x - \hbar D_y)$ and $(\hbar D_y - \hbar D_z)$ commute. By Lemma 4.3

$$\begin{aligned} &2(x - y) \cdot (y - z) + 2(\hbar D_x - \hbar D_y) \cdot (\hbar D_y - \hbar D_z) \\ &\leq \alpha C(x, y; \hbar D_x, \hbar D_y) + \frac{1}{\alpha} C(y, z; \hbar D_y, \hbar D_z), \end{aligned}$$

which concludes the proof of the third inequality. \square

Proof of Theorem 4.2 (a). By Theorem 2.12 in chapter 2 of [14], there exists an optimal coupling for $W_2(f, g)$, of the form $f(x, \xi) \delta_{\nabla \Phi(x, \xi)}(dyd\eta)$, where Φ is a convex function on $\mathbf{R}^d \times \mathbf{R}^d$. Let $Q \in \mathcal{C}(g, R_1)$ and set

$$P(x, \xi; dyd\eta) := f(x, \xi) \delta_{\nabla \Phi(x, \xi)}(dyd\eta) Q_g(y, \eta),$$

where Q_g is the disintegration of Q with respect to f obtained in Lemma 3.2. Then P is a nonnegative, self-adjoint operator-valued measure satisfying

$$\text{trace}_{\mathfrak{H}}(P(x, \xi; dyd\eta)) = f(x, \xi) \delta_{\nabla \Phi(x, \xi)}(dyd\eta)$$

while

$$\int P dx d\xi = (\nabla \Phi \# f)(y, \eta) dyd\eta Q_g(y, \eta) = g(y, \eta) Q_g(y, \eta) dyd\eta = Q(y, \eta) dyd\eta.$$

In particular

$$(10) \quad \int P(x, \xi; dyd\eta) = f(x, \xi) Q_g(\nabla \Phi(x, \xi)) \in \mathcal{C}(f, R_1).$$

Therefore

$$\mathcal{E}_h(f, R_1)^2 \leq \int \text{trace}_{\mathfrak{H}}(Q_g(\nabla \Phi(x, \xi))^{1/2} c_h(x, \xi) Q_g(\nabla \Phi(x, \xi))^{1/2}) f(x, \xi) dx d\xi.$$

By the first inequality in Lemma 4.4, one has

$$c_h(x, \xi; z, \hbar D_z) \leq (1 + \alpha) |(x, \xi) - \nabla \phi(x, \xi)|^2 + (1 + \frac{1}{\alpha}) c_h(\nabla \Phi(x, \xi); z, \hbar D_z)$$

for a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$ and all $\alpha > 0$. Since $g \in \mathcal{P}_2^{ac}(\mathbf{R}^d \times \mathbf{R}^d)$ and $R_1 \in \mathcal{D}_2(\mathfrak{H})$ and $Q \in \mathcal{C}(g, R_1)$, then

$$\begin{aligned} &\int \text{trace}_{\mathfrak{H}}(Q(y, \eta)^{1/2} c_h(y, \eta) Q(y, \eta)^{1/2}) dyd\eta \\ &= \int \text{trace}_{\mathfrak{H}}(Q_g(\nabla \Phi(x, \xi))^{1/2} c_h(\nabla \Phi(x, \xi)) Q_g(\nabla \Phi(x, \xi))^{1/2}) f(x, \xi) dx d\xi < \infty. \end{aligned}$$

For each $\epsilon > 0$, set

$$c_h^\epsilon(x, \xi; z, \hbar D_z) = (I + \epsilon c_h(x, \xi; z, \hbar D_z))^{-1} c_h(x, \xi; z, \hbar D_z) \leq c_h(x, \xi; z, \hbar D_z).$$

Then, for a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$ and each $\epsilon > 0$, one has

$$Q_g(\nabla\Phi(x, \xi))^{1/2} c_h(\nabla\Phi(x, \xi); z, \hbar D_z) Q_g(\nabla\Phi(x, \xi))^{1/2} \in \mathcal{L}^1(\mathfrak{H}),$$

and

$$\begin{aligned} & Q_g(\nabla\Phi(x, \xi))^{1/2} c_h^\epsilon(x, \xi; z, \hbar D_z) Q_g(\nabla\Phi(x, \xi))^{1/2} \\ & \leq (1 + \alpha) |(x, \xi) - \nabla\phi(x, \xi)|^2 Q_g(\nabla\Phi(x, \xi)) \\ & \quad + (1 + \frac{1}{\alpha}) Q_g(\nabla\Phi(x, \xi))^{1/2} c_h(\nabla\Phi(x, \xi); z, \hbar D_z) Q_g(\nabla\Phi(x, \xi))^{1/2}. \end{aligned}$$

Integrating both sides of this inequality with respect to the probability distribution $f(x, \xi)$, one finds

$$\begin{aligned} & \int \text{trace}_{\mathfrak{H}}(Q_g(\nabla\Phi(x, \xi))^{1/2} c_h^\epsilon(x, \xi) Q_g(\nabla\Phi(x, \xi))^{1/2}) f(x, \xi) dx d\xi \\ & \leq (1 + \alpha) \int |(x, \xi) - \nabla\phi(x, \xi)|^2 f(x, \xi) dx d\xi \\ & + (1 + \frac{1}{\alpha}) \int \text{trace}_{\mathfrak{H}}(Q_g(\nabla\Phi(x, \xi))^{1/2} c_h(\nabla\Phi(x, \xi)) Q_g(\nabla\Phi(x, \xi))^{1/2}) f(x, \xi) dx d\xi \\ & \leq (1 + \alpha) \text{dist}_{\text{MK},2}(f, g)^2 \\ & \quad + (1 + \frac{1}{\alpha}) \int \text{trace}_{\mathfrak{H}}(Q_g(y, \eta)^{1/2} c_h(y, \eta) Q_g(y, \eta)^{1/2}) g(y, \eta) dy d\eta \\ & \leq (1 + \alpha) \text{dist}_{\text{MK},2}(f, g)^2 + (1 + \frac{1}{\alpha}) \int \text{trace}_{\mathfrak{H}}(Q(y, \eta)^{1/2} c_h(y, \eta) Q(y, \eta)^{1/2}) dy d\eta. \end{aligned}$$

Minimizing the last right hand side of this inequality in $Q \in \mathcal{C}(g, R_1)$ shows that

$$\begin{aligned} & \int \text{trace}_{\mathfrak{H}}(Q_g(\nabla\Phi(x, \xi))^{1/2} c_h^\epsilon(x, \xi) Q_g(\nabla\Phi(x, \xi))^{1/2}) f(x, \xi) dx d\xi \\ & \leq (1 + \alpha) \text{dist}_{\text{MK},2}(f, g)^2 + (1 + \frac{1}{\alpha}) \mathcal{E}(g, R_1)^2. \end{aligned}$$

Passing to the limit as $\epsilon \rightarrow 0^+$ in the left hand side and applying Corollary 2.4 shows that

$$\begin{aligned} \mathcal{E}_h(f, R_1)^2 & \leq \int \text{trace}_{\mathfrak{H}}(Q_g(\nabla\Phi(x, \xi))^{1/2} c_h(x, \xi) Q_g(\nabla\Phi(x, \xi))^{1/2}) f(x, \xi) dx d\xi \\ & \leq (1 + \alpha) \text{dist}_{\text{MK},2}(f, g)^2 + (1 + \frac{1}{\alpha}) \mathcal{E}(g, R_1)^2, \end{aligned}$$

the first inequality being a consequence of the definition of \mathcal{E}_h according to (10).

Finally, minimizing the right hand side of this inequality as $\alpha > 0$, i.e. choosing $\alpha = \mathcal{E}_h(f, g) / \text{dist}_{\text{MK},2}(f, g)$ if $f \neq g$ a.e. on $\mathbf{R}^d \times \mathbf{R}^d$, or letting $\alpha \rightarrow +\infty$ if $f = g$, we arrive at the inequality

$$\begin{aligned} \mathcal{E}_h(f, R_1)^2 & \leq \text{dist}_{\text{MK},2}(f, g)^2 + \mathcal{E}_h(g, R_1)^2 + 2\mathcal{E}_h(g, R_1) \text{dist}_{\text{MK},2}(f, g) \\ & = (\text{dist}_{\text{MK},2}(f, g) + \mathcal{E}_h(g, R_1))^2, \end{aligned}$$

which is precisely the inequality (a). \square

Proof of Theorem 4.2 (b). Let $Q_1 \in \mathcal{C}(f, R_1)$ and $Q_3 \in \mathcal{C}(f, R_3)$. Let $Q_{1,f}$ and $Q_{3,f}$ be the disintegrations of Q_1 and Q_3 with respect to f obtained in Lemma 3.2. For each $\epsilon > 0$, set

$$(11) \quad C_h^\epsilon(x, z, \hbar D_x, \hbar D_z) = (I + \epsilon C_h(x, z, \hbar D_x, \hbar D_z))^{-1} C_h(x, z, \hbar D_x, \hbar D_z)$$

and observe that

$$0 \leq C_h^\epsilon(x, z, \hbar D_x, \hbar D_z) = C_h^\epsilon(x, z, \hbar D_x, \hbar D_z)^* \in \mathcal{L}(\mathcal{H} \otimes \mathfrak{H}).$$

By the second inequality in Lemma 4.4, for all $(y, \eta) \in \mathbf{R}^d \times \mathbf{R}^d$ and all $\alpha > 0$, one has

$$\begin{aligned} C_h^\epsilon(x, z, \hbar D_x, \hbar D_z) &\leq C_h(x, z, \hbar D_x, \hbar D_z) \\ &\leq (1 + \alpha)c_h(y, \eta; x, \hbar D_x) + (1 + \frac{1}{\alpha})c_h(y, \eta; z, \hbar D_z). \end{aligned}$$

Therefore, for a.e. $(y, \eta) \in \mathbf{R}^d \times \mathbf{R}^d$, one has

$$\begin{aligned} &(Q_{1,f}(y, \eta) \otimes Q_{3,f}(y, \eta))^{1/2} C_h^\epsilon(x, z, \hbar D_x, \hbar D_z) (Q_{1,f}(y, \eta) \otimes Q_{3,f}(y, \eta))^{1/2} \\ &\leq (1 + \alpha)(Q_{1,f}(y, \eta) \otimes Q_{3,f}(y, \eta))^{1/2} c_h(y, \eta; x, \hbar D_x) (Q_{1,f}(y, \eta) \otimes Q_{3,f}(y, \eta))^{1/2} \\ &\quad + (1 + \frac{1}{\alpha})(Q_{1,f}(y, \eta) \otimes Q_{3,f}(y, \eta))^{1/2} c_h(y, \eta; z, \hbar D_z) (Q_{1,f}(y, \eta) \otimes Q_{3,f}(y, \eta))^{1/2} \\ &= (1 + \alpha) (Q_{1,f}(y, \eta))^{1/2} c_h(y, \eta; x, \hbar D_x) Q_{1,f}(y, \eta)^{1/2} \otimes Q_{3,f}(y, \eta) \\ &\quad + (1 + \frac{1}{\alpha}) Q_{1,f}(y, \eta) \otimes (Q_{3,f}(y, \eta))^{1/2} c_h(y, \eta; z, \hbar D_z) Q_{1,f}(y, \eta)^{1/2}. \end{aligned}$$

Taking the trace in $\mathfrak{H} \otimes \mathfrak{H}$ of both sides of this inequality shows that

$$\begin{aligned} &\text{trace}_{\mathfrak{H} \otimes \mathfrak{H}} ((Q_{1,f} \otimes Q_{3,f}(y, \eta)) C_h^\epsilon(x, z, \hbar D_x, \hbar D_z)) \\ &= \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}} ((Q_{1,f} \otimes Q_{3,f}(y, \eta))^{1/2} C_h^\epsilon(x, z, \hbar D_x, \hbar D_z) (Q_{1,f} \otimes Q_{3,f}(y, \eta))^{1/2}) \\ &\leq (1 + \alpha) \text{trace}_{\mathfrak{H}} (Q_{1,f}(y, \eta)^{1/2} c_h(y, \eta; x, \hbar D_x) Q_{1,f}(y, \eta)^{1/2}) \\ &\quad + (1 + \frac{1}{\alpha}) \text{trace}_{\mathfrak{H}} (Q_{3,f}(y, \eta)^{1/2} c_h(y, \eta; z, \hbar D_z) Q_{1,f}(y, \eta)^{1/2}) \end{aligned}$$

for a.e. $(y, \eta) \in \mathbf{R}^d \times \mathbf{R}^d$. Integrating both sides of this inequality in (y, η) with respect to f shows that

$$\begin{aligned} &\text{trace}_{\mathfrak{H} \otimes \mathfrak{H}} \left(\left(\int (Q_{1,f} \otimes Q_{3,f}(y, \eta)) f((y, \eta)) dy d\eta \right) C_h^\epsilon(x, z, \hbar D_x, \hbar D_z) \right) \\ &\leq (1 + \alpha) \int \text{trace}_{\mathfrak{H}} (Q_{1,f}(y, \eta)^{1/2} c_h(y, \eta; x, \hbar D_x) Q_{1,f}(y, \eta)^{1/2}) f(y, \eta) dy d\eta \\ &\quad + (1 + \frac{1}{\alpha}) \int \text{trace}_{\mathfrak{H}} (Q_{3,f}(y, \eta)^{1/2} c_h(y, \eta; z, \hbar D_z) Q_{1,f}(y, \eta)^{1/2}) f(y, \eta) dy d\eta \\ &= (1 + \alpha) \int \text{trace}_{\mathfrak{H}} ((f Q_{1,f}(y, \eta))^{1/2} c_h(y, \eta; x, \hbar D_x) (f Q_{1,f}(y, \eta))^{1/2}) dy d\eta \\ &\quad + (1 + \frac{1}{\alpha}) \int \text{trace}_{\mathfrak{H}} ((f Q_{3,f}(y, \eta))^{1/2} c_h(y, \eta; z, \hbar D_z) (f Q_{1,f}(y, \eta))^{1/2}) dy d\eta \\ &= (1 + \alpha) \int \text{trace}_{\mathfrak{H}} (Q_1(y, \eta)^{1/2} c_h(y, \eta; x, \hbar D_x) Q_1(y, \eta)^{1/2}) dy d\eta \\ &\quad + (1 + \frac{1}{\alpha}) \int \text{trace}_{\mathfrak{H}} (Q_3(y, \eta)^{1/2} c_h(y, \eta; z, \hbar D_z) Q_3(y, \eta)^{1/2}) dy d\eta. \end{aligned}$$

By construction

$$P := \int (Q_{1,f} \otimes Q_{3,f}(y, \eta)) f((y, \eta)) dy d\eta \in \mathcal{C}(R_1, R_3);$$

on the other hand

$$\begin{aligned} &\int \text{trace}_{\mathfrak{H}} (Q_1(y, \eta)^{1/2} c_h(y, \eta; x, \hbar D_x) Q_1(y, \eta)^{1/2}) dy d\eta < \infty \\ &\int \text{trace}_{\mathfrak{H}} (Q_3(y, \eta)^{1/2} c_h(y, \eta; z, \hbar D_z) Q_3(y, \eta)^{1/2}) dy d\eta < \infty \end{aligned}$$

since $R_1, R_3 \in \mathcal{D}_2(\mathfrak{H})$ while $f \in \mathcal{P}_2^{ac}(\mathbf{R}^d \times \mathbf{R}^d)$. By Corollary 2.4

$$\text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}(PC_h^\epsilon(x, z, \hbar D_x, \hbar D_z)) \rightarrow \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}(P^{1/2}C_h(x, z, \hbar D_x, \hbar D_z)P^{1/2})$$

as $\epsilon \rightarrow 0^+$, so that

$$\begin{aligned} MK_h(R_1, R_3)^2 &\leq \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}(P^{1/2}C_h(x, z, \hbar D_x, \hbar D_z)P^{1/2}) \\ &\leq (1 + \alpha) \int \text{trace}_{\mathfrak{H}}(Q_1(y, \eta)^{1/2}c_h(y, \eta; x, \hbar D_x)Q_1(y, \eta)^{1/2}) dy d\eta \\ &\quad + (1 + \frac{1}{\alpha}) \int \text{trace}_{\mathfrak{H}}(Q_3(y, \eta)^{1/2}c_h(y, \eta; z, \hbar D_z)Q_3(y, \eta)^{1/2}) dy d\eta. \end{aligned}$$

Minimizing the right hand side of this inequality in $Q_1 \in \mathcal{C}(f, R_1)$ and in $Q_3 \in \mathcal{C}(f, R_3)$ shows that

$$MK_h(R_1, R_3)^2 \leq (1 + \alpha)\mathcal{E}_h(f, R_1)^2 + (1 + \frac{1}{\alpha})\mathcal{E}_h(f, R_3)^2.$$

Minimizing the right hand side of this inequality over $\alpha > 0$, i.e. taking

$$\alpha = \mathcal{E}_h(f, R_3)/\mathcal{E}_h(f, R_1)$$

(we recall that $\mathcal{E}_h(f, R_1) \geq \sqrt{d\hbar} > 0$), we arrive at

$$\begin{aligned} MK_h(R_1, R_3)^2 &\leq \mathcal{E}_h(f, R_1)^2 + \mathcal{E}_h(f, R_3)^2 + 2\mathcal{E}_h(f, R_1)\mathcal{E}_h(f, R_3) \\ &= (\mathcal{E}_h(f, R_1) + \mathcal{E}_h(f, R_3))^2, \end{aligned}$$

which is inequality (b). \square

The proofs of inequalities (c)-(e) are simpler because of the rank-one assumption on the intermediate point R_2 .

Proof of inequality (c). According to Lemma 3.1 (ii)

$$\begin{aligned} MK_h(R_1, R_2)^2 &= \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((R_1 \otimes R_2)^{1/2}C(x, y, \hbar D_x, \hbar D_y)(R_1 \otimes R_2)^{1/2}) \\ MK_h(R_2, R_3)^2 &= \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((R_2 \otimes R_3)^{1/2}C(y, z, \hbar D_y, \hbar D_z)(R_2 \otimes R_3)^{1/2}) \end{aligned}$$

since R_2 is a rank-one density. Applying the third inequality in Lemma 4.4 shows that

$$\begin{aligned} C_h^\epsilon(x, z, \hbar D_x, \hbar D_z) &\leq C_h(x, y, \hbar D_x, \hbar D_y) \\ &\leq (1 + \alpha)C_h(x, y, \hbar D_x, \hbar D_y) + (1 + \frac{1}{\alpha})C_h(y, z, \hbar D_y, \hbar D_z) \end{aligned}$$

so that

$$\begin{aligned} &(R_1 \otimes R_2 \otimes R_3)^{1/2}C_h^\epsilon(x, z, \hbar D_x, \hbar D_z)(R_1 \otimes R_2 \otimes R_3)^{1/2} \\ &\leq (1 + \alpha)(R_1 \otimes R_2 \otimes R_3)^{1/2}C_h(x, y, \hbar D_x, \hbar D_y)(R_1 \otimes R_2 \otimes R_3)^{1/2} \\ &\quad + (1 + \alpha)(R_1 \otimes R_2 \otimes R_3)^{1/2}C_h(y, z, \hbar D_y, \hbar D_z)(R_1 \otimes R_2 \otimes R_3)^{1/2} \\ &= (1 + \alpha)\left((R_1 \otimes R_2)^{1/2}C_h(x, y, \hbar D_x, \hbar D_y)(R_1 \otimes R_2)^{1/2}\right) \otimes R_3 \\ &\quad + (1 + \frac{1}{\alpha})R_1 \otimes \left((R_2 \otimes R_3)^{1/2}C_h(y, z, \hbar D_y, \hbar D_z)(R_2 \otimes R_3)^{1/2}\right). \end{aligned}$$

Taking the trace of both sides of this inequality in $\mathfrak{H} \otimes \mathfrak{H} \otimes \mathfrak{H}$

$$\begin{aligned}
& \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((R_1 \otimes R_3)C_h^\epsilon(x, z, \hbar D_x, \hbar D_z)) \\
&= \text{trace}_{\mathfrak{H} \otimes \mathfrak{H} \otimes \mathfrak{H}}((R_1 \otimes R_2 \otimes R_3)C_h^\epsilon(x, z, \hbar D_x, \hbar D_z)) \\
&= \text{trace}_{\mathfrak{H} \otimes \mathfrak{H} \otimes \mathfrak{H}}((R_1 \otimes R_2 \otimes R_3)^{1/2}C_h^\epsilon(x, z, \hbar D_x, \hbar D_z)(R_1 \otimes R_2 \otimes R_3)^{1/2}) \\
&\leq (1 + \alpha) \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((R_1 \otimes R_2)^{1/2}C_h(x, y, \hbar D_x, \hbar D_y)(R_1 \otimes R_2)^{1/2}) \\
&\quad + (1 + \frac{1}{\alpha}) \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((R_2 \otimes R_3)^{1/2}C_h(y, z, \hbar D_y, \hbar D_z)(R_2 \otimes R_3)^{1/2}) \\
&= (1 + \alpha)MK_h(R_1, R_2)^2 + (1 + \frac{1}{\alpha})MK_h(R_2, R_3)^2.
\end{aligned}$$

Passing to the limit as $\epsilon \rightarrow 0^+$ in the left hand side implies that

$$\begin{aligned}
MK_h(R_1, R_3)^2 &\leq \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((R_1 \otimes R_3)^{1/2}C_h(x, z, \hbar D_x, \hbar D_z)(R_1 \otimes R_3)^{1/2}) \\
&= \lim_{\epsilon \rightarrow 0^+} \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((R_1 \otimes R_3)C_h^\epsilon(x, z, \hbar D_x, \hbar D_z)) \\
&\leq (1 + \alpha)MK_h(R_1, R_2)^2 + (1 + \frac{1}{\alpha})MK_h(R_2, R_3)^2,
\end{aligned}$$

where the first inequality follows from the definition of MK_h and the fact that $R_1 \otimes R_3 \in \mathcal{C}(R_1, R_3)$, and the equality from Corollary 2.4.

Setting $\alpha := MK_h(R_2, R_3)/MK_h(R_1, R_2)$, we arrive at

$$\begin{aligned}
MK_h(R_1, R_3)^2 &\leq MK_h(R_1, R_2)^2 + MK_h(R_2, R_3)^2 + 2MK_h(R_1, R_2)MK_h(R_2, R_3) \\
&= (MK_h(R_1, R_2) + MK_h(R_2, R_3))^2
\end{aligned}$$

which is the inequality (c). \square

Proof of inequality (d). According to Lemma 3.1 (i)

$$\begin{aligned}
\mathcal{E}_h(f, R_2)^2 &= \int \text{trace}_{\mathfrak{H}}(R_2^{1/2}c_h(x, \xi)R_2^{1/2})f(x, \xi)dxd\xi \\
\mathcal{E}_h(g, R_2)^2 &= \int \text{trace}_{\mathfrak{H}}(R_2^{1/2}c_h(z, \zeta)R_2^{1/2})g(z, \zeta)dzd\zeta
\end{aligned}$$

since R_2 is a rank-one density. Applying the fourth inequality in Lemma 4.4 shows that

$$|x - z|^2 + |\xi - \zeta|^2 \leq (1 + \alpha)c_h(x, \xi; y, \hbar D_y) + (1 + \frac{1}{\alpha})c_h(z, \zeta; y, \hbar D_y),$$

so that

$$(|x - z|^2 + |\xi - \zeta|^2)R_2 \leq (1 + \alpha)R_2^{1/2}c_h(x, \xi)R_2^{1/2} + (1 + \frac{1}{\alpha})R_2^{1/2}c_h(z, \zeta)R_2^{1/2}$$

for all $x, z, \xi, \zeta \in \mathbf{R}^d$. Taking the trace of both sides of this inequality, and integrating in x, ξ, z, ζ after multiplying by $f(x, \xi)g(z, \zeta)$ shows that

$$\begin{aligned}
\text{dist}_{\text{MK}, 2}(f, g)^2 &\leq \int (|x - z|^2 + |\xi - \zeta|^2)f(x, \xi)g(z, \zeta)dxd\xi dzd\zeta \\
&= (1 + \alpha) \int \text{trace}_{\mathfrak{H}}(R_2^{1/2}c_h(x, \xi)R_2^{1/2})f(x, \xi)dxd\xi \\
&\quad + (1 + \frac{1}{\alpha}) \int \text{trace}_{\mathfrak{H}}(R_2^{1/2}c_h(z, \zeta)R_2^{1/2})g(z, \zeta)dzd\zeta \\
&= (1 + \alpha)\mathcal{E}_h(f, R_2)^2 + (1 + \frac{1}{\alpha})\mathcal{E}_h(g, R_2)^2,
\end{aligned}$$

since

$$\text{trace}_{\mathfrak{H}}(R_2) = \int f(x, \xi)dxd\xi = \int g(z, \zeta)dzd\zeta = 1.$$

The first inequality comes from the definition of the Monge-Kantorovich-Wasserstein distance $\text{dist}_{\text{MK},2}$ and the fact that $f \otimes g$ is a (nonoptimal) coupling of f and g . Choosing

$$\alpha = \mathcal{E}_h(g, R_2)/\mathcal{E}_h(f, R_2)$$

shows that

$$\begin{aligned} \text{dist}_{\text{MK},2}(f, g)^2 &\leq \mathcal{E}_h(f, R_2)^2 + \mathcal{E}_h(g, R_2)^2 + 2\mathcal{E}_h(f, R_2)\mathcal{E}_h(g, R_2) \\ &= (\mathcal{E}_h(f, R_2) + \mathcal{E}_h(g, R_2))^2 \end{aligned}$$

which is the inequality (d). \square

Proof of inequality (e). According to Lemma 3.1

$$\begin{aligned} \mathcal{E}_h(f, R_2)^2 &= \int \text{trace}_{\mathfrak{H}}(R_2^{1/2} c_h(x, \xi; y, \hbar D_y) R_2^{1/2}) f(x, \xi) dx d\xi \\ MK_h(R_2, R_3)^2 &= \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((R_2 \otimes R_3)^{1/2} C_h(y, z, \hbar D_y, \hbar D_z) (R_2 \otimes R_3)^{1/2}) \end{aligned}$$

since R_2 is a rank-one density. Applying the fifth inequality in Lemma 4.4 shows that

$$c_h(x, \xi; z, \hbar D_z) \leq (1 + \alpha) c_h(x, \xi; y, \hbar D_y) + (1 + \frac{1}{\alpha}) C_h(y, z, \hbar D_y, \hbar D_z)$$

so that, for each $\epsilon > 0$

$$0 \leq c_h^\epsilon(x, \xi; z, \hbar D_z) \leq (1 + \alpha) c_h(x, \xi; y, \hbar D_y) + (1 + \frac{1}{\alpha}) C_h(y, z, \hbar D_y, \hbar D_z)$$

with

$$\begin{aligned} c_h^\epsilon(x, \xi; z, \hbar D_z) &:= (I + \epsilon c_h(x, \xi; z, \hbar D_z))^{-1} c_h(x, \xi; z, \hbar D_z) \\ &= c_h^\epsilon(x, \xi; z, \hbar D_z)^* \in \mathcal{L}(\mathfrak{H}) \end{aligned}$$

for all $x, \xi \in \mathbf{R}^d$. Hence

$$\begin{aligned} &R_2 \otimes (R_3^{1/2} c_h^\epsilon(x, \xi; z, \hbar D_z) R_3^{1/2}) \\ &\leq (1 + \alpha) (R_2^{1/2} c_h(x, \xi; y, \hbar D_y) R_2^{1/2}) \otimes R_3 \\ &\quad + (1 + \frac{1}{\alpha}) (R_2 \otimes R_3)^{1/2} C_h(y, z, \hbar D_y, \hbar D_z) (R_2 \otimes R_3)^{1/2} \end{aligned}$$

for all $x, \xi \in \mathbf{R}^d$ and, taking the trace of both sides of this inequality leads to

$$\begin{aligned} (12) \quad &\text{trace}_{\mathfrak{H}}(R_3^{1/2} c_h^\epsilon(x, \xi; z, \hbar D_z) R_3^{1/2}) \\ &\leq (1 + \alpha) \text{trace}_{\mathfrak{H}}(R_2^{1/2} c_h(x, \xi; y, \hbar D_y) R_2^{1/2}) + (1 + \frac{1}{\alpha}) MK_h(R_2, R_3)^2. \end{aligned}$$

Multiplying both sides of this inequality by $f(x, \xi)$ and integrating in (x, ξ) shows that

$$\begin{aligned} &\int \text{trace}_{\mathfrak{H}}(R_3^{1/2} c_h^\epsilon(x, \xi; z, \hbar D_z) R_3^{1/2}) f(x, \xi) dx d\xi \\ &\leq (1 + \alpha) \mathcal{E}_h(f, R_2)^2 + (1 + \frac{1}{\alpha}) MK_h(R_2, R_3)^2 \\ &\quad = (\mathcal{E}_h(f, R_2) + MK_h(R_2, R_3))^2, \end{aligned}$$

with the choice

$$\alpha := MK_h(R_2, R_3)/\mathcal{E}_h(f, R_2).$$

Since the right-hand side of (12) is integrable with respect to $f(x, \xi) dx d\xi$, and therefore finite for $f(x, \xi) dx d\xi$ -a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$, one has

$$\text{trace}_{\mathfrak{H}}(R_3^{1/2} c_h^\epsilon(x, \xi; z, \hbar D_z) R_3^{1/2}) \rightarrow \text{trace}_{\mathfrak{H}}(R_3^{1/2} c_h(x, \xi; z, \hbar D_z) R_3^{1/2})$$

for $f(x, \xi)dxd\xi$ -a.e. $(x, \xi) \in \mathbf{R}^d \times \mathbf{R}^d$ by Corollary 2.4. By Fatou's lemma, observing that $f \otimes R_3 \in \mathcal{C}(f, R_3)$, one has

$$\begin{aligned} \mathcal{E}_\hbar(f, R_3) &\leq \int \text{trace}_{\mathfrak{H}}(R_3^{1/2} c_\hbar(x, \xi; z, \hbar D_z) R_3^{1/2}) f(x, \xi) dx d\xi \\ &\leq \varliminf_{\epsilon \rightarrow 0^+} \int \text{trace}_{\mathfrak{H}}(R_3^{1/2} c_\hbar^\epsilon(x, \xi; z, \hbar D_z) R_3^{1/2}) f(x, \xi) dx d\xi \\ &\leq (\mathcal{E}_\hbar(f, R_2) + MK_\hbar(R_2, R_3))^2, \end{aligned}$$

which is the inequality (e). \square

5. APPLICATIONS

One satisfying consequence of the triangle inequalities proved in the last section is the following statement, which confirms that MK_\hbar can indeed be thought of as a quantum deformation of the quadratic Monge-Kantorovich-Wasserstein distance.

Theorem 5.1. *Let R_\hbar, S_\hbar be families of density operators in $\mathcal{D}_2(\mathfrak{H})$, and let $f, g \in \mathcal{P}_2^{ac}(\mathbf{R}^d \times \mathbf{R}^d)$. Assume that*

$$\mathcal{E}_\hbar(f, R_\hbar) \rightarrow 0 \quad \text{and} \quad \mathcal{E}_\hbar(g, S_\hbar) \rightarrow 0$$

as $\hbar \rightarrow 0$. Then

$$\lim_{\hbar \rightarrow 0} MK_\hbar(R_\hbar, S_\hbar) = \text{dist}_{\text{MK},2}(f, g).$$

This statement is to be compared with the lower bound

$$MK_\hbar(R_\hbar, S_\hbar)^2 \geq \text{dist}_{\text{MK},2}(\widetilde{W}_\hbar(R_\hbar), \widetilde{W}_\hbar(S_\hbar))^2 - 2d\hbar,$$

which is Theorem 2.3 (2) in [6], and with the upper bound obtained in the special case of Töplitz operators

$$MK_\hbar(\text{OP}_\hbar^T((2\pi\hbar)^d \mu), \text{OP}_\hbar^T((2\pi\hbar)^d \nu))^2 \leq \text{dist}_{\text{MK},2}(\mu, \nu)^2 + 2d\hbar,$$

stated as Theorem 2.3 (1) in [6].

Proof. By Theorem 4.2 (a)-(b),

$$\begin{aligned} MK_\hbar(R_\hbar, S_\hbar) &\leq \mathcal{E}_\hbar(f, R_\hbar) + \mathcal{E}_\hbar(f, S_\hbar) \\ &\leq \mathcal{E}_\hbar(f, R_\hbar) + \text{dist}_{\text{MK},2}(f, g) + \mathcal{E}_\hbar(g, S_\hbar). \end{aligned}$$

Hence

$$\overline{\lim}_{\hbar \rightarrow 0^+} MK_\hbar(R_\hbar, S_\hbar) \leq \text{dist}_{\text{MK},2}(f, g).$$

By Theorem 2.4 (2) in [7]

$$\text{dist}_{\text{MK},2}(f, \widetilde{W}(R_\hbar))^2 \leq \mathcal{E}_\hbar(f, R_\hbar)^2 + d\hbar$$

(notice the slight change of normalization in the definition of \mathcal{E}_\hbar between [7] and the present paper), so that our assumption implies that

$$\text{dist}_{\text{MK},2}(f, \widetilde{W}(R_\hbar)) \rightarrow 0 \quad \text{and} \quad \text{dist}_{\text{MK},2}(g, \widetilde{W}(S_\hbar)) \rightarrow 0$$

as $\hbar \rightarrow 0$. From the inequality

$$\text{dist}_{\text{MK},2}(\widetilde{W}_\hbar(R_\hbar), \widetilde{W}_\hbar(S_\hbar))^2 \leq MK_\hbar(R_\hbar, S_\hbar)^2 + 2d\hbar,$$

(Theorem 2.3 (2) in [6]), we deduce that

$$\text{dist}_{\text{MK},2}(f, g) \leq \varliminf_{\hbar \rightarrow 0} MK_\hbar(R_\hbar, S_\hbar).$$

Notice that this last lower bound is a variant of the last inequality in Theorem 2.3 of [6], except that in the present case the assumption on R_h and S_h is slightly different (in other words, we have assumed that $\mathcal{E}_h(f, R_h) \rightarrow 0$ instead of assuming that $\widetilde{W}_h(R_h) \rightarrow f$ in $\mathcal{S}'(\mathbf{R}^d \times \mathbf{R}^d)$.) \square

6. KANTOROVICH DUALITY FOR \mathcal{E}_h

Theorem 6.1. *Let $S \in \mathcal{D}_2(\mathfrak{H})$ and let $p \equiv p(x, \xi)$ be a probability density on \mathbf{R}^{2d} such that*

$$\int_{\mathbf{R}^{2d}} (|x|^2 + |\xi|^2) p(x, \xi) dx d\xi < +\infty.$$

Then

$$\begin{aligned} \mathcal{E}_h(p, S)^2 &= \min_{Q \in \mathcal{C}(p, S)} \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(Q(x, \xi)^{1/2} c(x, \xi) Q(x, \xi)^{1/2}) dx d\xi \\ &= \sup_{\substack{a \in C_b(\mathbf{R}^{2d}), B = B^* \in \mathcal{L}(\mathfrak{H}) \\ a(x, \xi) I_{\mathfrak{H}} + B \leq c(x, \xi)}} \left(\int_{\mathbf{R}^{2d}} a(x, \xi) p(x, \xi) dx d\xi + \text{trace}_{\mathfrak{H}}(BS) \right). \end{aligned}$$

Notice that the duality theorem implies in particular the existence of at least one optimal coupling $Q \in \mathcal{C}(p, S)$.

Proof. The proof is split in several steps.

Step 1: the functions f and g . Consider the Banach space $E := C_b(\mathbf{R}^{2d}; \mathcal{L}(\mathfrak{H}))$, with

$$\|T\|_E := \sup_{(x, \xi) \in \mathbf{R}^{2d}} \|T(x, \xi)\|,$$

and set

$$f(T) := \begin{cases} 0 & \text{if } T(x, \xi) = T(x, \xi)^* \geq -c(x, \xi) \text{ for all } (x, \xi) \in \mathbf{R}^{2d}, \\ +\infty & \text{otherwise,} \end{cases}$$

while

$$g(T) := \begin{cases} \int_{\mathbf{R}^{2d}} ap(x, \xi) dx d\xi + \text{trace}_{\mathfrak{H}}(BS) & \text{if } T(x, \xi) = T(x, \xi)^* = a(x, \xi) I_{\mathfrak{H}} + B \\ & \text{for all } (x, \xi) \in \mathbf{R}^{2d}, \\ +\infty & \text{otherwise,} \end{cases}$$

The constraint $T(x, \xi) = T(x, \xi)^* \geq -c(x, \xi)$ means that, for each $(x, \xi) \in \mathbf{R}^{2d}$, one has

$$\langle \phi(x, \xi) | T(x, \xi) + c(x, \xi) | \phi(x, \xi) \rangle \geq 0$$

for all $\phi \in \text{Form-Dom}(c(x, \xi))$. On the other hand, the nullspace of the linear map

$$C_b(\mathbf{R}^{2d}) \times \mathcal{L}(\mathfrak{H}) \ni (a, B) \mapsto \Gamma(a, B) \equiv a(x, \xi) I_{\mathfrak{H}} + B \in E$$

is

$$\text{Ker}(\mathcal{L}) = \{(t, -tI_{\mathfrak{H}}), \quad t \in \mathbf{R}\}.$$

Since

$$g((a+t)I_{\mathfrak{H}} + (B-tI_{\mathfrak{H}})) = g(aI_{\mathfrak{H}} + B) + t \int_{\mathbf{R}^{2d}} p(x, \xi) dx d\xi - t \text{trace}_{\mathfrak{H}}(S) = g(aI_{\mathfrak{H}} + B),$$

the prescription above defines g on $\text{Ran}(\Gamma) \simeq (C_b(\mathbf{R}^{2d}) \times \mathcal{L}(\mathfrak{H}))/\text{Ker}(\Gamma)$. Observe that

$$\begin{aligned} g((aI_{\mathfrak{H}} + B)^*) &= g(\bar{a}I_{\mathfrak{H}} + B^*) = \int_{\mathbf{R}^{2d}} \overline{a(x, \xi)} p(x, \xi) dx d\xi + \text{trace}_{\mathfrak{H}}(B^* S) \\ &= \overline{\int_{\mathbf{R}^{2d}} a(x, \xi) p(x, \xi) dx d\xi} + \text{trace}_{\mathfrak{H}}((SB)^*) \\ &= \overline{\int_{\mathbf{R}^{2d}} a(x, \xi) p(x, \xi) dx d\xi} + \overline{\text{trace}_{\mathfrak{H}}(SB)} \\ &= \overline{g(aI_{\mathfrak{H}} + B)}, \end{aligned}$$

so that $(aI_{\mathfrak{H}} + B)^* = aI_{\mathfrak{H}} + B \implies g(aI_{\mathfrak{H}} + B) \in \mathbf{R}$. Thus the definition above implies that g takes its values in $(-\infty, +\infty]$.

The functions f and g are convex. Indeed, g is the extension by $+\infty$ of a \mathbf{R} -linear functional defined on the set of self-adjoint elements of $\text{Ran}(\Gamma)$, which is a linear subspace of E . As for f , it is the indicator function (in the sense of the definition in §4 of [11] on p. 28) of the convex set

$$\{T \in E \text{ s.t. } T(x, \xi) = T(x, \xi)^* \geq -c(x, \xi) \text{ for all } (x, \xi) \in \mathbf{R}^{2d}\}$$

and is therefore convex. Besides $f(0) = g(0) = 0$, and f is continuous at 0. Indeed, by the Heisenberg inequality

$$c(x, \xi) \geq d\hbar I_{\mathfrak{H}} \quad \text{for all } (x, \xi) \in \mathbf{R}^{2d},$$

so that, for each $T \in E$

$$\begin{aligned} T(x, \xi) = T(x, \xi)^* \text{ and } \|T(x, \xi)\| &< \frac{1}{2}d\hbar \text{ for all } (x, \xi) \in \mathbf{R}^{2d} \\ \implies T(x, \xi) &\geq -c(x, \xi) \text{ for all } (x, \xi) \in \mathbf{R}^{2d} \implies f(T) = 0. \end{aligned}$$

In particular f is continuous at 0.

Step 2: applying convex duality. By the Fenchel-Rockafellar convex duality theorem (Theorem 1.12 in [1])

$$\inf_{T \in E} (f(T) + g(T)) = \max_{\Lambda \in E'} (-f^*(-\Lambda) - g^*(\Lambda)).$$

Let us compute the Legendre duals f^* and g^* .

First

$$f^*(-\Lambda) = \sup_{T \in E} (\langle -\Lambda, T \rangle - f(T)) = \sup_{\substack{T \in E \\ T(x, \xi) = T(x, \xi)^* \geq -c(x, \xi)}} \langle -\Lambda, T \rangle.$$

If $\Lambda \in E'$ is not a nonnegative linear functional, there exists $T_0 \in E$ such that $T_0(x, \xi) = T_0(x, \xi)^* \geq 0$ such that $\langle \Lambda, T_0 \rangle = -\alpha < 0$. Since

$$nT_0(x, \xi) = nT_0(x, \xi)^* \geq 0 \geq -d\hbar I_{\mathfrak{H}} \geq -c(x, \xi) \text{ for all } (x, \xi) \in \mathbf{R}^{2d}$$

one has

$$f^*(-\Lambda) \geq \sup_{n \geq 1} \langle -\Lambda, nT_0 \rangle = \sup_{n \geq 1} (n\alpha) = +\infty.$$

For $\Lambda \in E'$ such that $\Lambda \geq 0$, we define

$$\langle \Lambda, c \rangle := \sup_{\substack{T \in E \\ T(x, \xi) = T(x, \xi)^* \leq c(x, \xi)}} \langle \Lambda, T \rangle \in [0, +\infty].$$

(Observe indeed that $T = 0$ satisfies the constraints since $c(x, \xi) = c(x, \xi)^* \geq 0$ for each $(x, \xi) \in \mathbf{R}^{2d}$.) With this definition, one has clearly

$$f^*(-\Lambda) := \begin{cases} \langle \Lambda, c \rangle & \text{if } \Lambda \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

Next

$$\begin{aligned} g^*(\Lambda) &= \sup_{T \in E} (\langle \Lambda, T \rangle - g(T)) \\ &= \sup_{\substack{T \in E \\ T(x, \xi) = T(x, \xi)^* = a(x, \xi)I_{\mathfrak{H}} + B}} \left(\langle \Lambda, T \rangle - \int_{\mathbf{R}^{2d}} a(x, \xi)p(x, \xi)dx d\xi - \text{trace}_{\mathfrak{H}}(BS) \right). \end{aligned}$$

If there exists $a \equiv a(x, \xi) \in C_b(\mathbf{R}^{2d}, \mathbf{R})$ and $B = B^* \in \mathcal{L}(\mathfrak{H})$ such that either

$$\langle \Lambda, aI_{\mathfrak{H}} + B \rangle > \int_{\mathbf{R}^{2d}} a(x, \xi)p(x, \xi)dx d\xi + \text{trace}_{\mathfrak{H}}(BS)$$

or

$$\langle \Lambda, aI_{\mathfrak{H}} + B \rangle < \int_{\mathbf{R}^{2d}} a(x, \xi)p(x, \xi)dx d\xi + \text{trace}_{\mathfrak{H}}(BS),$$

one has either

$$g(\Lambda) \geq \sup_{n \geq 1} \left(\langle \Lambda, n(aI_{\mathfrak{H}} + B) \rangle - n \int_{\mathbf{R}^{2d}} a(x, \xi)p(x, \xi)dx d\xi - n \text{trace}_{\mathfrak{H}}(BS) \right) = +\infty,$$

or

$$g(\Lambda) \geq \sup_{n \geq 1} \left(\langle \Lambda, n(-aI_{\mathfrak{H}} - B) \rangle + n \int_{\mathbf{R}^{2d}} a(x, \xi)p(x, \xi)dx d\xi + n \text{trace}_{\mathfrak{H}}(BS) \right) = +\infty.$$

Hence

$$g^*(\Lambda) := \begin{cases} 0 & \text{if } \langle \Lambda, aI_{\mathfrak{H}} + B \rangle = \int_{\mathbf{R}^{2d}} a(x, \xi)p(x, \xi)dx d\xi + \text{trace}_{\mathfrak{H}}(BS) \\ & \text{for each } a \equiv a(x, \xi) \in C_b(\mathbf{R}^{2d}, \mathbf{R}) \text{ and } B = B^* \in \mathcal{L}(\mathfrak{H}), \\ +\infty & \text{otherwise.} \end{cases}$$

Notice that the prescription

$$\langle \Lambda, aI_{\mathfrak{H}} + B \rangle = \int_{\mathbf{R}^{2d}} a(x, \xi)p(x, \xi)dx d\xi + \text{trace}_{\mathfrak{H}}(BS)$$

defines a unique linear functional on the set of $T \in \text{Ran } \Gamma$ such that $T(x, \xi)^* = T(x, \xi)$ for each $(x, \xi) \in \mathbf{R}^{2d}$ by the same argument as in Step 1.

Therefore, the Fenchel-Rockafellar duality theorem in this case results in the equality

$$\begin{aligned} \inf_{T \in E} (f(T) + g(T)) &= \inf_{\substack{a \in C_b(\mathbf{R}^{2d}, \mathbf{R}), B = B^* \\ a(x, \xi)I_{\mathfrak{H}} + B \geq -c(x, \xi)}} \left(\int_{\mathbf{R}^{2d}} a(x, \xi)p(x, \xi)dx d\xi + \text{trace}_{\mathfrak{H}}(SB) \right) \\ &= \max_{\Lambda \in E'} (f^*(-\Lambda) + g^*(\Lambda)) = \max_{\substack{0 \leq \Lambda \in E', \langle \Lambda, aI_{\mathfrak{H}} + B \rangle \\ = \int_{\mathbf{R}^{2d}} a(x, \xi)p(x, \xi)dx d\xi + \text{trace}_{\mathfrak{H}}(SB)}} -\langle \Lambda, c \rangle \end{aligned}$$

or, equivalently

$$\begin{aligned} \sup_{\substack{a \in C_b(\mathbf{R}^{2d}, \mathbf{R}), B = B^* \\ a(x, \xi)I_{\mathfrak{H}} + B \leq c(x, \xi)}} \left(\int_{\mathbf{R}^{2d}} a(x, \xi)p(x, \xi)dx d\xi + \text{trace}_{\mathfrak{H}}(SB) \right) \\ = \min_{\substack{0 \leq \Lambda \in E', \langle \Lambda, aI_{\mathfrak{H}} + B \rangle \\ = \int_{\mathbf{R}^{2d}} a(x, \xi)p(x, \xi)dx d\xi + \text{trace}_{\mathfrak{H}}(SB)}} \langle \Lambda, c \rangle. \end{aligned}$$

Step 3: representing the optimal Λ . Define a linear map $F_\Lambda : C_b(\mathbf{R}^{2d}) \rightarrow \mathcal{L}^1(\mathfrak{H})$ by the formula

$$\text{trace}_{\mathfrak{H}}(KF_\Lambda(a)) = \Lambda(aK), \quad \text{for each } K \in \mathcal{K}(\mathfrak{H}).$$

Indeed, since $K \mapsto \Lambda(aK)$ is a linear functional on $\mathcal{K}(\mathfrak{H})$ which is continuous for the norm topology, and since $\mathcal{K}(\mathfrak{H})' = \mathcal{L}^1(\mathfrak{H})$, this linear functional is represented by a trace-class operator $F_\Lambda(a)$. Since Λ is linear, the map F_Λ is linear.

Since $\Lambda \geq 0$, one has $F_\Lambda(a) = F_\Lambda(a)^* \geq 0$ for each $a \in C_b(\mathbf{R}^d)$ such that $a(x, \xi) \geq 0$ for each $(x, \xi) \in \mathbf{R}^{2d}$. Indeed, for $a \in C_b(\mathbf{R}^{2d}; \mathbf{R})$, set

$$T_1 := \frac{1}{2}(F_\Lambda(a) + F_\Lambda(a)^*), \quad T_2 := -\frac{1}{2}i(F_\Lambda(a) - F_\Lambda(a)^*).$$

Then, for each $K = K^* \in \mathcal{K}(\mathfrak{H})$, one has

$$\Lambda(aK) = \text{trace}_{\mathfrak{H}}(T_1K) + i \text{trace}_{\mathfrak{H}}(T_2K)$$

with

$$\overline{\text{Tr}_{\mathfrak{H}}(T_jK)} = \text{Tr}_{\mathfrak{H}}((T_jK)^*) = \text{Tr}_{\mathfrak{H}}(K^*T_j^*) = \text{Tr}_{\mathfrak{H}}(KT_j) = \text{Tr}_{\mathfrak{H}}(T_jK)$$

for $j = 1, 2$. Since $a \in C_b(\mathbf{R}^{2d}; \mathbf{R})$ and $K = K^* \in \mathcal{L}(\mathfrak{H})$, one has

$$-\|a\|_{L^\infty} \|K\|_{I_{\mathfrak{H}}} \leq aK \leq \|a\|_{L^\infty} \|K\|_{I_{\mathfrak{H}}}$$

so that

$$-\|a\|_{L^\infty} \|K\| \leq \Lambda(aK) \leq \|a\|_{L^\infty} \|K\| \quad \text{since } \Lambda(I_{\mathfrak{H}}) = \int_{\mathbf{R}^{2d}} p(x, \xi) dx d\xi = 1.$$

In particular, $\Lambda(aK) \in \mathbf{R}$, so that $\text{trace}_{\mathfrak{H}}(T_2K) = 0$ for each $K = K^* \in \mathcal{K}(\mathfrak{H})$. Since $T_2 = T_2^* \in \mathcal{L}^1(\mathfrak{H})$, specializing this identity to the case where K is the orthogonal projection on any eigenvector of T_2 shows that $T_2 = 0$. Thus

$$a \in C_b(\mathbf{R}^{2d}; \mathbf{R}) \implies F_\Lambda(a) = F_\Lambda(a)^*.$$

Moreover

$$a \in C_b(\mathbf{R}^{2d}; \mathbf{R}) \text{ and } a \geq 0 \implies \text{trace}_{\mathfrak{H}}(F_\Lambda(a)K) \geq 0 \text{ for each } K = K^* \geq 0 \text{ in } \mathcal{K}(\mathfrak{H})$$

and specializing this last inequality to the case where K is the orthogonal projection on any eigenvector of $F_\Lambda(a) = F_\Lambda(a)^* \in \mathcal{L}^1(\mathfrak{H})$ shows that all the eigenvalues of $F_\Lambda(a)$ are nonnegative, so that $F_\Lambda(a) \geq 0$.

Next we deduce from the defining identity for F_Λ , i.e.

$$\Lambda(aK) = \text{trace}_{\mathfrak{H}}(F_\Lambda(a)K) \text{ for each } a \in C_b(\mathbf{R}^{2d}; \mathbf{C}) \text{ and } K \in \mathcal{K}(\mathfrak{H})$$

that

$$\|F_\Lambda(a)\|_1 \leq \|\Lambda\| \|a\|_{L^\infty(\mathbf{R}^{2d})}.$$

Next we specialize this defining identity to the case where $a \geq 0$ on \mathbf{R}^{2d} while $K = \Pi_n$ is the orthogonal projection on $\text{span}\{e_1, \dots, e_n\}$, with (e_1, e_2, \dots) a complete orthonormal system in \mathfrak{H} . One has

$$\Lambda(a\Pi_n) = \text{trace}_{\mathfrak{H}}(F_\Lambda(a)\Pi_n) \rightarrow \text{trace}_{\mathfrak{H}}(F_\Lambda(a)) = \|F_\Lambda(a)\|_1 \quad \text{as } n \rightarrow \infty$$

while

$$a(I_{\mathfrak{H}} - \Pi_n) \geq 0 \text{ so that } \Lambda(a\Pi_n) \leq \Lambda(aI_{\mathfrak{H}}) = \int_{\mathbf{R}^{2d}} a(x, \xi) p(x, \xi) dx d\xi$$

so that

$$a \in C_b(\mathbf{R}^{2d}) \text{ and } a \geq 0 \implies \|F_\Lambda(a)\|_1 \leq \int_{\mathbf{R}^{2d}} a(x, \xi) p(x, \xi) dx d\xi.$$

More generally, for each $a \in C_b(\mathbf{R}^{2d}; \mathbf{R})$, one has $-|a| \leq a \leq |a|$ so that

$$|\text{trace}_{\mathfrak{H}}(F_\Lambda(a)|e_j\rangle\langle e_j|)| = |\Lambda(a|e_j\rangle\langle e_j|)| \leq \Lambda(|a||e_j\rangle\langle e_j|)$$

for each $j \geq 1$, where (e_1, e_2, \dots) is a complete orthonormal system of eigenvectors of $F_\Lambda(a) = F_\Lambda(a)^* \in \mathcal{L}^1(\mathfrak{H})$. Hence

$$\sum_{j=1}^n |\text{trace}_{\mathfrak{H}}(F_\Lambda(a)|e_j\rangle\langle e_j|)| \leq \Lambda\left(|a| \sum_{j=1}^n |e_j\rangle\langle e_j|\right) \leq \Lambda(|a|I_{\mathfrak{H}}),$$

and since

$$\sum_{j=1}^n |\text{trace}_{\mathfrak{H}}(F_\Lambda(a)|e_j\rangle\langle e_j|)| \rightarrow \|F_\Lambda(a)\|_1 \quad \text{as } n \rightarrow \infty$$

we conclude that

$$\|F_\Lambda(a)\|_1 \leq \Lambda(|a|I_{\mathfrak{H}}) = \int_{\mathbf{R}^{2d}} |a(x, \xi)| p(x, \xi) dx d\xi.$$

Since $C_b(\mathbf{R}^{2d})$ is dense in $L^1(\mathbf{R}^{2d}, p dx d\xi)$, this inequality, applied to the real and the imaginary part of a , shows that F_Λ is a continuous linear operator from $L^1(\mathbf{R}^{2d})$ to $\mathcal{L}^1(\mathfrak{H})$. Since $\mathcal{L}^1(\mathfrak{H})$ is separable and is the dual of the Banach space $\mathcal{K}(\mathfrak{H})$ (the norm closure in $\mathcal{L}(\mathfrak{H})$ of the set of finite rank operators), we conclude from the Dunford-Pettis theorem (Theorem 1 in §3 of chapter III in [5]) that $\mathcal{L}^1(\mathfrak{H})$ has the Radon-Nikodym property. By Theorem 5 in §1 of chapter III in [5], the operator F_Λ is Riesz-representable: in other words, there exists $q \in L^\infty(\mathbf{R}^{2d}, p dx d\xi; \mathcal{L}^1(\mathfrak{H}))$ such that

$$F_\Lambda(a) = \int_{\mathbf{R}^{2d}} a(x, \xi) q(x, \xi) p(x, \xi) dx d\xi, \quad \text{for all } a \in L^1(\mathbf{R}^{2d}, p dx d\xi).$$

Step 4: defining the optimal coupling. We have seen that

$$a \in C_b(\mathbf{R}^{2d}) \text{ and } a \geq 0 \implies F_\Lambda(a) = \int_{\mathbf{R}^{2d}} a(x, \xi) q(x, \xi) p(x, \xi) dx d\xi \geq 0.$$

This implies that $q(x, \xi) = q(x, \xi)^* \geq 0$ for a.e. $(x, \xi) \in \mathbf{R}^{2d}$.

Next, one has

$$\Lambda(K) = \text{trace}_{\mathfrak{H}}(F_\Lambda(1)K) = \text{trace}_{\mathfrak{H}}(KS), \quad K \in \mathcal{K}(\mathfrak{H}),$$

so that

$$F_\Lambda(1) = \int_{\mathbf{R}^{2d}} q(x, \xi) p(x, \xi) dx d\xi = S \in \mathcal{L}^1(\mathfrak{H}) = \mathcal{K}(\mathfrak{H})'.$$

On the other hand, for each $a \in C_b(\mathbf{R}^{2d})$ such that $a \geq 0$, one has

$$\begin{aligned} \text{trace}_{\mathfrak{H}}\left(P_n \int_{\mathbf{R}^{2d}} a(x, \xi) q(x, \xi) p(x, \xi) dx d\xi\right) &= \text{trace}_{\mathfrak{H}}(F_\Lambda(a)P_n) = \Lambda(aP_n) \\ &\leq \Lambda(aI_{\mathfrak{H}}) = \int_{\mathbf{R}^{2d}} a(x, \xi) p(x, \xi) dx d\xi \end{aligned}$$

where P_n is the orthogonal projection on $\text{span}\{e_1, \dots, e_n\}$, with (e_1, e_2, \dots) being a complete orthonormal system of eigenvectors of

$$\int_{\mathbf{R}^{2d}} a(x, \xi) q(x, \xi) p(x, \xi) dx d\xi \in \mathcal{L}^1(\mathfrak{H}).$$

Letting $n \rightarrow \infty$, one has

$$\begin{aligned} &\text{trace}_{\mathfrak{H}}\left(P_n \int_{\mathbf{R}^{2d}} a(x, \xi) q(x, \xi) p(x, \xi) dx d\xi\right) \\ &\rightarrow \text{trace}_{\mathfrak{H}}\left(\int_{\mathbf{R}^{2d}} a(x, \xi) q(x, \xi) p(x, \xi) dx d\xi\right), \end{aligned}$$

so that

$$\int_{\mathbf{R}^{2d}} a(x, \xi) \operatorname{trace}_{\mathfrak{H}}(q(x, \xi)) p(x, \xi) dx d\xi \leq \int_{\mathbf{R}^{2d}} a(x, \xi) d(x, \xi) dx d\xi.$$

Since this holds for each $a \in C_b(\mathbf{R}^{2d})$ such that $a \geq 0$, we conclude that

$$\operatorname{trace}_{\mathfrak{H}}(q(x, \xi)) \leq 1 \quad \text{for } p(x, \xi) dx d\xi\text{-a.e. } (x, \xi) \in \mathbf{R}^{2d}.$$

Moreover

$$\int_{\mathbf{R}^{2d}} (1 - \operatorname{trace}_{\mathfrak{H}}(q(x, \xi))) p(x, \xi) dx d\xi = 1 - \operatorname{trace}_{\mathfrak{H}}(S) = 0$$

so that

$$\operatorname{trace}_{\mathfrak{H}}(q(x, \xi)) = 1 \quad \text{for } p(x, \xi) dx d\xi\text{-a.e. } (x, \xi) \in \mathbf{R}^{2d}.$$

In other words, we have proved that $(x, \xi) \mapsto Q(x, \xi) = p(x, \xi)q(x, \xi)$ defines an element of $\mathcal{C}(p, S)$.

Step 5: extending the representation formula for Λ . For each $B \in E$, we define

$$\langle L, B \rangle := \langle \Lambda, B \rangle - \int_{\mathbf{R}^{2d}} \operatorname{trace}_{\mathfrak{H}}(B(x, \xi)Q(x, \xi)) dx d\xi.$$

Let us prove that

$$B \in E \text{ and } B(x, \xi) = B(x, \xi)^* \geq 0 \text{ for all } (x, \xi) \in \mathbf{R}^{2d} \implies \langle L, B \rangle \geq 0.$$

Pick $\epsilon > 0$, and let Q_ϵ be a simple $\mathcal{L}^1(\mathfrak{H})$ -valued function on \mathbf{R}^{2d} such that

$$\int_{\mathbf{R}^{2d}} \|Q(x, \xi) - Q_\epsilon(x, \xi)\|_1 dx d\xi < \epsilon.$$

Write

$$Q_\epsilon(x, \xi) = \sum_{j=1}^N \mathbf{1}_{\Omega_j}(x, \xi) Q_j, \quad 0 \leq Q_j = Q_j^* \in \mathcal{L}^1(\mathfrak{H}) \text{ for each } j = 0, \dots, N,$$

where Ω_j are bounded, pairwise disjoint measurable sets in \mathbf{R}^{2d} for $j = 1, \dots, N$. For each $j = 1, \dots, N$, let $(e_{j,1}, e_{j,2}, \dots)$ designate a complete orthonormal system of eigenvectors of Q_j , and let $P_{j,n}$ be the orthogonal projection on $\operatorname{span}\{e_{j,1}, \dots, e_{j,n}\}$. Define

$$\Pi_n(x, \xi) = \sum_{j=1}^N \mathbf{1}_{\Omega_j}(x, \xi) P_{j,n}.$$

One easily checks that $\Pi_n(x, \xi) = \Pi_n(x, \xi)^* = \Pi_n(x, \xi)^2$ for each $(x, \xi) \in \mathbf{R}^{2d}$. Then, for each $B \in E$ such that $B(x, \xi) = B(x, \xi)^* \geq 0$ for all $(x, \xi) \in \mathbf{R}^{2d}$, one has

$$\begin{aligned} 0 &\leq \langle \Lambda, (I_{\mathfrak{H}} - \Pi_n)B(I_{\mathfrak{H}} - \Pi_n) \rangle = \langle \Lambda, B \rangle - \langle \Lambda, \Pi_n B + B \Pi_n - \Pi_n B \Pi_n \rangle \\ &= \langle \Lambda, B \rangle - \int_{\mathbf{R}^{2d}} \operatorname{trace}_{\mathfrak{H}}((\Pi_n B + B \Pi_n - \Pi_n B \Pi_n)Q)(x, \xi) dx d\xi \\ &= \langle \Lambda, B \rangle - \int_{\mathbf{R}^{2d}} \operatorname{trace}_{\mathfrak{H}}((\Pi_n B + B \Pi_n - \Pi_n B \Pi_n)Q_\epsilon)(x, \xi) dx d\xi \\ &\quad + \int_{\mathbf{R}^{2d}} \operatorname{trace}_{\mathfrak{H}}((\Pi_n B + B \Pi_n - \Pi_n B \Pi_n)(Q_\epsilon - Q))(x, \xi) dx d\xi. \end{aligned}$$

By construction, keeping $\epsilon > 0$ fixed, one has

$$\begin{aligned} &\int_{\mathbf{R}^{2d}} \operatorname{trace}_{\mathfrak{H}}((\Pi_n B + B \Pi_n - \Pi_n B \Pi_n)Q_\epsilon)(x, \xi) dx d\xi \\ &= \int_{\mathbf{R}^{2d}} \operatorname{trace}_{\mathfrak{H}}(B \Pi_n Q_\epsilon \Pi_n)(x, \xi) dx d\xi \rightarrow \int_{\mathbf{R}^{2d}} \operatorname{trace}_{\mathfrak{H}}(B Q_\epsilon)(x, \xi) dx d\xi \end{aligned}$$

as $n \rightarrow \infty$, so that

$$\begin{aligned} 0 \leq \overline{\lim}_{n \rightarrow \infty} \langle \Lambda, (I_{\mathfrak{H}} - \Pi_n)B(I_{\mathfrak{H}} - \Pi_n) \rangle &= \langle \Lambda, B \rangle - \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(BQ)(x, \xi) dx d\xi \\ &+ \overline{\lim}_{n \rightarrow \infty} \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}((\Pi_n B + B\Pi_n - \Pi_n B\Pi_n)(Q_\epsilon - Q))(x, \xi) dx d\xi \\ &+ \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(B(Q - Q_\epsilon))(x, \xi) dx d\xi. \end{aligned}$$

On the other hand

$$\begin{aligned} &\left| \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}((\Pi_n B + B\Pi_n - \Pi_n B\Pi_n)(Q_\epsilon - Q))(x, \xi) dx d\xi \right| \\ &\leq \int_{\mathbf{R}^{2d}} |\text{trace}_{\mathfrak{H}}((\Pi_n B + B\Pi_n - \Pi_n B\Pi_n)(Q_\epsilon - Q))(x, \xi)| dx d\xi \\ &\leq \int_{\mathbf{R}^{2d}} \|(\Pi_n B + B\Pi_n - \Pi_n B\Pi_n)(x, \xi)\| \|(Q_\epsilon - Q)(x, \xi)\|_1 dx d\xi \\ &\leq 3 \sup_{(x, \xi) \in \mathbf{R}^{2d}} \|B(x, \xi)\| \int_{\mathbf{R}^{2d}} \|(Q_\epsilon - Q)(x, \xi)\|_1 dx d\xi \\ &\leq 3\epsilon \sup_{(x, \xi) \in \mathbf{R}^{2d}} \|B(x, \xi)\| \end{aligned}$$

while, by the same token,

$$\left| \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(B(Q - Q_\epsilon))(x, \xi) dx d\xi \right| \leq \epsilon \sup_{(x, \xi) \in \mathbf{R}^{2d}} \|B(x, \xi)\|.$$

Finally

$$\langle \Lambda, B \rangle - \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(BQ)(x, \xi) dx d\xi \geq -4\epsilon \sup_{(x, \xi) \in \mathbf{R}^{2d}} \|B(x, \xi)\|$$

and since this holds for each $\epsilon > 0$, we conclude that

$$B \in E \text{ and } B(x, \xi) = B(x, \xi)^* \geq 0 \text{ for all } (x, \xi) \in \mathbf{R}^{2d} \implies \langle L, B \rangle \geq 0.$$

By a classical argument, this implies that $\|L\| = \langle L, I_{\mathfrak{H}} \rangle$.

On the other hand

$$\begin{aligned} \langle L, I_{\mathfrak{H}} \rangle &= \langle L, I_{\mathfrak{H}} \rangle - \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(q(x, \xi))p(x, \xi) dx d\xi \\ &= \text{trace}_{\mathfrak{H}}(S) - \int_{\mathbf{R}^{2d}} p(x, \xi) dx d\xi = 0 \end{aligned}$$

so that $L = 0$. In other words, the representation formula

$$\langle \Lambda, B \rangle = \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(B(x, \xi)Q(x, \xi)) dx d\xi$$

holds for each $B \in E$, and not only for $B \in C_b(\mathbf{R}^{2d}; \mathcal{K}(\mathfrak{H}))$.

Step 6: computing $\langle \Lambda, c \rangle$. As explained in Step 2

$$\langle \Lambda, c \rangle = \sup_{\substack{T \in E \\ T(x, \xi) = T(x, \xi)^* \leq c(x, \xi)}} \langle \Lambda, T \rangle.$$

For each $n \geq 1$, set

$$c_n(x, \xi) := (I_{\mathfrak{H}} + \frac{1}{n}c(x, \xi))^{-1}c(x, \xi) \in \mathcal{L}(\mathfrak{H}),$$

so that

$$0 \leq c_1(x, \xi) = c_1(x, \xi)^* \leq \dots \leq c_n(x, \xi) = c_n(x, \xi)^* \leq \dots \leq c(x, \xi) = c(x, \xi)^*.$$

Thus, by definition

$$\langle \Lambda, c_n \rangle = \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(Q(x, \xi) c_n(x, \xi)) dx d\xi \leq \langle \Lambda, c \rangle$$

for each $n \geq 1$, so that, by Corollary 2.4

$$\begin{aligned} & \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(Q(x, \xi)^{1/2} c(x, \xi) Q(x, \xi)^{1/2}) dx d\xi \\ &= \lim_{n \rightarrow \infty} \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(Q(x, \xi) c_n(x, \xi)) dx d\xi \\ &\leq \langle \Lambda, c \rangle. \end{aligned}$$

On the other hand, let $(e_1(x, \xi), e_2(x, \xi), \dots)$ designate a complete orthonormal system in \mathfrak{H} of eigenfunctions of $c(x, \xi)$, with $c(x, \xi) e_j(x, \xi) = \lambda_j e_j(x, \xi)$ for $j \geq 1$. Since $c(x, \xi)$ is a phase space translate of the harmonic oscillator $H := \frac{1}{2}(|x|^2 - \hbar^2 \Delta_x)$, the eigenvalues λ_j are independent of (x, ξ) . Set

$$t_{kl}(x, \xi) := \langle e_k(x, \xi) | Q(x, \xi)^{1/2} | e_l(x, \xi) \rangle, \quad k, l \geq 1.$$

Since $(x, \xi) \mapsto Q(x, \xi)^{1/2} \in L^2(\mathbf{R}^{2d}; \mathcal{L}^2(\mathfrak{H}))$, one has

$$v_k(x, \xi) := \sum_{l \geq 1} t_{kl}(x, \xi) e_l(x, \xi) \in \text{Form-Dom}(c(x, \xi)) \quad \text{for a.e. } (x, \xi) \in \mathbf{R}^{2d}$$

and

$$\begin{aligned} \text{trace}_{\mathfrak{H}}(Q(x, \xi)^{1/2} c(x, \xi) Q(x, \xi)^{1/2}) &= \sum_{k, l \geq 1} \lambda_l |t_{kl}(x, \xi)|^2 \\ &= \sum_{k \geq 1} \langle v_k(x, \xi) | c(x, \xi) | v_k(x, \xi) \rangle < \infty \end{aligned}$$

for a.e. $(x, \xi) \in \mathbf{R}^{2d}$, since

$$\int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(Q(x, \xi)^{1/2} c(x, \xi) Q(x, \xi)^{1/2}) dx d\xi < \infty.$$

Taking this last inequality for granted, we conclude as follows. Let $a \equiv a(x, \xi) \in C_b(\mathbf{R}^{2d})$ and $B = B^* \in \mathcal{L}(\mathfrak{H})$ satisfy the constraint

$$a(x, \xi) I_{\mathfrak{H}} + B \leq c(x, \xi), \quad (x, \xi) \in \mathbf{R}^{2d}$$

in the sense that

$$a(x, \xi) \|\phi\|_{\mathfrak{H}}^2 + \langle \phi | B | \phi \rangle \leq \langle \phi | c(x, \xi) | \phi \rangle \quad \text{for each } \phi \in \text{Form-Dom}(c(x, \xi)).$$

Since $v_k(x, \xi) \in \text{Form-Dom}(c(x, \xi))$ for a.e. $(x, \xi) \in \mathbf{R}^{2d}$ and each $k \geq 1$

$$\begin{aligned} & a(x, \xi) p(x, \xi) + \text{trace}_{\mathfrak{H}}(Q(x, \xi) B) \\ &= a(x, \xi) \text{trace}_{\mathfrak{H}}(Q(x, \xi)) + \text{trace}_{\mathfrak{H}}(Q(x, \xi)^{1/2} B Q(x, \xi)^{1/2}) \\ &= a(x, \xi) \sum_{k \geq 1} \langle v_k(x, \xi) | v_k(x, \xi) \rangle + \sum_{k \geq 1} \langle v_k(x, \xi) | B | v_k(x, \xi) \rangle \\ &\leq \sum_{k \geq 1} \langle v_k(x, \xi) | c(x, \xi) | v_k(x, \xi) \rangle = \text{trace}_{\mathfrak{H}}(Q(x, \xi)^{1/2} c(x, \xi) Q(x, \xi)^{1/2}). \end{aligned}$$

Integrating in (x, ξ) shows that

$$\begin{aligned} & \int_{\mathbf{R}^{2d}} a(x, \xi) p(x, \xi) dx d\xi + \text{trace}_{\mathfrak{H}}(SB) \\ &\leq \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(Q(x, \xi)^{1/2} c(x, \xi) Q(x, \xi)^{1/2}) dx d\xi \end{aligned}$$

since, by construction,

$$\int_{\mathbf{R}^{2d}} Q(x, \xi) dx d\xi = S.$$

Thus

$$\begin{aligned} \langle \Lambda, c \rangle &= \sup_{\substack{a \in C_b(\mathbf{R}^{2d}), B = B^* \in \mathcal{L}(\mathfrak{H}) \\ a(x, \xi) I_{\mathfrak{H}} + B \leq c(x, \xi)}} \left(\int_{\mathbf{R}^{2d}} a(x, \xi) p(x, \xi) dx d\xi + \text{trace}_{\mathfrak{H}}(SB) \right) \\ &\leq \int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(Q(x, \xi)^{1/2} c(x, \xi) Q(x, \xi)^{1/2}) dx d\xi \leq \langle \Lambda, c \rangle, \end{aligned}$$

where the first equality follows from convex duality as explained in Step 2, while the last inequality has been obtained above at the beginning of Step 6. This completes the proof.

It remains to prove that

$$\int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(Q(x, \xi)^{1/2} c(x, \xi) Q(x, \xi)^{1/2}) dx d\xi < \infty.$$

Since

$$c(x, \xi) \leq (|x|^2 + |\xi|^2) I_{\mathfrak{H}} + H$$

one has

$$v_k(x, \xi) \in \text{Form-Dom}(H) \implies v_k(x, \xi) \in \text{Form-Dom}(c(x, \xi))$$

and

$$\langle v_k(x, \xi) | c(x, \xi) | v_k(x, \xi) \rangle \leq (|x|^2 + |\xi|^2) \|v_k(x, \xi)\|_{\mathfrak{H}}^2 + \langle v_k(x, \xi) | H | v_k(x, \xi) \rangle.$$

Let (h_1, h_2, \dots) be a complete orthonormal system of eigenvectors of H in \mathfrak{H} (the Hermite functions), with eigenvalues μ_j . Since

$$\sum_{k \geq 1} \overline{t_{km}(x, \xi)} t_{kn}(x, \xi) = \langle e_m(x, \xi) | Q(x, \xi) | e_n(x, \xi) \rangle$$

by definition of $t_{kl}(x, \xi)$, one has

$$\begin{aligned} \sum_{k \geq 1} |\langle v_k(x, \xi) | h_j \rangle|^2 &= \sum_{k \geq 1} \sum_{m, n \geq 1} \overline{t_{km}(x, \xi)} t_{kn}(x, \xi) \langle e_m(x, \xi) | h_j \rangle \langle h_j | e_n(x, \xi) \rangle \\ &= \sum_{m, n \geq 1} \langle e_m(x, \xi) | Q(x, \xi) | e_n(x, \xi) \rangle \langle e_m(x, \xi) | h_j \rangle \langle h_j | e_n(x, \xi) \rangle = \langle h_j | Q(x, \xi) | h_j \rangle. \end{aligned}$$

Hence

$$\begin{aligned} \int_{\mathbf{R}^{2d}} \sum_{k \geq 1} \langle v_k(x, \xi) | H | v_k(x, \xi) \rangle dx d\xi &= \sum_{j \geq 1} \mu_j \int_{\mathbf{R}^{2d}} \langle h_j | Q(x, \xi) | h_j \rangle \\ &= \sum_{j \geq 1} \mu_j \langle h_j | S | h_j \rangle = \text{trace}_{\mathfrak{H}}(S^{1/2} |H| S^{1/2}) < \infty, \end{aligned}$$

and since

$$\sum_{k \geq 1} \|v_k(x, \xi)\|_{\mathfrak{H}}^2 = \text{trace}_{\mathfrak{H}}(Q(x, \xi)) = p(x, \xi),$$

one concludes that

$$\begin{aligned} &\int_{\mathbf{R}^{2d}} \text{trace}_{\mathfrak{H}}(Q(x, \xi)^{1/2} c(x, \xi) Q(x, \xi)^{1/2}) dx d\xi \\ &\leq \int_{\mathbf{R}^{2d}} (|x|^2 + |\xi|^2) p(x, \xi) dx d\xi + \text{trace}_{\mathfrak{H}}(S^{1/2} H S^{1/2}) < \infty. \end{aligned}$$

□

7. APPLICATIONS OF DUALITY FOR \mathcal{E}_h I: INEQUALITIES BETWEEN MK_h , \mathcal{E}_h AND $\text{dist}_{\text{MK},2}$.

Theorem 7.1. *Let $R, S \in \mathcal{D}_2(\mathfrak{H})$ and p be a probability density on \mathbf{R}^{2d} . Then*

$$\begin{aligned}\mathcal{E}_h(\widetilde{W}_h(R), S)^2 &\geq \text{dist}_{\text{MK},2}(\widetilde{W}_h[R], \widetilde{W}_h[S]) - d\hbar, \\ MK_h(R, S)^2 &\geq \mathcal{E}_h(\widetilde{W}_h(R), S)^2 - d\hbar, \\ MK_h(R, S)^2 &\geq \text{dist}_{\text{MK},2}(\widetilde{W}_h[R], \widetilde{W}_h[S]) - 2d\hbar.\end{aligned}$$

Proof. The first inequality and the third inequality (also a consequence of the two others) were proved in Theorem 2.4 (2) of [7] and Theorem 2.3 (2) of [6] respectively.

The second inequality is proved along the same lines as Theorem 2.3 (2) of [6]. Let $a \equiv a(x, \xi)$ in $C_b(\mathbf{R}^{2d}, \mathbf{R})$ and $B = B^* \in \mathcal{L}(\mathfrak{H})$ satisfy

$$a(x, \xi)I_{\mathfrak{H}} + B \leq c(x, \xi) \quad \text{for a.e. } (x, \xi) \in \mathbf{R}^{2d}.$$

Then

$$a(x, \xi)|x, \xi\rangle\langle x, \xi| \otimes I_{\mathfrak{H}} + |x, \xi\rangle\langle x, \xi| \otimes B \leq |x, \xi\rangle\langle x, \xi| \otimes c(x, \xi)$$

for a.e. $(x, \xi) \in \mathbf{R}^{2d}$, so that

$$\begin{aligned}\text{OP}_h^T((2\pi\hbar)^d a) \otimes I_{\mathfrak{H}} + I_{\mathfrak{H}} \otimes B &\leq \frac{1}{(2\pi\hbar)^d} \int_{\mathbf{R}^{2d}} |x, \xi\rangle\langle x, \xi| \otimes c(x, \xi) dx d\xi \\ &= C + d\hbar I_{\mathfrak{H} \otimes \mathfrak{H}}.\end{aligned}$$

Thus, for each $\mathcal{Q} \in \mathcal{C}(R, S)$, one has

$$\begin{aligned}\text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}(\mathcal{Q}^{1/2} C \mathcal{Q}^{1/2}) + d\hbar &\geq \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}(\mathcal{Q}^{1/2} (\text{OP}_h^T((2\pi\hbar)^d a) \otimes I_{\mathfrak{H}} + I_{\mathfrak{H}} \otimes B) \mathcal{Q}^{1/2}) \\ &= \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}(\mathcal{Q} (\text{OP}_h^T((2\pi\hbar)^d a) \otimes I_{\mathfrak{H}} + I_{\mathfrak{H}} \otimes B)) \\ &= \text{trace}_{\mathfrak{H}}(R \text{OP}_h^T((2\pi\hbar)^d a) + SB) \\ &= \int_{\mathbf{R}^{2d}} a(x, \xi) \widetilde{W}_h(R)(x, \xi) dx d\xi + \text{trace}_{\mathfrak{H}}(SB).\end{aligned}$$

In particular

$$\begin{aligned}MK_h(R, S)^2 + d\hbar &= \inf_{\mathcal{Q} \in \mathcal{C}(R, S)} \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}(\mathcal{Q}^{1/2} C \mathcal{Q}^{1/2}) + d\hbar \\ &\geq \sup_{\substack{a \in C_b(\mathbf{R}^{2d}, \mathbf{R}), B = B^* \in \mathcal{L}(\mathfrak{H}) \\ a(x, \xi)I_{\mathfrak{H}} + B \leq c(x, \xi)}} \left(\int_{\mathbf{R}^{2d}} a(x, \xi) \widetilde{W}_h(R)(x, \xi) dx d\xi + \text{trace}_{\mathfrak{H}}(SB) \right) \\ &= \mathcal{E}_h(\widetilde{W}_h(R), S)^2.\end{aligned}$$

□

8. APPLICATIONS OF DUALITY FOR \mathcal{E}_h II: “TRIANGLE” INEQUALITIES

Theorem 8.1. *Let $R, S, T \in \mathcal{D}_2(\mathfrak{H})$ and let $f, g \in \mathcal{P}_2(\mathbf{R}^{2d})$. Then*

(i) *one has*

$$\begin{aligned}\text{dist}_{\text{MK},2}(f, g) &\leq \sqrt{\mathcal{E}_h(f, S)^2 + d\hbar} + \sqrt{\mathcal{E}_h(g, S)^2 + d\hbar} \\ &< \mathcal{E}_h(f, S) + \mathcal{E}_h(g, S) + d\hbar;\end{aligned}$$

(ii) *one has*

$$\begin{aligned}\mathcal{E}_h(f, T) &\leq \text{dist}_{\text{MK},2}(f, \widetilde{W}_h(S)) + \mathcal{E}_h(\widetilde{W}_h(S), T) \\ &\leq \sqrt{\mathcal{E}_h(f, S)^2 + d\hbar} + \sqrt{MK_h(S, T)^2 + d\hbar} \\ &< \mathcal{E}_h(f, S) + MK_h(S, T) + d\hbar;\end{aligned}$$

(iii) one has

$$\begin{aligned} MK_{\hbar}(R, T) &\leq \mathcal{E}_{\hbar}(\widetilde{W}_{\hbar}(S), R) + \mathcal{E}_{\hbar}(\widetilde{W}_{\hbar}(S), T) \\ &\leq \sqrt{MK_{\hbar}(R, S)^2 + d\hbar} + \sqrt{MK_{\hbar}(S, T)^2 + d\hbar} \\ &< MK_{\hbar}(R, S) + MK_{\hbar}(S, T) + d\hbar. \end{aligned}$$

Proof. The triangle inequality for $\text{dist}_{\text{MK},2}$ implies that

$$\text{dist}_{\text{MK},2}(f, g) \leq \text{dist}_{\text{MK},2}(f, \widetilde{W}_{\hbar}(S)) + \text{dist}_{\text{MK},2}(\widetilde{W}_{\hbar}(S), g).$$

Then, Theorem 2.4 (2) of [7] implies that

$$\begin{aligned} \text{dist}_{\text{MK},2}(f, \widetilde{W}_{\hbar}(S)) &\leq \sqrt{\mathcal{E}_{\hbar}(f, S) + d\hbar}, \\ \text{dist}_{\text{MK},2}(\widetilde{W}_{\hbar}(S), g) &\leq \sqrt{\mathcal{E}_{\hbar}(g, S) + d\hbar}. \end{aligned}$$

This implies the first inequality in (i). As for the second inequality, for each $X, Y > 0$, one has the obvious elementary inequality

$$\sqrt{X + Y} < X + \frac{1}{2}Y.$$

This inequality obviously applies to the present case since $\mathcal{E}_{\hbar}(f, S) \geq d\hbar$ and $\mathcal{E}_{\hbar}(g, S) \geq d\hbar$ by Theorem 2.4 (2) of [7]. This proves (i).

Observe that the first inequality in (ii) is inequality (a) in Theorem 4.2 with $g = \widetilde{W}_{\hbar}(S)$ and $R_1 = T$. Then Theorem 2.4 (2) of [7] implies that

$$\text{dist}_{\text{MK},2}(f, \widetilde{W}_{\hbar}(S)) \leq \sqrt{\mathcal{E}_{\hbar}(f, S)^2 + d\hbar},$$

while Theorem 7.1 implies that

$$\mathcal{E}_{\hbar}(\widetilde{W}_{\hbar}(S), T) \leq \sqrt{MK_{\hbar}(S, T)^2 + d\hbar},$$

and this implies the second inequality in (ii). The third inequality is obtained as in (i).

Finally, the first inequality in (iii) is inequality (b) in Theorem 4.2 with $R_1 = R$, while $R_3 = T$ and $f = \widetilde{W}_{\hbar}(S)$. Then, Theorem 7.1 implies that

$$\begin{aligned} \mathcal{E}_{\hbar}(\widetilde{W}_{\hbar}(S), R) &\leq \sqrt{MK_{\hbar}(R, ST)^2 + d\hbar}, \\ \mathcal{E}_{\hbar}(\widetilde{W}_{\hbar}(S), T) &\leq \sqrt{MK_{\hbar}(S, T)^2 + d\hbar}, \end{aligned}$$

which gives the second inequality in (iii). Finally, the third inequality is obtained as in (i). \square

Remark. It is interesting to compare the inequality (iii) above with the “generalized triangle inequality” in [4]. Let us recall that DePalma and Trevisan have constructed a pseudo-distance on density operators on \mathfrak{H} which is similar to ours to some extent. The DePalma-Trevisan distance D is defined through a different notion of coupling than in [6]; specifically, their notion of couplings is based on “quantum channels” (completely positive linear maps on the set of density operators): see Definition 1 in [4]. While the transport cost in formula (19) of [4] is in some sense reminiscent of the transport cost used in [6], these two costs are in fact significantly different. For instance, the transport cost used in the definition of MK_{\hbar} in [6], and in the present paper, has compact resolvent, and therefore its spectrum consists of eigenvalues only. On the contrary, the cost operator in [4] in the case of Gaussian quantum systems has continuous spectrum on $[0, +\infty)$.

In Theorem 2 of [4], DePalma and Trevisan prove what they call a “triangle inequality” for their distance D , of the form

$$D(R, T) \leq D(R, S) + D(S, S) + D(S, T)$$

(inequality (35) in [4]). Of course, if D was a real distance, $D(S, S) = 0$, and the inequality above coincides with the usual triangle inequality. In [4], there is an explicit formula for $D(S, S)$ in terms of the canonical purification of S (Corollary 1, formula (34) in [4]).

With the distance MK_h defined in [6], one has

$$MK_h(R, S) \geq 2d\hbar, \quad \text{for all } R, S \in \mathcal{D}_2(\mathfrak{H}),$$

so that Theorem 8.1 (iii) implies that

$$MK_h(R, T) < MK_h(R, S) + MK_h(S, S) + MK_h(S, T).$$

In other words, MK_h satisfies the same “generalized triangle inequality” as the DePalma-Trevisan distance D , with a strict inequality.

9. APPLICATIONS OF DUALITY FOR \mathcal{E}_h III: CLASSICAL/QUANTUM OPTIMAL TRANSPORT AND SEMIQUANTUM LEGENDRE TRANSFORM

9.1. A classical/quantum optimal transport. Let r be a probability density on \mathbf{R}^{2d} and S a density operator on $L^2(\mathbf{R}^d)$.

We suppose that an optimal operator \tilde{B} and an optimal function \tilde{a} exists for the Kantorovich duality formulation of $\mathcal{E}_h(r, S)$, as in Theorem 6.1, and that $\tilde{a} \in C_b(\mathbf{R}^{2d})$ and $\tilde{B} \in \mathcal{L}(\mathfrak{H})$. That is to say that

$$\tilde{a}(q, p) + \tilde{B} \leq (Z - z)^2 \text{ and } \mathcal{E}_h(r, S)^2 = \int_{\mathbf{R}^{2d}} \tilde{a}(z) r(z) dz + \text{trace}_{L^2(\mathbf{R}^d)}(\tilde{B}S).$$

Here we have used the notation $z = (q, p)$, $dz = dqdp$, and $Z = (Q, P)$.

Let us denote by $\Pi(z)$ an optimal coupling of r, S and let us define

$$\begin{aligned} a(z) &:= \frac{1}{2}(|z|^2 - \tilde{a}(z)) \\ B &:= \frac{1}{2}(|Z|^2 - \tilde{B}). \end{aligned}$$

One has

$$(a(z) + B - z \cdot Z) \geq 0 \text{ and } \text{trace}_{L^2(\mathbf{R}^d)} \int_{\mathbf{R}^{2d}} \Pi(z)^{\frac{1}{2}} (a(z) + B - z \cdot Z) \Pi(z)^{\frac{1}{2}} dz = 0,$$

Therefore, since $\Pi(z)^{\frac{1}{2}} (a(z) + B - z \cdot Z) \Pi(z)^{\frac{1}{2}} \geq 0$

$$\Pi(z)^{\frac{1}{2}} (a(z) + B - z \cdot Z) \Pi(z)^{\frac{1}{2}} = 0 \text{ a.e.,}$$

In other words,

$$\Pi(z)^{\frac{1}{2}} (a(z) + B - z \cdot Z)^{\frac{1}{2}} \left(\Pi(z)^{\frac{1}{2}} (a(z) + B - z \cdot Z)^{\frac{1}{2}} \right)^* = 0$$

which implies that

$$(a(z) + B - z \cdot Z)^{\frac{1}{2}} \Pi(z)^{\frac{1}{2}} = 0 \text{ a.e.}$$

and (forgetting the “a.e.” in the sequel)

$$(13) \quad (a(z) + B - z \cdot Z) \Pi(z) = 0.$$

Hence, the range of Π consists in functions $\mathbb{R}^{2d} \ni z \mapsto \psi_z \in L^2(\mathbb{R}^d)$ such that

$$(14) \quad (a(z) + B - z \cdot Z) \psi_z = 0 \iff (B - z \cdot Z) \psi_z = -a(z) \psi_z :$$

the vectors ψ_z are the eigenvectors of $B - z \cdot Z$ with eigenvalue $-a(z)$.

But $B + a(z) - z \cdot Z \geq 0$. Therefore

$$-a(z) \text{ is the lowest eigenvalue of } B - 2z \cdot Z.$$

From now on, we will suppose that the fundamental of $B - z \cdot Z$ is non degenerate. This means that $\Pi(z)$ is proportional to $|\psi_z\rangle\langle\psi_z|$ and therefore, since $\Pi(z)$ is a coupling between r and S ,

$$\Pi(z) = r(z)|\psi_z\rangle\langle\psi_z|$$

and

$$S = \int_{\mathbf{R}^{2d}} r(z)|\psi_z\rangle\langle\psi_z|dz.$$

We just prove the following result.

Theorem 9.1. *Let B be a bounded optimal Kantorovich operator of $\mathcal{E}_h(r, S)$. Let moreover, for each $z \in \mathbf{R}^{2d}$, ψ_z be the ground state of $B - z \cdot Z$.*

Then S admits the following representation

$$S = \int_{\mathbf{R}^{2d}} r(z)|\psi_z\rangle\langle\psi_z|dz.$$

Theorem 9.1 suggests to associate to any probability density μ the following operator

$$(15) \quad \mu \longrightarrow \text{OP}_h^{r, S}[\mu] := \int_{\mathbf{R}^{2d}} |\psi_z\rangle\langle\psi_z| \mu(dz).$$

The arrow in (15) can be seen as the ‘‘optimal transport’’, from classical probability densities to quantum density matrices, transporting r to S .

Note that, for any density μ ,

$$\text{trace OP}_h^{r, S}[\mu] = \int_{\mathbf{R}^{2d}} \mu(dz).$$

Finally, using (13), we easily show, by analogy with the proof of Theorem 2.6 (b) in [2], that, when $a \in C^1(\mathbf{R}^{2d})$, $(\nabla a)r \in C_b(\mathbf{R}^{2d})$ and, e.g., $\psi_z \in \text{Dom}(\frac{1}{i\hbar}[Z, B])$ for all $z \in \text{supp}(r)$,

$$\begin{aligned} 0 &= \Pi(z) \frac{1}{i\hbar} [Z, (a(z) + B - z \cdot Z) \Pi(z)] = \Pi_z \frac{1}{i\hbar} [Z, a(z) + B - z \cdot Z] \Pi(z) \\ &= \Pi(z) ([Z, B] - z) \Pi(z) \end{aligned}$$

and

$$\begin{aligned} 0 &= \Pi(z) |Z, (a(z) + B - z \cdot Z) \Pi(z)\rangle = \Pi(z) \{Z, a(z) + B - z \cdot Z\} \Pi(z) \\ &= \Pi(z) (\nabla a(z) - Z) \Pi(z). \end{aligned}$$

Therefore the (classical and quantum) ‘‘gradient’’ aspect appears in the following expressions

$$\begin{aligned} \langle\psi_z|Z|\psi_z\rangle &= \nabla a(z) \\ z &= \langle\psi_z|Z|\nabla^Q B\psi_z\rangle \end{aligned}$$

where $\nabla^Q := \frac{1}{i\hbar}[JZ, \cdot]$ with J the symplectic matrix defined by $\{f, g\} = \nabla f \cdot J \nabla g$, as introduced and motivated in [2, Section 1].

Let us finish this section by an example. Suppose that

$$S = \text{OP}_\hbar^T((2\pi\hbar)^d r).$$

In this case, one knows, [7, Theorem 2.4 (1)] (note a difference of normalization: in [7], $E_\hbar^2 = \frac{1}{2}\mathcal{E}_\hbar^2$),

$$\mathcal{E}_\hbar(r, S)^2 = d\hbar = \int_{\mathbf{R}^{2d}} \tilde{a}(z)r(z)dz + \text{trace}(\tilde{B}S) \text{ with } \tilde{a} = 0, \tilde{B} = d\hbar I_{\mathfrak{S}}.$$

Since $(q-x)^2 + (p+i\hbar\nabla_x)^2 \geq d\hbar I_{\mathfrak{S}} = \tilde{a}(z)I_{\mathfrak{S}} + \tilde{B}$, \tilde{a} and \tilde{B} are optimal and

$$a(q, p) = \frac{1}{2}|z|^2 \text{ and } B = \frac{1}{2}(|Z|^2 - d\hbar).$$

Hence

$$a(z) + B - z \cdot Z = \frac{1}{2}(-\nabla_x + x - (q + ip))(\nabla_x + x - (q - ip)),$$

the solution of (14) is

$$\psi_z = (\pi\hbar)^{-d/4} e^{-\frac{(x-q)^2}{2\hbar}} e^{i\frac{p \cdot x}{\hbar}}$$

and Theorem 9.1 expresses back that $S = \text{OP}_\hbar^T((2\pi\hbar)^d r)$ and

$$\text{OP}_\hbar^{\mu, \text{OP}_\hbar^T(\mu)} = \text{OP}_\hbar^T$$

for any probability density μ .

9.2. A semiquantum Legendre transform. As we have seen, $-a(z)$ is the fundamental of the operator $B - z \cdot Z$. Therefore, by the variational characterization of the lowest eigenvalue,

$$-a(z) = \inf_{\substack{\phi \in \text{Dom}(B) \\ \|\phi\|_{\mathfrak{S}}=1}} (\langle \phi | B | \phi \rangle - z \cdot \langle \phi | Z | \phi \rangle),$$

to be faced to the classical definition of the Legendre transform

$$a(z) = \sup_{z'} (z \cdot z' - b(z')).$$

Let us define the semiquantum Legendre transform by

$$B^{sq*} := \sup_{\substack{\phi \in \text{Dom}(B) \\ \|\phi\|_{\mathfrak{S}}=1}} (z \cdot \langle \phi | Z | \phi \rangle - \langle \phi | B | \phi \rangle).$$

Theorem 9.2. *Let $a(z) = \frac{1}{2}(|z|^2 - \tilde{a}(z))$, $B = \frac{1}{2}(|Z|^2 - \tilde{B})$ where $\tilde{a}(z)$ and \tilde{B} are bounded optimal Kantorovich potentials for $\mathcal{E}_\hbar(r, S)$. Then*

$$a = B^{sq*}.$$

Proof. We just recall the variational argument.

Let $A \geq 0$ and $A|\phi_0\rangle = 0$. Then,

$$\langle \phi_0 | A | \phi_0 \rangle \leq \inf_{\substack{\phi \in \text{Dom}(B) \\ \|\phi\|_{\mathfrak{S}}=1}} \langle \phi | A | \phi \rangle$$

and

$$\langle \phi_0 + \delta\phi_0 | A | \phi_0 + \delta\phi_0 \rangle = \langle \delta\phi | A | \delta\phi \rangle.$$

□

REFERENCES

- [1] Brezis, H.: “Functional Analysis, Sobolev Spaces and Partial Differential Equations”, Springer Science + Business Media 2011.
- [2] Caglioti, E., Golse, F., Paul, T.: *Toward Optimal Transport for Quantum Densities*, preprint [arXiv:2101.03256](https://arxiv.org/abs/2101.03256) [math-ph].
- [3] Caglioti, E., Golse, F., Paul, T.: *Quantum Optimal Transport is Cheaper*, J. Statist. Phys. **181** (2020), 149–162.
- [4] DePalma, G., Trevisan, D.: *Quantum optimal transport with quantum channels*, preprint [arXiv:1911.00803](https://arxiv.org/abs/1911.00803) [math-ph].
- [5] Diestel, J., Uhl Jr, J.J.: “Vector Measures”, Amer. Math. Soc., Providence, RI, 1977.
- [6] Golse, F., Mouhot, C., Paul, T.: *On the mean-field and classical limits of quantum mechanics*, Commun. Math. Phys. **343** (2016), 165–205.
- [7] Golse, F., Paul, T.: *The Schrödinger Equation in the Mean-Field and Semiclassical Regime*, Arch. Rational Mech. Anal. **223** (2017), 57–94.
- [8] Golse, F., Paul, T.: *Semiclassical evolution with low regularity*, preprint [arXiv:2011.14884](https://arxiv.org/abs/2011.14884) [math.AP], to appear in J. Math. Pures et Appl..
- [9] Reed, M., Simon, B.: “Methods of Modern Mathematical Physics I. Functional Analysis” Acad. Press., Inc., 1980.
- [10] Reed, M., Simon, B.: “Methods of Modern Mathematical Physics IV. Analysis of Operators” Acad. Press., Inc., 1978.
- [11] Rockafellar, R.T. : “Convex Analysis”, 2nd printing, Princeton University Press, 1972.
- [12] Rudin, W.: “Functional Analysis”, 2nd ed., McGraw Hill, Inc., 1991.
- [13] Simon, B.: “Trace Ideals and their Applications”, 2nd ed., Amer. Math. Soc. Providence, 2005.
- [14] Villani, C.: “Topics in Optimal Transportation”, Amer. Math. Soc., Providence, 2003.

(F.G.) CMLS, ÉCOLE POLYTECHNIQUE, CNRS, UNIVERSITÉ PARIS-SACLAY , 91128 PALAISEAU CEDEX, FRANCE

E-mail address: francois.golse@polytechnique.edu

(T.P.) CNRS & LJLL SORBONNE UNIVERSITÉ 4 PLACE JUSSIEU 75005 PARIS, FRANCE

E-mail address: thierry.paul@upmc.fr