



Towards an automatic detection of charcoal production platforms in airborne LiDAR images

Cláudia Oliveira, Stéphanie Aravecchia, L. May, Cedric Pradalier, Vincent Robin, Simon Devin

► To cite this version:

Cláudia Oliveira, Stéphanie Aravecchia, L. May, Cedric Pradalier, Vincent Robin, et al.. Towards an automatic detection of charcoal production platforms in airborne LiDAR images. 5ème colloque des Zones Ateliers - CNRS, Nov 2020, Blois, France. hal-03137113

HAL Id: hal-03137113

<https://hal.science/hal-03137113>

Submitted on 10 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

TOWARDS AN AUTOMATIC DETECTION OF CHARCOAL PRODUCTION PLATFORMS IN AIRBORNE LIDAR IMAGES

Oliveira C^{1*}, Aravecchia S², May L³, Pradalier C², Robin V¹, Devin S¹

¹LIEC, Université de Lorraine, Metz, France; *claudia.oliveira@univ-lorraine.fr

² Georgia Tech Lorraine, Metz, France

³ Georgia Institute of Technology, Atlanta, Georgia, USA

INTRODUCTION

Charcoal produced from woodland exploitation was a key source of energy for the (proto-) industrial development from the Middle-Ages to the emergence of fossil energy sources during the 19th century. Charcoal production platforms (or kilns) are some evidences of these activities and they are spread and preserved, at some extent, in forest areas. These structures with a characteristic circular to oval shape (Fig. 1) are recognizable in airborne images such as LiDAR (Light Detection and Ranging). Frequently, the density of these kilns in woodlands is so high that the visual inspection is insufficient. Methods of automatic detection based on machine learning for image analysis are being implemented to detect and quantify these structures, remaining traces of ancient forest exploitation.




Fig. 1 | Examples of charcoal kilns in southern Meuse with the aspect on LiDAR-derived slope images.

OBJECTIVES


Development of an **expedite** and **reliable** methodology to routinely perform analysis in airborne images to increment the **detection** and quantification of kilns in order to better understand the history and evolution of forest resources' exploitation.

METHODS

- charcoal kilns' detection by visual inspection of images (1) and fieldwork (2)



- charcoal kilns' detection using deep learning techniques



STUDY AREA




Fig. 2 | Study area in southern Meuse (NE France). The forest patches (black contour, 57.3km²) and ancient industrial activities (smithies) displayed in the LiDAR-derived Digital Elevation Model.

RESULTS





Fig. 4 | Results of the visual and automatically-detected kilns in a selected area.

PERSPECTIVES

Future work will focus on (1) **assessing** of detection in the field (planned for mid-November 2020), (2) **improving** the charcoal kiln detector and, with the feedback of the fieldwork, fine-tune our model and (3) **using** our charcoal kiln detector in a new unstudied area.