



HAL
open science

An uncommon geomorphological crisis - Dominica Coastal systems response and adjustments in the Aftermath of Hurricane Maria (September 2017 episode)

Samuel Battut, Tony Rey, Stéphanie Defossez, Frédéric Leone

► **To cite this version:**

Samuel Battut, Tony Rey, Stéphanie Defossez, Frédéric Leone. An uncommon geomorphological crisis - Dominica Coastal systems response and adjustments in the Aftermath of Hurricane Maria (September 2017 episode). Oceanext international interdisciplinary conference second edition, Jul 2019, Nantes, France. hal-03137389

HAL Id: hal-03137389

<https://hal.science/hal-03137389>

Submitted on 12 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An uncommon geomorphological crisis — Dominica Coastal system response and adjustments in the aftermath of Hurricane Maria (September 2017 episode)

S. Battut*, T. Rey, S. Defossez & F. Leone - Université Paul Valéry - Montpellier 3 - UMR GRED

1. Objectives

Analysing the physical impact of Hurricane Maria on the Dominican coastline is relevant to the TIREX 2017 research project funded by the ANR OURAGANS - a national grant devoted to the study of hurricanes. TIREX (*Transfert des apprentissages de Retours d'Expériences scientifiques pour le renforcement des capacités de réponse et d'adaptation individuelles et collectives dans un contexte de changement climatique*) refers to the use of scientific feedback to encourage better response and adjustment from the civilian population as well as the local and national authorities in a context of climate change ; it focuses on the north area of the West Indies during 2017 hurricane season.




Figure 1 - Maria Hurricane track with winds speed and watch or warning levels for Dominica. Sources : ibtracs /NHC, NOAA, 2018 - GADM, 2019 (BATTUT, 2019)

2. The event

The 2017 Atlantic hurricane season is undoubtedly one of the most active and devastating ever recorded. Hurricane Maria ranks as the deadliest of the six strongest hurricanes of the season causing about 3,000 fatalities as recently reported. It started as a tropical wave off the West African coast before turning into a category 5 hurricane in less than 18 hours just before making landfall on the island of Dominica in the evening of the 18th of September 2017 (fig. 1). It stroked the island going from the south-east to the north-west and wrecked considerably this small island developing state (1.3 billion American Dollars loss or 224% of the 2016 gross domestic product). It affected seriously a population already made vulnerable by the passing of Hurricane Erika in 2015.

3. A diachronic and comparative approach for sedimentary responses analysis

In Dominica, hurricane Maria impacts have been studied through a comparative and diachronic approach on several sediment cells and watersheds across the island (fig. 5) displaying different signs of anthropisation (fig. 4). Their resilience was assessed a year and a half after the event. The use of aerial footage (fig. 3) and topomorphological data analysis made on field (fig. 2) have been put forward thanks to a cartographic translation in order to emphasize morphosedimentary response at different moments in time.




Figure 2 - Geomorphological cartography of Whiter river mouth in Pointe Mulâtre made with field observations and high resolution aerial views (BATTUT, 2019).



Figure 3 - Coastal evolutions in Colihaut between 2005 and 2018 based on coastline monitoring. (BATTUT, 2019)

Sedimentary cell	Morphological nature	Average sediment size	Human occupation level of seafloor
Grand Bay	Bay	Pebble	Few houses
Colihaut	Small bay	Pebble	Coastal village
Mahaut	Small bay	Pebble	Coastal village
Pointe Mulâtre	Small bay	Pebble	Very few houses
Petite Savanne Bay	Creek	Block	No occupation
Fort Saint Jean	Small bay	Block	Small coastal village
Number One beach	Beach at the upper end of a bay	Sand	No occupation
Portsmouth Bay	Large bay	Sand	Second island city
Scott's Head	Small bay with tombolo	Pebble	Coastal village

Sites show a large range of shapes, materials and occupation levels so as to be well representative of Dominican coasts.

Figure 4 - Sites under study and their general features.




Figure 5 - Geomorphological map of Dominica and sites under study location (BATTUT, 2019). 1 - Portsmouth bay 2 - Colihaut 3 - Mahaut 4 - Scott's Head 5 - Grand bay 6 - Fond Saint Jean 7 - Petite Savanne bay 8 - Pointe Mulâtre 9 - Number one beach.

4. Binary sedimentary responses to Maria hurricane: from accretion to erosion




Figure 6 - Sediment balance in 8 different sites between 2005 and 2018 based on evolution surface (m2) (BATTUT, 2019).

The results show that coast evolutions are mainly due to fluvial processes rather than marine dynamics. Heavy rains caused major landslides (fig. 10) and flooding while enabling organic materials and debris to flow downstream towards the mouths. This phenomenon of extreme sedimentation accounts for major morphological alterations such as land elevation, sedimentary progradation, and river avulsion. Contrary to the idea according to which hurricanes generally make coastlines recede, in the case of Dominica island, large amounts of input materials of heterogeneous granulometry have generally contributed to a considerable accretion in sediment cells (fig. 6). Besides, large amounts of wooden debris washed away into the sea and reshaped by the cyclonic swell intensified the impact of the waves on the coastline and accumulated on supratidal parts of beaches.

Only two studied sites showed erosive processes signs (fig. 6). These locations are low lying sandy beaches with estuaries. Contrary to most of the sites, they include low gradient rivers. Erosion was mainly due to cyclonic waves that transported sediments towards the subtidal zone, causing the beach profile to retreat (fig. 9).

6. Coastal sedimentary accretion as a result of upstream extreme sedimentation processes




Figure 10 - Sedimentation processes in Fond Saint Jean (west) and Petite Savanne (east) watersheds. (BATTUT, 2019).



Figure 11 - Morphological evolutions in Fond Saint Jean bay between 2005 and 2017. (BATTUT, 2019).



Figure 12 - Coarse deposits in Petite Savanne. (BATTUT, 2019).




Figure 13 - Large block on Pointe Mulâtre river mouth.

5. Erosive dynamics appearing on low lying sites with low gradient rivers and fine materials




Figure 7 - Erosion induced by Hurricane Maria in Lagoon, Portsmouth (BATTUT, 2019)




Figure 8 - Erosive processes and damages on seafront urbanization in Scott's Head and Picard (BATTUT, 2019).



Figure 9 - Washover deposits on Number one beach (BATTUT, 2019).

Erosive signs were observed in Portsmouth bay and Number one beach (fig. 7 & 8). Their orographic settings are rare on the island of Dominica where slopes are particularly steep, even in coastal areas. Large low plains at the back of the seafloor determine rivers declivity and their sediment load. On these sites, hurricane impacts were mainly induced by marine processes like waves impact and sediment erosion. Urbanized seafronts have suffered very high destructions as reflective erosion damaged sea walls and buildings (fig. 8).

7. Coastal resilience



From left to right : Figure 14 - Maria and Post Maria sediment balances in the studied sites (L is the length of the sediment cell, S/L is the evolution surface related to the length of the cell (expressed in m² per linear meter) - Figure 15 - Post Maria coastline adjustments in Fond Saint Jean - Figure 16 - Post Maria adjustments on Lagoon beach, Portsmouth (BATTUT, 2019)

8. Conclusion

Paroxysmic events like hurricane Maria are able to disrupt coastal morphological equilibria and set a new one that is different from the previous one. Impacts are strongly related to the coast morphological settings and to human occupation. In dominica, extreme sedimentation processes mainly implied significant accretion on coastal systems. Longer observations (several years after the event) would allow to determine if these tendencies will be sustained. Some indicators show that coasts tend to restore previous morphological settings one year and a half after the event. This evolution depends on the next hurricane seasons and future events intensity.