

A chemomechanobiological model of the long-term healing response of arterial tissue to a clamping injury

Lauranne Maes, Julie Vastmans, Stéphane Avril, Nele Famaey

▶ To cite this version:

Lauranne Maes, Julie Vastmans, Stéphane Avril, Nele Famaey. A chemomechanobiological model of the long-term healing response of arterial tissue to a clamping injury. Frontiers in Bioengineering and Biotechnology, 2021, 8, pp.589889. 10.3389/fbioe.2020.589889. hal-03139778

HAL Id: hal-03139778 https://hal.science/hal-03139778

Submitted on 12 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1

A chemomechanobiological model of the long-term healing response of arterial tissue to a clamping injury

Lauranne Maes¹, Julie Vastmans¹, Stéphane Avril² and Nele Famaey^{1,*}

¹Biomechanics Section, Department of Mechanical Engineering, KU Leuven, Leuven, Belaium ² Mines Saint-Etienne, Université de Lyon, Université Jean Monnet, INSERM, U 1059 SAINBIOSE, F-42023 Saint-Etienne, France

Correspondence*: Nele Famaey nele.famaey@kuleuven.be

ABSTRACT 2

7

8

Vascular clamping often causes injury to arterial tissue, leading to a cascade of cellular and 3 extracellular events. A reliable *in silico* prediction of these processes following vascular injury 4 5 could help us to increase our understanding thereof, and eventually optimize surgical techniques or drug delivery to minimize the amount of long-term damage. However, the complexity and 6 interdependency of these events make translation into constitutive laws and their numerical implementation particularly challenging.

9 We introduce a finite element simulation of arterial clamping taking into account acute endothelial denudation, damage to extracellular matrix and smooth muscle cell loss. The model captures 10 how this causes tissue inflammation and deviation from mechanical homeostasis, both triggering 11 12 vascular remodeling. A number of cellular processes are modelled, aiming at restoring this homeostasis, *i.e.* smooth muscle cell phenotype switching, proliferation, migration and the 13 production of extracellular matrix. We calibrated these damage and remodeling laws by comparing 14 our numerical results to in vivo experimental data of clamping and healing experiments. In these 15 same experiments, the functional integrity of the tissue was assessed through myograph tests, 16 which were also reproduced in the present study through a novel model for vasodilator and 17 -constrictor dependent smooth muscle contraction. 18

The simulation results show a good agreement with the *in vivo* experiments. The computational 19 model was then also used to simulate healing beyond the duration of the experiments in order 20 to exploit the benefits of computational model predictions. These results showed a significant 21 sensitivity to model parameters related to smooth muscle cell phenotypes, highlighting the 22 pressing need to further elucidate the biological processes of smooth muscle cell phenotypic 23 switching in the future. 24

25 Keywords: Smooth muscle cells (SMC), Phenotype switch, Vascular remodeling, Vascular clamping, Myograph, Finite elements

1 INTRODUCTION

Multiple studies indicate that arterial occlusion by almost any type of clamp systematically leads to intimal 26 injury at the site of application. For example, endothelial denudation is a widely known effect of clamping 27 (Geenens et al., 2016a; Slayback et al., 1976; Barone et al., 1989; Margovsky et al., 1997, 1999; Hangler 28 et al., 2008; Vural et al., 2008; Famaey et al., 2010; Geenens et al., 2016b). Several studies also report 29 damage to the extracellular matrix (ECM) in the media with flattened elastic lamellae (Geenens et al., 30 2016a; Barone et al., 1989; Margovsky et al., 1999; Famaey et al., 2010). Moreover, healing after arterial 31 clamping usually implies some degree of inflammation and subsequent tissue remodeling (Geenens et al., 32 2016a). 33

34 The tonicity of vascular smooth muscle cells (SMC) after clamping was also studied extensively (Barone et al., 1989; Famaey et al., 2010; Geenens et al., 2016b). Experimental data on the effects of arterial 35 clamping were collected in mice (Geenens et al., 2016a,b). In this study, descending thoracic aortas were 36 37 clamped at different levels of loading. Then, the aorta was either excised immediately or excised after 38 a fixed duration of healing. After excision, rings were cut and tested with a myograph to measure the 39 vascular tone under vasoconstriction and vasodilatation stimulations, followed by histological analyses. An 40 acute decline of endothelium-dependent vasodilatation was observed just after clamping, but the functional response was restored after one month (Geenens et al., 2016b). Arterial clamping was also followed by an 41 inflammatory response leading to some degree of fibrosis. 42

The role of mechanobiology in the response to arterial clamping is not clearly understood. It is known that 43 in many conditions, vascular remodeling is mediated by the mechanical stimuli sensed by vascular SMCs, 44 permitting to maintain wall stresses at homeostasis (Humphrey, 2002). SMCs modulate their phenotype in 45 response to changing local environmental cues (Epstein et al., 1994), possibly performing biosynthetic, 46 proliferative, and contractile roles in the vessel wall (Thyberg et al., 1995). Contractile SMCs react to 47 environmental changes on the short term by contracting and relaxing to restore a homeostatic state. On 48 the longer term, biosynthetic vascular SMC produce and degrade the extracellular matrix, thus enabling 49 growth and remodeling (Owens et al., 2004). 50

51 In order to decipher the role of mechanobiology in the response to arterial clamping, in silico predictive 52 models can be helpful. A number of computational models for damage through overloading of soft tissues have been developed and tested by Balzani et al. (2006); Rodríguez et al. (2006); Peña (2011); 53 54 Gasser (2011); Balzani et al. (2012); Sáez et al. (2012); Forsell et al. (2013); Famaey et al. (2013); 55 Peña (2014); Schmidt and Balzani (2016); Li and Holzapfel (2019). Most of these models are based on continuum damage mechanics (Kachanov, 1986; Simo and Ju, 1987), where the amount of damaged tissue 56 57 is determined by a damage parameter. These models were successfully applied to predict acute damage after arterial clamping. However, most of them focused on short term fiber damage and modeled neither 58 the active behavior of vascular SMCs nor the healing process occurring on the longer term. 59

Modeling the active behavior of vascular SMCs has been a topic of extensive investigation (Murtada 60 et al., 2017), combining continuum mechanics (Murtada et al., 2010, 2012) and the kinetics of pathways 61 involved in the active behavior (Hai and Murphy, 1988; Schmitz and Böl, 2011; Bouklas et al., 2018; Böl 62 et al., 2012; Liu, 2014; Adachi and Kida, 2014; Ferreira et al., 2018), including phosphorylation of myosin 63 light chain, variations of intracellular calcium concentration and membrane depolarisation (Sharifimajd 64 and Stålhand, 2014). However, to our best knowledge, none of these models depend on the concentration 65 of specific vasoreactive agents used in myograph testing: phenylephrine (PE), acetylcholine (ACh) and 66 sodium nitroprusside (SNP) as nitric oxide (NO) donor. 67

Modeling vascular healing is also rather recent. Comellas et al. (2016) presented a computational model 68 69 of tissue healing after mechanical overload, in which temporal evolutions of damage are homeostasis-driven. However, no discrimination was made between the different tissue constituents (elastin, collagen, cells) in 70 71 terms of damage and mechanical behavior. This can be addressed by microstructurally-motivated growth 72 and remodeling models based on the constrained mixture model introduced by Humphrey and Rajagopal (2002). In the constrained mixture theory, the different constituents of the tissue are constrained to move 73 together in a mixture but all have different biologically relevant stress-free states. Tissue remodeling is 74 75 governed by laws of production and degradation for each constituent based on stress states. This type 76 of model has been used to predict different tissue adaptations such as aneurysm growth for instance by Mousavi et al. (2019); Latorre and Humphrey (2018b); Famaey et al. (2018); Braeu et al. (2017); Cyron 77 et al. (2016); Watton and Hill (2009); Zeinali-Davarani and Baek (2012); Baek et al. (2006); Valentín et al. 78 (2013); Alberto Figueroa et al. (2009) or wound healing by Zuo et al. (2020). However, to the best of our 79 knowledge, the constrained mixture theory has never been used to model healing after arterial clamping. 80

In the present work, we aim to computationally capture the mechanobiological effects of arterial clamping. Therefore, we introduce a chemomechanical model in a constrained mixture framework, considering inflammation, collagen deposition, SMC proliferation, SMC active response as well as SMC switch from contractile to synthetic phenotype, all depending on the mechanical and chemical environment. After introducing the details of the model, we simulate the response to arterial clamping after one and two months of healing and compare the results to experimental data.

2 MATERIALS AND METHODS

87 2.1 Mouse experiments

As reported by Geenens et al. (2016a), 108 wildtype mice were subjected to a surgical procedure, in 88 which the descending thoracic aorta was isolated and clamped in vivo with a non-serrated, 2 mm wide 89 90 clamp at either a loading level of 0.0 N (control group), 0.6 N or 1.27 N. The clamped tissue was then either immediately excised, or *in vivo* healing was allowed for 6 hours, 2 weeks or 1 month. After these four 91 time points, histological analyses were carried out to assess the structural integrity of the tissue through 92 CD105, CD45, Verhoeff's-Van Gieson and osteopontin - α -SMA stainings. After the immediate excision 93 or after 1 month, myograph tests were carried out to assess the functional integrity of the tissue. The aorta 94 segment was mounted onto two rods in an organ bath and, upon stretching of the tissue, a stable preload 95 of 20 mN was reached. Afterwards, the vasoactive substances PE, ACh and SNP were subsequentially 96 97 added to the solution to assess endothelium dependent and independent vasodilation. In total, all mice that underwent surgery were divided into eighteen groups corresponding to a particular condition, depending 98 on the clamping force and the healing time, and on the type of assessment, *i.e.* histology or myograph. 99 100 More details on these animal experiments are given in Geenens et al. (2016a).

101 2.2 Constitutive model

102 2.2.1 Passive material behavior

103 The anisotropic and nonlinear passive mechanical behavior of arterial tissue is often represented by 104 a Gasser-Ogden-Holzapfel (Gasser et al., 2006) hyperelastic formulation. The deviatoric strain energy 105 function is decomposed in an isotropic Neo-Hookean part, representing the elastin fibers in the tissue, 106 and an exponential, anisotropic part, representing two collagen fiber families running in two symmetric 107 directions. Assuming a fully incompressible material and ignoring the volumetric contribution, the strain 108 energy function of the elastin and collagen contribution is respectively written as

$$\hat{\Psi}^{elas} = C_{10}(\bar{I}_1^{elas} - 3),$$

$$\hat{\Psi}_i^{coll} = \frac{k_1}{2k_2} \exp\left\{k_2[(\kappa \bar{I}_1^{coll} + (1 - 3\kappa)\bar{I}_i^{coll}) - 1]^2\right\} - 1, \quad i = 4, 6,$$
(1)

109 where C_{10} and k_1 represent the stiffness of elastin and collagen. k_2 determines the exponential collagen 110 behavior and κ quantifies the fiber dispersion. \bar{I}_1^{elas} and \bar{I}_1^{coll} are the first invariants or traces of the 111 deviatoric right Cauchy-Green stretch tensors $\operatorname{tr}(\bar{C}^{elas}) = \operatorname{tr}(J^{-2/3} F^{elas^T} F^{elas})$ and $\operatorname{tr}(\bar{C}^{coll}) =$ 112 $\operatorname{tr}(J^{-2/3} F^{coll^T} F^{coll})$, where F^{elas} and F^{coll} are the deformation gradients of elastin and collagen 113 respectively and J is the Jacobian of the deformation gradient F. More information on these different 114 deformation gradients follows in section 2.2.4. \bar{I}_4^{coll} and \bar{I}_6^{coll} are the fourth and sixth invariants of \bar{C}^{coll} 115 and M_i , representing the stretch along the preferred fiber direction, written as

$$\bar{I}_i^{coll} = \boldsymbol{M}_i \cdot (\bar{C}^{coll} \boldsymbol{M}_i), \quad i = 4, 6,$$
(2)

with M_i the undeformed fiber vector defined by the fiber angle α_i with respect to the circumferential direction. Therefore, $M_i = [0 \cos \alpha_i \sin \alpha_i]^T$, assuming that the radial direction is the first direction, the circumferential direction the second and the axial the third.

119 2.2.2 Active material behavior

120 Contractile SMCs in the media actively generate vascular tone. An active component to the strain energy121 function, as described by Murtada et al. (2010) and used by Famaey et al. (2013) takes the form

$$\hat{\Psi}^{csmc} = \frac{\mu_{smc}}{2} \left(n_3 + n_4 \right) \left(\sqrt{\bar{I}_4^{smc}} + u_{rs} - 1 \right)^2, \tag{3}$$

where μ_{smc} is a stiffness-like material parameter, n_3 and n_4 together are the fractions of the smooth muscle 122 filaments in the force-producing states. u_{rs} represents the normalized sliding between the filaments arising 123 124 from the difference between the stress in the surrounding matrix P_{mat} and the driving stresses of the crossbridges of the filaments P_{smc} . Murtada et al. (2010) give an in-depth explanation of these variables. This is 125 also further elaborated in section 2.5. \bar{I}_4^{smc} is the fourth invariant of M_{smc} and $\bar{C}^{smc} = J^{-2/3} F^{smcT} F^{smc}$, 126 the deviatoric right Cauchy-Green stretch tensor of the smooth muscle fibers associated with the smooth 127 muscle deformation gradient F^{smc} . \bar{I}_4^{smc} can be written similarly to equation 2, where M_{smc} represents 128 the orientation of the cells. Assuming that the cells are aligned along the circumferential direction, we 129 write $M_{smc} = [0 \ 1 \ 0]^T$. 130

131 2.2.3 Strain energy function

Similarly to Famaey et al. (2018), the overall strain energy density stored in the material is calculatedwith a mass-averaged rule as

$$\Psi = \Psi^{elas} + \Psi^{coll} + \Psi^{csmc}$$

$$= \rho^{elas} (k) \hat{\Psi}^{elas} (k) + \sum_{\tau=0}^{k} \rho^{coll} (k,\tau) \hat{\Psi}^{coll}_{i} (k,\tau) + \rho^{csmc} (k) \hat{\Psi}^{csmc} (k) , \qquad (4)$$

134 where $\rho^{elas}(k)$ represents the elastin density at the current time step k. $\rho^{csmc}(k)$ is the density of SMCs in 135 their contractile phenotype and $\rho^{coll}(k, \tau)$ is the density of collagen cohort τ . These considered densities and constituent specific strain energy densities relate to the reference configuration. The deposition ofcollagen is discretized, such that different collagen cohorts can be identified, depending on the time of

138 production. We consider all the initially present collagen as one cohort deposited at k = 0. On top of that,

139 at every discrete time step, one cohort per collagen family is produced. At every time step, all existing

140 cohorts, for example previously deposited at time step τ , are degraded through a slowly decaying survival 141 fraction. In the present study, we consider two symmetric fiber families, each divided into k + 1 cohorts at

142 every time step k.

143 2.2.4 Deformation gradient

144 The strain energy defined in equation 4 depends on the deformation gradients of the considered 145 constituents. According to the constrained mixture growth and remodeling theory, the total deformation 146 gradient of elastin F^{elas} is written as

$$F^{elas} = FG^{elas},\tag{5}$$

147 where G^{elas} is a deformation gradient containing the deposition stretches of elastin in the *in vivo* 148 homeostatic reference state of the artery and F represents any deformation of the mixture as a whole with 149 respect to this reference.

150 The total deformation gradient of a certain collagen cohort τ is

$$\boldsymbol{F}^{coll}\left(\tau\right) = \boldsymbol{F} \boldsymbol{F}_{dep}^{coll}\left(\tau\right)^{-1} \boldsymbol{G}^{coll}.$$
(6)

151 G^{coll} represents the deformation of the collagen cohort at deposition. $F^{coll}_{dep}(\tau)$ is the deformation of the 152 mixture at the time of deposition with respect to the homeostatic reference state and F is the current 153 deformation. In a steady state regime, the deformation at deposition of all collagen still present is equal 154 to the current deformation, such that $F^{coll}_{dep} = F$ and that F^{coll} is simply equal to G^{coll} , the collagen 155 deposition stretch tensor. Collagen is assumed to be deposited at a constant stretch g^{coll} (Bellini et al., 156 2014) along the main fiber direction. Therefore, for a particular fiber direction M (Cyron et al., 2016),

$$\boldsymbol{G}^{coll} = g^{coll} \boldsymbol{M} \otimes \boldsymbol{M} + \frac{1}{\sqrt{g^{coll}}} (\boldsymbol{I} - \boldsymbol{M} \otimes \boldsymbol{M}).$$
(7)

157 Contractile SMCs are assumed to only feel the deformation with respect to the state at which they were158 deposited. Therefore, their deformation gradient is

$$\boldsymbol{F}^{csmc}\left(\tau\right) = \boldsymbol{F}\boldsymbol{F}_{dep}^{csmc}\left(\tau\right)^{-1},\tag{8}$$

159 where $F_{dep}^{csmc}(\tau)$ is the deformation gradient of the mixture at the time of deposition τ of the considered 160 cohort.

All deformations are considered fully incompressible. Moreover, volumetric changes due to mass additionor loss are neglected, such that no deformation is observed as a result of growth.

163 2.3 Damage model

Figure 1 gives an overview of the considered damage effects. A short-term damage model for contractile SMCs and collagen inspired by Famaey et al. (2013) and Balzani et al. (2006) is considered. The fraction of damaged cells is modeled as a damage parameter d^{csmc} , calculated as

$$d^{csmc} = 1 - \exp\left(-\beta/m^{csmc}\right),\tag{9}$$

167 where m^{csmc} is a damage constant and

$$\beta = \max abs \left(\lambda_{\theta\theta} - 1\right),\tag{10}$$

where $\lambda_{\theta\theta}$ is the local circumferential stretch with respect to the *in vivo* reference stretch, assuming that the deformation gradient is known in a predefined local coordinate system whose axes are aligned with the

170 local radial, circumferential and axial directions.

171 The fraction of damaged collagen d^{coll} becomes

$$d^{coll} = 1 - \exp\left(-\Delta\zeta/m^{coll}\right),\tag{11}$$

172 where again m^{coll} is a constant. $\Delta \zeta$ is calculated as the difference between the current and homeostatic 173 fiber stresses (see also equation 13).

We assume that endothelium can be damaged as a result of overloading of the inner elastin of the media, since the endothelium itself bears almost no load. As stated by Jufri et al. (2015), endothelial cells react differently to physiological and pathological ranges of mechanical stretch, where the latter may induce apoptosis (Kou et al., 2009). We therefore assume that the local endothelium dies if a certain threshold m^{ec} of the local β is exceeded.

179 2.4 Remodeling model

A remodeling algorithm is defined considering six main components. In the following sections, remodeling pathways are introduced for the two main passive load-bearing constituents elastin and collagen. Two SMC phenotypes are considered, active load-bearing contractile cells and non-load-bearing synthetic cells that produce extracellular matrix. We also consider the healing of the endothelium and the infiltration of inflammatory agents. The scheme on figure 2 is an overview of the remodeling pathways with the corresponding equations introduced in the following sections.

186 2.4.1 Elastin

187 We assume that the production or degradation of elastin is negligible over the considered time frame 188 and that new elastin cannot be produced. The elastin density at each time point is therefore equal to the 189 homeostatic density ρ_0^{elas} .

190 2.4.2 Collagen

191 The density at time step k of a collagen cohort deposited at time τ is (Famaey et al., 2018; Valentín et al., 192 2013)

$$\rho^{coll}(k,\tau) = m^{coll}(\tau) q^{coll}(k,\tau).$$
(12)

193 $m^{coll}(\tau)$ represents the amount of collagen of the specified cohort at the time of deposition τ and $q^{coll}(k, \tau)$ 194 is the fraction is this cohort that survives until time k.

The degradation of collagen depends on the current fiber stress. Upon discretization of equation 8 in
Famaey et al. (2018) or 53 in Valentín et al. (2013), the survival fraction of a certain collagen cohort is

$$q^{coll}(k,\tau) = \exp\left(-\sum_{\tilde{\tau}=\tau}^{k} K_{qh}^{coll} \Delta t \left(1 + \Delta \zeta \left(k,\tilde{\tau}\right)^{2}\right)\right),\tag{13}$$

197 where K_{qh}^{coll} is the homeostatic decay constant and $\Delta \zeta (k, \tilde{\tau})$ represents the difference between the current 198 and homeostatic fiber stresses as defined in Famaey et al. (2018); Valentín et al. (2013). 199 The production of new collagen cohorts is proportional to the current density of synthetic cells. The rate 200 at which they produce collagen depends on the presence of contractile cells and on the mechanical stimulus 201 $\Delta\lambda$ felt by these latter cells. The production rate at time τ is written as

$$m^{coll}\left(\tau\right) = m_{0}^{coll}\Gamma\left(\tau\right),\tag{14}$$

202 where

$$\Gamma(\tau) = \left(1 + \frac{\rho^{csmc}(\tau)}{\rho_0^{csmc}} K_m^{coll} \Delta \lambda\right) \frac{\rho^{ssmc}(\tau)}{\rho_0^{ssmc}}.$$
(15)

203 ρ^{csmc} and ρ_0^{csmc} are the current and homeostatic densities of contractile cells and ρ_0^{ssmc} and ρ_0^{ssmc} are the 204 corresponding densities of the synthetic cells. K_m^{coll} is a remodeling parameter and the mechanical criterion 205 for remodeling is written as

$$\Delta \lambda = \lambda_{\theta\theta} - 1. \tag{16}$$

Note that $\Delta \lambda$ is very similar to β^{csmc} . It can however be positive or negative for circumferential stretch or compression respectively.

208 2.4.3 Contractile smooth muscle cells

Contractile SMCs dedifferentiate into synthetic cells upon mechanical triggering, for example as observed
by Wang et al. (2018) or when losing grip to the surrounding extracellular matrix. We assume that these
cells react to stretch in the circumferential direction as

$$\frac{d\rho^{csmc}}{dt} = -\Delta\lambda K_{dd}^{smc}\rho^{csmc}.$$
(17)

212 Through numerical integration, this becomes

$$\rho^{csmc}\left(k\right) = \rho^{csmc}\left(k-1\right)\left(1 - \Delta\lambda K_{dd}^{smc}\Delta t\right),\tag{18}$$

213 where K_{dd}^{smc} is a rate parameter for cell differentiation and Δt the considered time step. Regardless of 214 this equation, the maximum relative amount of contractile cells $\rho^{csmc}/\rho_0^{csmc}$ is bounded by the relative 215 amount of elastin $\rho^{elas}/\rho_0^{elas}$ since we assume that cells cannot be contractile if they are unable to grip the 216 extracellular environment. The increase of contractile SMCs is also bounded by the available amount of 217 synthetic cells to differentiate from.

218 2.4.4 Synthetic smooth muscle cells

Whereas contractile cells are quiescent in their normal state, synthetic SMCs are more proliferative (Hao et al., 2003). We therefore assume that their density can increase through dedifferentiation or proliferation based on the mechanical environment (Mantella et al., 2015). Moreover, their proliferation also increases as a reaction to inflammation (Yang et al., 2018). We write the evolution law of these cells as

$$\frac{d\rho^{ssmc}}{dt} = \left(\Delta\lambda K_{pl}^{smc} + K_{ic}^{smc}\phi^{ic}\right)\rho^{ssmc} + \Delta\lambda K_{dd}^{smc}\rho^{csmc}.$$
(19)

223 Therefore, in discretized form, the current synthetic cell density is

$$\rho^{ssmc}\left(k\right) = \rho^{ssmc}\left(k-1\right) \left(\Delta\lambda K_{pl}^{smc}\Delta t + 1 + K_{ic}^{smc}\Delta t\phi^{ic}\right) + \rho^{csmc}\left(k-1\right) - \rho^{csmc}\left(k\right),\tag{20}$$

where K_{pl}^{smc} and K_{ic}^{smc} are rate parameters related to the mechanical and inflammatory stimulus respectively and ϕ^{ic} is the current fraction of inflammation.

226 2.4.5 Endothelial cells

227 After degradation, the endothelium heals following the logistic growth law

$$\frac{d\phi^{ec}}{dt} = K^{ec} \left(1 - \phi^{ec}\right) \phi^{ec} \tag{21}$$

228 or

$$\phi^{ec}(k) = (K^{ec}\Delta t (1 - \phi^{ec} (k - 1)) + 1) \phi^{ec} (k - 1), \qquad (22)$$

in which K^{ec} is a rate parameter and ϕ^{ec} is the total fraction of endothelium present. $\phi^{ec} = 1$ means that the endothelium is fully recovered. If $\phi^{ec} = 0$ and no endothelial cells are present at all, no recovery is possible.

232 2.4.6 Inflammation

We model inflammation provoked when platelets and leukocyte adhere to the de-endothelialized artery and send inflammatory agents in the tissue. We therefore assume that the inflammation is directly related to the fraction of intact endothelium:

$$\phi^{ic} = 1 - \phi^{ec},\tag{23}$$

236 where ϕ^{ic} represents a relative level of inflammation with a maximum of 1.

237 2.5 Contractility model

Equation 3 describes the energy generated by SMCs. As stated before, this energy depends on the muscle filament sliding and on the fraction of the filaments in their force-producing states.

Similarly to Famaey et al. (2013); Murtada et al. (2010), the driving equation for the evolution of the relative sliding of the myofilaments u_{rs} is

$$\dot{u}_{rs} = \frac{1}{\eta} \left(P_{smc} - P_{mat} \right), \tag{24}$$

242 with

$$P_{mat} = \frac{\partial \Psi^{csmc}}{\partial \lambda_{\theta}} = J^{-2/3} \mu^{csmc} \left(n_3 + n_4 \right) \left(\sqrt{\bar{I}_4^{csmc}} + u_{rs} - 1 \right), \tag{25}$$

243 and

$$P_{smc} = \begin{cases} \kappa_c n_3, & \text{for } P_{mat} < \kappa_c n_3 \\ P_{mat}, & \text{for } \kappa_c (n_3 + n_4) \ge P_{mat} \ge \kappa_c n_3 \\ \kappa_c (n_3 + n_4), & \text{for } P_{mat} > \kappa_c (n_3 + n_4) \end{cases}$$
(26)

In a steady-state or homeostatic condition, u_{rs} is evolved to a situation where $P_{smc} = P_{mat}$. Mathematically, three situations can be discerned. Potentially, the u_{rs} value from a previous state already allows $\kappa_c (n_3 + n_4) \ge P_{mat} \ge \kappa_c n_3$ to be true in the new steady-state, such that $P_{smc} = P_{mat}$ already holds, as can be seen from equation 26. u_{rs} then does not evolve further in the new steady-state. The previous u_{rs} may also cause P_{mat} to be smaller than $\kappa_c n_3$. In that case, u_{rs} evolves until $P_{mat} = \kappa_c n_3$. At the final steady-state, u_{rs} is then written as

$$u_{rs} = \frac{\kappa_c n_3}{\mu_{smc} \left(n_3 + n_4\right)} + 1 - \sqrt{\bar{I}_4^{csmc}}.$$
(27)

250 Alternatively, if the previous u_{rs} causes P_{mat} to be greater than $\kappa_c (n_3 + n_4)$, u_{rs} evolves to

$$u_{rs} = \frac{\kappa_c}{\mu_{smc}} + 1 - \sqrt{\bar{I}_4^{csmc}}.$$
(28)

251 The steady-state configuration of the muscle filaments can therefore be calculated at any level of 252 deformation, quantified by \bar{I}_4 .

As defined by Murtada et al. (2010); Hai and Murphy (1988), the myofilaments switch between their states n_1 , n_2 , n_3 or n_4 by the set of differential equations

where n_1 and n_2 represent the fractions of myofilaments in their detached state, while n_4 and n_3 represent the fractions of attached filaments, dephosphorylated and phosphorylated, respectively. As explained by Murtada et al. (2010), the rate parameters k_1 and k_6 are dependent on the calcium concentration $[Ca^{2+}]$ using Michaelis-Menten kinetics as

$$k_1 = k_6 = \frac{[CaCaM]^2}{[CaCaM]^2 + K_{CaCaM}^2} s^{-1}, [CaCaM] = \alpha_{Ca}[Ca^{2+}].$$
(30)

259 The second equation represents the formation of Calcium-Calmodulin complex, where α is a positive 260 constant. K_{CaCaM} in the first equation, inspired by Yang et al. (2003) is a CaCaM-dependent 261 phosphorylation rate parameter.

We assume that $[Ca^{2+}]$ represents the intracellular calcium contraction. This concentration can be influenced by vasoactive agents, such as the vasodilator NO and the vasoconstrictor PE. The response to these agents is normalized with respect to the maximal possible response by the Hill equation:

$$R_{NO} = \frac{[NO]}{[NO] + K_{NO}} \tag{31}$$

265 and

$$R_{PE} = \frac{[PE]}{[PE] + K_{PE}},\tag{32}$$

where [NO] and [PE] are the respective extracellular concentrations of NO and PE. K_{NO} and K_{PE} determine the inflection points of the Hill curves. PE causes an increase of the calcium concentration, whereas NO decreases it. The current intracellular calcium concentration is determined as

$$[Ca^{2+}] = [Ca^{2+}]_{hom} + \alpha_{PE}R_{PE} - \alpha_{NO}R_{NO},$$
(33)

where $[Ca^{2+}]_{hom}$ is the homeostatic intracellular calcium concentration and α_{PE} is the maximum extra calcium concentration in response to PE. α_{PE} is the maximal calcium concentration that is removed in response to NO. Plugging the resulting $[Ca^{2+}]$ back into equation 30, a NO and PE dependency of the rate parameters k_1 and k_6 is observed. k_2 and k_5 , that define the rate of dephosphorylation, are also directly affected by the NO concentration since NO activates myosin light chain phosphatase (Carvajal et al., 2000). We write

$$k_2 = k_5 = k_{2,hom} + \alpha_2 R_{NO}, \tag{34}$$

where $k_{2,hom}$ is the homeostatic rate of dephosphorylation and α_2 is the maximal increase in response to NO.

NO is produced by a healthy endothelium in response to, for example, the vasodilating agent acetylcholine
(ACh) or wall shear stress (WSS). Cohen et al. (1997) measured the NO concentration in response to ACh.
Their findings are approximated with the equation

$$[NO]_{ec} = 2.8 \cdot 10^{-7} \cdot \frac{\log_{10} \left([ACh]/1M \right) + 8.2}{0.9 + \log_{10} \left([ACh]/1M \right) + 8.2} \cdot 1M,$$
(35)

where $[NO]_{ec}$ is the extracellular concentration of NO produced by the endothelium, [ACh] is the extracellular concentration of ACh and 1M refers to a concentration of one molar.

In summary, the whole dependency of the contractile state of the contractile cells on the vasoactive agent PE, NO and ACh is schematically represented in figure 3.

284 2.6 Finite element model

The material, damage and remodeling models explained above are used for an *in silico* reproduction of the experiments carried out by Geenens et al. (2016a) *in silico*. A finite element model is set up in Abaqus/Standard 2017 to represent a mouse aorta. The diastolic geometry of the aorta is represented as a half cylinder with inner diameter 0.65 mm and thickness 0.04 mm (Bersi et al., 2016). Due to symmetry, only a length of 0.04 mm of the cylinder is modeled. The geometry consists of 12852 full integration, hexahedral, hybrid elements (C3D8H). The simulation goes through a number of steps explained below, according to the steps followed in the actual experiments. An overview is also shown in figure 4.

292 2.6.1 Homeostasis

In order to model the *in vivo*, mechanobiologically homeostatic condition of the mouse aorta, the prestressing algorithm explained in Mousavi and Avril (2017); Famaey et al. (2018); Maes et al. (2019) is used. This algorithm looks for a suitable deposition stretch deformation gradient for elastin G^{elas} in order to balance the diastolic *in vivo* reference geometry with the intraluminal diastolic pressure p = 10 kPa, while the top and bottom of the arterial section are fixed in axial direction. The collagen deposition stretch g^{coll} and the axial elastin deposition stretch g^{elas}_{ax} are fixed as prior knowledge. This simulation is shown in figure 4 as step 1.

300 2.6.2 Clamping

As shown in step 2 of figure 4, the clamping of the aorta is simulated using two undeformable parallel 301 plates, similarly to Famaey et al. (2013) and as shown figure 4. Self-contact of the inner surface of the artery 302 is defined, as well as contact between the plates and the outer surface of the artery. The plates first move 303 towards each other until reaching the desired clamping force, while no damage to the material is allowed. 304 Two different clamp forces are discerned: 0.6 N or 1.27 N per 2 mm length of the clamp, respectively 305 named load 1 and load 2. In a few next steps, the clamps are held at a constant distance and the damage 306 model explained in section 2.3 is activated. When the damage to the endothelium and contractile SMC 307 has stabilized, the damage is held constant again, while the clamp plates are removed. During the whole 308 process the intraluminal pressure is fixed at mouse a rtic level p = 10 kPa. 309

310 2.6.3 Remodeling

After the releasing of the clamp, the remodeling algorithm explained in section 2.4 is activated, while keeping the pressure constant. This is shown as step 3 in figure 4. The local collagen and SMC densities are initialized based on the previously calculated local damage. The initial value of ϕ^{ec} is defined as the percentage of intact endothelial layer in the entire considered segment. Therefore, this is not a locally defined variable, and the same value is attributed to every integration point.

316 Due to the initial loss of contractile SMC and collagen, a dilatation of the artery is observed. This 317 nonhomeostatic mechanical state drives remodeling. Every remodeling step corresponds to one day of 318 remodeling. During this process, all nodes are constrained to only move in the radial direction in order to 319 avoid excessive shearing between the layers and failure of the simulation.

320 2.6.4 Myograph test

After 31 days of remodeling (cases R1 and R2 for loads 1 and 2), immediately after clamping (cases A1 and A2 for loads 1 and 2) or immediately after obtaining the homeostatic configuration (cases A0 and R0), a myograph test is simulated, as in the mouse experiments explained in section 2.1. For further reference, an overview of these six cases is given in table 1.

325 During the simulation, first, the axial boundary condition is released, as well as the intraluminal pressure 326 to simulated excision, as shown by step 4 in figure 4. Step 5 depicts the simulation of a myograph 327 experiment as explained in section 2.1 similarly to Famaey et al. (2013). An undeformable rod with a 328 radius of 0.15 mm in Abaqus is pushed into the arterty until the approximate required preload of 0.0133 N 329 per mm length of the arterial segment is reached, assuming that the preload was set at 0.02 N for a length of approximately 1.5 mm in the actual experiments carried out by Geenens et al. (2016a). The rod is then fixed 330 at this position, while 10^{-6} M PE, 10^{-5} M ACh or 10^{-6} M NO is virtually added through field variables. 331 A contact definition is prescribed between the rod and the inner surface of the artery. Preconstriction due to 332 333 PE and relaxation due to ACh or NO can then be observed according to the smooth muscle contraction model described in section 2.5. 334

335 2.6.5 Remodeling beyond 31 days

In order to further examine the behavior of the remodeling model, the finite element analysis of section 2.6.3 was extended to a remodeling period of 91 days. The effect of slight adaptations to the model was investigated as well. The first adaptation is the assumption that synthetic SMCs do not redifferentiate into their contractile phenotype, such that equations 17 and 19 only hold when $\Delta\lambda$ is greater than or equal to zero. In the opposite case, only the inflammation level influences the SMC densities:

$$\frac{d\rho^{csmc}}{dt} = 0$$

$$\frac{d\rho^{ssmc}}{dt} = K_{ic}^{smc} \phi^{ic} \rho^{ssmc}.$$
(36)

The second adaptation is the assumption that collagen production is solely dependent on the amount
of synthetic SMCs, while their production rate is not directly affected by the mechanical environment.
Equation 15 then simplifies to

$$\Gamma\left(\tau\right) = \frac{\rho^{ssmc}\left(\tau\right)}{\rho_{0}^{ssmc}}.$$
(37)

344 2.7 Model parameters

An overview of all used parameter values is given in tables 2 and 3. A code number from 1 to 5 is attributed to every parameter, explaining the way its value was determined. We either used an exact value from the specified reference (1), or used a representative value from the reference, when for example a range was given based on the results of tests on multiple samples (2). Some parameter values are estimated (5), or the parameter is manually fitted to the experimental myograph data (see section 2.1), either with an idea of the order of magnitude from literature (3), or without (4).

3 RESULTS

351 3.1 Damage due to clamping

Figure 5 shows the distribution of β (see equation 10), defining the local loss of contractile SMC due to clamping. The highest damage is concentrated at the inner side of the wall at the edge of the clamp.

Table 4 gives an overview of the relative collagen, contractile SMC and endothelium content acutely after clamping at the three different clamp loads (cases A0, A1 and A2). From this table it can be concluded that the difference between clamping at 0.6 N and 1.27 N is small in terms of acute damage. There is approximately 70% loss of endothelium, 9% collagen loss and 28% contractile SMC loss in both cases. The small difference in damage is due to a minimal required clamp displacement to increase the reaction force from 0.6 N to 1.27 N, yielding only small stretch differences.

360 3.2 Remodeling

Table 4 also shows the situation after the simulated *in vivo* healing period of 31 days (cases R0, R1 and R2) using the presented remodeling model, taking into account cell differentiation, ECM production by synthetic cells and inflammation after clamping injury.

Figure 6 shows the evolution of the total content of each constituent relative to its normal amount over a remodeling time of 31 days after damage due to clamping at 1.27N. In other words, it shows the evolution from case A2 to case R2. Due to the similar level of damage at cases A1 and A2, as is clear from table 4, the evolution from cases A1 to R1 resembles the one depicted in this figure.

There is an initial dedifferentiation of the cells from their contractile to synthetic phenotype due to an initial overstretching of the wall. Along, with the high initial inflammation level, this also causes the synthetic cells to proliferate, such that the collagen content increases. At about 14 days, the initial stiffness loss is compensated, and the collagen, synthetic and contractile cell contents slowly return to their normal levels.

373 3.3 Myograph test

Figure 7 shows the normalized reaction force in the simulated rod while it moves towards the preload position before the addition of vasoactive substances. It indicates the overall stiffness of each material for each case. There was no discernible difference between cases A1 and A2 on the one hand and cases R1 and R2 on the other hand, as can be observed in figure 7. Furthermore, the simulations do not show differences between cases A0 and R0.

Figure 8 gives an overview of the results of the simulated myograph experiments upon the addition of vasoactive substances, compared to the results obtained on mouse arteries, as explained in section 2.1. The figure shows how the isometric force changes upon addition of PE, NO and ACh. PE drives an increased phosphorylation rate $k_1 = k_6$ of the myofilaments through an increased intracellular calcium level, inducing a vasoconstrictive effect. NO has the reverse effect on calcium and it also increases the dephosphorylation rate $k_2 = k_5$. ACh does not act directly on the contractile SMC, but triggers the endothelium to produce NO. Therefore, the vasodilating effect of ACh is smaller than that of NO.

386 3.4 Remodeling beyond 31 days

The evolution of relative collagen, synthetic cells and contractile cells density over a remodeling period of 91 days is shown in figure 9, for the original remodeling model (A) and two adapted models (B and C) as explained in section 2.6.5. Beyond one month, unnatural periodic behavior emerges when using the original model, caused by the initial extra loss of contractile SMC upon overstretching, causing an extra stiffness loss, and a delay in the increased collagen production through the proliferation of synthetic cells.

When synthetic cells do not redifferentiate into the contractile phenotype, a more rapid stabilization of the remodeling is observed (figure 9 B). However, the loss of contractile cells due to clamping overload will never be compensated in this case.

The results of the last variant of the remodeling model are shown in figure 9 C and show an increased oscillation of the synthetic cell density. Collagen production is not dependent on a mechanical stimulus anymore, such that a bigger increase in synthetic cell count is required to restore the collagen density, and along with it, restore the homeostatic mechanical environment.

4 DISCUSSION

The aim of this study is to introduce a computational model predicting healing in arterial tissues subjected to mechanical overloading and damage, for instance after clamping. Three models are introduced for the *in silico* simulation of the experiments carried out by Geenens et al. (2016a): a damage model for clamping, a remodeling model to predict healing and a contractility model to simulate myograph experiments.

403 The contractility model is original as it is the first to take the vasoactive substances PE, NO and ACh 404 into account. Their respective influence on the rate of phosphorylation and dephosphorylation of myosin 405 light chain leads to a reliable response in the simulation of a myograph experiment, as shown in figure 8. The model is based on signaling pathways on the cellular level, dependent as well as independent on 406 407 intracellullar calcium, as shown in figure 3. The approach is different from the recent model presented by 408 Murtada et al. (2016), in which the smooth muscle tone prediction was based on a structurally motivated model of the contractile unit. In their implementation, the response to an external factor, such as a change 409 410 in loading or in the concentration of a vasoactive agent, is modeled as an evolving scaling factor for 411 the myosin filament length. Before us, the continuum mechanics-based model of Murtada et al. (2017) was the only one that accounted for the dependency of the phophorylation rates on the diffusion of the 412 vasoconstrictor potassium chloride (KCl) from the adventitia, although diffusion itself is neglected in the 413 414 present study.

415 The remodeling model includes novel aspects of cell differentiation upon mechanical stimulus and the production of extracellular matrix by synthetic SMCs. This production is also dependent on a certain 416 level of tissue inflammation, as for example done by Latorre and Humphrey (2018b). In the present 417 418 approach however, the inflammation level is directly related to the damage and healing of the endothelium. 419 Inflammation increases the synthetic cell proliferation, thus indirectly enhancing collagen production (Davis et al., 2003), as summarized in figure 2. Hence, our remodeling model includes all the relevant 420 biological processes and pathways, in contrast to more phenomenological models, where collagen turnover 421 422 is directly related to a mechanical stimulus, such as in Mousavi et al. (2019); Famaey et al. (2018); Braeu et al. (2017); Cyron et al. (2016); Baek et al. (2006); Valentín et al. (2013); Alberto Figueroa et al. (2009). 423

424 This more detailed description of SMC behavior in vascular healing and remodeling comes at an increased 425 computational cost. Moreover, figure 9 shows stability issues of the model in the form of unnatural temporal oscillations of the densities at longer time scales. A solution could be to neglect the transient effects and 426 only consider the steady state, such as done by Latorre and Humphrey (2018a). Alternatively, we can 427 428 include damping in the model to obtain a critical or overdamped dynamic system in order to avoid unnatural periodic behavior. From a mathematical point of view, the main limitation is the high number of parameters, 429 as summarized in tables 2 and 3. Some parameters are determined based on previous works or based 430 on their physical meaning, others were set in order to match experimental findings, mainly based on the 431 tissue properties at 0 and 31 days of healing, which were however not sufficient to uniquely determine the 432 parameter values. 433

434 Unfortunately, the currently available experimental data is not sufficient to proof its pilot application. It is likely that other parameter combinations would amount to the same results as shown in figure 8. 435 436 Nevertheless, the phenomenological nature of this new model is strongly reduced as compared to state-437 of-the-art models. A high number of parameters can be qualified as physics-based, such that their values 438 can be obtained through the design of dedicated biochemical experimental set-ups. This will allow these 439 parameter values to be measured with more certainty, or with smaller confidence intervals, capturing the 440 individual differences and differences between tissue types, allowing a better focus of the parameter fitting 441 process.

442 Constrained mixture models are generally computationally expensive due to its high memory use, inversely related to the length of the time step. To ensure the feasibility, we chose to use a time step of one 443 444 day, where a time convergence study showed errors of less than five percent with respect to the situation with a time step of half a day. Also in an attempt to limit the computational cost, only a very short segment 445 of artery is modeled and the defined boundary conditions cause a plane strain situation. Considering a 446 longer segment, possibly along with a more realistic patient-specific geometry, would improve the reliability 447 of the model, mainly near the edges of the clamp and near the edges of the excised sample during the 448 myograph simulation. Using this very short artery segment allows to use a non-localized variable ϕ^{ec} that 449 represents the overall intactness of the endothelium in the segment. Localizing the endothelial damage 450 would greatly affect the complexity of the model, since diffusion of inflammatory agents and NO would 451 need to be integrated. Similarly, taking into account the migration of SMC as an important mechanism in 452 vascular remodeling, would increase the complexity as the remodeling in a certain integration point would 453 be affected, not only by all variables defined in that specific location, but also by its surroundings. In a 454 similar way, one could also consider re-endothelialization as a non-localized process of proliferation and 455 migration of nearby endothelial cells. Including all these processes would increase the biofidelity of the 456 model, although it is unclear to what extent, given the already many unknowns in the present version. 457

Furthermore, to further improve the remodeling model, an improved understanding of biological and 458 biochemical phenomena is required. To this day, some unknowns, uncertainties and controversies remain. 459 For example, it is unclear to what kind of mechanical stimulus cells react. There are indications that SMCs 460 and fibroblasts have a preferred structural stiffness of the extracellular matrix and react based on deviations 461 from this ideal value (Humphrey, 2008). On the other hand, certain signaling pathways are thought to be 462 triggered by so-called baroreceptors, sensitive to mechanical stretch (Lacolley et al., 2017). Multiple studies 463 have investigated the effects of cyclic straining of arterial tissue, as reviewed by Mantella et al. (2015). 464 Some apparently contradictory results emerge. For example, Chang et al. (2003) observed an increased 465 SMC proliferation under in vitro cyclic strain, while Morrow et al. (2005); Guha et al. (2011) observed 466 a decreased proliferation, potentially due to a different experimental design that mimics *in vivo* loading 467

468 conditions better (Mantella et al., 2015). The widely accepted theory that precursor cells differentiate into
469 synthetic cells and subsequently become fully differentiated contractile cells has been challenged recently
470 as well, given that both phenotypes can be present in healthy tissues while maintaining vascular tone and
471 tissue architecture (Rensen et al., 2007).

In summary, the presented models provide a detailed description of vascular SMC behavior under conditions of damage as well as at different concentrations of vasoactive agents. This allows us to study tissue healing and the effects of, for example, vasoactive or anti-proliferative drugs. However, there are still many unknowns regarding these phenomena, which is why more detailed and carefully designed experiments are needed in order to fully capture SMC behavior in all its aspects.

477 To conclude, in this study, a damage model, as well as a remodeling and cell contractility model were introduced, taking into account endothelial damage and healing, tissue inflammation, mechanosensing, 478 479 extracellular matrix production and phenotype switching of SMCs. Using these models, in vivo clamping 480 tests on mice aortas and subsequent healing and myograph tests, were simulated through finite element 481 modeling. The results of the simulated myograph tests showed great resemblance to the results of the actual 482 experiments. This detailed mechanobiological description of vascular SMC behavior can be clinically 483 relevant to enable *in silico* investigations of drug effects. However, the results show that there is still a need for an improved biological and biochemical fundamental understanding to reliably capture vascular SMC 484 485 mechanobiology at all the relevant spatio-temporal scales.

CONFLICT OF INTEREST STATEMENT

The authors declare that the research was conducted in the absence of any commercial or financialrelationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

LM, SA and NF conceived the presented ideas. LM developed and implemented the presented algorithms.
JV, SA and NF reviewed the computational methods. LM wrote the manuscript. JV, SA and NF reviewed
the manuscript. SA and NF supervised the project.

FUNDING

This work was supported by the Research Foundation-Flanders (FWO) through a travel grant (V414519N)
and a doctoral fellowship (11A6519N) to LM, a doctoral grant strategic basic research (SB 1S35316N)
to JV and a postdoctoral fellowship (PDO/012) to NF. It was also supported by the ERC through the
ERC-2014-CoG BIOLOCHANICS grant to SA.

ACKNOWLEDGMENTS

The authors would like to thank Prof. Erik Verbeken, Dr. Rachel Geenens and Ir. Yelliz Mattheus (KULeuven, Belgium) for their interesting insights in this work.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without unduereservation.

REFERENCES

- 499 Adachi, T. and Kida, N. (2014). Numerical analysis of arterial contraction regulated by smooth muscle
- stretch and intracellular calcium ion concentration. *Journal of Biomechanical Science and Engineering*doi:10.1299/jbse.2014jbse0002
- Alberto Figueroa, C., Baek, S., Taylor, C. A., and Humphrey, J. D. (2009). A computational framework for
 fluid-solid-growth modeling in cardiovascular simulations. *Computer Methods in Applied Mechanics and Engineering* 198, 3583–3602. doi:10.1016/j.cma.2008.09.013
- Baek, S., Rajagopal, K. R., and Humphrey, J. D. (2006). A Theoretical Model of Enlarging Intracranial
 Fusiform Aneurysms. *Journal of Biomechanical Engineering* 128, 142–9. doi:10.1115/1.2132374
- Balzani, D., Brinkhues, S., and Holzapfel, G. A. (2012). Constitutive framework for the modeling of
 damage in collagenous soft tissues with application to arterial walls. *Computer Methods in Applied Mechanics and Engineering* 213-216, 139–151. doi:10.1016/j.cma.2011.11.015
- Balzani, D., Schröder, J., and Gross, D. (2006). Simulation of discontinuous damage incorporating residual
 stresses in circumferentially overstretched atherosclerotic arteries. *Acta Biomaterialia* 2, 609–618.
 doi:10.1016/j.actbio.2006.06.005
- Barone, G. W., Conerly, J. M., Farley, P. C., Flanagan, T. L., and Kron, I. L. (1989). Assessing clamp-related
 vascular injuries by measurement of associated vascular dysfunction. *Surgery* 105, 465–71
- Bellini, C., Ferruzzi, J., Roccabianca, S., Di Martino, E. S., and Humphrey, J. D. (2014). A
 Microstructurally Motivated Model of Arterial Wall Mechanics with Mechanobiological Implications. *Annals of Biomedical Engineering* 42, 488–502. doi:10.1007/s10439-013-0928-x
- Bersi, M. R., Bellini, C., Di Achille, P., Humphrey, J. D., Genovese, K., and Avril, S. (2016). Novel
 methodology for characterizing regional variations in the material properties of murine aortas. *Journal of Biomechanical Engineering* 138, 0710051–07100515. doi:10.1115/1.4033674
- Böl, M., Schmitz, A., Nowak, G., and Siebert, T. (2012). A three-dimensional chemo-mechanical
 continuum model for smooth muscle contraction. *Journal of the Mechanical Behavior of Biomedical Materials* 13, 215–229. doi:10.1016/j.jmbbm.2012.05.015
- Bouklas, N., Sakar, M. S., and Curtin, W. A. (2018). A model for cellular mechanotransduction and
 contractility at finite strain. *ZAMM Zeitschrift fur Angewandte Mathematik und Mechanik* 98, 1754–1770.
 doi:10.1002/zamm.201890001
- Braeu, F. A., Seitz, A., Aydin, R. C., and Cyron, C. J. (2017). Homogenized constrained mixture models
 for anisotropic volumetric growth and remodeling. *Biomechanics and Modeling in Mechanobiology* 16,
 889–906. doi:10.1007/s10237-016-0859-1
- Carvajal, J. A., Germain, A. M., Huidobro-Toro, J. P., and Weiner, C. P. (2000). Molecular mechanism of
 cGMP-mediated smooth muscle relaxation. *Journal of Cellular Physiology* 184, 409–420. doi:10.1002/
 1097-4652(200009)184:3(409::AID-JCP16)3.0.CO;2-K
- Chang, H., Shyu, K. G., Wang, B. W., and Kuan, P. (2003). Regulation of hypoxia-inducible factor-1α
 by cyclical mechanical stretch in rat vascular smooth muscle cells. *Clinical Science* 105, 447–56.
 doi:10.1042/CS20030088
- Cohen, R. A., Plane, F., Najibi, S., Huk, I., Malinski, T., and Garland, C. J. (1997). Nitric oxide is the
 mediator of both endothelium-dependent relaxation and hyperpolarization of the rabbit carotid artery.
- Proceedings of the National Academy of Sciences of the United States of America 94, 4193–4198.
 doi:10.1073/pnas.94.8.4193
- 540 Comellas, E., Gasser, T. C., Bellomo, F. J., and Oller, S. (2016). A homeostatic-driven turnover remodelling
- constitutive model for healing in soft tissues. *Journal of the Royal Society Interface* 13. doi:10.1098/rsif.
 2015.1081

- 543 Cyron, C. J., Aydin, R. C., and Humphrey, J. D. (2016). A homogenized constrained mixture (and mechanical analog) model for growth and remodeling of soft tissue. *Biomechanics and Modeling in* 545 *Mechanobiology* 15, 1389–1403. doi:10.1007/s10237-016-0770-9
- Davis, C., Fischer, J., Ley, K., and Sarembock, I. J. (2003). The role of inflammation in vascular injury and
 repair. *journal of thrombosis and heamostasis* 1, 1699–1709. doi:10.1046/j.1538-7836.2003.00292.x
- Epstein, F. H., Gibbons, G. H., and Dzau, V. J. (1994). The emerging concept of vascular remodeling. *New England Journal of Medicine* 330, 1431–8. doi:10.1056/NEJM199405193302008
- Famaey, N., Vander Sloten, J., and Kuhl, E. (2013). A three-constituent damage model for arterial
 clamping in computer-assisted surgery. *Biomechanics and Modeling in Mechanobiology* 12, 123–136.
 doi:10.1007/s10237-012-0386-7
- Famaey, N., Vastmans, J., Fehervary, H., Maes, L., Vanderveken, E., Rega, F., et al. (2018). Numerical simulation of arterial remodeling in pulmonary autografts. *ZAMM Journal of Applied Mathematics and Mechanics / Zeitschrift für Angewandte Mathematik und Mechanik* 98, 2239–2257. doi:10.1002/zamm.201700351
- Famaey, N., Verbeken, E., Vinckier, S., Willaert, B., Herijgers, P., and Vander Sloten, J. (2010). In vivo
 soft tissue damage assessment for applications in surgery. *Medical Engineering and Physics* 32, 437–43.
 doi:10.1016/j.medengphy.2010.04.002
- Ferreira, J. P., Parente, M. P., and Natal Jorge, R. M. (2018). Continuum mechanical model for cross-linked
 actin networks with contractile bundles. *Journal of the Mechanics and Physics of Solids* 110, 100–117.
 doi:10.1016/j.jmps.2017.09.009
- Forsell, C., Swedenborg, J., Roy, J., and Gasser, T. C. (2013). The quasi-static failure properties of the
 abdominal aortic aneurysm wall estimated by a mixed experimental-numerical approach. *Annals of Biomedical Engineering* 41, 1554–66. doi:10.1007/s10439-012-0711-4
- Gasser, C. (2011). An irreversible constitutive model for fibrous soft biological tissue: A 3-D microfiber
 approach with demonstrative application to abdominal aortic aneurysms. *Acta Biomaterialia* 7, 2457–
 2466. doi:10.1016/j.actbio.2011.02.015
- Gasser, T. C., Ogden, R. W., and Holzapfel, G. A. (2006). Hyperelastic modelling of arterial layers with
 distributed collagen fibre orientations. *Journal of The Royal Society Interface* 3, 15–35. doi:10.1098/rsif.
 2005.0073
- Geenens, R., Famaey, N., Gijbels, A., Verhelle, S., Vinckier, S., Vander Sloten, J., et al. (2016a).
 Atherosclerosis alters loading-induced arterial damage: Implications for robotic surgery. *PLoS ONE* 11.
 doi:10.1371/journal.pone.0156936
- Geenens, R., Famaey, N., Gijbels, A., Verhulst, V., Vinckier, S., Vander Sloten, J., et al. (2016b). Arterial
 vasoreactivity is equally affected by in vivo cross-clamping with increasing loads in young and middleaged mice aortas. *Annals of Thoracic and Cardiovascular Surgery* 15, 1389–1403. doi:10.5761/atcs.oa.
 15-00225
- Guha, S., Cullen, J. P., Morrow, D., Colombo, A., Lally, C., Walls, D., et al. (2011). Glycogen
 synthase kinase 3 beta positively regulates Notch signaling in vascular smooth muscle cells: Role in cell
 proliferation and survival. *Basic Research in Cardiology* 106. doi:10.1007/s00395-011-0189-5
- Hai, C. M. and Murphy, R. A. (1988). Cross-bridge phosphorylation and regulation of latch state in smooth
 muscle. *American Journal of Physiology Cell Physiology* 254, C99–106. doi:10.1152/ajpcell.1988.254.
 1.C99
- Hangler, H., Mueller, L., Ruttmann, E., Antretter, H., and Pfaller, K. (2008). Shunt or Snare: Coronary
 Endothelial Damage due to Hemostatic Devices for Beating Heart Coronary Surgery. *Annals of Thoracic*
- 587 Surgery 86, 1873–1877. doi:10.1016/j.athoracsur.2008.06.047

- Hao, H., Gabbiani, G., and Bochaton-Piallat, M. L. (2003). Arterial smooth muscle cell heterogeneity:
 Implications for atherosclerosis and restenosis development. *Arteriosclerosis, Thrombosis, and Vascular Biology* 23, 1510–20. doi:10.1161/01.ATV.0000090130.85752.ED
- Humphrey, J. D. (2002). Cardiovascular Solid Mechanics (New York: Springer). doi:10.1007/
 978-0-387-21576-1
- Humphrey, J. D. (2008). Vascular adaptation and mechanical homeostasis at tissue, cellular, and sub-cellular
 levels. *Cell Biochemistry and Biophysics* 50, 53–78. doi:10.1007/s12013-007-9002-3
- Humphrey, J. D. and Rajagopal, K. R. (2002). A Constrained Mixture Model For Growth And Remodelling
 Of Soft Tissues. *Mathematical Models and Methods in Applied Sciences* 12, 407–430. doi:10.1142/
 S0218202502001714
- Jufri, N. F., Mohamedali, A., Avolio, A., and Baker, M. S. (2015). Mechanical stretch: Physiological
 and pathological implications for human vascular endothelial cells. *Vascular Cell* 7. doi:10.1186/
 s13221-015-0033-z
- Kachanov, L. M. (1986). *Introduction to Continuum Damage Mechanics* (Dordrecht: Springer Netherlands).
 doi:10.1007/978-94-017-1957-5
- Kou, B., Zhang, J., and Singer, D. R. (2009). Effects of cyclic strain on endothelial cell apoptosis
 and tubulogenesis are dependent on ROS production via NAD(P)H subunit p22phox. *Microvascular Research* 77, 125–133. doi:10.1016/j.mvr.2008.08.001
- Lacolley, P., Regnault, V., Segers, P., and Laurent, S. (2017). Vascular smooth muscle cells and arterial
 stiffening: Relevance in development, aging, and disease. *Physiological Reviews* 97, 1555–1617.
 doi:10.1152/physrev.00003.2017
- Latorre, M. and Humphrey, J. D. (2018a). A mechanobiologically equilibrated constrained mixture model
 for growth and remodeling of soft tissues. *ZAMM Zeitschrift fur Angewandte Mathematik und Mechanik*98, 2048–2071. doi:10.1002/zamm.201700302
- Latorre, M. and Humphrey, J. D. (2018b). Modeling mechano-driven and immuno-mediated aortic
 maladaptation in hypertension. *Biomechanics and Modeling in Mechanobiology* 17, 1497–1511. doi:10.
 1007/s10237-018-1041-8
- Li, K. and Holzapfel, G. A. (2019). Multiscale modeling of fiber recruitment and damage with a
 discrete fiber dispersion method. *Journal of the Mechanics and Physics of Solids* 126, 226–244.
 doi:10.1016/j.jmps.2019.01.022
- Liu, T. (2014). A constitutive model for cytoskeletal contractility of smooth muscle cells. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* 470. doi:10.1098/rspa.2013.0771
- Maes, L., Fehervary, H., Vastmans, J., Mousavi, S., Avril, S., and Famaey, N. (2019). Constrained mixture
 modeling affects material parameter identification from planar biaxial tests. *Journal of the Mechanical Behavior of Biomedical Materials* 95, 124–135. doi:10.1016/j.jmbbm.2019.03.029
- 623 Mantella, L. E., Quan, A., and Verma, S. (2015). Variability in vascular smooth muscle cell stretch-induced
- 624 responses in 2D culture. *Vascular Cell* 7. doi:10.1186/s13221-015-0032-0
- Margovsky, A. I., Chambers, A. J., and Lord, R. S. A. (1999). The effect of increasing clamping forces
 on endothelial and arterial wall damage: An experimental study in the sheep. *Vascular* 7, 457–63.
 doi:10.1177/096721099900700413
- Margovsky, A. I., Lord, R. S., and Chambers, A. J. (1997). The effect of arterial clamp duration on
 endothelial injury: An experimental study. *Australian and New Zealand Journal of Surgery* 67, 448–451.
 doi:10.1111/j.1445-2197.1997.tb02012.x
- Morrow, D., Sweeney, C., Birney, Y. A., Cummins, P. M., Walls, D., Redmond, E. M., et al. (2005). Cyclic
 strain inhibits notch receptor signaling in vascular smooth muscle cells in vitro. *Circulation Research*

- 633 96, 567–75. doi:10.1161/01.RES.0000159182.98874.43
- Mousavi, S. J. and Avril, S. (2017). Patient-specific stress analyses in the ascending thoracic aorta using
 a finite-element implementation of the constrained mixture theory. *Biomechanics and Modeling in Mechanobiology* 16, 1765–1777. doi:10.1007/s10237-017-0918-2
- Mousavi, S. J., Farzaneh, S., and Avril, S. (2019). Patient-specific predictions of aneurysm growth
 and remodeling in the ascending thoracic aorta using the homogenized constrained mixture model. *Biomechanics and Modeling in Mechanobiology* 18, 1895–1913. doi:10.1007/s10237-019-01184-8
- Murtada, S. C., Arner, A., and Holzapfel, G. A. (2012). Experiments and mechanochemical modeling
 of smooth muscle contraction: Significance of filament overlap. *Journal of Theoretical Biology* 297,
 176–86. doi:10.1016/j.jtbi.2011.11.012
- Murtada, S. I., Humphrey, J. D., and Holzapfel, G. A. (2017). Multiscale and Multiaxial Mechanics of
 Vascular Smooth Muscle. *Biophysical Journal* 113, 714–727. doi:10.1016/j.bpj.2017.06.017
- Murtada, S. I., Kroon, M., and Holzapfel, G. A. (2010). A calcium-driven mechanochemical model for
 prediction of force generation in smooth muscle. *Biomechanics and Modeling in Mechanobiology* 9,
 749–62. doi:10.1007/s10237-010-0211-0
- Murtada, S. I., Lewin, S., Arner, A., and Humphrey, J. D. (2016). Adaptation of active tone in the
 mouse descending thoracic aorta under acute changes in loading. *Biomechanics and Modeling in Mechanobiology* 15, 579–592. doi:10.1007/s10237-015-0711-z
- Owens, G. K., Kumar, M. S., and Wamhoff, B. R. (2004). Molecular regulation of vascular smooth muscle
 cell differentiation in development and disease. *Physiological Reviews* 84, 767–801. doi:10.1152/
 physrev.00041.2003
- Peña, E. (2011). A rate dependent directional damage model for fibred materials: Application to soft
 biological tissues. *Computational Mechanics* 48, 407–420. doi:10.1007/s00466-011-0594-5
- Peña, E. (2014). Computational aspects of the numerical modelling of softening, damage and permanent set
 in soft biological tissues. *Computers and Structures* 130, 57–72. doi:10.1016/j.compstruc.2013.10.002
- Rensen, S. S., Doevendans, P. A., and Van Eys, G. J. (2007). Regulation and characteristics of vascular
 smooth muscle cell phenotypic diversity. *Netherlands Heart Journal* 15, 100–108. doi:10.1007/
 BF03085963
- Rodríguez, J. F., Cacho, F., Bea, J. A., and Doblaré, M. (2006). A stochastic-structurally based three
 dimensional finite-strain damage model for fibrous soft tissue. *Journal of the Mechanics and Physics of Solids* 54, 864–886. doi:10.1016/j.jmps.2005.10.005
- Sáez, P., Alastrué, V., Peña, E., Doblaré, M., and Martínez, M. A. (2012). Anisotropic microsphere-based
 approach to damage in soft fibered tissue. *Biomechanics and Modeling in Mechanobiology* 11, 595–608.
 doi:10.1007/s10237-011-0336-9
- Schmidt, T. and Balzani, D. (2016). Relaxed incremental variational approach for the modeling of damage induced stress hysteresis in arterial walls. *Journal of the Mechanical Behavior of Biomedical Materials* 58, 149–162. doi:10.1016/j.jmbbm.2015.08.005
- Schmitz, A. and Böl, M. (2011). On a phenomenological model for active smooth muscle contraction.
 Journal of Biomechanics 44, 2090–2095. doi:10.1016/j.jbiomech.2011.05.020
- Sharifimajd, B. and Stålhand, J. (2014). A continuum model for excitation-contraction of smooth muscle
 under finite deformations. *Journal of Theoretical Biology* 355, 1–9. doi:10.1016/j.jtbi.2014.03.016
- 674 Simo, J. C. and Ju, J. W. (1987). Strain- and stress-based continuum damage models-II. Computational
- aspects. International Journal of Solids and Structures 23, 841–869. doi:10.1016/0020-7683(87)
 90084-9

- Slayback, J. B., Bowen, W. W., and Hinshaw, D. B. (1976). Intimal injury from arterial clamps. *The American Journal of Surgery* 132, 183–8. doi:10.1016/0002-9610(76)90045-3
- Thyberg, J., Blomgren, K., Hedin, U., and Dryjski, M. (1995). Phenotypic modulation of smooth muscle
 cells during the formation of neointimal thickenings in the rat carotid artery after balloon injury: an
 electron-microscopic and stereological study. *Cell & Tissue Research* 281, 421–433. doi:10.1007/
 BF00417860
- Utz, J., Eckert, R., and Trautwein, W. (1999). Changes of intracellular calcium concentrations by
 phenylephrine in renal arterial smooth muscle cells. *Pflugers Archiv European Journal of Physiology*438, 725–731. doi:10.1007/s004240051099
- Valentín, A., Humphrey, J. D., and Holzapfel, G. A. (2013). A finite element-based constrained mixture
 implementation for arterial growth, remodeling, and adaptation: Theory and numerical verification.
 International Journal for Numerical Methods in Biomedical Engineering 29, 822–849. doi:10.1002/
 cnm.2555
- Vural, A. H., Yalcinkaya, S., Türk, T., Yümün, G., Gül, N., Yalcinkaya, U., et al. (2008). Intracoronary
 Shunt Versus Bulldog Clamp in Off-Pump Bypass Surgery. Endothelial Trauma: Shunt Versus Clamp.
 Journal of Surgical Research 150, 261–265. doi:10.1016/j.jss.2007.12.774
- Wang, Y., Cao, W., Cui, J., Yu, Y., Zhao, Y., Shi, J., et al. (2018). Arterial wall stress induces phenotypic
 switching of arterial smooth muscle cells in vascular remodeling by activating the YAP/TAZ signaling
 pathway. *Cellular Physiology and Biochemistry* 51, 842–853. doi:10.1159/000495376
- Watton, P. N. and Hill, N. A. (2009). Evolving mechanical properties of a model of abdominal aortic
 aneurysm. *Biomechanics and Modeling in Mechanobiology* 8, 25–42. doi:10.1007/s10237-007-0115-9
- Yang, D., Sun, C., Zhang, J., Lin, S., Zhao, L., Wang, L., et al. (2018). Proliferation of vascular smooth
 muscle cells under inflammation is regulated by NF-κB p65/microRNA-17/RB pathway activation. *International journal of molecular medicine* 41, 43–50. doi:10.3892/ijmm.2017.3212
- Yang, J., Clark, J. W., Bryan, R. M., and Robertson, C. (2003). The myogenic response in isolated rat cerebrovascular arteries: Smooth muscle cell model. *Medical Engineering and Physics* 25, 691–709. doi:10.1016/S1350-4533(03)00100-0
- Zeinali-Davarani, S. and Baek, S. (2012). Medical image-based simulation of abdominal aortic aneurysm
 growth. *Mechanics Research Communications* 42, 107–117. doi:10.1016/j.mechrescom.2012.01.008
- Zuo, D., Avril, S., Yang, H., Jamaleddin Mousavi, S., Hackl, K., and He, Y. (2020). Three-dimensional
 numerical simulation of soft-tissue wound healing using constrained-mixture anisotropic hyperelasticity
- and gradient-enhanced damage mechanics. *Journal of the Royal Society Interface* 17. doi:10.1098/rsif.
 2019.0708

FIGURE CAPTIONS

Table 1. Clarification of the codes of the six cases for which a myograph experiment is modeled: A0, R0, A1, R1, A2 and R2.

	Immediately after clamping	After one month healing
Clamp load of 0.0N	A0	R0
Clamp load of 0.6N	A1	R 1
Clamp load of 1.27N	A2	R2



Figure 1. Schematic representation of the damage effects presented in section 2.3.



Figure 2. Schematic representation of the remodeling pathways presented from equation 13 to 22.



Figure 3. Schematic representation of the pathways presented from equation 29 to 35. A green arrow with a plus sign represents a positive influence, a red arrow with a minus sign represents a negative influence.



Figure 4. Overview of all simulation steps. Step 1: obtaining homeostatic configuration, step 2: clamping the artery while allowing damage to the constituents, step 3: remodeling of the tissue after damage, step 4: releasing intraluminal pressure and axial tension, step 5: simulation of a myograph experiment. Step 4 and 5 are done after step 1 (cases A0 and R0), after step 2 (cases A1 and A2) and after step 3 (cases R1 and R2).



Figure 5. Distribution of β during clamping at a load level of 1.27N, causing contractile SMC and endothelial damage.



Figure 6. 31 days evolution of the relative content of all considered constituents in the arterial wall during healing after damage due to clamping at load level 2.



Figure 7. Normalized force versus rod displacement for the six cases. Three different zones are discerned. From 0 to approximately 0.35 mm of displacement, the rod does not touch the sample yet, such that the force is zero. From approximately 0.35 to 0.44 mm, the cylindrical shape of the sample is straightened out. Finally, after approximately 0.44 mm, the sample is further stretched.



Figure 8. Overview of the results of the simulated (red squares) and experimental (boxplot) myograph results. (A) representative scheme of the isometric force measured in the simulated myograph upon addition of vasoactive substances, (B) force increase due to PE addition, (C) and (D) subsequent force decrease after addition of NO and ACh respectively. All forces are normalized with the axial length of the sample. The boxplots show the median values, the 25th and 75th percentiles, the total extent of the measurements without outliers (whiskers) and the outliers (crosses).



Figure 9. Evolution of the relative density of collagen, synthetic SMC and contractile SMC during 91 days of healing for the original remodeling model (A) and two adapted versions (B and C). (A) Original model as described in section 2.4. (B) The mechanical trigger that regulates the number of synthetic cells can only act by increasing the number of cells, and is otherwise deactivated. (C) The production of collagen is only related to the number of synthetic cells, not to the mechanical environment.

Table 2. Overview of parameter values for constituent densities, passive and material properties, prestretch values, damage parameters and remodeling parameters. (1) The exact value from the reference is used. (2) A representative value from the reference is used. (3) The parameter is manually fitted in the same order of mangitude as the reference. (4) The parameter is manually fitted. (5) The parameter is estimated.

Parameter	Value	Source		
Initial densities				
$ ho_0^{elas}$	0.35	Bersi et al. (2016) (2)		
ρ_0^{coll}	0.30	Bersi et al. (2016) (2)		
ρ_0^{csmc}	0.30	Bersi et al. (2016) (2)		
ρ_0^{8smc}	0.05	Bersi et al. (2016) (2)		
Passive material parameters				
C_{10} –	0.04 MPa	Bersi et al. (2016) (2)		
k_1	1.0 MPa	Bersi et al. (2016) (2)		
k_2	1.5	Bersi et al. (2016) (2)		
κ	0.1	(5)		
α	$\pi/8 \text{ rad}$	(5)		
SMC parameters				
μ^{csmc}	0.42 MPa	Murtada et al. (2010) (3)		
κ_c	1.55 MPa	Murtada et al. (2010) (3)		
Prestretches				
g_{ax}	1.67	Bersi et al. (2016) (1)		
g_c	1.1	Bellini et al. (2014) (2)		
Damage parameters				
m^{csmc}	1.0	(4)		
m^{coll}	20.0	(4)		
m^{ec}	0.38	(4)		
Remodeling parameters				
K_{ah}^{coll}	$\log\left(2.0 ight)/100~\mathrm{day}^{-1}$	(4)		
K_m^{coll}	26.64	(4)		
K_{pl}^{smc}	$4.0~\mathrm{day}^{-1}$	(4)		
$\dot{K^{ec}}$	$0.08~\mathrm{day}^{-1}$	(4)		
K_{dd}^{smc}	$1.6 day^{-1}$	(4)		
$K_{ic}^{\widetilde{smc}}$	$0.01~\mathrm{day}^{-1}$	(4)		

Table 3. Overview of parameter values for the SMC contractility model. (1) The exact value from the reference is used. (2) A representative value from the reference is used. (3) The parameter is manually fitted in the same order of mangitude as the reference. (4) The parameter is manually fitted. (5) The parameter is estimated.

Parameter	Value	Value Source		
SMC contractility parameters				
k_3	$0.4 \ { m s}^{-1}$	Hai and Murphy (1988) (1)		
k_4	$0.1 \ { m s}^{-1}$	Hai and Murphy (1988) (1)		
k_7	$0.01 \ { m s}^{-1}$	Hai and Murphy (1988) (1)		
η	60.0 MPa·s	Murtada et al. (2010) (1)		
K_{NO}	8.0e-8 M	(4)		
K_{PE}	2.0e-7 M	(4)		
$[Ca^{2+}]_{hom}$	2.7e-7 M	Utz et al. (1999) (2)		
α_{NO}	1.4e-7 M	(4)		
$lpha_{PE}$	1.28e-7 M	(4)		
$lpha_{Ca}$	0.24	(4)		
K_{CaCaM}	1.78e-7 M	Murtada et al. (2010) (1)		
$k_{2,hom}$	$0.5 \ { m s}^{-1}$	Hai and Murphy (1988) (1)		
<u>α2</u>	$0.1 \ { m s}^{-1}$	(4)		

Table 4. The fraction of overall elastin, collagen, contractile SMC (cSMC), synthetic SMC (sSMC) and endothelium content with respect to their normal content. Note that the level of inflammation is zero in the normal artery wall state, and has a maximal value of 1. Cases A0, A1 and A2 refer to the acute situation after clamping at three load levels (0.0N, 0.6N and 1.27N). R0, R1 and R2 correspond to the respective cases after 31 healing days.

0						
Case	A0	R0	A1	R1	A2	R2
Elastin	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
Collagen	1.0000	1.0000	0.9076	1.6123	0.9071	1.6157
cSMČ	1.0000	1.0000	0.7170	0.6539	0.7190	0.6575
sSMC	1.0000	1.0000	1.0000	1.7225	1.0000	1.7253
Endothelium	1.0000	1.0000	0.3113	0.8463	0.2719	0.8184
Inflammation	0.0000	0.0000	0.6887	0.1537	0.7281	0.1816