



HAL
open science

Reflected BSDEs in non-convex domains

Jean-François Chassagneux, Sergey Nadtochiy, Adrien Richou

► **To cite this version:**

Jean-François Chassagneux, Sergey Nadtochiy, Adrien Richou. Reflected BSDEs in non-convex domains. Probability Theory and Related Fields, In press. hal-03140279

HAL Id: hal-03140279

<https://hal.science/hal-03140279>

Submitted on 12 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

REFLECTED BSDES IN NON-CONVEX DOMAINS

JEAN-FRANÇOIS CHASSAGNEUX, SERGEY NADTOCHIY, ADRIEN RICHOUE

ABSTRACT. This paper establishes the well-posedness of reflected backward stochastic differential equations in the non-convex domains that satisfy a weaker version of the star-shaped property. The main results are established (i) in a Markovian framework with Hölder-continuous generator and terminal condition and (ii) in a general setting under a smallness assumption on the input data. We also investigate the connections between this well-posedness result and the theory of martingales on manifolds.

1. INTRODUCTION

Backward stochastic differential equations (BSDEs), originally introduced in [2] and fully developed in [39, 37], can be viewed as the probabilistic analogues of semi-linear partial differential equations (PDEs). In particular, BSDEs are used to describe the solutions of stochastic control problems (see, among many others, [38, 14, 23]). If the control variable of such an optimization problem has a discrete component – e.g., an option to switch the state process to a different regime or to terminate the process and obtain an instantaneous payoff – then, the associated PDE obtains a free-boundary feature and the associated BSDE becomes reflected: i.e., its solution lives inside a given domain and is reflected at the boundary of this domain. The theory of reflected BSDEs in dimension one, i.e. when the reflected process is one-dimensional, is well developed in a very high generality: see, e.g. [13, 9, 12, 21, 22]. However, the multidimensional case presents significant additional challenges (e.g., due to the lack of the comparison principle), and, to date, the well-posedness of multidimensional reflected BSDEs (or, systems of reflected BSDEs) has only been established in the case of convex reflection domains: see, e.g., [20, 32, 8, 17]. The systems of reflected BSDEs in convex domains appear in certain types of stochastic control problems, such as the switching problems: see, among others, [24, 27, 7, 6, 35, 1]. On the other hand, in a certain class of control-stopping stochastic differential games, the associated equilibria are described by the systems of reflected BSDEs in non-convex domains, as shown, e.g., in [19] (see also [18] for the convex case). We

⁰Authors would like to thank Marc Arnaudon for the enlightening discussions about martingales on manifolds. The authors also thank the Illinois Institute of Technology for hosting the meetings during which this research was initiated. Partial support from the NSF CAREER grant 1855309 is acknowledged.

also refer to [4], which considers another case of a system of reflected BSDEs in a non-convex domain. This paper presents the first general well-posedness result for the systems of reflected BSDEs in non-convex domains under the assumption of a weak star-shape property, see Assumption 1.1 below.

In addition to the stochastic control-stopping games, the reflected BSDEs in non-convex domains have a direct connection to the theory of martingales on manifolds. We refer to [15] for an introduction and an overview of this theory. One of the key questions therein is the following: given a random variable ξ with values in a manifold M , is it possible to define a martingale Y in M such that the terminal value of this martingale (at time $T > 0$) is given by ξ (i.e., $Y_T = \xi$), and is such a martingale unique? A positive answer to this question, in particular, allows one to extend the notion of conditional expectation and gives one possible way to define a barycenter on a manifold (see e.g. [16, 42]). We refer to [30, 31, 41, 10] for other applications. As explained in [10], it is possible to give a positive answer to this question by solving a BSDE with quadratic non-linearities with respect to the z -variable, stated in \mathbb{R}^d – the Euclidean space in which the manifold is embedded. It turns out that for a certain class of non-convex reflection domains \mathcal{D} , the reflected BSDE in \mathcal{D} gives rise to a martingale on the manifold $\partial\mathcal{D}$, see Section 5. In particular, our results provide a new proof of the existence and uniqueness of a martingale with a prescribed terminal value in a given strict sub-sector of a hemisphere of \mathbb{S}^{d-1} , in the Markovian framework or under additional smallness assumptions (see the example in Section 5).

On a technical level, our analysis is connected to the theory of BSDEs with quadratic growth in the z -variable. This connection is made precise in Section 3, but it can also be seen if one attempts to map a given non-convex domain into a convex one – the resulting reflected BSDE in a convex domain will have quadratic terms in z . Thus, the reflected BSDEs in non-convex domains can be viewed as the quadratic reflected BSDEs in convex domains. This observation also explains the additional challenges of the case of a non-convex domain, relative to a convex one: the mathematical difficulties in the former case are similar to those arising in the well-posedness theory for the systems of quadratic BSDEs [43, 28, 44, 25]. The present work uses some of the results developed in the latter theory: in particular, the results of [44] are crucial for our analysis.

Another important connection is to the methods of [33, 34], which establish the well-posedness of the forward (or, regular) stochastic differential equations (SDEs) reflected at the boundary of a given domain. In particular, we use the arguments of [34] to establish the stability of the solutions to the reflected BSDEs considered herein, see Section 2. It is important to mention, however, that many crucial arguments used in the proof of the well-posedness of a reflected (forward) SDE cannot be applied to the case of a reflected BSDE due to the adaptedness issues which, in particular, prohibit the application of the Skorokhod's mapping, used in [34], and of the standard localization methods.

The remainder of this paper is organized as follows. Section 1.1 states the reflected BSDE (equation (1.2)) and the main assumptions (Assumptions 1.1 and 1.2) which hold throughout the paper. Section 2 describes various auxiliary properties and a priori estimates, as well as the stability (Proposition 2.2) and uniqueness (Corollary 2.2) of the solutions to the target reflected BSDE in a certain class. Section 3 describes a sequence of penalized quadratic BSDEs in a Markovian framework, shows that their solutions converge to a solution of the target reflected BSDE, and verifies that this solution belongs to the class in which the uniqueness holds, thus establishing the well-posedness of the target reflected BSDE in a Markovian framework (Theorem 3.1). In Section 4, we approximate a general reflected BSDE by the Markovian ones, to obtain the well-posedness of the former (Theorem 4.2) under an additional smallness assumption (Assumption 2.1). Finally, Section 5 provides a more detailed description of the connection between the reflected BSDEs in non-convex domains and the martingales on manifolds, which, in particular, illustrates the sharpness of some of our assumptions.

1.1. The setup and main assumptions. Let \mathcal{D} be a subset of \mathbb{R}^d given by

$$\mathcal{D} = \{y \in \mathbb{R}^d : \phi(y) < 0\},$$

with a function $\phi : \mathbb{R}^d \rightarrow \mathbb{R}$. We denote by ∇ the gradient, and by ∇^2 the Hessian, of a given function. For any subset A of an Euclidean space, we denote its closure by \bar{A} and, when $A \neq \emptyset$, we denote by $d(\cdot, A)$ the distance function to A .

Assumption 1.1. *We assume that ϕ satisfies the following:*

- (Compactness) *There exists $R > 0$, s.t. $\phi(y) > 0$ for all $|y| \geq R$.*
- (Smoothness) *$\phi \in C^2(\mathbb{R}^d)$, $|\nabla\phi(y)| > 0$ for all $y \in \partial\mathcal{D}$, and $\nabla^2\phi$ is locally Lipschitz.*
- (Weak star-shape property) *There exists a non-empty open convex set $\mathcal{C} \subset \mathcal{D}$ such that*
 - $0 \in \mathcal{C}$,
 - *there exists a convex function $\phi_{\mathcal{C}} : \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying $\phi_{\mathcal{C}} \in C^2(\mathbb{R}^d)$,*

$$\mathcal{C} = \{y \in \mathbb{R}^d : \phi_{\mathcal{C}}(y) < 0\},$$

$\phi_{\mathcal{C}} \geq \phi_{\mathcal{C}}(0)$ and $\phi_{\mathcal{C}}(y) = |y - \mathfrak{P}_{\bar{\mathcal{C}}}(y)|$ for all $y \in \mathbb{R}^d \setminus \mathcal{C}$ where $\mathfrak{P}_{\bar{\mathcal{C}}}$ stands for the projection function onto $\bar{\mathcal{C}}$,

- *it holds that*

$$(1.1) \quad \gamma := \inf_{y \in \partial\mathcal{D}} \nabla\phi_{\mathcal{C}}(y) \cdot \frac{\nabla\phi(y)}{|\nabla\phi(y)|} > 0.$$

Remark 1.1. *If \mathcal{D} is a strictly star-shaped domain w.r.t. 0, i.e., if it satisfies*

$$\inf_{y \in \partial\mathcal{D}} \frac{y}{|y|} \cdot \frac{\nabla\phi(y)}{|\nabla\phi(y)|} > 0,$$

then the weak star-shape property is also satisfied, with \mathcal{C} being a ball of radius $\varepsilon > 0$ centered at 0, and with

$$\phi_{\mathcal{C}}(y) = \varrho_{\varepsilon}(|y| - \varepsilon),$$

where $\varrho_{\varepsilon} : \mathbb{R} \rightarrow \mathbb{R}$ is a convex increasing function satisfying $\varrho_{\varepsilon} \in C^2(\mathbb{R}, \mathbb{R})$, $\varrho_{\varepsilon}(x) = -\varepsilon/2$ for $x < -\varepsilon$ and $\varrho_{\varepsilon}(x) = x$ for $x > 0$.

All stochastic processes and random variables, appearing in this paper, are constructed on a fixed stochastic basis $(\Omega, \mathbb{F}, \mathbb{P})$, with the filtration \mathbb{F} being a completion of the natural filtration of a multidimensional Brownian motion W in \mathbb{R}^d on a time interval $[0, T]$.

For $p \geq 1$, we denote by \mathcal{L}^p the space of (classes of equivalence of)¹ \mathcal{F}_T -measurable random variables ξ (with values in a Euclidean space), s.t. $\|\xi\|_{\mathcal{L}^p} := \mathbb{E}[|\xi|^p]^{1/p} < \infty$. The space \mathcal{L}^{∞} stands for all \mathcal{F}_T -measurable essentially bounded random variables. We also define \mathcal{H}^2 as the space of progressively measurable processes (with values in a Euclidean space) Z , s.t. $\|Z\|_{\mathcal{H}^2} := \mathbb{E}\left[\int_0^T |Z_t|^2 dt\right]^{1/2} < \infty$. Next, for $p \geq 1$, we define \mathcal{M}^p as the space of all continuous local martingales M with $\|M\|_{\mathcal{M}^p} := \mathbb{E}\left[\langle M \rangle_T^{p/2}\right]^{1/p} < \infty$. For $p \in [1, \infty]$, we denote by \mathcal{S}^p the set of continuous adapted process U such that $\|\sup_{t \in [0, T]} |U_t|\|_{\mathcal{L}^p} < \infty$. We also denote by $\text{Var}_t(K)$ the variation of a process K (with values in a Euclidean space) on the time interval $[0, t]$ and by \mathcal{K}^p , for $p \in [1, \infty]$, the set of finite-variation process K such that $\|\text{Var}_{[0, T]}(K)\|_{\mathcal{L}^p} < \infty$ and $K_0 = 0$. Finally, we denote by \mathcal{B}^2 the set of processes $V \in \mathcal{H}^2$, satisfying

$$\|V\|_{\mathcal{B}^2} := \left\| \sup_{t \in [0, T]} \mathbb{E} \left[\int_t^T |V_s|^2 ds \middle| \mathcal{F}_t \right] \right\|_{\mathcal{L}^{\infty}}^{1/2} < +\infty.$$

Let us remark that $V \in \mathcal{B}^2$ implies that the martingale $\int_0^\cdot V_s dW_s$ is a BMO martingale, and $\|V\|_{\mathcal{B}^2}$ is the BMO norm of $\int_0^\cdot V_s dW_s$. We refer to [29] for further details about BMO martingales.

We are investigating the well-posedness of the following reflected BSDE $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$

$$(1.2) \quad \begin{cases} (i) Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T dK_s - \int_t^T Z_s dW_s, & 0 \leq t \leq T, \\ (ii) Y_t \in \bar{\mathcal{D}} \text{ a.s.}, & K_t = \int_0^t \mathbf{n}(Y_s) d\text{Var}_s(K), & 0 \leq t \leq T, \end{cases}$$

¹We drop this clarification in further definitions.

where \mathbf{n} is the unit outward normal to $\partial\mathcal{D}$, extended as zero into \mathcal{D} :

$$\mathbf{n}(y) = \frac{\nabla\phi(y)}{|\nabla\phi(y)|}, \quad \forall y \in \partial\mathcal{D} \quad \text{and} \quad \mathbf{n}(y) = 0, \quad \forall y \in \mathcal{D}.$$

Assumption 1.2. *We assume that ξ takes values in $\bar{\mathcal{D}}$, $f(\cdot, y, z)$ is progressively measurable, $f(t, \cdot, \cdot)$ is globally Lipschitz ($K_{f,y}$ -Lipschitz in y and $K_{f,z}$ -Lipschitz in z), uniformly in (t, ω) , and $\|f(\cdot, 0, 0)\|_{\mathcal{L}^\infty} < \infty$. In addition, w.l.o.g. (in view of the boundedness of \mathcal{D}), there exists a compact $\mathcal{K} \subset \mathbb{R}^d$, s.t. $f(t, y, z) = 0$ whenever $y \notin \mathcal{K}$.*

Assumptions 1.1 and 1.2 hold throughout the rest of the paper even if not cited explicitly.

2. GEOMETRIC PROPERTIES AND A PRIORI ESTIMATES

In this section, we derive useful geometric properties of the domain \mathcal{D} , expressed via the corresponding properties of the function ϕ , and construct an auxiliary function ψ which is used in the next section to define a sequence of approximating equations to (1.2). We also present some key *a priori* estimates and properties of the solutions to the RBSDEs (1.2).

2.1. Absolute continuity of the process K . As noticed in [20], we can take advantage of the smoothness of \mathcal{D} to show that the process K is absolutely continuous with respect to the Lebesgue measure.

Lemma 2.1. *Assume that $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ solves (1.2). Then, almost every path of K is absolutely continuous with respect to the Lebesgue measure.*

Proof. Applying Itô's formula to $t \mapsto \phi(Y_t)$, we obtain

$$(2.1) \quad \begin{aligned} d\phi(Y_t) &= \left(-\nabla\phi(Y_t) \cdot f(t, Y_t, Z_t) + \frac{1}{2} \text{Tr}[Z_t^\top \nabla^2\phi(Y_t) Z_t] \right) dt \\ &+ \nabla\phi(Y_t) \cdot dK_t + \nabla\phi(Y_t) \cdot Z_t dW_t \end{aligned}$$

Then, the Itô-Tanaka formula applied to the positive part of the semi-martingale $-\phi(Y_t)$ reads

$$(2.2) \quad d[-\phi(Y_t)]^+ = \mathbf{1}_{\{-\phi(Y_t) > 0\}} d[-\phi(Y_t)] + \frac{1}{2} dL_t^0,$$

where L^0 is the local time of the semi-martingale $-\phi(Y)$ at zero. Since $\phi(Y_t) \leq 0$, we have $d[-\phi(Y_t)]^+ = -d\phi(Y_t)$ which yields, combining (2.1)–(2.2),

$$\begin{aligned} &\mathbf{1}_{\{\phi(Y_t)=0\}} \left(-\nabla\phi(Y_t) \cdot f(t, Y_t, Z_t) + \frac{1}{2} \text{Tr}[Z_t^\top \nabla^2\phi(Y_t) Z_t] \right) dt + |\nabla\phi(Y_t)| d\text{Var}_t(K) \\ &+ \mathbf{1}_{\{\phi(Y_t)=0\}} \nabla\phi(Y_t) \cdot Z_t dW_t + \frac{1}{2} dL_t^0 = 0. \end{aligned}$$

In particular, we deduce that

$$(2.3) \quad |\nabla\phi(Y_t)|d\text{Var}_t(K) \leq \mathbf{1}_{\{\phi(Y_t)=0\}} \left[\nabla\phi(Y_t) \cdot f(t, Y_t, Z_t) - \frac{1}{2}\text{Tr}[Z_t^\top \nabla^2\phi(Y_t)Z_t] \right]^+ dt,$$

which proves the absolute continuity of K . \square

2.2. The exterior sphere property. The following lemma states the well known observation that, for any boundary point of a smooth domain, there exists a small enough tangent external sphere, see e.g. [34].

Lemma 2.2. *There exists $R_0 > 0$, s.t.*

$$(2.4) \quad (y - y') \cdot \mathbf{n}(y) + \frac{1}{2R_0}|y - y'|^2 \geq 0, \quad \forall y \in \partial\mathcal{D}, y' \in \bar{\mathcal{D}}.$$

Proof. Due to the smoothness of ϕ , for any $y \in \partial\mathcal{D}$ and $y' \in \bar{\mathcal{D}}$, there exists $\lambda \in [0, 1]$, s.t.

$$(2.5) \quad 0 \geq \phi(y') = \phi(y) + (y' - y) \cdot \mathbf{n}(y)|\nabla\phi(y)| + \frac{1}{2}(y - y')^\top \nabla^2\phi(\lambda y + (1 - \lambda)y')(y - y'),$$

It only remains to notice that: $\phi = 0$ and $|\nabla\phi|$ is bounded away from zero on $\partial\mathcal{D}$, and $|\nabla^2\phi|$ is bounded from above on $\bar{\mathcal{D}}$. Thus, we obtain the statement of the lemma. \square

Using the above lemma, we can define the projection operator that is used in the subsequent sections. To this end, we first define the set

$$Q = \{y \in \mathbb{R}^d : d(y, \mathcal{D}) < R_0\},$$

and the set-valued projection operator

$$\mathfrak{P}(y) = \text{argmin}_{x \in \bar{\mathcal{D}}} |x - y|, \quad y \in \mathbb{R}^d.$$

Corollary 2.1. *For any $y \in Q$, $\mathfrak{P}(y)$ is a singleton.*

Proof. It is easy to see that, for a ball $B_r(y) \subset \mathbb{R}^d$, with radius $r > 0$ and center at y , we have:

$$(2.6) \quad (x - x') \cdot \frac{y - x}{|y - x|} + \frac{1}{2r}|x - x'|^2 = 0, \quad \forall x, x' \in \partial B_r(y).$$

Next, assume that there exist $y \in \mathbb{R}^d \setminus \bar{\mathcal{D}}$ and $x \neq x' \in \bar{\mathcal{D}}$, s.t.

$$|x - y| = |x' - y| = \text{argmin}_{z \in \bar{\mathcal{D}}} |z - y|.$$

Then, it is clear that $x, x' \in \partial B_r \cap \partial\mathcal{D}$, with $r = \min_{z \in \bar{\mathcal{D}}} |z - y| < R_0$, and the equations (2.4), (2.6) yield a contradiction. \square

W.l.o.g., we will identify the value of $\mathfrak{P}(y)$ with its only element, for any $y \in Q$.

Remark 2.1. Thanks to (2.5) and to (2.3) we easily deduce for any solution $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ to (1.2) that

$$d\text{Var}_t(K) \leq \mathbf{1}_{\{\phi(Y_t)=0\}} \left(\left[\frac{\nabla\phi(Y_t)}{|\nabla\phi(Y_t)|} \cdot f(t, Y_t, Z_t) \right]^+ + \frac{1}{2R_0} |Z_t|^2 \right) dt,$$

with R_0 satisfying (2.4).

The following lemma provides an alternative to (1.2)(ii), and it becomes useful in the subsequent sections.

Lemma 2.3. Assume that $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ solves (1.2)(i) and $Y_t \in \bar{\mathcal{D}}$ a.s. for all $t \in [0, T]$. Then

$$K_t = \int_0^t \mathbf{n}(Y_s) d\text{Var}_s(K), \quad t \in [0, T],$$

holds if and only if there exists a constant $c > 0$, depending only on \mathcal{D} , such that for all essentially bounded continuous adapted process V in \mathcal{D} , we have

$$(2.7) \quad \int_0^T ((Y_s - V_s) + c|Y_s - V_s|^2 \mathbf{n}(Y_t)) dK_s \geq 0.$$

Proof. One implication is a direct consequence of Lemma 2.2. The other implication is a mere generalization of Lemma 2.1 in [20]. \square

2.3. The pseudo-distance function. In this subsection, we modify the function ϕ in order to construct a new smooth function ψ which satisfies the inequality (1.1) in $\mathbb{R}^d \setminus \mathcal{D}$ instead of $\partial\mathcal{D}$. We denote by $\vartheta : \mathbb{R} \rightarrow [0, 1]$ an infinitely smooth nondecreasing function which is equal to zero on $(-\infty, 0]$ and to one on $[1, \infty)$. We also choose a large enough $R > 1$, s.t. $\mathcal{D} \subset B_{R-1}(0)$, and a small enough $\epsilon \in (0, 1)$, s.t., for all $y \in B_{R+1}(0) \setminus \mathcal{D}$, we have:

$$\phi(y) \leq \epsilon \quad \Rightarrow \quad y \in B_R(0), \quad \nabla\phi_{\mathcal{C}}(y) \cdot \nabla\phi(y) > 0.$$

Then, we define

$$(2.8) \quad \tilde{\phi}(y) := \phi^+(y)(1 - \vartheta(|y| - R - 1)) + \vartheta(|y| - R), \quad \psi(y) := \tilde{\phi}(y) + \kappa|y|\vartheta(\tilde{\phi}(y)/\epsilon),$$

for an arbitrary constant $\kappa > 0$.

We refer to ψ as the pseudo-distance function.

Notice that

$$\begin{aligned} \nabla\phi_{\mathcal{C}}(y) \cdot \nabla\psi(y) &= \nabla\phi_{\mathcal{C}}(y) \cdot \nabla\tilde{\phi}(y) + \kappa\nabla\phi_{\mathcal{C}}(y) \cdot \frac{y}{|y|}\vartheta(\tilde{\phi}(y)/\epsilon) \\ &\quad + \kappa\nabla\phi_{\mathcal{C}}(y) \cdot \nabla\tilde{\phi}(y)|y|\vartheta'(\tilde{\phi}(y)/\epsilon)/\epsilon \\ &= \nabla\phi_{\mathcal{C}}(y) \cdot \nabla\tilde{\phi}(y) \left(1 + \kappa|y|\vartheta'(\tilde{\phi}(y)/\epsilon)/\epsilon \right) + \kappa\nabla\phi_{\mathcal{C}}(y) \cdot \frac{y}{|y|}\vartheta(\tilde{\phi}(y)/\epsilon). \end{aligned}$$

It is clear that $\psi \in C^2(\mathbb{R}^d \setminus \bar{\mathcal{D}})$ and that its derivatives up to the second order are locally Lipschitz-continuous on $\mathbb{R}^d \setminus \mathcal{D}$. It is also easy to see that $\tilde{\phi}(y) \in (0, \epsilon]$ if and only if $y \in B_{R+1}(0) \setminus \mathcal{D}$ and $\phi(y) \leq \epsilon$, in which case $y \in B_R(0)$, $\tilde{\phi}(y) = \phi(y)$, $\nabla \tilde{\phi}(y) = \nabla \phi(y)$, and

$$\nabla \phi_{\mathcal{C}}(y) \cdot \nabla \psi(y) \geq \nabla \phi_{\mathcal{C}}(y) \cdot \nabla \phi(y) > 0,$$

where we also observed that $\inf_{y \in \mathbb{R}^d \setminus \mathcal{D}} \nabla \phi_{\mathcal{C}}(y) \cdot y/|y| > 0$, which follows from the convexity of \mathcal{C} and from the fact that $0 \in \mathcal{C}$. If $\tilde{\phi}(y) \leq 0$, then $y \in \bar{\mathcal{D}}$. If $\tilde{\phi}(y) > \epsilon$, then

$$\nabla \phi_{\mathcal{C}}(y) \cdot \nabla \psi(y) = \nabla \phi_{\mathcal{C}}(y) \cdot \nabla \tilde{\phi}(y) + \kappa \nabla \phi_{\mathcal{C}}(y) \cdot \frac{y}{|y|},$$

which can be made positive for all $y \in \mathbb{R}^d \setminus \mathcal{D}$ by choosing large enough $\kappa > 0$, as $|\nabla \tilde{\phi}|$ is bounded on $\mathbb{R}^d \setminus \mathcal{D}$ and $\inf_{y \in \mathbb{R}^d \setminus \mathcal{D}} \nabla \phi_{\mathcal{C}}(y) \cdot y/|y| > 0$.

The following lemma summarizes the above properties of ψ and states several additional properties which can be easily verified.

Lemma 2.4. *There exist constants $R, \epsilon, \kappa > 0$, s.t. the function ψ defined in (2.8) satisfies the following properties.*

- (1) ψ is globally Lipschitz-continuous in \mathbb{R}^d .
- (2) There exist constants $c, C > 0$, s.t. $c d(y, \mathcal{D}) \leq \psi(y) \leq C d(y, \mathcal{D})$ for $y \in \mathbb{R}^d$.
- (3) $\psi \in C^2(\mathbb{R}^d \setminus \bar{\mathcal{D}})$, and its derivatives up to the second order are globally Lipschitz-continuous in $\mathbb{R}^d \setminus \mathcal{D}$.
- (4) $\inf_{y \in \mathbb{R}^d \setminus \mathcal{D}} \nabla \phi_{\mathcal{C}}(y) \cdot \nabla \psi(y) > 0$.
- (5) $\inf_{y \in \mathbb{R}^d \setminus \mathcal{D}} |\nabla \psi(y)| > 0$.
- (6) $\psi(y) = \phi(y) = 0$, $\nabla \psi(y) = \nabla \phi(y)$, and $\nabla^2 \psi(y) = \nabla^2 \phi(y)$, for $y \in \partial \mathcal{D}$.

In the remainder of the paper, we fix (R, ϵ, κ) as in the above lemma and consider the associated pseudo-distance function ψ . For convenience, we also extend the vector-valued function \mathbf{n} to \mathbb{R}^d as follows:

$$\mathbf{n}(y) = \frac{1}{|\nabla \psi(y)|} \nabla \psi(y) \mathbf{1}_{\{\mathbb{R}^d \setminus \mathcal{D}\}}(y).$$

2.3.1. *Asymptotic convexity of the squared pseudo-distance.* Due to Lemma 2.4, the Hessian of ψ^2 , denoted $\nabla^2 \psi^2$, is well defined in $\mathbb{R}^d \setminus \mathcal{D}$ (it is extended to the boundary of the latter set by continuity). The following lemma shows that $\nabla^2 \psi^2$, viewed as a bilinear form, becomes positive semidefinite close to \mathcal{D} .

Lemma 2.5. *There exists a constant $C > 0$, s.t., for all $y \in \mathbb{R}^d \setminus \mathcal{D}$ and $z \in \mathbb{R}^d$,*

$$z^\top \nabla^2 \psi^2(y) z \geq -C \psi(y) |z|^2.$$

Proof. Notice that, for $y \in \mathbb{R}^d \setminus \mathcal{D}$ and $z \in \mathbb{R}^d$,

$$\nabla^2 \psi^2(y) = 2 \nabla \psi(y) \nabla^\top \psi(y) + 2 \psi(y) \nabla^2 \psi(y),$$

$$z^\top \nabla^2 \psi^2(y) z = 2(\nabla \psi(y) \cdot z)^2 + 2\psi(y) z^\top \nabla^2 \psi(y) z \geq 2\psi(y) z^\top \nabla^2 \psi(y) z.$$

Using the fact that $\nabla^2 \psi$ is bounded (cf. the third property in Lemma 2.4) and the second property in Lemma 2.4, we complete the proof. \square

2.4. *A priori* estimates. In this subsection, we prove *a priori* estimates in the case of general terminal condition ξ and generator f . We first introduce the appropriate “smallness assumption”.

Assumption 2.1. *We assume that at least one of the following four conditions is fulfilled with some $\theta \geq 1$:*

- (i) $|\phi_{\mathcal{C}}^+(\xi)|_{\mathcal{L}^\infty} < \frac{\gamma R_0}{\theta}$ and $\nabla \phi_{\mathcal{C}}(y) \cdot f(s, y, z) \leq 0$, $\forall s, y, z \in [0, T] \times \bar{\mathcal{D}} \setminus \mathcal{C} \times \mathbb{R}^{d \times d'}$,
- (ii) or $\sup_{x \in \mathcal{D}} \phi_{\mathcal{C}}^+(x) < \frac{\gamma R_0}{\theta}$,
- (iii) or \mathcal{C} is the Euclidean ball centered at 0 with radius $\lambda > 0$, and

$$|\xi|_{\mathcal{L}^\infty}^2 < \lambda^2 + \frac{2R_0^2}{\theta}, \quad \nabla \phi_{\mathcal{C}}(y) \cdot f(s, y, z) \leq 0, \quad \forall s, y, z \in [0, T] \times \bar{\mathcal{D}} \setminus \mathcal{C} \times \mathbb{R}^{d \times d'},$$

- (iv) or \mathcal{C} is the Euclidean ball centered at 0 with radius $\lambda > 0$, and

$$\sup_{x \in \mathcal{D}} |x|^2 < \lambda^2 + \frac{2R_0^2}{\theta},$$

with R_0 satisfying (2.4) and γ appearing in Assumption 1.1.

It is worth mentioning that Assumption 2.1 is not our standing assumption and is cited explicitly whenever it is invoked. In particular, our well-posedness results in the Markovian framework do not require the smallness assumption, see Section 3.

Next, we consider the following class of solution:

Definition 2.1. *For any $\theta \geq 1$, we denote by $\mathcal{U}(\theta, \xi, f, T)$ the set of all solutions $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ to (1.2) s.t.*

$$(2.9) \quad \mathbb{E} \left[e^{\frac{\theta p}{R_0} \text{Var}_T(K)} \right] < \infty,$$

with some $p > 1$ and with R_0 satisfying (2.4).

In the sequel, we will generally drop (ξ, f, T) in the notation for the class \mathcal{U} . Note also that we will mainly consider $\theta = 1$ or $\theta = 2$.

The following proposition clarifies the link between Assumption 2.1 and the class $\mathcal{U}(\theta)$.

Proposition 2.1. *Let $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ be a solution to the RBSDE (1.2). Then, $Z \in \mathcal{B}^2$. Moreover, if Assumption 2.1 holds for some $\theta \geq 1$, then, there exist constants C and $p > 1$, which depend only on $K_{f,y}$, $K_{f,z}$, γ ,*

λ , $\sup_{y \in \mathcal{D}} |y|$, $\sup_{y \in \bar{\mathcal{D}}} \phi_{\mathcal{C}}^+(y)$, $\|\phi_{\mathcal{C}}^+(\xi)\|_{\mathcal{L}^\infty}$, $\|f(\cdot, 0, 0)\|_{\mathcal{L}^\infty}$ and R_0 (recall Assumption 1.1 and (2.4)), such that

$$(2.10) \quad \mathbb{E} \left[e^{\frac{\theta p}{R_0} \text{Var}_T(K)} \right] \leq C.$$

Thus, under Assumption 2.1, any solution (Y, Z, K) belongs to $\mathcal{U}(\theta, \xi, f, T)$.

Proof. 1. We start by applying Itô-Tanaka's formula to $\phi_{\mathcal{C}}^+(Y_t)$ (note that $\phi_{\mathcal{C}}^+$ is convex): for all $t \leq t'$,

$$(2.11) \quad \mathbb{E}_t \left[\int_t^{t'} \nabla \phi_{\mathcal{C}}^+(Y_s) \cdot dK_s \right] \leq \mathbb{E}_t \left[\phi_{\mathcal{C}}^+(Y_{t'}) + \int_t^{t'} \nabla \phi_{\mathcal{C}}^+(Y_s) \cdot f(s, Y_s, Z_s) ds \right].$$

In the equation above and the proofs below, we use the shorter notation $\mathbb{E}_t[\cdot]$ for $\mathbb{E}[\cdot | \mathcal{F}_t]$.

Recalling Assumption 1.1, we also have

$$(2.12) \quad \gamma \mathbb{E}_t \left[\int_t^{t'} d\text{Var}_s(K) \right] \leq \mathbb{E}_t \left[\int_t^{t'} \nabla \phi_{\mathcal{C}}^+(Y_s) \cdot \mathbf{n}(Y_s) d\text{Var}_s(K) \right] = \mathbb{E}_t \left[\int_t^{t'} \nabla \phi_{\mathcal{C}}^+(Y_s) \cdot dK_s \right].$$

This yields, for $t' = T$, that

$$(2.13) \quad \gamma \mathbb{E}_t \left[\int_t^T d\text{Var}_s(K) \right] \leq \mathbb{E}_t \left[\phi_{\mathcal{C}}^+(\xi) + \int_t^T \nabla \phi_{\mathcal{C}}^+(Y_s) \cdot f(s, Y_s, Z_s) ds \right].$$

Next, we consider an arbitrary $\varepsilon > 0$ and apply Itô's formula to $\varepsilon |Y_t|^2$ between t and t' , to obtain:

$$\begin{aligned} \varepsilon \mathbb{E}_t \left[\int_t^{t'} |Z_s|^2 ds \right] &\leq \varepsilon \mathbb{E}_t [|Y_{t'}|^2] + C_1 \varepsilon \mathbb{E}_t \left[\int_t^{t'} (1 + |f(s, 0, 0)| + |Z_s|) ds \right] \\ &\quad + \varepsilon C_1 \mathbb{E} \left[\int_t^{t'} d\text{Var}_s(K) \right], \end{aligned}$$

where we used that $|Y|$ is bounded. The above inequality implies

$$(2.14) \quad \frac{\varepsilon}{2} \mathbb{E}_t \left[\int_t^{t'} |Z_s|^2 ds \right] \leq \varepsilon \mathbb{E}_t [|Y_{t'}|^2] + C_\varepsilon \mathbb{E}_t \left[\int_t^{t'} (1 + |f(s, 0, 0)|) ds \right] + \varepsilon C_1 \mathbb{E} \left[\int_t^{t'} d\text{Var}_s(K) \right].$$

Setting $t' = T$ and $\varepsilon = 1$ in the previous inequality and combining it with (2.13), we obtain

$$(2.15) \quad \begin{aligned} \frac{1}{2} \mathbb{E}_t \left[\int_t^T |Z_s|^2 ds \right] &\leq \mathbb{E}_t \left[|\xi|^2 + \frac{C_1}{\gamma} \phi_{\mathcal{C}}^+(\xi) \right] + C \mathbb{E}_t \left[\int_t^T (1 + |f(s, 0, 0)|) ds \right] \\ &+ \frac{C}{\gamma} \mathbb{E} \left[\int_t^T \nabla \phi_{\mathcal{C}}^+(Y_s) \cdot f(s, Y_s, Z_s) ds \right]. \end{aligned}$$

Now, we observe that

$$\frac{C_1}{\gamma} \mathbb{E} \left[\int_t^T \nabla \phi_{\mathcal{C}}^+(Y_s) \cdot f(s, Y_s, Z_s) ds \right] \leq C \mathbb{E}_t \left[\int_t^{t'} (1 + |f(s, 0, 0)|) ds \right] + \frac{1}{4} \mathbb{E}_t \left[\int_t^T |Z_s|^2 ds \right]$$

Inserting the previous estimate back into (2.15), we get

$$\frac{1}{4} \mathbb{E}_t \left[\int_t^T |Z_s|^2 ds \right] \leq \mathbb{E}_t \left[|\xi|^2 + \frac{C}{\gamma} \phi_{\mathcal{C}}^+(\xi) \right] + C \mathbb{E}_t \left[\int_t^T (1 + |f(s, 0, 0)|) ds \right].$$

This proves that $Z \in \mathcal{B}^2$.

2. We now turn to the estimation of the exponential moments of $\text{Var}_T(K)$, under the smallness Assumption 2.1.

2.a First, combining Assumption 2.1(i) with (2.13), we obtain

$$(2.16) \quad \frac{\theta}{R_0} \left\| \sup_{t \in [0, T]} \mathbb{E}_t \left[\int_t^T d\text{Var}_s(K) \right] \right\|_{\mathcal{L}^\infty} < 1.$$

Then, we apply the energy inequalities for non-decreasing processes with bounded potential (see, e.g., (105.1)-(105.2) in [11]) to obtain (2.10) in this case.

2.b Let now Assumption 2.1-(ii) hold. Using (2.11)-(2.12) and recalling that $|Y|$ is bounded, we obtain, for all $0 \leq t < t' \leq T$ and for any $\varepsilon > 0$,

$$\gamma \mathbb{E}_t \left[\int_t^{t'} d\text{Var}_s(K) \right] \leq \sup_{y \in \mathcal{D}} \phi_{\mathcal{C}}^+(y) + C'_\varepsilon (t' - t) (1 + |f(\cdot, 0, 0)|_{\mathcal{L}^\infty}) + \frac{\varepsilon}{2} \mathbb{E}_t \left[\int_t^{t'} |Z_s|^2 ds \right].$$

Using the above inequality and (2.14) (with the same $\varepsilon > 0$), we obtain:

$$(\gamma - C\varepsilon) \mathbb{E}_t \left[\int_t^{t'} d\text{Var}_s(K) \right] \leq \varepsilon \sup_{y \in \mathcal{D}} |y|^2 + \sup_{y \in \mathcal{D}} \phi_{\mathcal{C}}^+(y) + C''_\varepsilon (t' - t) (1 + |f(\cdot, 0, 0)|_{\mathcal{L}^\infty}).$$

In particular, by taking ε small enough, we conclude that, for any $\varepsilon' > 0$, there exists $C_{\varepsilon'} > 0$ such that

$$(2.17) \quad \mathbb{E}_t \left[\int_t^{t'} d\text{Var}_s(K) \right] \leq \frac{\sup_{y \in \mathcal{D}} \phi_{\mathcal{C}}^+(y)}{\gamma} (1 + \varepsilon') + C_{\varepsilon'} (t' - t).$$

Next, using (2.17) and Assumption 2.1(ii), we conclude that there exist $0 < \varepsilon'' < 1$, $p > 1$, and $N \geq 1$, depending only on $K_{f,y}$, $K_{f,z}$, γ , $\sup_{y \in \mathcal{D}} |y|$, $\sup_{y \in \bar{\mathcal{D}}} \phi_{\mathcal{C}}(y)^+$, $\|f(\cdot, 0, 0)\|_{\mathcal{L}^\infty}$ and R_0 , such that, a.s.:

$$(2.18) \quad \mathbb{E}_t \left[\int_t^{T(k+1)/N} d\text{Var}_s(K) \right] \leq \frac{R_0}{\theta p} (1 - \varepsilon''), \quad \forall 0 \leq k < N, \quad \forall t \in [Tk/N, T(k+1)/N].$$

Then, we apply the energy inequalities for non-decreasing processes with bounded potential (see, e.g., (105.1)-(105.2) in [11]), to obtain, for all $0 \leq k < N$,

$$(2.19) \quad \mathbb{E}_{Tk/N} \left[e^{\frac{\theta p}{R_0} \int_{Tk/N}^{T(k+1)/N} d\text{Var}_s(K)} \right] \leq \tilde{C},$$

with \tilde{C} that depends only on $K_{f,y}$, $K_{f,z}$, γ , $\sup_{y \in \mathcal{D}} |y|$, $\sup_{y \in \bar{\mathcal{D}}} \phi_{\mathcal{C}}(y)^+$, $\|f(\cdot, 0, 0)\|_{\mathcal{L}^\infty}$ and R_0 . We now observe that

$$\begin{aligned} \mathbb{E} \left[e^{\frac{\theta p}{R_0} \text{Var}_T(K)} \right] &= \mathbb{E} \left[e^{\frac{\theta p}{R_0} \text{Var}_{T(N-1)/N}(K)} \mathbb{E}_{T(N-1)/N} \left[e^{\frac{\theta p}{R_0} \int_{T(N-1)/N}^T d\text{Var}_s(K)} \right] \right] \\ &\leq \tilde{C} \mathbb{E} \left[e^{\frac{\theta p}{R_0} \text{Var}_{T(N-1)/N}(K)} \right], \end{aligned}$$

where we used (2.19) with $k = N - 1$ to obtain the last inequality. Iterating the above procedure concludes the proof of this case.

2.c Next, we let Assumption 2.1(iii) hold. Using (2.3), the linear growth of f , and Young's inequality, we have, for all $\varepsilon > 0$,

$$(2.20) \quad \text{Var}_T(K) \leq C_\varepsilon + \frac{1 + \varepsilon}{2R_0} \int_0^T \mathbf{1}_{\{\phi(Y_t)=0\}} |Z_t|^2 dt.$$

Moreover, we apply Itô-Tanaka formula to $(|Y_s|^2 - \lambda^2)^+$ to obtain, for all $t \leq t'$,

$$(2.21) \quad \begin{aligned} \mathbb{E}_t \left[\int_t^{t'} \mathbf{1}_{\{\phi(Y_s)=0\}} |Z_s|^2 ds \right] &\leq \mathbb{E}_t \left[\int_t^{t'} \mathbf{1}_{\{\phi_{\mathcal{C}}(Y_s)>0\}} |Z_s|^2 ds \right] \\ &\leq \mathbb{E}_t \left[(|Y_{t'}|^2 - \lambda^2)^+ + 2 \int_t^{t'} \mathbf{1}_{\{\phi_{\mathcal{C}}(Y_s)>0\}} |Y_s| \nabla \phi_{\mathcal{C}}(Y_s) \cdot f(s, Y_s, Z_s) ds \right], \end{aligned}$$

where we also recall that $\mathbf{1}_{\{\phi_{\mathcal{C}}(Y_s)>0\}} |Y_s| \nabla \phi_{\mathcal{C}}(Y_s) = \mathbf{1}_{\{\phi_{\mathcal{C}}(Y_s)>0\}} Y_s$ since \mathcal{C} is a Euclidean ball centered at zero. Then, by taking $t' = T$ in (2.21) and using Assumption 2.1(iii), we obtain, for $\varepsilon > 0$ and $p > 1$ small enough,

$$\frac{\theta p(1 + \varepsilon)}{2R_0^2} \left\| \sup_{t \in [0, T]} \mathbb{E}_t \left[\int_t^T \mathbf{1}_{\{\phi(Y_s)=0\}} |Z_s|^2 ds \right] \right\|_{\mathcal{L}^\infty} < 1.$$

It remains to apply the John-Nirenberg inequality for BMO Martingales (see Theorem 2.2 in [29]) and recall (2.20), to conclude that

$$\mathbb{E} \left[e^{\frac{\theta p}{R_0} \text{Var}_T(K)} \right] \leq C_\varepsilon \mathbb{E} \left[e^{\frac{\theta p(1+\varepsilon)}{2R_0^2} \int_0^T \mathbf{1}_{\{\phi(Y_s)=0\}} |Z_s|^2 ds} \right] < +\infty,$$

which yields (2.10).

2.d Finally, the proof in the case of Assumption 2.1(iv) follows from (2.20) and (2.21), by partitioning $[0, T]$ into small time intervals as in step 2.b. To avoid the redundant calculations, we skip the details. \square

2.5. Stability and uniqueness in $\mathcal{U}(\theta)$. Using the a priori estimates established in the previous subsection, we prove the following stability property of the solutions to (1.2).

Proposition 2.2. *Let us consider $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ (resp. $(Y', Z', K') \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$) which solve the RBSDE (1.2) with a domain \mathcal{D} (resp. \mathcal{D}'), with a terminal condition ξ (resp. ξ'), and with a generator f (resp. f'). Assume, moreover, that there exists $p > 1$ such that*

$$(2.22) \quad \kappa := \mathbb{E} \left[e^{\frac{p}{R_0} (\text{Var}_T(K) + \text{Var}_T(K'))} \right] < +\infty,$$

with R_0 satisfying (2.4) for \mathcal{D} and \mathcal{D}' . Let us denote by $\bar{\mathfrak{P}}$ (resp. $\bar{\mathfrak{P}}'$) a measurable selection of the projection operator onto \mathcal{D} (resp. \mathcal{D}'). Then, the following stability result holds: there exists a constant $C > 0$, which depends only on $K_{f,y}$, $K_{f',y}$, $K_{f,z}$, $K_{f',z}$ (recall Assumption 1.2), $\sup_{y \in \mathcal{D} \cup \mathcal{D}'} |y|$, R_0 , and on κ , and is such that

$$\begin{aligned} & \|Y - Y'\|_{\mathcal{S}^2} + \|Z - Z'\|_{\mathcal{H}^2} + \|K - K'\|_{\mathcal{K}^1} \\ & \leq C \mathbb{E} [|\xi - \xi'|^{2p/(p-1)}]^{(p-1)/(2p)} + C \mathbb{E} \left[\left(\int_0^T |f(s, Y_s, Z_s) - f'(s, Y_s, Z_s)| ds \right)^{2p/(p-1)} \right]^{(p-1)/(2p)} \\ & \quad + C \mathbb{E} \left[\sup_{s \in [0, T]} |Y_s - \bar{\mathfrak{P}}'(Y_s)|^{p/(p-1)} \right]^{(p-1)/(2p)} + C \mathbb{E} \left[\sup_{s \in [0, T]} |Y'_s - \bar{\mathfrak{P}}(Y'_s)|^{p/(p-1)} \right]^{(p-1)/(2p)}. \end{aligned}$$

Proof. We apply Itô's formula to the process

$$e^{\beta t + \frac{1}{R_0} (\text{Var}_t(K) + \text{Var}_t(K'))} |Y_t - Y'_t|^2,$$

with the constant β to be determined later on. By denoting

$$\delta f_t := f(t, Y_t, Z_t) - f'(t, Y'_t, Z'_t), \quad \delta \xi := \xi - \xi',$$

$$\Gamma_t := e^{\beta t + \frac{1}{R_0} (\text{Var}_t(K) + \text{Var}_t(K'))}, \quad \delta Y := Y - Y', \quad \delta Z = Z - Z',$$

we obtain

$$\begin{aligned}
& \Gamma_t |\delta Y_t|^2 + \int_t^T \Gamma_s |\delta Z_s|^2 ds \\
&= \Gamma_T |\delta \xi|^2 + 2 \int_t^T \Gamma_s \delta Y_s \cdot \delta f_s ds - 2 \int_t^T \Gamma_s \delta Y_s \cdot dK_s + 2 \int_t^T \Gamma_s \delta Y_s \cdot dK'_s \\
&\quad - \beta \int_t^T \Gamma_s |\delta Y_s|^2 ds - \frac{1}{R_0} \int_t^T \Gamma_s |\delta Y_s|^2 d\text{Var}_s(K) - \frac{1}{R_0} \int_t^T \Gamma_s |\delta Y_s|^2 d\text{Var}_s(K') \\
(2.23) \quad & - 2 \int_t^T \Gamma_s \delta Y_s \cdot \delta Z_s dW_s.
\end{aligned}$$

Using the BDG inequality, the fact that $|\delta Y|$ is bounded, Hölder inequality (with $q = p/(p-1) > 1$ being the conjugate exponent), we obtain:

$$\begin{aligned}
\mathbb{E} \left[\sup_{t \in [0, T]} \left| \int_0^t \Gamma_s \delta Y_s \cdot \delta Z_s dW_s \right| \right] &\leq C \mathbb{E} \left[\left(\int_0^T |\Gamma_s \delta Z_s|^2 ds \right)^{\frac{1}{2}} \right] \\
&\leq C \mathbb{E}[(\Gamma_T)^p]^{\frac{1}{p}} \mathbb{E} \left[\left(\int_0^T |\delta Z_s|^2 ds \right)^{[q/2]} \right]^{\frac{1}{q}} < \infty,
\end{aligned}$$

where the last inequality is due to (2.22) and to the Energy Inequality (since $Z, Z' \in \mathcal{B}^2$). Hence, we conclude that the local martingale term in the right hand side of (2.23) is a true martingale.

Next, we estimate the second term in the right hand side of (2.23) using the Lipschitz property of f' :

$$\delta Y_s \cdot \delta f_s \leq |\delta Y_s| |f(s, Y_s, Z_s) - f'(s, Y_s, Z_s)| + \beta |\delta Y_s|^2 + \frac{1}{4} |\delta Z_s|^2,$$

provided $\beta > 0$ is large enough. In addition, the condition (1.2)(ii) and the exterior sphere property (recall (2.4)) yield

$$\begin{aligned}
& - 2 \int_t^T \Gamma_s \delta Y_s \cdot dK_s - \frac{1}{R_0} \int_t^T \Gamma_s |\delta Y_s|^2 d\text{Var}_s(K) \\
&= - 2 \int_t^T \Gamma_s (\bar{\mathfrak{P}}(Y'_s) - Y'_s) \cdot dK_s - 2 \int_t^T \Gamma_s (Y_s - \bar{\mathfrak{P}}(Y'_s)) \cdot dK_s - \frac{1}{R_0} \int_t^T \Gamma_s |\delta Y_s|^2 d\text{Var}_s(K) \\
&= - 2 \int_t^T \Gamma_s (Y_s - \bar{\mathfrak{P}}(Y'_s)) \cdot dK_s - \frac{1}{R_0} \int_t^T \Gamma_s |Y_s - \bar{\mathfrak{P}}(Y'_s)|^2 d\text{Var}_s(K) \\
&\quad + \frac{1}{R_0} \int_t^T \Gamma_s (|\bar{\mathfrak{P}}(Y'_s) - Y'_s|^2 - |\delta Y_s|^2) d\text{Var}_s(K) - 2 \int_t^T \Gamma_s (\bar{\mathfrak{P}}(Y'_s) - Y'_s) \cdot dK_s \\
&\leq C \int_t^T \Gamma_s |\bar{\mathfrak{P}}(Y'_s) - Y'_s| d\text{Var}_s(K) \leq C \Gamma_T \sup_{s \in [0, T]} |\bar{\mathfrak{P}}(Y'_s) - Y'_s|,
\end{aligned}$$

where for the last inequality, we used: $\int_t^T \exp\left(\frac{\text{Var}_s(K)}{R_0}\right) d\text{Var}_s(K) \leq R_0 \exp\left(\frac{\text{Var}_T(K)}{R_0}\right)$.
By the same arguments we obtain

$$2 \int_t^T \Gamma_s \delta Y_s \cdot dK'_s - \frac{1}{R_0} \int_t^T \Gamma_s |\delta Y_s|^2 d\text{Var}_s(K') \leq C \Gamma_T \sup_{s \in [0, T]} |\bar{\mathfrak{P}}'(Y_s) - Y_s|.$$

Using the above estimates, we take expectations on both sides of 2.23, with $t = 0$, and apply Hölder inequality to obtain

$$\begin{aligned} \|\Gamma^{1/2} \delta Z\|_{\mathcal{H}^2} &\leq \mathbb{E} \left[\Gamma_T |\delta \xi|^2 + 2 \int_0^T \Gamma_s |\delta Y_s| |f(s, Y_s, Z_s) - f'(s, Y_s, Z_s)| ds \right]^{1/2} \\ &\quad + \mathbb{E} \left[\Gamma_T \left(\sup_{s \in [0, T]} |\bar{\mathfrak{P}}'(Y_s) - Y_s| + |\bar{\mathfrak{P}}(Y'_s) - Y'_s| \right) \right]^{1/2} \\ &\leq C \mathbb{E}[|\delta \xi|^{2q}]^{1/(2q)} \\ &\quad + 2 \mathbb{E} \left[\sup_{s \in [0, T]} (\Gamma_s^{1/2} |\delta Y_s|) \Gamma_T^{1/2} \int_0^T |f(s, Y_s, Z_s) - f'(s, Y_s, Z_s)| ds \right]^{1/2} \\ (2.24) \quad &\quad + C \mathbb{E} \left[\sup_{s \in [0, T]} (|\bar{\mathfrak{P}}'(Y_s) - Y_s| + |\bar{\mathfrak{P}}(Y'_s) - Y'_s|)^q \right]^{1/(2q)}. \end{aligned}$$

Using (2.23) and (2.24), we apply BDG, Hölder and Young inequalities to obtain

$$\begin{aligned} \|\Gamma^{1/2} \delta Y\|_{\mathcal{H}^2} &\leq C \mathbb{E}[|\delta \xi|^q]^{1/(2q)} \\ &\quad + C \mathbb{E} \left[\sup_{s \in [0, T]} (\Gamma_s^{1/2} |\delta Y_s|) \Gamma_T^{1/2} \int_0^T |f(s, Y_s, Z_s) - f'(s, Y_s, Z_s)| ds \right]^{1/2} \\ &\quad + C \mathbb{E} \left[\sup_{s \in [0, T]} (|\bar{\mathfrak{P}}'(Y_s) - Y_s| + |\bar{\mathfrak{P}}(Y'_s) - Y'_s|)^q \right]^{1/(2q)} \\ &\leq C \mathbb{E}[|\delta \xi|^q]^{1/(2q)} + \frac{1}{2} \|\Gamma^{1/2} \delta Y\|_{\mathcal{H}^2} + C \mathbb{E} \left[\left(\int_0^T |f(s, Y_s, Z_s) - f'(s, Y_s, Z_s)| ds \right)^{2q} \right]^{1/(2q)} \\ (2.25) \quad &\quad + C \mathbb{E} \left[\sup_{s \in [0, T]} (|\bar{\mathfrak{P}}'(Y_s) - Y_s| + |\bar{\mathfrak{P}}(Y'_s) - Y'_s|)^q \right]^{1/(2q)}. \end{aligned}$$

Then, combining (2.24), Young inequality, and (2.25), yields

$$\begin{aligned}
& \|Y - Y'\|_{\mathcal{S}^2} + \|Z - Z'\|_{\mathcal{H}^2} \leq \|\Gamma^{1/2}\delta Y\|_{\mathcal{S}^2} + \|\Gamma^{1/2}\delta Z\|_{\mathcal{H}^2} \\
& \leq C\mathbb{E}[|\xi - \xi'|^{2q}]^{1/(2q)} + C\mathbb{E}\left[\left(\int_0^T |f(s, Y_s, Z_s) - f'(s, Y_s, Z_s)|ds\right)^{2q}\right]^{1/(2q)} \\
(2.26) \quad & + C\mathbb{E}\left[\sup_{s \in [0, T]} (|\bar{\mathfrak{P}}'(Y_s) - Y_s| + |\bar{\mathfrak{P}}(Y'_s) - Y'_s|)^q\right]^{1/(2q)}.
\end{aligned}$$

Finally, we recall that

$$K_t - K'_t = \delta Y_t - \delta Y_0 + \int_0^t f(s, Y_s, Z_s) - f'(s, Y'_s, Z'_s)ds - \int_0^t \delta Z_s dW_s.$$

Then, the BDG inequality, the Lipschitz property of f' , as well as (2.26), yield

$$\begin{aligned}
\|K - K'\|_{\mathcal{S}^2} & \leq C\mathbb{E}[|\xi - \xi'|^{2q}]^{1/(2q)} + C\mathbb{E}\left[\left(\int_0^T |f(s, Y_s, Z_s) - f'(s, Y_s, Z_s)|ds\right)^{2q}\right]^{1/(2q)} \\
& + C\mathbb{E}\left[\sup_{s \in [0, T]} (|\bar{\mathfrak{P}}'(Y_s) - Y_s| + |\bar{\mathfrak{P}}(Y'_s) - Y'_s|)^q\right]^{1/(2q)},
\end{aligned}$$

which completes the proof of the proposition. \square

In a general non-Markovian framework, we obtain the following uniqueness result as a direct consequence of Proposition 2.2.

Corollary 2.2. *The reflected BSDE (1.2) has at most one solution in the class $\mathcal{U}(2)$.*

Proof. Indeed, it suffices to check that, for any two solutions in the class $\mathcal{U}(2)$, (2.22) holds. This follows directly from the Cauchy-Schwarz inequality. \square

This uniqueness result is improved in the Markovian setting: see Theorem 3.1 and Remark 3.5.

3. WELL-POSEDNESS IN A MARKOVIAN FRAMEWORK

In this section, we establish the existence and uniqueness of the solution to (3.3) under the assumption that the terminal condition and the generator of the reflected BSDE are functions of a Markov diffusion process X in \mathbb{R}^d :

$$(3.1) \quad X_t = x + \int_0^t b(s, X_s)ds + \int_0^t \sigma(s, X_s)dW_s, \quad x \in \mathbb{R}^d.$$

Namely, we make the following assumptions.

Assumption 3.1. *We assume that (b, σ) are bounded measurable functions, uniformly Lipschitz with respect to x , and such that $\sigma^\top \sigma$ is uniformly positive definite (i.e., uniformly elliptic), which implies in particular that σ is invertible.*

Assumption 3.2. *We assume that*

$$\xi := g(X_T) \quad \text{and} \quad f(t, y, z) := F(t, X_t, y, z),$$

where g is α -Hölder and $\bar{\mathcal{D}}$ -valued, F is measurable in all variables, globally Lipschitz in (y, z) , and s.t. $|F(\cdot, \cdot, 0, 0)|$ is bounded.

Recall that Assumptions 1.1 and 1.2 hold throughout the paper, even if they are not cited explicitly.

3.1. Penalized equation. We begin by noticing that $\psi^2 \in C^1(\mathbb{R}^d)$ and denote

$$\Psi(y) := \frac{1}{2} \nabla \psi(y)^2 = \psi(y) \nabla \psi(y), \quad y \in \mathbb{R}^d,$$

where we extend (naturally) $\nabla \psi$ to \mathcal{D} by zero. We also extend $\nabla^2 \psi^2$ to \mathcal{D} by zero.

It is useful to note that there exist constants c, C , s.t.

$$(3.2) \quad 0 < c\psi \leq |\Psi| \leq C\psi.$$

Next, we consider the following penalized equation:

$$(3.3) \quad Y_t^n = \xi + \int_t^T f(s, Y_s^n, Z_s^n) ds - \int_t^T n\Psi(Y_s^n)(1 + |Z_s^n|^2) ds - \int_t^T Z_s^n dW_s.$$

Let us remark that, contrarily to the convex framework tackled in [20], it is natural (and necessary) to add a $|z|^2$ inside the penalization term due to (2.3). For convenience, we introduce the notation:

$$(3.4) \quad \begin{aligned} \Phi_t^n &:= \int_0^t n\Psi(Y_s^n) ds, & \Theta_t^n &:= \int_0^t n\Psi(Y_s^n)|Z_s^n|^2 ds, \\ K_t^n &:= \Phi_t^n + \Theta_t^n. \end{aligned}$$

3.2. Existence of a solution to the penalized equation. We start by considering the following family of approximating BSDEs, indexed by a pair of positive integers $M = (M_1, M_2)$:

$$(3.5) \quad Y_t^{n,M} = g(X_T) + \int_t^T F^{n,M}(s, X_s, Y_s^{n,M}, Z_s^{n,M}) ds - \int_t^T Z_s^{n,M} dW_s,$$

with

$$F^{n,M}(t, x, y, z) := f(t, x, y, z) - n\rho_{M_1}(\psi(y))\nabla\psi(y)(1 + \rho_{M_2}(|z|^2)), \quad \rho_k(x) := x \wedge k.$$

The above BSDE has a globally Lipschitz generator and, therefore, is known to have a unique Markovian solution $(Y^{n,M}, Z^{n,M}) \in \mathcal{S}^2 \times \mathcal{H}^2$ (see, e.g., Theorem

4.1 in [14]). The following Proposition uses the weak star-shape property of \mathcal{D} , stated in Assumption 1.1, to establish a uniform estimate on $(Y^{n,M}, Z^{n,M})$.

Lemma 3.1. *There exists a constant $C > 0$, s.t., for any $n \geq 1$, any $M = (M_1, M_2)$, and any $t \in [0, T]$, the following holds a.s.:*

$$(3.6) \quad |Y_t^{n,M}|^2 + \mathbb{E}_t \left[\int_t^T |Y_s^{n,M}|^2 + |Z_s^{n,M}|^2 ds \right] \leq C \mathbb{E}_t \left[|\xi|^2 + \int_t^T (1 + |f(s, 0, 0)|^2) ds \right],$$

$$(3.7) \quad \mathbb{E}_t \left[\int_t^T n \rho_{M_1}(\psi(Y_s^{n,M})) (1 + \rho_{M_2}(|Z_s^{n,M}|^2)) ds \right] \leq C \mathbb{E}_t \left[|\xi|^2 + \int_t^T (1 + |f(s, 0, 0)|^2) ds \right].$$

Proof. W.l.o.g., we assume that ϕ_C attains its minimum at zero. Then, we consider arbitrary $t \in [0, T]$ and constants $\alpha > 0$, $\beta > 0$, to be fixed later, and define

$$[t, T] \times \mathbb{R}^d \ni (s, y) \mapsto h(s, y) := e^{\beta(s-t)} (\alpha |y|^2 + (\phi_C(y) - \phi_C(0))^2) \in \mathbb{R}.$$

We observe that $(\phi_C - \phi_C(0))^2$ is convex and $h(s, y) \leq e^{\beta(T-s)} c_0 |y|^2$, for some positive constant c_0 . Then, we apply Itô's formula to the process $h(s, Y_s^{n,M})$ (recalling (3.3)), to obtain

$$(3.8) \quad \begin{aligned} \alpha |Y_t^{n,M}|^2 &\leq h(t, Y_t^{n,M}) \leq h(T, \xi) \\ &+ 2 \int_t^T e^{\beta(s-t)} (\alpha Y_s^{n,M} + (\phi_C(Y_s^{n,M}) - \phi_C(0)) \nabla \phi_C(Y_s^{n,M})) \cdot f(s, Y_s^{n,M}, Z_s^{n,M}) ds \\ &- \int_t^T n \rho_{M_1}(\psi(Y_s^{n,M})) \nabla_y h(s, Y_s^{n,M}) \cdot \nabla \psi(Y_s^{n,M}) (1 + \rho_{M_2}(|Z_s^{n,M}|^2)) ds \\ &- 2 \int_t^T \nabla_y h(s, Y_s^{n,M}) \cdot Z_s^{n,M} dW_s - \alpha \int_t^T e^{\beta(s-t)} |Z_s^{n,M}|^2 ds - \beta \int_t^T e^{\beta(s-t)} |Y_s^{n,M}|^2 ds. \end{aligned}$$

As $Y^{n,M} \in \mathcal{S}^2$ and $Z^{n,M} \in \mathcal{H}^2$, the local martingale in the above representation is in \mathcal{M}^1 and, hence, is a true martingale.

Next, we notice that the fourth property in Lemma 2.4 implies the existence of a constant $c_1 > 0$, s.t.

$$\nabla \phi_C(Y_s^{n,M}) \cdot \nabla \psi(Y_s^{n,M}) \geq c_1 \mathbf{1}_{\{Y_s^{n,M} \notin \mathcal{D}\}}.$$

Then, there exist constants $c_2, c_3 > 0$ such that

$$\begin{aligned}
& - \int_t^T n \rho_{M_1}(\psi(Y_s^{n,M})) \nabla_y h(s, Y_s^{n,M}) \cdot \nabla \psi(Y_s^{n,M}) (1 + \rho_{M_2}(|Z_s^{n,M}|^2)) ds \\
\leq & - 2 \int_t^T n e^{\beta(s-t)} \rho_{M_1}(\psi(Y_s^{n,M})) (c_1(\phi_C(Y_s^{n,M}) - \phi_C(0)) - \alpha c_2 |Y_s^{n,M}|) (1 + \rho_{M_2}(|Z_s^{n,M}|^2)) ds \\
(3.9) \quad & \leq - 2 \int_t^T n e^{\beta(s-t)} \rho_{M_1}(\psi(Y_s^{n,M})) [c_1(\phi_C(Y_s^{n,M}) - \phi_C(0)) \\
& \quad - \alpha c_2(\phi_C(Y_s^{n,M}) + |\mathfrak{P}_{\bar{C}}(Y_s^{n,M})|)] (1 + \rho_{M_2}(|Z_s^{n,M}|^2)) ds \\
\leq & - c_3 \int_t^T n e^{\beta(s-t)} \rho_{M_1}(\psi(Y_s^{n,M})) (1 + \rho_{M_2}(|Z_s^{n,M}|^2)) ds,
\end{aligned}$$

provided α is small enough. In the rest of the proof, we assume that α is chosen so that the above inequality holds.

Next, we remark that

$$\begin{aligned}
& \left| 2 \int_t^T e^{\beta(s-t)} (\alpha Y_s^{n,M} + (\phi_C(Y_s^{n,M}) - \phi_C(0)) \nabla \phi_C(Y_s^{n,M})) \cdot f(s, Y_s^{n,M}, Z_s^{n,M}) ds \right| \\
& \leq C_1 \int_t^T e^{\beta(s-t)} ((\alpha + 1) |Y_s^{n,M}| - \phi_C(0)) (|f(s, 0, 0)| + C_2 |Y_s^{n,M}| + C_2 |Z_s^{n,M}|) ds \\
& \leq \int_t^T e^{\beta(s-t)} \left(C_3 |Y_s^{n,M}|^2 + C_3 + |f(s, 0, 0)|^2 + \frac{\alpha}{2} |Z_s^{n,M}|^2 \right) ds.
\end{aligned}$$

Combining the above estimates and (3.8), we conclude that, for a large enough $\beta > 0$, there exists a constant $C_4 > 0$, s.t.

$$\begin{aligned}
\alpha |Y_t^{n,M}|^2 + \mathbb{E}_t \left[c_3 \int_t^T n \rho_{M_1}(\psi(Y_s^{n,M})) (1 + \rho_{M_2}(|Z_s^{n,M}|^2)) ds + \frac{\alpha}{2} \int_t^T |Y_s^{n,M}|^2 + |Z_s^{n,M}|^2 ds \right] \\
\leq e^{\beta(T-t)} \mathbb{E}_t \left[c_0 |\xi|^2 + \int_t^T (C_4 + |f(s, 0, 0)|^2) ds \right],
\end{aligned}$$

which yields the statement of the lemma. \square

Proposition 3.1. *Under Assumptions 3.1 and 3.2, for any $n \geq 1$, the BSDE (3.3) has a Markovian solution (Y^n, Z^n) . In particular, there exists a measurable function u^n such that $Y_t^n = u^n(t, X_t)$. Moreover, the estimates (3.6)–(3.7) hold with $(Y^{n,M}, Z^{n,M})$ and ρ_{M_i} replaced, respectively, by any solution (Y^n, Z^n) of (3.3) and by the identity function.*

Proof. The main statement of the proposition follows from Theorem 2.8 in [44] (without the localization). To be able to apply the latter theorem, we first

consider the following auxiliary BSDE, which can be viewed as a middle ground between (3.3) and (3.5):

$$(3.10) \quad \tilde{Y}_t^{n,M_1} = g(X_T) + \int_t^T \tilde{F}^{n,M_1}(s, X_s, \tilde{Y}_s^{n,M_1}, \tilde{Z}_s^{n,M_1}) ds - \int_t^T \tilde{Z}_s^{n,M_1} dW_s,$$

with

$$\tilde{F}^{n,M_1}(t, x, y, z) := f(t, x, y, z) - n\rho_{M_1}(\psi(y))\nabla\psi(y)(1 + |z|^2)$$

and $\rho_{M_1}(x) = x \wedge M_1$. We claim that the unique solution $(Y^{n,M}, Z^{n,M})$ of (3.5) converges (along a subsequence) to a Markovian solution $(\tilde{Y}^{n,M_1}, \tilde{Z}^{n,M_1})$ of (3.10), as $M_2 \rightarrow \infty$. Indeed, this claim follows directly from Theorem 2.8 in [44]. To verify the assumptions of the latter theorem, we first notice that, due to (3.6), there exists a constant $c > 0$ such that $|Y_t^{n,M}| \leq c$, for all $t \in [0, T]$ and n, M . Moreover, for large enough $C > 0$ (independent of n and M), $h(y) := C(\alpha|y|^2 + (\phi_C(y) - \phi_C(0))^2)$ is a global c -Lyapunov function for $(F^{n,M})_M$, in the sense of Definition 2.3 in [44], where α is the constant chosen in the proof of Lemma 3.1. Indeed, there exists a large enough $C > 0$, s.t., for all $|y| \leq c$, we have:

$$\begin{aligned} & \frac{1}{2}C \operatorname{Tr}[(z\sigma)^\top (\nabla^2 h(y)) z\sigma] - C\nabla h(y) \cdot F^{n,M} \\ & \geq C\alpha \operatorname{Tr}[(z\sigma)^\top z\sigma] - 2C[\alpha y + (\phi_C(y) - \phi_C(0))\nabla\phi_C(y)] \cdot f(t, x, y, z) \\ & \quad + 2Cn[\alpha y + (\phi_C(y) - \phi_C(0))\nabla\phi_C(y)] \cdot \nabla\psi(y)\rho_{M_1}(\psi(y))(1 + \rho_{M_2}(|z|^2)) \\ & \geq |z|^2 - C', \end{aligned}$$

where we used the uniform ellipticity of $\sigma^\top\sigma$, Assumption 1.2, and the fourth property in Lemma 2.4, and repeated the estimates used in (3.9). In addition, we have $|F^{n,M}(t, x, y, z)| \leq C + C_n|z|^2$, with the constants (C, C_n) independent of M_2 . Observing that $F^{n,M}$ converges to \tilde{F}^{n,M_1} locally uniformly, as $M_2 \rightarrow \infty$, we conclude that the assumptions of Theorem 2.8 in [44] are satisfied and that (3.10) has a Markovian solution $(\tilde{Y}^{n,M_1}, \tilde{Z}^{n,M_1})$ which is a limit point of $\{(Y^{n,M}, Z^{n,M})\}_{M_2}$.

Next, we recall that, due to (3.6), $|Y^{n,M}|$ is bounded uniformly over M . Hence, $|\tilde{Y}^{n,M_1}|$ can be bounded uniformly over $M_1 \geq 1$, and, in turn, $(\tilde{Y}^{n,M_1}, \tilde{Z}^{n,M_1})$ solve (3.3) for any large enough $M_1 > 0$.

The estimates (3.6)–(3.7) are obtained by repeating the proof of Lemma 3.1 for the equation (3.3) in place of (3.5). \square

3.3. A priori estimates. The following result relies on the asymptotic convexity of the squared pseudo-distance function, stated in Lemma 2.5.

Lemma 3.2. *Under Assumptions 3.1 and 3.2, there exists a constant $C > 0$, s.t., for any $n \geq 1$, any solution (Y^n, Z^n) of (3.3), and any $t \in [0, T]$, the following*

holds a.s.:

$$\begin{aligned} n\psi^2(Y_t^n) + \mathbb{E}_t \left[\int_t^T n^2 |\Psi(Y_s^n)|^2 (1 + |Z_s^n|^2) ds \right] \\ \leq C \mathbb{E}_t \left[|\xi|^2 + \int_t^T |f(s, 0, 0)|^2 ds \right], \end{aligned}$$

and, in particular,

$$d(Y_t^n, \mathcal{D}) \leq Cn^{-1/2}.$$

Proof. We begin by applying Itô's formula to $|\psi(Y_t^n)|^2$, to obtain

$$\begin{aligned} (3.11) \quad \psi^2(Y_t^n) &= 2 \int_t^T \Psi(Y_s^n) \cdot f(s, Y_s^n, Z_s^n) ds - 2 \int_t^T n |\Psi(Y_s^n)|^2 ds \\ &\quad - 2 \int_t^T n |\Psi(Y_s^n)|^2 |Z_s^n|^2 ds - 2 \int_t^T \Psi(Y_s^n) \cdot Z_s^n dW_s \\ &\quad - \frac{1}{2} \int_t^T \text{Tr}[(Z_s^n)^\top \nabla^2 \psi^2(Y_s^n) Z_s^n] ds \end{aligned}$$

Remark 3.1. Note that the Hessian of ψ^2 has a discontinuity at $\partial\mathcal{D}$. To justify the use of Itô's formula, we approximate ψ^2 by a sequence of C^2 functions $\{g^m\}$, so that g^m , ∇g^m and $\nabla^2 g^m$ converge, respectively, to ψ^2 , $\nabla \psi^2$ and $\nabla^2 \psi^2$ everywhere in \mathbb{R}^d , and $|\nabla g^m|$, $|\nabla^2 g^m|$ are locally bounded uniformly over m . To construct such a sequence, we first define

$$\hat{\phi}(y) := \phi(y)(1 - \vartheta(|y| - R - 1)) + \vartheta(|y| - R), \quad \hat{\psi}(y) := \hat{\phi}(y) + \kappa|y|\vartheta(\hat{\phi}(y)/\epsilon), \quad y \in \mathbb{R}^d,$$

where we recall the original function ϕ , appearing in Assumption 1.1, and use the same ϑ , R , and ϵ , as the ones used in Subsection 2.3 to define ψ (see (2.8)). It is clear that $\hat{\psi}(y) = \psi(y)$, for $y \in \mathbb{R}^d \setminus \mathcal{D}$, and that $\hat{\psi}(y) = \phi(y)$, for $y \in \mathcal{D}$. Thus, $\hat{\psi}$ is a smooth extension of ψ into \mathcal{D} . Next, we consider an infinitely smooth nondecreasing function $\rho: \mathbb{R} \rightarrow \mathbb{R}$, such that $\rho(x) = -1$ for $x \leq -1$ and $\rho(x) = x$ for $x \geq 0$, and define

$$g^m(y) := \frac{1}{m^2} \rho^2 \left(m \hat{\psi}(y) \right), \quad y \in \mathbb{R}^d.$$

It is easy to check by a direct computation that $g^m(y)$, $\nabla g^m(y)$ and $\nabla^2 g^m(y)$ converge to zero as $m \rightarrow \infty$, for any $y \in \mathcal{D}$. On the other hand, $g^m(y)$ and its first two derivatives coincide with $\psi^2(y)$ and with its respective derivatives, for all $y \in \mathbb{R}^d \setminus \mathcal{D}$ and all m . Thus, we obtain the desired sequence $\{g^m\}$. Applying Itô's formula to $g^m(Y_t^n)$ and using the dominated convergence theorem to pass to the limit as $m \rightarrow \infty$, we establish (3.11).

As $|\Psi|$ is linearly bounded (see Lemma 2.4), we conclude, as in the proof of Lemma 3.1, that the local martingale in the above representation is a true martingale.

Next, we note that

$$2\Psi(Y_s^n) \cdot f(s, Y_s^n, Z_s^n) \leq n|\Psi(Y_s^n)|^2 + n^{-1}|f(s, Y_s^n, Z_s^n)|^2,$$

and use Lemma 3.1, to obtain:

$$(3.12) \quad \mathbb{E}_t \int_t^T 2\Psi(Y_s^n) \cdot f(s, Y_s^n, Z_s^n) ds \leq \mathbb{E}_t \int_t^T n|\Psi(Y_s^n)|^2 ds + Cn^{-1} \mathbb{E}_t \left[|\xi|^2 + \int_t^T (1 + |f(s, 0, 0)|^2) ds \right].$$

In addition, Lemmas 2.4 and 2.5 yield

$$\text{Tr}[(Z_s^n)^\top \nabla^2 \psi^2(Y_s^n) Z_s^n] \geq -C\Psi(Y_s^n)|Z_s^n|^2.$$

Then,

$$(3.13) \quad -n \left(\frac{1}{2} \text{Tr}[(Z_s^n)^\top \nabla^2 \psi^2(Y_s^n) Z_s^n] + 2n|\Psi(Y_s^n)|^2 |Z_s^n|^2 \right) \leq (Cn|\Psi(Y_s^n)| - cn^2|\Psi(Y_s^n)|^2) |Z_s^n|^2$$

Next, we observe that

$$(3.14) \quad (Cn|\Psi(Y_s^n)| - Cn^2|\Psi(Y_s^n)|^2) |Z_s^n|^2 \leq C|Z_s^n|^2.$$

Collecting (3.13)–(3.14) and using Lemma 3.1, we obtain

$$(3.15) \quad -\mathbb{E}_t \int_t^T (\text{Tr}[(Z_s^n)^\top \nabla^2 \psi^2(Y_s^n) Z_s^n] + 2n|\Psi(Y_s^n)|^2 |Z_s^n|^2) ds \leq Cn^{-1} \mathbb{E}_t \int_t^T |Z_s^n|^2 ds \leq Cn^{-1} \mathbb{E}_t \left[|\xi|^2 + \int_t^T (1 + |f(s, 0, 0)|^2) ds \right].$$

Taking the conditional expectation in (3.11), multiplying both sides by n , and using (3.12), (3.15), we complete the proof. \square

The following proposition improves the rate of convergence of Y^n to \mathcal{D} .

Proposition 3.2. *Under Assumptions 3.1 and 3.2, there exist $\mathfrak{N}, C > 0$, s.t. for any $n \geq \mathfrak{N}$, any solution (Y^n, Z^n) of (3.3), and any $t \in [0, T]$, the following holds a.s.:*

$$n\psi(Y_t^n) \leq C$$

Proof. First, we denote by $\|\nabla^2\psi(y)\|_*$ the maximum absolute value across all negative parts of the entries of the matrix $\nabla^2\psi(y)$. Next, we fix arbitrary $\epsilon, \varepsilon > 0$ satisfying

$$\epsilon \leq \left(\varepsilon + \sup_{y \in \partial\mathcal{D}} \frac{\|\nabla^2\psi(y)\|_*}{|\nabla\psi(y)|^2} \right)^{-1}, \quad \epsilon \leq \left(\varepsilon + 2 \sup_{y \in \partial\mathcal{D}, z \in \mathbb{R}^{d \times m}, s \in [0, T]} \frac{\|\mathbf{n}(y) \cdot f(s, y, z)\|_{\mathcal{L}^\infty}}{|\nabla\psi(y)|} \right)^{-1},$$

and define

$$\Psi^n(y) := (\psi(y) - 1/(\epsilon n))^+ \nabla\psi(y) = \frac{1}{2} \nabla \left((\psi(y) - 1/(\epsilon n))^+ \right)^2,$$

$$\tilde{H}(y) := \nabla^2 \left((\psi(y) - 1/(\epsilon n))^+ \right)^2 = 2\nabla\psi(y) \nabla^\top \psi(y) \mathbf{1}_{\psi \geq 1/(\epsilon n)} + 2(\psi(y) - 1/(\epsilon n))^+ \nabla^2\psi(y).$$

Next, we apply Itô's formula to $((\psi(Y_t^n) - 1/(\epsilon n))^+)^2$ (the validity of Itô's formula for the function $((\psi - 1/(\epsilon n))^+)^2$ is justified similarly to Remark 3.1), to obtain

$$\begin{aligned} ((\psi(Y_t^n) - 1/(\epsilon n))^+)^2 &= 2 \int_t^T \Psi^n(Y_s^n) \cdot f(s, Y_s^n, Z_s^n) ds - 2 \int_t^T n |\Psi(Y_s^n)| |\Psi^n(Y_s^n)| ds \\ (3.16) \quad &- 2 \int_t^T n |\Psi(Y_s^n)| |\Psi^n(Y_s^n)| |Z_s^n|^2 ds - 2 \int_t^T \Psi^n(Y_s^n) \cdot Z_s^n dW_s \\ &- \frac{1}{2} \int_t^T \text{Tr}[(Z_s^n)^\top \tilde{H}(Y_s^n) Z_s^n] ds. \end{aligned}$$

As $|\Psi^n|$ is linearly bounded (see Lemma 2.4), we conclude, as in the proof of Proposition 3.1, that the local martingale in the above representation is a true martingale.

Next, we note that

$$\begin{aligned} &2\Psi^n(Y_s^n) \cdot f(s, Y_s^n, Z_s^n) - n |\Psi(Y_s^n)| |\Psi^n(Y_s^n)| \\ &\leq |\Psi^n(Y_s^n)| (2|\mathbf{n}(Y_s^n) \cdot f(s, Y_s^n, Z_s^n)| - n\psi(Y_s^n) |\nabla\psi(Y_s^n)|). \end{aligned}$$

Notice that, whenever $\Psi^n(Y_s^n) > 0$, we have $\psi(Y_s^n) \geq 1/(\epsilon n)$ and, hence, $n\psi(Y_s^n) \geq 1/\epsilon$. Then, since $\epsilon > 0$ satisfies

$$\epsilon \leq \left(\varepsilon + 2 \sup_{y \in \partial\mathcal{D}, z \in \mathbb{R}^{d \times m}, s \in [0, T]} \frac{\|\mathbf{n}(y) \cdot f(s, y, z)\|_{\mathcal{L}^\infty}}{|\nabla\psi(y)|} \right)^{-1},$$

and since Y_s^n is close to \mathcal{D} for large enough n (due to Lemma 3.2), we conclude that

$$\begin{aligned} n\psi(Y_s^n) |\nabla\psi(Y_s^n)| &\geq |\nabla\psi(Y_s^n)| \left(\varepsilon + 2 \sup_{y \in \partial\mathcal{D}, z \in \mathbb{R}^{d \times m}, s \in [0, T]} \frac{\|\mathbf{n}(y) \cdot f(s, y, z)\|_{\mathcal{L}^\infty}}{|\nabla\psi(y)|} \right) \\ &\geq |\nabla\psi(Y_s^n)| \varepsilon/2 + 2|\mathbf{n}(Y_s^n) \cdot f(s, Y_s^n, Z_s^n)| \end{aligned}$$

and, in turn,

$$(3.17) \quad 2\Psi^n(Y_s^n) \cdot f(s, Y_s^n, Z_s^n) - n|\Psi(Y_s^n)||\Psi^n(Y_s^n)| \leq 0.$$

Next, we recall that

$$\frac{1}{2}\mathrm{Tr}[(Z_s^n)^\top \tilde{H}(Y_s^n) Z_s^n] \geq (\psi(Y_s^n) - 1/(\epsilon n))^+ \mathrm{Tr}[(Z_s^n)^\top \nabla^2 \psi(Y_s^n) Z_s^n]$$

and, hence,

$$-\frac{1}{2}\mathrm{Tr}[(Z_s^n)^\top \tilde{H}(Y_s^n) Z_s^n] \leq \|\nabla^2 \psi(Y_s^n)\|_* (\psi(Y_s^n) - 1/(\epsilon n))^+ |Z_s^n|^2.$$

In addition,

$$-|\Psi(Y_s^n)||\Psi^n(Y_s^n)||Z_s^n|^2 = -|\nabla \psi(Y_s^n)|^2 \psi(Y_s^n) (\psi(Y_s^n) - 1/(\epsilon n))^+ |Z_s^n|^2.$$

Collecting the two equations above, we deduce

$$\begin{aligned} -\frac{1}{2}\mathrm{Tr}[(Z_s^n)^\top \tilde{H}(Y_s^n) Z_s^n] - n|\Psi(Y_s^n)||\Psi^n(Y_s^n)||Z_s^n|^2 \\ \leq (\psi(Y_s^n) - 1/(\epsilon n))^+ |Z_s^n|^2 (\|\nabla^2 \psi(Y_s^n)\|_* - |\nabla \psi(Y_s^n)|^2 n \psi(Y_s^n)). \end{aligned}$$

Recall that, whenever $\psi(Y_s^n) \geq 1/(\epsilon n)$, we have $n\psi(Y_s^n) \geq 1/\epsilon$. Then, since $\epsilon > 0$ satisfies

$$\epsilon \leq \left(\epsilon + \sup_{y \in \partial \mathcal{D}} \frac{\|\nabla^2 \psi(y)\|_*}{|\nabla \psi(y)|^2} \right)^{-1},$$

and since Y_s^n is close to \mathcal{D} for large enough n , we conclude that

$$\begin{aligned} |\nabla \psi(Y_s^n)|^2 n \psi(Y_s^n) &\geq |\nabla \psi(Y_s^n)|^2 \left(\epsilon + \sup_{y \in \partial \mathcal{D}} \frac{\|\nabla^2 \psi(y)\|_*}{|\nabla \psi(y)|^2} \right) \\ &\geq |\nabla \psi(Y_s^n)|^2 \epsilon/2 + \|\nabla^2 \psi(Y_s^n)\|_* \end{aligned}$$

and, in turn,

$$(3.18) \quad -\frac{1}{2}\mathrm{Tr}[(Z_s^n)^\top \tilde{H}(Y_s^n) Z_s^n] - n|\Psi(Y_s^n)||\Psi^n(Y_s^n)||Z_s^n|^2 \leq 0.$$

Taking the conditional expectation in (3.16), we make use of equations (3.17) and (3.18), and of the fact that $|\Psi^n| \leq |\Psi|$, to obtain

$$((\psi(Y_t^n) - 1/(\epsilon n))^+)^2 + \mathbb{E}_t \int_t^T n |\Psi^n(Y_s^n)|^2 (1 + |Z_s^n|^2) ds \leq 0$$

and complete the proof. \square

Using Proposition 3.2, we can improve the statement of Proposition 3.1 and deduce that the Hölder norm of the Markovian solution of the penalized BSDE is bounded uniformly over n .

Corollary 3.1. *Under Assumptions 3.1 and 3.2, there exist constants $\mathfrak{N} \geq 1$, $\alpha' \in (0, 1]$, and $C > 0$ (independent of n), s.t., for any $n \geq \mathfrak{N}$, the BSDE (3.3) has a Markovian solution (Y^n, Z^n) , with $Y_t^n = u^n(t, X_t)$, and any such solution satisfies*

$$(3.19) \quad \sup_{(t,x) \neq (t',x')} \frac{|u^n(t, x) - u^n(t', x')|}{|t - t'|^{\alpha'/2} + |x - x'|^{\alpha'}} \leq C.$$

Proof. The statement of the corollary follows from Theorem 2.5 in [44] (without the localization). To verify the assumptions of the latter theorem, we consider the following capped version of (3.3):

$$(3.20) \quad \hat{Y}_t^{n,N} = g(X_T) + \int_t^T \hat{F}^{n,N}(s, X_s, \hat{Y}_s^{n,N}, \hat{Z}_s^{n,N}) ds - \int_t^T \hat{Z}_s^{n,N} dW_s,$$

with

$$\hat{F}^{n,N}(t, x, y, z) := f(t, x, y, z) - \rho_N(n\psi(y)) \nabla \psi(y) (1 + |z|^2)$$

and $\rho_N(x) = x \wedge N$. Propositions 3.1 and 3.2 imply the existence of (large enough) $N, \mathfrak{N} > 0$, s.t., for every $n \geq \mathfrak{N}$, there exists a Markovian solution (Y^n, Z^n) of (3.3), with $Y_t^n = u^n(t, X_t)$, and any such solution also solves (3.20). Moreover, there exists $c > 0$ such that $|u^n| \leq c$ for all n .

Next, we fix N as above and verify easily (as in the proof of Proposition 3.1) that, for large enough $C > 0$ and small enough $\alpha > 0$ (independent of n), $C(\alpha|y|^2 + (\phi_C(y) - \phi_C(0))^2)$ is a global c -Lyapunov function for $(\hat{F}^{n,N})_n$, in the sense of Definition 2.3 in [44]. In addition, $|\hat{F}^{n,N}(t, x, y, z)| \leq C + C_N|z|^2$, with the constants (C, C_N) independent of n . Thus, Theorem 2.5 in [44] yields the uniform boundedness of the Hölder norm of u^n . \square

W.l.o.g. we assume that the statements of Proposition 3.2 and Corollary 3.1 hold with $\mathfrak{N} = 1$. From Corollary 3.1, we deduce that there exists a subsequence of $\{u^n\}_{n \geq 1}$ converging locally uniformly to a function u satisfying (3.19). To alleviate the notation, this subsequence is still denoted $(u^n)_{n \geq 1}$. Recalling that $Y_t^n = u^n(t, X_t)$ and introducing $Y_t := u(t, X_t)$, for $t \in [0, T]$, we observe that

$$(3.21) \quad \mathbb{E} \left[\sup_{t \in [0, T]} |Y_t^n - Y_t|^2 \right] \xrightarrow{n \rightarrow +\infty} 0,$$

since $t \mapsto (t, X_t)$ is a.s. continuous and $\{|Y^n|\}$ is bounded uniformly by a constant, see Lemma 3.2.

We conclude this section with the following lemma, which is used in the next section. This lemma provides a uniform upper bound on the second moment of the auxiliary process

$$\Gamma_t^{n,m} := \exp \left(\int_0^t \left(1 + |\dot{K}_s^n| + |\dot{K}_s^m| \right) ds \right), \quad t \in [0, T],$$

where we recall (3.4).

Lemma 3.3. *Under Assumptions 3.1 and 3.2, for any $\varepsilon > 0$, there exists $N \geq 1$ (independent of n) such that, for all $n \geq 1$ and all $0 \leq k < N$, we have a.s.:*

$$\mathbb{E}_t \left[\int_t^{T(k+1)/N} |Z_s^n|^2 + |\dot{K}_s^n| ds \right] \leq \varepsilon, \quad \forall t \in [Tk/N, T(k+1)/N].$$

In particular, for any $\beta > 0$, there exists a constant $C = C(\beta)$ (independent of (n, m)), s.t.

$$\mathbb{E}[(\Gamma_T^{n,m})^\beta] \leq C,$$

for all $n, m \geq 1$.

Remark 3.2. *It is worth noticing that the constant C , appearing in Lemma 3.3, does not depend on the initial value x of the diffusion X , as follows from the proof of the lemma.*

Proof. The proof of the first statement of the lemma is an improvement of the estimates in the proof of Lemma 3.1, with the use of Corollary 3.1. We fix $t < t' \in [0, T]$, $\beta' > 0$ and $\alpha > 0$, and apply Itô's formula to the process $(e^{\beta'(s-t)}(\alpha|Y_s^n|^2 + (\phi_C(Y_s^n) - \phi_C(0))^2)_{s \in [t, t']}$ (recall (3.3)) to obtain, as in the proof of Lemma 3.1,

$$|Y_t^n|^2 + c\mathbb{E}_t \left[\int_t^{t'} |Z_s^n|^2 + |\dot{K}_s^n| ds \right] \leq \mathbb{E}_t \left[e^{\beta'(t'-t)} |Y_{t'}^n|^2 + C \int_t^{t'} e^{\beta'(s-t)} (1 + |F(s, X_s, 0, 0)|^2) ds \right],$$

which holds for large enough β' and small enough α .

Then, by using the upper bounds on $|Y^n|$, see Proposition 3.2 and on $|F(\cdot, \cdot, 0, 0)|$, see Assumption 3.2, we obtain:

$$\begin{aligned} & \mathbb{E}_t \left[\int_t^{t'} |Z_s^n|^2 + |\dot{K}_s^n| ds \right] \\ & \leq \mathbb{E}_t \left[e^{\beta'(t'-t)} |Y_{t'}^n|^2 - |Y_t^n|^2 + C \int_t^{t'} e^{\beta'(s-t)} (1 + |F(s, X_s, 0, 0)|^2) ds \right] \\ & \leq \mathbb{E}_t \left[(e^{\beta'(t'-t)} - 1) |Y_{t'}^n|^2 + |Y_{t'}^n + Y_t^n| |u^n(t', X_{t'}) - u^n(t, X_t)| + \frac{C}{\beta'} (e^{\beta'(t'-t)} - 1) \right] \\ & \leq C(\beta')(t' - t) + C\mathbb{E}_t \left[(t' - t)^{\alpha'/2} + |X_{t'} - X_t|^{\alpha'} \right] \leq C'(\beta')(t' - t)^{\alpha'/2}, \end{aligned}$$

where C' is independent of n , and we made use of Jensen's inequality and of standard SDE estimates on X in the last inequality. The above proves the first statement of the lemma.

To prove the second statement, we fix an arbitrary $\beta > 0$ and consider N corresponding to $\varepsilon = 1/(8\beta)$. Then, the first statement of the lemma and the John-Nirenberg inequality yield:

$$\begin{aligned} \mathbb{E} \left[e^{2\beta \int_0^T |\dot{K}_s^n| + |\dot{K}_s^m| ds} \right] &\leq \mathbb{E} \left[e^{2\beta \int_0^{T(N-1)/N} |\dot{K}_s^n| + |\dot{K}_s^m| ds} \mathbb{E}_{T(N-1)/N} \left[e^{2\beta \int_{T(N-1)/N}^T |\dot{K}_s^n| + |\dot{K}_s^m| ds} \right] \right] \\ &\leq 2\mathbb{E} \left[e^{2\beta \int_0^{T(N-1)/N} |\dot{K}_s^n| + |\dot{K}_s^m| ds} \right]. \end{aligned}$$

Iterating the above, we obtain the desired estimate. \square

3.4. Existence and uniqueness. We denote by $\{(Y^n, Z^n)\}_{n \geq 1}$ a sequence of Markovian solutions to (3.3) satisfying (3.21) (whose existence is established in the previous subsection). The goal of this subsection is to establish that $\{(Y^n, Z^n, K^n) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1\}_{n \geq 1}$ (with K^n defined in (3.4))² converges to a solution of the reflected BSDE (1.2) and that this solution is unique in the appropriate class.

Theorem 3.1. *Let Assumptions 3.1 and 3.2 hold. Then, there exists a triplet $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$, such that*

$$\lim_{n \rightarrow \infty} (\|Y^n - Y\|_{\mathcal{S}^2}, \|Z^n - Z\|_{\mathcal{H}^2}, \|K^n - K\|_{\mathcal{S}^2}) = 0.$$

which solves (1.2). The process K is absolutely continuous and satisfies, for all $\beta > 0$,

$$(3.22) \quad \mathbb{E}[e^{\beta \text{Var}_T(K)}] < \infty.$$

Moreover, this solution is unique in the class $\mathcal{U}(1)$ (recall Definition 2.1).

Remark 3.3. *If, in addition to Assumptions 3.1 and 3.2, g and F are globally Lipschitz in x (i.e., $\alpha = 1$ in Assumption 3.2), then there exists a constant C such that*

$$|Z| \leq C, \quad dt \times d\mathbb{P}\text{-a.e.}$$

Indeed, using the same arguments as in the proof of Corollary 3.1, we conclude that the conditions of Theorem 2.16 in [25] are satisfied. The latter theorem yields the existence of a constant C , s.t. $|Z_t^n| \leq C$ for a.e. (t, ω) and for all n . Then, it follows that $|Z| \leq C$.

Remark 3.4. *It is worth noticing that every exponential moment of $\text{Var}_T(K)$ can be bounded by a constant that does not depend on the initial value x of the diffusion X , as follows from Remark 3.2 and from the proof of Theorem 3.1.*

²The fact that $K^n \in \mathcal{K}^1$ follows from the inequality (3.7) and the second statement of Proposition 3.1.

Proof. 1.a We first prove the uniqueness of the solution in the desired class. For any solution (Y', Z', K') in $\mathcal{U}(1)$, we have

$$(3.23) \quad \mathbb{E} \left[e^{\frac{p'}{R_0} \text{Var}_T(K')} \right] < +\infty,$$

for some $p' > 1$. Setting $1 < p := (1 + p')/2 < p'$, $q' := p'/p > 1$ and $q = q'/(q' - 1)$, we obtain, using Hölder inequality,

$$(3.24) \quad \mathbb{E} \left[e^{\frac{p}{R_0} (\text{Var}_T(K) + \text{Var}_T(K'))} \right] \leq \mathbb{E} \left[e^{\frac{qp}{R_0} \text{Var}_T(K)} \right]^{\frac{1}{q}} \mathbb{E} \left[e^{\frac{q'p}{R_0} \text{Var}_T(K')} \right]^{\frac{1}{q'}}.$$

By (3.23), we have $\mathbb{E} \left[e^{\frac{q'p}{R_0} \text{Var}_T(K')} \right] = \mathbb{E} \left[e^{\frac{p'}{R_0} \text{Var}_T(K')} \right] < \infty$. Then using (3.22), which is proved below, we obtain

$$(3.25) \quad \mathbb{E} \left[e^{\frac{p}{R_0} (\text{Var}_T(K) + \text{Var}_T(K'))} \right] < +\infty.$$

Proposition 2.2, then, yields the uniqueness stated in the theorem.

1.b The fact that K is absolutely continuous is proved in Lemma 2.1.

2. Turning to the existence part of the proof, we have already obtained the convergence of $\{Y^n\}$ – recall (3.21). Moreover, it follows easily from Proposition 3.2 that, with probability one, Y_t takes values in $\bar{\mathcal{D}}$ for all $t \in [0, T]$.

We now turn to the convergence of $\{Z^n\}$. For $n, m \geq 1$, we denote

$$\begin{aligned} \delta f_t &:= f(t, Y_t^n, Z_t^n) - f(t, Y_t^m, Z_t^m), \quad \delta K := K^n - K^m, \\ \delta Y &:= Y^n - Y^m, \quad \delta Z := Z^n - Z^m. \end{aligned}$$

Applying Itô's formula to $(e^{\beta' s} |\delta Y_s|^2)_{s \in [t, T]}$, we obtain

$$(3.26) \quad \begin{aligned} |\delta Y_t|^2 + \int_t^T e^{\beta'(s-t)} |\delta Z_s|^2 ds &= 2 \int_t^T e^{\beta'(s-t)} \delta Y_s \cdot \delta f_s ds - 2 \int_t^T e^{\beta'(s-t)} \delta Y_s \cdot \delta \dot{K}_s ds \\ &- 2 \int_t^T e^{\beta'(s-t)} \delta Y_s \cdot \delta Z_s dW_s - \beta' \int_t^T e^{\beta'(s-t)} |\delta Y_s|^2 ds. \end{aligned}$$

Choosing a large enough $\beta' > 0$ and using the standard estimates, we deduce

$$(3.27) \quad \mathbb{E} \left[\int_0^T |\delta Z_s|^2 ds \right] \leq C \mathbb{E} \left[\int_0^T |\delta Y_s \cdot \delta \dot{K}_s| ds \right].$$

Note that Lemma 3.3 yields the existence of a constant C , s.t. $\mathbb{E} \left[\left(\int_0^T |\delta \dot{K}_s| ds \right)^2 \right] \leq C$, for all n, m . Then, using the Cauchy-Schwartz inequality, we obtain

$$\mathbb{E} \left[\int_0^T |\delta Y_s \cdot \delta \dot{K}_s| ds \right] \leq \mathbb{E} \left[\sup_{s \in [0, T]} |\delta Y_s|^2 \right]^{\frac{1}{2}} \left[\mathbb{E} \left(\int_0^T |\delta \dot{K}_s| ds \right)^2 \right]^{\frac{1}{2}}.$$

The above estimate, along with (3.27) and (3.21), implies that $\{Z^n\}_{n \geq 1}$ is a Cauchy sequence. Thus, there exists $(Y, Z) \in \mathcal{S}^2 \times \mathcal{H}^2$ such that $(Y^n, Z^n) \rightarrow (Y, Z)$.

Next, we recall that

$$K_t^n = Y_t^n - Y_0^n + \int_0^t f(s, Y_s^n, Z_s^n) ds - M_t^n, \quad M_t^n := \int_0^t Z_s^n dW_s.$$

Doob's maximal inequality implies that $\{M^n\}$ converges in \mathcal{S}^2 to M , with $M_t := \int_0^t Z_s dW_s$. As $f(t, \cdot, \cdot)$ is Lipschitz, we conclude that

$$(3.28) \quad \|K^n - K\|_{\mathcal{S}^2} \rightarrow 0,$$

with the continuous process K defined as

$$K_t := Y_t - Y_0 + \int_0^t f(s, Y_s, Z_s) ds - \int_0^t Z_s dW_s.$$

We now prove that $K \in \mathcal{K}^1$, and that dK_t is directed along \mathbf{n} and is active only when Y touches the boundary. To this end, we define the auxiliary nondecreasing processes

$$\hat{K}_t^n := \int_0^t n\psi(Y_s^n)(1 + |Z_s^n|^2) ds, \quad t \in [0, T].$$

From Lemma 3.1 we deduce the existence of a constant C , s.t. $\mathbb{E}[\hat{K}_T^n] \leq C$ for all n . Then, using Proposition 3.4 in [5], we know that there exists a nondecreasing nonnegative process \hat{K} , two sequences of integers $\{p \leq N_p\}$, with $p \rightarrow \infty$, and a family of numbers $\{\lambda_r^p\}$, with $\sum_{r=p}^{N_p} \lambda_r^p = 1$, such that

$$(3.29) \quad \mathbb{P}\left(\hat{K}_t^p := \sum_{r=p}^{N_p} \lambda_r^p \hat{K}_t^r \rightarrow \hat{K}_t, \forall t \in [0, T]\right) = 1.$$

The above implies that the measure induced by $d^p \hat{K}_t$ on $[0, T]$ converges a.s. to $d\hat{K}_t$. Then, for any bounded continuous process χ and any $0 \leq t_1 < t_2 \leq T$,

$$(3.30) \quad \eta^p(t_1, t_2) := \int_{t_1}^{t_2} \chi_t \sum_{r=p}^{N_p} \lambda_r^p \psi(Y_t^r)(1 + |Z_t^r|^2) dt = \int_{t_1}^{t_2} \chi_t d^p \hat{K}_t \rightarrow \int_0^T \chi_t d\hat{K}_t, \quad a.s.$$

From the first statement of Lemma 3.3 (with the use of Proposition 3.2), we conclude that, for any $\varepsilon > 0$, there exists $N \geq 1$ (independent of n) such that, for all p and all $0 \leq k < N$, we have a.s.:

$$\mathbb{E}_d[|\eta^p(t, T(k+1)/N)|] \leq \varepsilon, \quad \forall t \in [Tk/N, T(k+1)/N].$$

Then, repeating the last part of the proof of Lemma 3.3, we conclude that, for any $\beta > 0$, there exists a constant C , s.t.

$$\mathbb{E}[e^{\beta\eta^p(0,T)}] \leq C, \quad \forall p.$$

Thus, the family $\{\exp(\beta\eta^p(0,T))\}_p$ is uniformly integrable. The latter implies, in particular, that the convergence in (3.30) holds in \mathcal{L}^1 and that all exponential moments of \hat{K}_T are finite.

Next, we define

$${}^pK_t := \sum_{r=p}^{N_p} \lambda_r^p K_t^r, \quad t \in [0, T].$$

We also denote by $\widetilde{\nabla}\psi$ a Lipschitz extension of $\nabla\psi$ into \mathcal{D} (constructed as in Remark 3.1). Then, for any event A and any $t \in [0, T]$, we have:

$$\begin{aligned} \mathbb{E}[{}^pK_t \mathbf{1}_A] &= \mathbb{E}\left[\int_0^t \sum_{r=p}^{N_p} \lambda_r^p \widetilde{\nabla}\psi(Y_s^r) r \psi(Y_s^r) (1 + |Z_s^r|^2) ds \mathbf{1}_A\right] \\ &= \mathbb{E}\left[\int_0^t \widetilde{\nabla}\psi(Y_s) \sum_{r=p}^{N_p} \lambda_r^p r \psi(Y_s^r) (1 + |Z_s^r|^2) ds \mathbf{1}_A\right] \\ &\quad + O\left(\mathbb{E}\left[\sum_{r=p}^{N_p} \lambda_r^p \sup_{s \in [0,t]} |Y_s^r - Y_s| \int_0^t (1 + |Z_s^r|^2) ds\right]\right) \\ &\rightarrow \mathbb{E}\left[\int_0^t \widetilde{\nabla}\psi(Y_s) d\hat{K}_s ds \mathbf{1}_A\right], \end{aligned}$$

where we used (3.30), and its \mathcal{L}^1 version, and the estimate

$$\mathbb{E}\left[\sup_{s \in [0,t]} |Y_s^r - Y_s| \int_0^t (1 + |Z_s^r|^2) ds\right] \leq C \|Y^r - Y\|_{\mathcal{S}^2} \mathbb{E}\left[e^{\int_0^t (1 + |Z_s^r|^2) ds}\right]^{1/2} \leq C \|Y^r - Y\|_{\mathcal{S}^2},$$

which follows from Lemma 3.3.

On the other hand, as K^n converges to K in \mathcal{S}^2 , $\mathbb{E}[{}^pK_t \mathbf{1}_A]$ converges to $\mathbb{E}[K_t \mathbf{1}_A]$, and since A is arbitrary and K is continuous, we conclude:

$$(3.31) \quad \mathbb{P}\left(K_t = \int_0^t \widetilde{\nabla}\psi(Y_s) d\hat{K}_s ds, \quad \forall t \in [0, T]\right) = 1.$$

Note that the integrability of \hat{K}_T and the above representation, in particular, imply $K \in \mathcal{K}^1$.

It only remains to show that

$$(3.32) \quad \int_0^T \mathbf{1}_{\mathcal{D}}(Y_t) d\hat{K}_t = 0.$$

To this end, we choose an arbitrary Lipschitz f supported in \mathcal{D} and any event A , to obtain:

$$\begin{aligned} \mathbb{E}\left[\int_0^T f(Y_t)d\hat{K}_t \mathbf{1}_A\right] &= \lim_{n \rightarrow \infty} \mathbb{E}\left[\int_0^T f(Y_t) \sum_{r=p}^{N_p} \lambda_r^p r \psi(Y_t^r)(1 + |Z_t^r|^2)dt \mathbf{1}_A\right] \\ &= \lim_{n \rightarrow \infty} \mathbb{E}\left[\int_0^T \sum_{r=p}^{N_p} \lambda_r^p f(Y_t^r) r \psi(Y_t^r)(1 + |Z_t^r|^2)dt \mathbf{1}_A\right] \\ &\quad + O\left(\mathbb{E}\left[\sum_{r=p}^{N_p} \lambda_r^p \sup_{t \in [0, T]} |Y_t^r - Y_t| \int_0^T (1 + |Z_t^r|^2)dt\right]\right) = 0. \end{aligned}$$

As A is arbitrary, we conclude that, for any Lipschitz f supported in \mathcal{D} , we have $\int_0^T f(Y_t)d\hat{K}_t = 0$ a.s.. Approximating $\mathbf{1}_{\mathcal{D}}$ with a sequence of such f , we, e.g., use the monotone convergence theorem to deduce (3.32). Combining the latter with (3.31), we obtain (1.2)(ii) and conclude the proof of the first part of Theorem 3.1. \square

Remark 3.5. *Theorem 3.1 implies that, under Assumptions 3.1, 3.2 and 2.1 with $\theta = 1$, there exists a unique solution to (1.2) in $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$.*

4. WELL-POSEDNESS BEYOND MARKOVIAN FRAMEWORK

4.1. Discrete path-dependent framework. In this subsection, we extend the existence and uniqueness result obtained in a Markovian framework, see Theorem 3.1, to a discrete path-dependent framework.

Assumption 4.1. *Let ℓ be an arbitrary strictly positive integer and consider the partition $0 = t_0 < t_1 < \dots < t_\ell = T$ of $[0, T]$. We assume that*

$$\xi = g(X_{t_1}, \dots, X_{t_\ell}) \quad \text{and} \quad f(s, y, z) = F(s, X_{t_1 \wedge s}, \dots, X_{t_\ell \wedge s}, y, z)$$

where

- (i) g is α -Hölder and takes values in $\bar{\mathcal{D}}$,
- (ii) F is measurable in all variables, globally Lipschitz in (y, z) , uniformly over (x_1, \dots, x_ℓ) , and globally α -Hölder in (x_1, \dots, x_ℓ) , uniformly over (y, z) , and $|F(\cdot, \dots, 0, 0)|$ is bounded.

We note that $\ell = 1$ corresponds to the Markovian framework with an extra regularity assumption on the generator with respect to x . We also recall that Assumptions 1.1 and 1.2 hold throughout Section 4 even if not cited explicitly.

Theorem 4.1. *Let Assumptions 3.1 and Assumption 4.1 hold. Then, there exists a triplet $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ that solves (1.2). Moreover, all exponential moments of $\text{Var}_T(K)$ are finite, and this solution is unique in the class $\mathcal{U}(1)$ (recall Definition 2.1).*

Proof. Once the finiteness of the exponential moments of $\text{Var}_T(K)$ is proven, the uniqueness of the solution in the class $\mathcal{U}(1)$ follows from the same arguments as in step 1.a of the proof of Theorem 3.1. Let us now prove the existence part of the theorem. To this end, we use the backward recursion to construct a solution on each interval $[t_i, t_{i+1}]$ for $0 \leq i \leq \ell - 1$.

Since the case $\ell = 1$ corresponds to the already treated Markovian framework, we assume that $\ell > 1$ and we consider the interval time $[t_{\ell-1}, T]$. For any $(t, x) \in [0, T] \times \mathbb{R}^d$, we denote by $X^{t,x}$ the unique solution of (3.1) on $[t, T]$, which starts from x at time t . We reserve the notation X for the original diffusion started at time zero. For any $\mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_{\ell-1}) \in (\mathbb{R}^d)^{\ell-1}$, we denote by $(Y^{\mathbf{x}}, Z^{\mathbf{x}}, K^{\mathbf{x}})$ the solution of (1.2) on $[t_{\ell-1}, T]$, with the terminal condition $g(\mathbf{x}, X_T^{t_{\ell-1}, \mathbf{x}_{\ell-1}})$ and with the generator $F(\cdot, \mathbf{x}, X^{t_{\ell-1}, \mathbf{x}_{\ell-1}}, \cdot, \cdot)$, whose existence is ensured by Theorem 3.1 and whose uniqueness in an appropriate class follows from Theorem 2.2.

Next, we denote by $(Y^{\mathbf{x},n}, Z^{\mathbf{x},n})$ a Markovian solution of the penalized BSDE (3.3) on $[t_{\ell-1}, T]$, whose existence is ensured by Proposition 3.1. In particular, there exist measurable functions $u^n(\mathbf{x}, \cdot, \cdot)$ and $v^n(\mathbf{x}, \cdot, \cdot)$ such that

$$Y_t^{\mathbf{x},n} = u^n(\mathbf{x}, t, X_t^{t_{\ell-1}, \mathbf{x}_{\ell-1}}), \quad Z_t^{\mathbf{x},n} = v^n(\mathbf{x}, t, X_t^{t_{\ell-1}, \mathbf{x}_{\ell-1}}).$$

By considering a sequence of Lipschitz approximations of (3.3), given by (3.5), we apply Theorem 5.4 in [26] and, passing to the limit for the Lipschitz approximations as in the proof of Proposition 3.1, we conclude that a Markovian solution to (3.3) can be constructed so that u^n and v^n are jointly measurable in all variables. Passing to the limit in n along a subsequence, we use Theorem 3.1 and the uniform Hölder estimate in Corollary 3.1, to deduce the existence of jointly measurable functions u and v satisfying

$$(4.1) \quad Y_t^{\mathbf{x}} = u(\mathbf{x}, t, X_t^{t_{\ell-1}, \mathbf{x}_{\ell-1}}), \quad Z_t^{\mathbf{x}} = v(\mathbf{x}, t, X_t^{t_{\ell-1}, \mathbf{x}_{\ell-1}}).$$

Then, by denoting $\mathbf{X} = (X_{t_1}, \dots, X_{t_{\ell-1}})$, we consider the progressively measurable processes $(Y_t^{\mathbf{X}}, Z_t^{\mathbf{X}})_{t \in [t_{\ell-1}, T]}$ and define

$$K_t^{\mathbf{X}} := Y_t^{\mathbf{X}} - Y_{t_{\ell-1}}^{\mathbf{X}} + \int_{t_{\ell-1}}^t F(s, \mathbf{X}, X_s^{t_{\ell-1}, X_{t_{\ell-1}}}, Y_s^{\mathbf{X}}, Z_s^{\mathbf{X}}) ds - \int_{t_{\ell-1}}^t Z_s^{\mathbf{X}} dW_s, \quad t_{\ell-1} \leq t \leq T.$$

We note that $X_s^{t_{\ell-1}, X_{t_{\ell-1}}} = X_s$ and that $(Y_t^{\mathbf{X}}, Z_t^{\mathbf{X}}, K_t^{\mathbf{X}})_{t \in [t_{\ell-1}, T]}$ is a solution of (1.2) on the time interval $[t_{\ell-1}, T]$ satisfying $K_{t_{\ell-1}}^{\mathbf{X}} = 0$.

In order to iterate this construction and to extend the solution to the time interval $[t_{\ell-2}, t_{\ell-1}]$, we have to ensure that the associated terminal condition $Y_{t_{\ell-1}}^{\mathbf{X}}$ of the reflected BSDE (1.2) on $[t_{\ell-2}, t_{\ell-1}]$ is an α -Hölder function of \mathbf{X} . To this end, we recall the function u in (4.1) and define, for all $\tilde{\mathbf{x}} = (\mathbf{x}_1, \dots, \mathbf{x}_{\ell-2}) \in (\mathbb{R}^d)^{\ell-2}$ and $\mathbf{x}_{\ell-1} \in \mathbb{R}^d$, the deterministic function

$$\tilde{g}(\tilde{\mathbf{x}}, \mathbf{x}_{\ell-1}) := u(\tilde{\mathbf{x}}, \mathbf{x}_{\ell-1}, t_{\ell-1}, \mathbf{x}_{\ell-1}) = Y_{t_{\ell-1}}^{\tilde{\mathbf{x}}, \mathbf{x}_{\ell-1}}.$$

Let us prove that this function is α -Hölder. Indeed, for any $\mathbf{x} := (\tilde{\mathbf{x}}, \mathbf{x}_{\ell-1}) \in (\mathbb{R}^{d'})^{\ell-1}$ and $\mathbf{x}' := (\tilde{\mathbf{x}}', \mathbf{x}'_{\ell-1}) \in (\mathbb{R}^{d'})^{\ell-1}$, Proposition 2.2 with $p = 2$ yields

$$\begin{aligned} & |\tilde{g}(\mathbf{x}) - \tilde{g}(\mathbf{x}')| \leq \|Y^{\mathbf{x}} - Y^{\mathbf{x}'}\|_{\mathcal{S}^2} \\ & \leq C \mathbb{E} \left[\left| g(\mathbf{x}, X_T^{t_{\ell-1}, \mathbf{x}_{\ell-1}}) - g(\mathbf{x}', X_T^{t_{\ell-1}, \mathbf{x}'_{\ell-1}}) \right|^4 \right]^{1/4} \\ & \quad + C \mathbb{E} \left[\left(\int_0^T |F(s, \mathbf{x}, X_s^{t_{\ell-1}, \mathbf{x}_{\ell-1}}, Y_s^{\mathbf{x}}, Z_s^{\mathbf{x}}) - F(s, \mathbf{x}', X_s^{t_{\ell-1}, \mathbf{x}'_{\ell-1}}, Y_s^{\mathbf{x}}, Z_s^{\mathbf{x}})| ds \right)^4 \right]^{1/4} \\ & \leq C \left(|\mathbf{x} - \mathbf{x}'|^\alpha + \mathbb{E} \left[\sup_{0 \leq s \leq T} |X_s^{t_{\ell-1}, \mathbf{x}_{\ell-1}} - X_s^{t_{\ell-1}, \mathbf{x}'_{\ell-1}}|^{4\alpha} \right]^{1/4} \right), \end{aligned}$$

with a constant C that does not depend on x (see Remark 3.4). Then, the Jensen's inequality and the standard SDE estimates yield

$$|\tilde{g}(\mathbf{x}) - \tilde{g}(\mathbf{x}')| \leq C |\mathbf{x} - \mathbf{x}'|^\alpha,$$

which gives us the α -Hölder property of \tilde{g} . Considering the reflected BSDE (1.2) on $[t_{\ell-2}, t_{\ell-1}]$, with the terminal condition $Y_{t_{\ell-1}}^{\mathbf{x}} = \tilde{g}(X_{t_1}, \dots, X_{t_{\ell-1}})$ and with the generator

$$F(s, X_{t_1 \wedge s}, \dots, X_{t_{\ell-2} \wedge s}, X_{t_{\ell-1} \wedge s}, X_{t_{\ell-1} \wedge s}, y, z),$$

we deduce, as in the first part of the proof, that it has a solution in the form (4.1).

Finally, iterating the above construction, we concatenate the “ Y ” and “ Z ” parts of the solutions constructed in individual sub-intervals, and we sum up the “ K ” parts (assuming that every individual “ K ” part is extended continuously as a constant to the left and to the right of the associated sub-interval). It is easy to see that the resulting process $(Y, Z, \tilde{K}) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ is a solution of (1.2) on $[0, T]$. \square

4.2. General case.

Theorem 4.2. *Let Assumption 2.1 hold with $\theta = 2$. Then, there exists a triplet $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ that solves (1.2), and this solution is unique in the class $\mathcal{U}(2)$.*

Proof. The uniqueness part of the theorem is a direct consequence of Proposition 2.1 and Corollary 2.2. Let us now prove the existence part: To do so, we shall construct a Cauchy sequence of approximating reflected BSDEs.

First, we observe that the terminal condition ξ can be approximated by a sequence of random variables of the form $\xi^n := g_n(W_{t_1}, \dots, W_{t_n})$, where g_n is infinitely differentiable. The sequence $(\xi^n)_{n \in \mathbb{N}^*}$ can be chosen so that it converges to ξ in \mathcal{L}^q , for any $q \geq 1$ (see, e.g., [36]). In particular,

$$(4.2) \quad \lim_{n \rightarrow \infty} \mathbb{E} [|\xi - \xi^n|^{2p/(p-1)}] = 0,$$

with $p > 1$ appearing in Proposition 2.1. Replacing g_n by $g_n \wedge \|\xi\|_{\mathcal{L}^\infty}$, we can assume $\|\xi^n\|_{\mathcal{L}^\infty} \leq \|\xi\|_{\mathcal{L}^\infty}$. We observe that ξ^n satisfies Assumption 4.1(i) for $X = W$.

Second, to approximate the generator, for every $n \in \mathbb{N}^*$, we denote by \mathcal{K}^n the closed ball in $\mathbb{R}^{d \times d'}$ of radius n centered at zero, and choose a sequence of numbers $\epsilon_n \downarrow 0$. We set

$$\ell_n := \|f(\cdot, 0, 0)\|_{\mathcal{L}^\infty} + K_{f,y} \sup_{y \in \bar{\mathcal{D}}} |y| + nK_{f,z},$$

recalling Assumption 1.2. For each n , we introduce \mathfrak{L}^n the compact convex subset of $\mathcal{C}(\bar{\mathcal{D}} \times \mathcal{K}^n)$ (the space of continuous function endowed with the uniform norm denoted $\|\cdot\|_s$) consisting of Lipschitz functions, with the Lipschitz coefficients $K_{f,y}$ in the variable $y \in \bar{\mathcal{D}}$ and $K_{f,z}$ in the variable $z \in \mathcal{K}^n$, and whose (uniform) norm is bounded by ℓ_n . Note that $f|_{\mathcal{K}^n}$ is valued in \mathfrak{L}^n . We are now going to build an approximation of $f|_{\bar{\mathcal{D}} \times \mathcal{K}^n}$ in \mathfrak{L}^n satisfying Assumption 4.1 (for $X = W$).

Let $\{\phi_n^m\}_{m=1}^{M_n}$ be an ϵ_n -cover of the compact set \mathfrak{L}^n with M_n a positive integer. We denote by $\tilde{f}^n(t, \cdot)$ the (measurable selection of the) proximal projection of $f|_{\bar{\mathcal{D}} \times \mathcal{K}^n}(t, \cdot)$ on $\{\phi_n^m\}_{m=1}^{M_n}$. It satisfies

$$\tilde{f}^n(t, \cdot) = \sum_{m=1}^{M_n} \phi_n^m(\cdot) (\tilde{\eta}_t^n)^m =: \phi_n(\cdot) \tilde{\eta}_t^n \quad \text{and} \quad \|\tilde{f}^n(t, \cdot) - f|_{\bar{\mathcal{D}} \times \mathcal{K}^n}(t, \cdot)\|_s \leq \epsilon_n \quad \text{a.s.}$$

with $\tilde{\eta}^n$ a progressively measurable process which takes its value on the (non zero) extremal points of $\mathcal{S}_{M_n} := \{x \in \mathbb{R}^{M_n} \mid 0 \leq x^m \leq 1, \sum_{m=1}^{M_n} x^m \leq 1\}$. Then, using the dominated convergence theorem, we have

$$(4.3) \quad \mathbb{E} \left[\int_0^T \|f|_{\bar{\mathcal{D}} \times \mathcal{K}^n}(t, \cdot) - \tilde{f}^n(t, \cdot)\|_s^{2p/(p-1)} dt \right] \leq T(\epsilon_n)^{2p/(p-1)}.$$

We now classically approximate $(\tilde{\eta}_t^n)_{t \in [0, T]}$ by an adapted process $(\hat{\eta}_t^n)_{t \in [0, T]}$ piecewise constant on a grid $\Pi_n := \{t_0 = 0 < \dots < t_k^n < \dots < t_{\kappa_n}^n = T\}$. This process can be chosen to be \mathcal{S}_{M_n} -valued and satisfying

$$\mathbb{E} \left[\int_0^T |\tilde{\eta}_t^n - \hat{\eta}_t^n|^{2p/(p-1)} dt \right] \leq \frac{\epsilon_n}{(M_n \ell_n^2)^{p/(p-1)}}.$$

Setting

$$(4.4) \quad \hat{f}^n(t, \cdot) = \sum_{k=0}^{\kappa_n-1} \phi_n(\cdot) \hat{\eta}_{t_k^n}^n \mathbf{1}_{(t_k^n, t_{k+1}^n]}(t),$$

which is \mathfrak{L}^n -valued, as a random convex combination of the $\{\phi_n^m\}_{m=1}^{M_n}$, we compute

$$(4.5) \quad \mathbb{E} \left[\int_0^T \|\tilde{f}^n(t, \cdot) - \hat{f}^n(t, \cdot)\|_s^{2p/(p-1)} dt \right] \leq \epsilon_n.$$

Then, we apply the approximation result of [36] for each $\hat{\eta}_{t_k}^n$. Introducing, if necessary, a finer grid $\mathfrak{R}_n \subset \Pi_n$, we set

$$\eta_{t_k}^n := \mathfrak{P}_{\mathcal{S}} \left[r_k^n \left((W_r)_{r \in \mathfrak{R}_n, r \leq t_k^n} \right) \right],$$

where r_k^n is a smooth function with values in \mathbb{R}^{M_n} and $\mathfrak{P}_{\mathcal{S}}$ the (orthogonal) projection onto \mathcal{S}_{M_n} . We can chose r_k^n such that

$$(4.6) \quad \mathbb{E} \left[|\eta_{t_k}^n - \hat{\eta}_{t_k}^n|^{2p/(p-1)} \right] \leq \frac{\epsilon_n}{(M_n \ell_n^2)^{p/(p-1)}}.$$

Setting $f^n(t, \cdot) = \sum_{k=0}^{k_n-1} \phi_n(\cdot) \eta_{t_k}^n \mathbf{1}_{(t_k^n, t_{k+1}^n]}(t)$, which belongs to \mathfrak{L}^n , we have

$$(4.7) \quad \mathbb{E} \left[\int_0^T \|\hat{f}^n(t, \cdot) - f^n(t, \cdot)\|_s^{2p/(p-1)} dt \right] \leq T \epsilon_n.$$

Collecting the above, we conclude that

$$(4.8) \quad \lim_{n \rightarrow \infty} \mathbb{E} \int_0^T \sup_{y \in \bar{D}, z \in \mathcal{K}^n} |f(t, y, z) - f^n(t, y, z)|^{2p/(p-1)} dt = 0.$$

We extend $f^n(t, y, \cdot)$ to $\mathbb{R}^{d \times d} \setminus \mathcal{K}^n$ as a constant in each radial direction, so that its uniform norm and Lipschitz coefficient do not change.

When, moreover, f satisfies Assumption 2.1-(i) (resp. Assumption 2.1-(iii)), the above construction allow to build an approximating sequence f^n having the same properties. One simply works with $\tilde{\mathfrak{L}}^n$ instead of \mathfrak{L}^n , where $\tilde{\mathfrak{L}}^n$ is the closed convex subset of \mathfrak{L}^n whose function satisfies Assumption 2.1-(i) (resp. Assumption 2.1-(iii)).

Now, since, for any $n \in \mathbb{N}^*$, ξ^n and f^n satisfy Assumption 4.1, we can invoke Theorem 4.1 to obtain an unique solution $(Y^n, Z^n, K^n) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ to (1.2) associated with this data. Thanks to Proposition 2.1, we are allowed to apply Proposition 2.2: for all $n, m \in \mathbb{N}^*$,

$$(4.9) \quad \begin{aligned} & \|Y^n - Y^m\|_{\mathcal{S}^2} + \|Z^n - Z^m\|_{\mathcal{H}^2} + \|K^n - K^m\|_{\mathcal{S}^2} \\ & \leq C \mathbb{E} \left[|\xi^n - \xi^m|^{2p/(p-1)} \right]^{(p-1)/(2p)} \\ & + C \mathbb{E} \left[\left(\int_0^T |f^n(s, Y_s^n, Z_s^n) - f^m(s, Y_s^m, Z_s^m)| ds \right)^{2p/(p-1)} \right]^{(p-1)/(2p)}, \end{aligned}$$

with a constant C that does not depend on n and m .

Applying the Cauchy-Schwartz, Jensen's, and Chebyshev's inequalities, we obtain

$$\begin{aligned} & \mathbb{E} \left[\left(\int_0^T (1 + |Z_t^n|) \mathbf{1}_{\{|Z_t^n| > n\}} dt \right)^{2p/(p-1)} \right] \\ & \leq T^{p/(p-1)-1/2} \mathbb{E} \left[\left(\int_0^T (1 + |Z_t^n|)^2 dt \right)^{2p/(p-1)} \right]^{1/2} \mathbb{E} \left[\int_0^T \mathbf{1}_{\{|Z_t^n| > n\}} dt \right]^{1/2} \\ & \leq \frac{T^{p/(p-1)-1/2}}{n} \mathbb{E} \left[\left(\int_0^T (1 + |Z_t^n|)^2 dt \right)^{2p/(p-1)} \right]^{1/2} \mathbb{E} \left[\int_0^T |Z_t^n|^2 dt \right]^{1/2}. \end{aligned}$$

Using Proposition 2.1 and the energy inequality for BMO martingales, we can estimate $\mathbb{E} \left(\int_0^T (1 + |Z_t^n|)^2 dt \right)^{2p/(p-1)}$ uniformly over n . Then, for all $m \geq n$, we obtain from the above estimate:

$$\begin{aligned} & \mathbb{E} \left[\left(\int_0^T |f^n(t, Y_t^n, Z_t^n) - f^m(t, Y_t^n, Z_t^n)| dt \right)^{2p/(p-1)} \right]^{(p-1)/(2p)} \\ & \leq C \mathbb{E} \left[\left(\int_0^T (1 + |Z_t^n|) \mathbf{1}_{\{|Z_t^n| > n\}} dt \right)^{2p/(p-1)} \right]^{(p-1)/(2p)} \\ (4.10) \quad & + C \mathbb{E} \left[\left(\int_0^T |f^n(t, Y_t^n, Z_t^n) - f^m(t, Y_t^n, Z_t^n)| \mathbf{1}_{\{|Z_t^n| \leq n\}} dt \right)^{2p/(p-1)} \right]^{(p-1)/(2p)} \\ & \leq \frac{C}{n^{(p-1)/(2p)}} + C \mathbb{E} \left[\int_0^T |f^n(t, Y_t^n, Z_t^n) - f(t, Y_t^n, Z_t^n)|^{2p/(p-1)} \mathbf{1}_{\{Y_t^n \in \bar{\mathcal{D}}, |Z_t^n| \leq n\}} dt \right]^{(p-1)/(2p)} \\ & + C \mathbb{E} \left[\int_0^T |f^m(t, Y_t^n, Z_t^n) - f(t, Y_t^n, Z_t^n)|^{2p/(p-1)} \mathbf{1}_{\{Y_t^n \in \bar{\mathcal{D}}, |Z_t^n| \leq m\}} dt \right]^{(p-1)/(2p)}. \end{aligned}$$

In view of (4.8), the right hand side of the above vanishes as $n, m \rightarrow \infty$. Collecting (4.2), (4.9) and (4.10), we conclude:

$$\|Y^n - Y^m\|_{\mathcal{S}^2} + \|Z^n - Z^m\|_{\mathcal{H}^2} + \|K^n - K^m\|_{\mathcal{S}^2} \xrightarrow{n, m \rightarrow +\infty} 0.$$

In other words, $(Y^n, Z^n, K^n)_{n \in \mathbb{N}^*}$ is a Cauchy sequence in $\mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{S}^2$. Then, there exists $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{S}^2$ such that $(Y^n, Z^n, K^n) \xrightarrow{n \rightarrow +\infty} (Y, Z, K)$. Moreover, Y takes values on $\bar{\mathcal{D}}$. As (Y^n, Z^n, K^n) is the unique solution to (1.2)

associated to the terminal condition ξ^n and the generator f^n , and since we have

$$\begin{aligned} & \mathbb{E} \left[\int_t^T |f^n(s, Y_s^n, Z_s^n) - f(s, Y_s, Z_s)| ds \right] \\ & \leq \mathbb{E} \left[\int_t^T |f^n(s, Y_s, Z_s) - f(s, Y_s, Z_s)| ds \right] + C \mathbb{E} \left[\int_t^T |Y_s^n - Y_s| + |Z_s^n - Z_s| ds \right], \end{aligned}$$

we can easily pass to the limit in (1.2)(i) to show that (Y, Z, K) satisfies (1.2)(i).

It remains to prove that $K \in \mathcal{K}^1$, that dK_t is directed along $\mathbf{n}(Y_t)$ and that it is active only when Y touches the boundary (the latter two properties will be shown via the alternative characterization given by Lemma 2.3). Repeating the derivation of (2.11)-(2.12) for (Y^n, Z^n, K^n) , but without taking the conditional expectations, and with $\beta = 0$, we obtain:

$$\int_0^T d\text{Var}_s(K^n) \leq C \left(|\xi^n|^2 + \int_0^T 2Y_s^n \cdot f(s, Y_s^n, Z_s^n) ds - \int_0^T 2Y_s^n Z_s^n dW_s \right),$$

where the constant C does not depend on n . The right hand side of the above inequality converges in probability, hence it also converges a.s. up to a subsequence which we still denote $\{(Y^n, Z^n, K^n)\}$. Then, $\{\text{Var}_T(K^n)\}_{n \in \mathbb{N}^*}$ is a.s. bounded uniformly over n , and Fatou's lemma yields that $\text{Var}_T(K)$ is a.s. bounded – i.e. K is a bounded variation process. Thanks to Proposition 2.1, $\{\text{Var}_T(K^n)\}_{n \in \mathbb{N}^*}$ is uniformly integrable and, hence, $K \in \mathcal{K}^1$. As (Y^n, Z^n, K^n) solve (1.2) with the terminal condition ξ^n and the generator f^n , Lemma 2.3 yields the existence of a constant c , independent of n , such that, for all continuous adapted process V with values in $\bar{\mathcal{D}}$, we have

$$\int_0^T (Y_s^n - V_s) dK_s^n + c|Y_s^n - V_s|^2 \mathbf{n}(Y_s^n) dK_s^n \geq 0 \quad \text{a.s.}$$

Finally, we use Lemma 5.8 in [20] to pass to the limit in the above inequality, to obtain

$$\int_0^T (Y_s - V_s) dK_s + c|Y_s - V_s|^2 \mathbf{n}(Y_s) dK_s \geq 0 \quad \text{a.s.}$$

and complete the proof of the theorem via another application of Lemma 2.3. \square

5. CONNECTION TO BROWNIAN Γ -MARTINGALES

It turns out that the solutions to reflected BSDEs in non-convex domains, defined via (1.2) and constructed in the previous sections, are naturally connected to the notion of martingales on manifolds (a.k.a. Γ -martingales – see [15]). In this section, we investigate this connection more closely, in particular, discovering a new proof of the existence and uniqueness of a Brownian martingale with a prescribed terminal value on a section of a sphere and showing the sharpness of the weak star-shape property in Assumption 1.1.

The connection to martingales on manifolds is made precise by the following proposition, which states that, under certain assumptions, the Y -component of the solution to (1.2) always stays on the boundary of the domain \mathcal{D} . Treating $\partial\mathcal{D}$ as a manifold and expressing dK_t via $\nabla^2\phi(Y_t)$ and Z_t , we discover that Y satisfies the definition of a Brownian Γ -martingale on the manifold $\partial\mathcal{D}$ given in [15].

Proposition 5.1. *Assume the following:*

- *there exists a convex domain \mathcal{A} , satisfying $\bar{\mathcal{A}} \cap \bar{\mathcal{D}} \subset \partial\mathcal{D}$,*
- *$\mathbf{1}_{\{y \in \partial\mathcal{D} \setminus \bar{\mathcal{A}}\}} \nabla d(y, \mathcal{A}) \cdot \nabla\phi(y) \geq 0$,*
- *$f \equiv 0$ and $\xi \in \bar{\mathcal{A}} \cap \partial\mathcal{D}$ almost surely,*
- *$(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ solve (1.2).*

Then, $Y \in \bar{\mathcal{A}} \cap \bar{\mathcal{D}} \subset \partial\mathcal{D}$ almost surely. Moreover, we have

$$(5.1) \quad d\text{Var}_t(K) = \left[-\frac{1}{2} \text{Tr}[Z_t^\top \nabla^2\phi(Y_t) Z_t] \right]^+ dt.$$

Finally, Y is a Γ -martingale, with the prescribed terminal value ξ , in the manifold $\partial\mathcal{D}$ endowed with the Riemannian structure inherited from \mathbb{R}^d and its canonical connection Γ , as defined in [15].

Remark 5.1. *It is worth mentioning that the assumptions made in Proposition 5.1 imply that the set \mathcal{A} cannot be smooth. To obtain an intuitive understanding of what the set \mathcal{A} may look like, we refer the reader to the example that follows.*

Proof. We apply Itô's formula for general convex functions (in the form of an inequality, as in [3]) to the process $d(Y_t, \mathcal{A})$ to obtain

$$0 \leq d(Y_t, \mathcal{A}) \leq \mathbb{E}_t \left[d(\xi, \mathcal{A}) - \int_t^T \mathbf{1}_{\{Y_s \in \partial\mathcal{D} \setminus \bar{\mathcal{A}}\}} \nabla d(Y_s, \mathcal{A}) dK_s \right] \leq 0, \quad t \in [0, T],$$

which gives us $Y \in \bar{\mathcal{A}} \cap \mathcal{D} \subset \partial\mathcal{D}$. Applying Itô's formula to $\phi(Y_t)$ yields (5.1). Finally, using (5.1), the fact that dK_t is orthogonal to the tangent space of $\partial\mathcal{D}$ at the point Y_t , as well as (4.9), (4.10), (5.6)(ii) from [15], we conclude that Y is a Γ -martingale on $\partial\mathcal{D}$. \square

In the remainder of this section, we assume $f = 0$ and present a simple example of the domains \mathcal{D} and \mathcal{A} for which the assumptions of Proposition 5.1 hold. This example allows us to obtain an alternative proof of a known result on Γ -martingales using the reflected BSDEs and, on the other hand, to illustrate the sharpness of the weak star-shape assumption (see Assumption 1.1) using the martingales on manifolds.

In this example, we construct the functions ϕ and ϕ_C , which define the domains \mathcal{D} and \mathcal{C} as in Assumption 1.1, first, in the plane $\mathcal{P} := \mathbb{R} \times \{0\}^{d-2} \times \mathbb{R}$ of \mathbb{R}^d . These functions are built symmetric with respect to the y_d -axis, as the domains are themselves symmetric, see the precise description below. Then, we extend


them to \mathbb{R}^d via

$$\phi(y) = \phi((r(y), 0, \dots, 0, y_d)), \quad \phi_{\mathcal{C}}(y) = \phi_{\mathcal{C}}((r(y), 0, \dots, 0, y_d)),$$

with $r(y) := \left(\sum_{i=1}^{d-1} |y^i|^2\right)^{1/2}$. By an abuse of notation we denote by the same names the associated domains \mathcal{D} and \mathcal{C} constructed in \mathbb{R}^d and their intersections with \mathcal{P} .

We consider the three parameters $\alpha \in (0, \pi/2)$, $\eta > 0$, $\varepsilon \in (0, \pi/2 - \alpha)$, and the domains $\mathcal{D}_{\alpha, \eta, \varepsilon}$, $\mathcal{C}_{\alpha, \eta}$, $\mathcal{A}_{\alpha, \varepsilon}$ given in Figure 1, which satisfy the following properties.

- $\mathcal{C}_{\alpha, \eta}$ is a square centered at $(0, -1 - \eta - \sin(\alpha))$, with length $2 \sin(\alpha) + 2\eta$, with the edges parallel to axes and with rounded corners (obtained by modifying the square in the η -neighborhoods of these corners), such that $\partial \mathcal{C}_{\alpha, \eta}$ is a C^2 curve and $\mathcal{C}_{\alpha, \eta}$ is convex,
- $\mathcal{D}_{\alpha, \eta, \varepsilon}$ is symmetric w.r.t. the axis y_d .
- $\partial \mathcal{D}_{\alpha, \eta, \varepsilon}$ is C^2 and is made up of the following pieces:
 - the arc \mathcal{S}_{α} of angle 2α , symmetric w.r.t. the axis y_d , of the circle centered at zero and with the radius 1,
 - the arc of angle 2α , symmetric w.r.t. the axis y_d , of the circle centered at zero and with the radius $(2 \sin(\alpha) + 2\eta + 1)/\cos(\alpha)$,
 - and two smooth curves \mathcal{L}_1 and \mathcal{L}_2 , symmetric to each other w.r.t. the axis y_d , which connect the two arcs described above forming a C^2 closed curve that does not intersect itself nor $\mathcal{C}_{\alpha, \eta}$.
- We denote by A^1 (respectively, A^2) the end point of the curve \mathcal{S}_{α} that belongs to the right (respectively, left) half-plane w.r.t. the axis y_d .
- Let us assume that \mathcal{L}_1 (respectively, \mathcal{L}_2) belongs to the right (respectively, left) half-plane w.r.t. the axis y_d . We also assume that the curve \mathcal{L}_1 is constructed so that, in its natural parameterization with the starting point A^1 , the slope of its tangent vector has exactly one change of monotonicity. Namely, we assume that there exists a point B_{ε}^1 , such that the angle between B_{ε}^1 and A^1 relative to zero is ε and such that the derivative of the slope of the aforementioned tangent vector is continuous, nonincreasing, and equal to zero at B_{ε}^1 . The curve \mathcal{L}_2 , then, satisfies the analogous property due to symmetry, with the associated point B_{ε}^2 .
- As the curve $\partial \mathcal{D}_{\alpha, \eta, \varepsilon}$ is C^2 , closed, and without self-intersections, we construct ϕ as the signed distance to $\partial \mathcal{D}_{\alpha, \eta, \varepsilon}$ in a neighborhood of $\partial \mathcal{D}_{\alpha, \eta, \varepsilon}$ and, then, extend it in a smooth way to \mathbb{R}^2 . $\phi_{\mathcal{C}}$ is constructed similarly.
- We define $\mathcal{S}_{\alpha, \varepsilon}$ as the concatenation of the curves $B_{\varepsilon}^2 A^2$, \mathcal{S}_{α} , $A^1 B_{\varepsilon}^1$, and we define $\mathcal{A}_{\alpha, \varepsilon}$ as the interior of the convex hull of $\mathcal{S}_{\alpha, \varepsilon}$.
- Finally, we always assume that $\eta > 0$ is small enough, such that $\mathcal{C}_{\alpha, \eta}$ is included in the triangle with vertices P^1 , 0 and P^2 , as shown in Figure 1. This ensures that $\tilde{\mathcal{C}}_{\alpha, \eta} \subset \mathcal{D}_{\alpha, \eta, \varepsilon}$ for any $\varepsilon > 0$.

FIGURE 1. Domains $\mathcal{D}_{\alpha,\eta,\varepsilon}$, $\mathcal{C}_{\alpha,\eta}$, $\mathcal{A}_{\alpha,\varepsilon}$

Let us now consider a terminal condition $\xi \in \mathcal{S}_\alpha$ and verify that $\mathcal{D}_{\alpha,\eta,\varepsilon}$, $\mathcal{A}_{\alpha,\varepsilon}$ and ξ satisfy the desired assumptions. We easily deduce that $R_0 = 1$. Then, for any $\alpha \in (0, \pi/2)$ and $\eta > 0$, there exists $\varepsilon_0 \in (0, \pi/2 - \alpha)$, such that, for all $0 < \varepsilon < \varepsilon_0$, the condition (1.1) holds up to a shift of coordinates in \mathbb{R}^d that maps the origin to $a_{\alpha,\eta} := (0, \dots, 0, -1 - \eta - \sin(\alpha))$. The other conditions of Assumption 1.1 follow easily.

Next, we notice that, in the discrete path-dependent framework and under Assumption 4.1, we can apply Theorem 4.1 to conclude that there exists a unique (in $\mathcal{U}(1)$) triplet $(Y^\varepsilon, Z^\varepsilon, K^\varepsilon) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ that solves (1.2) in the domain $\mathcal{D} = \mathcal{D}_{\alpha,\eta,\varepsilon}$ (we suppress the dependence of the solution on η and α as they are fixed in what follows). Proposition 5.1 applied to $\mathcal{D} = \mathcal{D}_{\alpha,\eta,\varepsilon}$ and $\mathcal{A} = \mathcal{A}_{\alpha,\varepsilon}$ (whose assumptions are satisfied by the construction of $\mathcal{A}_{\alpha,\varepsilon}$, \mathcal{L}_1 and \mathcal{L}_2) states that Y^ε lives in $\mathcal{S}_{\alpha,\varepsilon}$. Then, the stability result of Proposition 2.2 yields that $\{(Y^{1/n}, Z^{1/n}, K^{1/n})\}_{n=1}^\infty$ is a Cauchy sequence and, hence, has a limit (Y, Z, K) . It is clear that Y lives on \mathcal{S}_α . Then, applying the arguments similar to those used in the proof of Theorem 3.1, one can deduce that (Y, Z, K) solve the reflected BSDE (1.2) in the domain $\mathcal{D} = \mathcal{D}_{\alpha,\eta,\varepsilon'}$, for any $\varepsilon' \in (0, \varepsilon_0)$. Applying Proposition 5.1 once more and recalling that Y lives on \mathcal{S}_α , we conclude that Y is a Γ -martingale on the manifold \mathcal{S}_α with the terminal condition ξ . The uniqueness part of Theorem 4.1 yields that such a Γ -martingale is unique (in $\mathcal{U}(1)$).

Now, let us come back to a general terminal condition. We first notice that Proposition 2.1 holds for any solution $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ of (1.2) that lives in \mathcal{S}_α and satisfies Assumption 2.1(i) with γ replaced by

$$\gamma_\alpha := \inf_{y \in \mathcal{S}_\alpha} \nabla \phi_{\mathcal{C}}(y) \cdot \frac{\nabla \phi(y)}{|\nabla \phi(y)|}.$$

We can easily compute $\gamma_\alpha = \cos(\alpha)$. Moreover, we also have $R_0 = 1$ and

$$|\phi_{\mathcal{C}}^+(\xi)|_{\mathcal{L}^\infty} \leq 1 - \cos(\alpha).$$

Thus, we conclude that Assumption 2.1(i) is fulfilled with $\theta = 2$ as long as $\cos(\alpha) > 2/3$. Considering a sequence of discrete path-dependent terminal conditions that approximate the given (general) terminal condition and take values in \mathcal{S}_α , we repeat the proof of Theorem 4.2 obtaining a unique (in $\mathcal{U}(2)$) triplet $(Y, Z, K) \in \mathcal{S}^2 \times \mathcal{H}^2 \times \mathcal{K}^1$ that solves (1.2) in the domain $\mathcal{D} = \mathcal{D}_{\alpha, \eta, \varepsilon'}$, for any $\varepsilon' \in (0, \varepsilon_0)$, and is such that Y lives in \mathcal{S}_α . Applying Proposition 5.1 once more, we conclude that Y is a Γ -martingale on the manifold \mathcal{S}_α with the terminal condition ξ . The uniqueness part of Theorem 4.1 yields that such a Γ -martingale is unique in $\mathcal{U}(2)$.

To sum up, the above construction proves the existence and uniqueness result for a Brownian Γ -martingale with a prescribed discrete path-dependent terminal condition ξ , satisfying Assumption 4.1, on any sector of the sphere \mathbb{S}^{d-1} (we understand a sector as an intersection of a sphere and a half-space) that is strictly contained in a hemisphere. For a general terminal condition ξ , we are only able to tackle the case $\alpha < \arccos(2/3)$. Thus, our results provide an alternative proof of a particular case of [30, 41], where the existence and uniqueness for any $\alpha < \pi/2$ is established. Considering the case $\alpha = \pi/2$, we notice that, for any \mathcal{D} that is included in the outside of a sphere and whose boundary contains a hemisphere, it is impossible to find a convex domain $\mathcal{C} \subset \mathcal{D}$ that can “see” all points on the boundary of this hemisphere with a strictly positive angle: in other words, (1.1) can not be fulfilled. Therefore, our existence and uniqueness results fail for the case of a hemisphere. On the other hand, considering directly the problem of existence and uniqueness of a Brownian Γ -martingale with a prescribed terminal condition on a closed hemisphere of \mathbb{S}^{d-1} , we notice the major challenge that is due to the non-uniqueness of geodesics, when $d \geq 3$. Indeed, assume that ξ takes its values in the set $\{z_1, z_2\}$ consisting of two antipodes on the sphere (i.e., the line connecting the two points goes through the center of the sphere) and note that $\mathcal{S}_{\pi/2}$ does contain such points. Then, for any shortest arc $\widehat{z_1 z_2} \subset \mathcal{S}_{\pi/2}$, there exists a Γ -martingale on the manifold $\widehat{z_1 z_2}$ with the terminal condition ξ . As the arc $\widehat{z_1 z_2}$ is a geodesic, we conclude that the resulting Γ -martingale is also a Γ -martingale in the larger manifold $\mathcal{S}_{\pi/2}$. Assuming that ξ takes each of its two values with strictly positive probability and recalling that there are infinitely many geodesic arcs $\widehat{z_1 z_2}$ on $\mathcal{S}_{\pi/2}$, we conclude that the uniqueness of a Γ -martingale on \mathbb{S}^{d-1} with

the terminal condition ξ does not hold. This observation, in particular, illustrates the sharpness of the weak star-shape property in Assumption 1.1 (condition (1.1)) in the case of a general terminal condition and general $d \geq 2$.

Let us also mention that the non-uniqueness described above does not occur for $d = 2$, which indicates that it may be possible to relax our assumptions for reflected BSDEs in planar non-convex domains. In particular, we refer to [40] for a complete treatment of Γ -martingales on \mathbb{S}^1 . The latter result also yields the existence and uniqueness of a solution to the reflected BSDE in the domain $\mathcal{D} = \{y \in \mathbb{R}^2, 1 < |y| < 2\}$, which does not possess the weak star-shape property, with zero generator and with a terminal condition satisfying $|\xi| = 1$.

Moreover, in Section 3 of [41], Picard was able to prove the existence and uniqueness of a Brownian Γ -martingale with a prescribed terminal condition in a closed hemisphere of \mathbb{S}^{d-1} , and in an even bigger domain, for a small enough T and under a smoothness assumption on the terminal condition³. The latter indicates that in a smooth Markovian or discrete path-dependent framework, under an additional smallness assumption, it may also be possible to relax the requirement of a weak star-shape property even for $d > 2$.

Finally, let us give a simple example showing that the *a priori* estimates of Proposition 2.1 are not sharp.⁴ Mimicking [40], we consider a \mathcal{F}_T -measurable random variable ν with values in $[-\alpha, \alpha]$, where $0 < \alpha < \pi/2$ is a given parameter, and let $(\theta_t, \eta_t)_{t \in [0, T]}$ be the solution of the BSDE $\theta_t = \nu - \int_t^T \eta_s dW_s$ for $t \in [0, T]$. We set $Y_t = (\cos(\theta_t), \sin(\theta_t))^\top$ for all $t \in [0, T]$ and we easily check that Y is a solution to the BSDE

$$Y_t = \xi + \int_t^T \frac{|Z_s|^2}{2} Y_s ds - \int_t^T Z_s dW_s, \quad 0 \leq t \leq T,$$

where $\xi = (\cos(\nu), \sin(\nu))^\top$ and $Z_t = (-\eta_t \sin(\theta_t), \eta_t \cos(\theta_t))^\top$. Notice that this multidimensional quadratic BSDE can also be seen as a reflected BSDE in the domain $\mathcal{D}_{\alpha, \eta, \varepsilon}$, with sufficiently small $\eta, \varepsilon > 0$, rotated by $\pi/2$. Indeed, Y lives on (rotated) \mathcal{S}_α and the reflecting term is always pointing along the outer normal vector to (rotated) \mathcal{S}_α . Recall that $\mathcal{D}_{\alpha, \eta, \varepsilon}$ satisfies the weak star-shape property and note that $d\text{Var}_t(K) = \frac{|\eta_t|^2}{2}$. Then, an application of Itô's formula to θ_t^2 yields

$$\mathbb{E}_t \left[\int_t^T d\text{Var}_s(K) \right] = \frac{1}{2} \mathbb{E}_t \left[\int_t^T |\eta_s|^2 ds \right] = \frac{1}{2} \mathbb{E}_t \left[\nu^2 - (\mathbb{E}_t \nu)^2 \right] \leq \frac{\alpha^2}{2}.$$

Moreover, the above becomes an equality for $t = 0$ and $\nu = \text{sign}(W_T)\alpha$. Then, recalling that $R_0 = 1$ for $\mathcal{D}_{\alpha, \eta, \varepsilon}$, we deduce from the John-Nirenberg inequality

³To be precise, it is assumed that the process Z , defined by $\xi = \mathbb{E}[\xi] + \int_0^T Z_s dW_s$, has sufficiently small $\int_0^T \text{ess sup}_\Omega |Z_s|^2 ds$.

⁴Note that these estimates are not needed in a Markovian or discrete path-dependent case.

that

$$\mathbb{E} \left[e^{\frac{2p}{R_0} \text{Var}_T(K)} \right] < \infty,$$

for some $p > 1$, provided $\alpha < 1$, which is weaker than the condition $\alpha < \arccos(2/3) < 1$ required by Assumption 2.1(i) with $\theta = 2$, as computed earlier in this subsection.

REFERENCES

- [1] C. Benezet, J.-F. Chassagneux, and A. Richou. Randomised switching problems and obliquely reflected BSDEs.
- [2] J.-M. Bismut. Théorie probabiliste du contrôle des diffusions. *Mem. Amer. Math. Soc.*, 4(167):xiii+130, 1976.
- [3] N. Bouleau. Formules de changement de variables. *Ann. Inst. H. Poincaré Probab. Statist.*, 20(2):133–145, 1984.
- [4] P. Briand and H. Higon. Particles systems for mean reflected BSDEs. *Stochastic Process. Appl.*, 131:253–275, 2021.
- [5] L. Campi and W. Schachermayer. A super-replication theorem in kabanov’s model of transaction costs. *Finance and Stochastics*, 10(4):579–596, 2006.
- [6] J.-F. Chassagneux, R. Elie, and I. Kharroubi. A note on existence and uniqueness for solutions of multidimensional reflected BSDEs. *Electron. Commun. Probab.*, 16:120–128, 2011.
- [7] J.-F. Chassagneux, R. Elie, and I. Kharroubi. Discrete-time approximation of multidimensional BSDEs with oblique reflections. *Ann. Appl. Probab.*, 22(3):971–1007, 2012.
- [8] J.-F. Chassagneux and A. Richou. Obliquely reflected backward stochastic differential equations. *Ann. Inst. Henri Poincaré Probab. Stat.*, 56(4):2868–2896, 2020.
- [9] J. Cvitanic and I. Karatzas. Backward stochastic differential equations with reflection and Dynkin games. *The Annals of Probability*, pages 2024–2056, 1996.
- [10] R. W. R. Darling. Constructing gamma-martingales with prescribed limit, using backwards SDE. *Ann. Probab.*, 23(3):1234–1261, 1995.
- [11] C. Dellacherie and P.-A. Meyer. *Probabilités et potentiel. Chapitres V à VIII*, volume 1385 of *Actualités Scientifiques et Industrielles [Current Scientific and Industrial Topics]*. Hermann, Paris, revised edition, 1980. Théorie des martingales. [Martingale theory].
- [12] R. Dumitrescu, M.-C. Quenez, and A. Sulem. Generalized Dynkin games and doubly reflected BSDEs with jumps. *Electronic Journal of Probability*, 21, 2016.
- [13] N. El Karoui, C. Kapoudjian, É. Pardoux, S. Peng, and M.-C. Quenez. Reflected solutions of backward SDE’s, and related obstacle problems for PDE’s. *the Annals of Probability*, pages 702–737, 1997.
- [14] N. El Karoui, S. Peng, and M. C. Quenez. Backward stochastic differential equations in finance. *Math. Finance*, 7(1):1–71, 1997.
- [15] M. Émery. *Stochastic calculus in manifolds*. Universitext. Springer-Verlag, Berlin, 1989. With an appendix by P.-A. Meyer.
- [16] M. Émery and G. Mokobodzki. Sur le barycentre d’une probabilité dans une variété. In *Séminaire de Probabilités, XXV*, volume 1485 of *Lecture Notes in Math.*, pages 220–233. Springer, Berlin, 1991.
- [17] I. Fakhouri, Y. Ouknine, and Y. Ren. Reflected backward stochastic differential equations with jumps in time-dependent random convex domains. *Stochastics*, 90(2):256–296, 2018.
- [18] R. Gayduk and S. Nadtochiy. Endogenous formation of Limit Order Books: dynamics between trades. *SICON*, 56(3):1577–1619, 2018.

- [19] R. Gayduk and S. Nadtochiy. Control-stopping games for market microstructure and beyond. *Mathematics of Operations Research*, 45(4):1193–1620, 2020.
- [20] A. Gégout-Petit and É. Pardoux. Équations différentielles stochastiques rétrogrades réfléchies dans un convexe. *Stochastics Stochastics Rep.*, 57(1-2):111–128, 1996.
- [21] M. Grigorova, P. Imkeller, E. Offen, Y. Ouknine, and M.-C. Quenez. Reflected BSDEs when the obstacle is not right-continuous and optimal stopping. *Ann. Appl. Probab.*, 27(5):3153–3188, 2017.
- [22] M. Grigorova, P. Imkeller, Y. Ouknine, and M.-C. Quenez. Doubly reflected BSDEs and \mathcal{E}^f -Dynkin games: beyond the right-continuous case. *Electron. J. Probab.*, 23:Paper No. 122, 38, 2018.
- [23] S. Hamadène, J.-P. Lepeltier, and S. Peng. BSDEs with continuous coefficients and stochastic differential games. In *Backward stochastic differential equations*, pages 115–128. Harlow: Longman, 1997.
- [24] S. Hamadène and J. Zhang. Switching problem and related system of reflected backward SDEs. *Stochastic Process. Appl.*, 120(4):403–426, 2010.
- [25] J. Harter and A. Richou. A stability approach for solving multidimensional quadratic BSDEs. *Electron. J. Probab.*, 24:Paper No. 4, 51, 2019.
- [26] Y. Hu and J. Ma. Nonlinear Feynman-Kac formula and discrete-functional-type BSDEs with continuous coefficients. *Stochastic Process. Appl.*, 112(1):23–51, 2004.
- [27] Y. Hu and S. Tang. Multi-dimensional BSDE with oblique reflection and optimal switching. *Probab. Theory Related Fields*, 147(1-2):89–121, 2010.
- [28] Y. Hu and S. Tang. Multi-dimensional backward stochastic differential equations of diagonally quadratic generators. *Stochastic Process. Appl.*, 126(4):1066–1086, 2016.
- [29] N. Kazamaki. *Continuous exponential martingales and BMO*, volume 1579 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1994.
- [30] W. S. Kendall. Probability, convexity, and harmonic maps with small image. I. Uniqueness and fine existence. *Proc. London Math. Soc. (3)*, 61(2):371–406, 1990.
- [31] W. S. Kendall. Convex geometry and nonconfluent Γ -martingales. I. Tightness and strict convexity. In *Stochastic analysis (Durham, 1990)*, volume 167 of *London Math. Soc. Lecture Note Ser.*, pages 163–178. Cambridge Univ. Press, Cambridge, 1991.
- [32] T. Klimsiak, A. Rozkosz, and L. Słomiński. Reflected BSDEs in time-dependent convex regions. *Stochastic Process. Appl.*, 125(2):571–596, 2015.
- [33] P.-L. Lions, J.-L. Menaldi, and A.-S. Sznitman. Construction de processus de diffusion réfléchis par pénalisation du domaine. *CR Acad. Sc. Paris*, 229:459–462, 1981.
- [34] P.-L. Lions and A.-S. Sznitman. Stochastic differential equations with reflecting boundary conditions. *Comm. Pure Appl. Math.*, 37(4):511–537, 1984.
- [35] R. Martyr. Finite-horizon optimal multiple switching with signed switching costs. *Math. Oper. Res.*, 41(4):1432–1447, 2016.
- [36] D. Nualart. *The Malliavin calculus and related topics*. Probability and its Applications (New York). Springer-Verlag, Berlin, second edition, 2006.
- [37] É. Pardoux and S. Peng. Backward stochastic differential equations and quasilinear parabolic partial differential equations. In *Stochastic partial differential equations and their applications (Charlotte, NC, 1991)*, volume 176 of *Lecture Notes in Control and Inform. Sci.*, pages 200–217. Springer, Berlin, 1992.
- [38] S. Peng. Backward stochastic differential equations and applications to optimal control. *Appl. Math. Optim.*, 27(2):125–144, 1993.
- [39] S. G. Peng. Probabilistic interpretation for systems of quasilinear parabolic partial differential equations. *Stochastics Stochastics Rep.*, 37(1-2):61–74, 1991.

- [40] J. Picard. Martingales sur le cercle. In *Séminaire de Probabilités, XXIII*, volume 1372 of *Lecture Notes in Math.*, pages 147–160. Springer, Berlin, 1989.
- [41] J. Picard. Martingales on Riemannian manifolds with prescribed limit. *J. Funct. Anal.*, 99(2):223–261, 1991.
- [42] J. Picard. Barycentres et martingales sur une variété. *Ann. Inst. H. Poincaré Probab. Statist.*, 30(4):647–702, 1994.
- [43] R. Tevzadze. Solvability of backward stochastic differential equations with quadratic growth. *Stochastic Process. Appl.*, 118(3):503–515, 2008.
- [44] H. Xing and G. Žitković. A class of globally solvable Markovian quadratic BSDE systems and applications. *Ann. Probab.*, 46(1):491–550, 2018.

UFR DE MATHÉMATIQUES & LPSM, UNIVERSITÉ DE PARIS.

Email address: `chassagneux@lpsm.paris`

DEPARTMENT OF APPLIED MATHEMATICS, ILLINOIS INSTITUTE OF TECHNOLOGY, CHICAGO, IL 60616.

Email address: `snadtochiy@iit.edu`

UNIVERSITÉ DE BORDEAUX, IMB, UMR 5251, F-33400 TALENCE, FRANCE.

Email address: `adrien.richou@math.u-bordeaux.fr`