



HAL
open science

A General Framework for the Derandomization of PAC-Bayesian Bounds

Paul Viallard, Pascal Germain, Amaury Habrard, Emilie Morvant

► **To cite this version:**

Paul Viallard, Pascal Germain, Amaury Habrard, Emilie Morvant. A General Framework for the Derandomization of PAC-Bayesian Bounds. 2021. hal-03143025v1

HAL Id: hal-03143025

<https://hal.science/hal-03143025v1>

Preprint submitted on 16 Feb 2021 (v1), last revised 18 Sep 2023 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PAUL VIALARD¹, PASCAL GERMAIN², AMAURY HABRARD¹, and EMILIE MORVANT¹

¹Univ Lyon, UJM-Saint-Etienne, CNRS, Institut d Optique Graduate School,
Laboratoire Hubert Curien UMR 5516, F-42023, SAINT-ETIENNE, France

²Département d’informatique et de génie logiciel, Université Laval, Québec, Canada

Abstract

PAC-Bayesian bounds are known to be tight and informative when studying the generalization ability of randomized classifiers. However, when applied to some family of deterministic models such as neural networks, they require a loose and costly derandomization step. As an alternative to this step, we introduce three new PAC-Bayesian generalization bounds that have the originality to be *pointwise*, meaning that they provide guarantees over one single hypothesis instead of the usual *averaged* analysis. Our bounds are rather general, potentially parameterizable, and provide novel insights for various machine learning settings that rely on randomized algorithms. We illustrate the interest of our theoretical result for the analysis of neural network training.

1 Introduction

PAC-Bayesian theory—introduced by [Shawe-Taylor and Williamson \(1997\)](#); [McAllester \(1998\)](#)—provides a powerful framework for analyzing tightly the generalization ability of famous machine learning models such as linear classifiers ([Germain et al., 2009](#)), SVM ([Ambroladze et al., 2006](#)) or neural networks ([Dziugaite and Roy, 2017](#)). PAC-Bayesian analyses take usually the form of bounds on the average risk of a randomized classifier with respect to a learned posterior distribution given a chosen prior distribution defined over a set of hypotheses (before observing the learning sample). While such bounds are very effective for analyzing stochastic classifiers, some machine learning approaches need nevertheless guarantees on deterministic models. Then, in this situation, a *derandomization step* of the bound is required. Different forms of derandomization have been introduced in the literature for specific settings. Among them, [Langford and Shawe-Taylor \(2002\)](#); [Langford \(2005\)](#); [Germain et al. \(2009\)](#) propose a derandomization step for linear classifiers; it consists in finding a closed-form solution for the average risk of the randomized linear classifiers from which a bound on the risk of a new deterministic classifier is deduced. Hence, for this classifier, the bound is not randomized anymore. From a similar analysis [Letarte](#)

[et al. \(2019\)](#) present a PAC-Bayes bound for a very specific deterministic network architecture using sign functions as activations. Another line of works derandomizes neural networks ([Neyshabur et al., 2018](#); [Nagarajan and Kolter, 2019a](#)). While being technically different, it consists of starting from PAC-Bayesian guarantees on the randomized classifier and use an “output perturbation” bound to convert a guarantee from a random classifier to the mean classifier.

Above all, these works mainly highlight the need of a general PAC-Bayesian framework for the derandomization of classic PAC-Bayesian bounds. In this paper, our objective is thus twofold: *(i)* to derive new PAC-Bayesian bounds that directly derandomize (any type of classifiers) without any other additional step, *(ii)* to obtain a derandomization with (almost) no impact on the guarantee. To do so, we derive three novels derandomized bounds based on a so-called *pointwise* approach: instead of bounding the *average risk of a randomized classifier* with respect to the posterior distribution, our analyses bound the *risk of a sampled* (unique) classifier from the posterior distribution. Note that there are pointwise bounds that have been introduced as an intermediate step to derive PAC-Bayesian bounds ([Blanchard and Fleuret, 2007](#); [Rivasplata et al., 2020](#)). To the best of our knowledge, our work appears to be the first tentative to provide a general framework for the derandomization of PAC-Bayesian bounds with a new strategy: *(a)* we sample the classifier according to a posterior distribution, *(b)* we provide a PAC-Bayesian guarantee on the sampled classifier.

The paper is organized as follows. Section 2 introduces the notations we follow, and recalls some basics on generalization bounds. In Section 3, we derive our main contribution that relies on three *pointwise* PAC-Bayesian bounds. Then, we illustrate the derandomization on deterministic neural networks in Section 4. We conclude in Section 5.

2 Setting and Basics

General Notations. We stand in a supervised classification setting, where \mathcal{X} is the *input space*, \mathcal{Y} is the *label set*, and \mathcal{D} is an unknown *data distribution* over $\mathcal{X} \times \mathcal{Y} = \mathcal{Z}$. An *example* is denoted by $z = (\mathbf{x}, y) \in \mathcal{Z}$, and the *learning sample* by $\mathcal{S} = \{z_i\}_{i=1}^m$ constituted by m examples drawn *i.i.d.*

from \mathcal{D} ; the distribution of such a m -sample being \mathcal{D}^m . We consider a *hypothesis set* \mathcal{H} of functions $h : \mathcal{X} \rightarrow \mathcal{Y}$. The learner aims to find $h : \mathcal{X} \rightarrow \mathcal{Y}$ from \mathcal{H} that assigns a label y to a new input x as correctly as possible. Given an example $z \in \mathcal{Z}$ and a hypothesis $h \in \mathcal{H}$, we assess the quality of the prediction of h with a *loss function* $\ell : \mathcal{H} \times \mathcal{Z} \rightarrow [0, 1]$ that evaluates to which extent the prediction is accurate. From the loss ℓ , we define the *true risk* $R_{\mathcal{D}}(h)$ of $h \in \mathcal{H}$, which corresponds to the expected loss over the data distribution \mathcal{D} , and its empirical counterpart $R_{\mathcal{S}}(h)$ estimated on \mathcal{S} as

$$R_{\mathcal{D}}(h) \triangleq \mathbb{E}_{z \sim \mathcal{D}} \ell(h, z), \quad \text{and} \quad R_{\mathcal{S}}(h) \triangleq \frac{1}{m} \sum_{i=1}^m \ell(h, z_i).$$

Then, we want to find the hypothesis from \mathcal{H} that minimizes the true risk on the distribution \mathcal{D} . However, this value is not computable since we do not know the distribution \mathcal{D} . In practice we can deal with the Empirical Risk Minimization (ERM) principle that consists in finding a hypothesis h that minimizes $R_{\mathcal{S}}(h)$. Nevertheless, we have no guarantee that such h will generalize well on new unseen data from \mathcal{D} . We thus need to bring theoretical guarantees to quantify how much the empirical risk is a good estimator of the true risk. The statistical machine learning theory (e.g., Vapnik, 2000) studies the conditions of consistency and therefore convergence of ERM towards the true value. This kind of result is usually called generalization bounds, often referred to as PAC (Probably Approximately Correct) bounds (Valiant, 1984). These bounds take the following form

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m} \left[|R_{\mathcal{D}}(h) - R_{\mathcal{S}}(h)| \leq \epsilon \left(\frac{1}{\delta}, \frac{1}{m}, \mathcal{H} \right) \right] \geq 1 - \delta.$$

Put into words, with high probability $(1-\delta)$ on the random choice of \mathcal{S} , we obtain good generalization properties when the deviation between the true risk $R_{\mathcal{D}}(h)$ and its empirical estimate $R_{\mathcal{S}}(h)$ is low, i.e., $\epsilon \left(\frac{1}{\delta}, \frac{1}{m}, \mathcal{H} \right)$ should be as small as possible. The function ϵ depends on three entities: (i) the number of examples m , the more we have, the more precise $R_{\mathcal{S}}(h)$ is, (ii) the hypothesis set \mathcal{H} with the idea that some specificities of it play a role in its generalization power, (iii) the parameter confidence δ , the most probable is the bound (δ close to 0), the larger is ϵ . We now recall three families of such bounds with a strong emphasis on the PAC-Bayes theory at the heart of our contributions.

Uniform Convergence Bound. A first classical type of generalization bounds is referred as *Uniform Convergence* bounds that are based on a measure of complexity of the hypothesis set \mathcal{H} and stand for all the hypotheses of \mathcal{H} . Among the most renowned complexity measure, we can cite the VC-dimension or the Rademacher Complexity. This type of bound usually takes the form

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m} \left[\sup_{h \in \mathcal{H}} |R_{\mathcal{D}}(h) - R_{\mathcal{S}}(h)| \leq \epsilon \left(\frac{1}{\delta}, \frac{1}{m}, \mathcal{H} \right) \right] \geq 1 - \delta.$$

This bound holds for all $h \in \mathcal{H}$, including the best, but also the worst. This *worst-case* analysis makes hard to obtain a non-vacuous bound with $\epsilon \left(\frac{1}{\delta}, \frac{1}{m}, \mathcal{H} \right) < 1$. The ability of such bounds to explain the generalization of deep learning has been recently challenged (Nagarajan and Kolter, 2019b).

Algorithmic-Dependent Bound. Other bounds depend on the learning algorithm (Bousquet and Elisseeff, 2002; Xu and Mannor, 2012). These bounds directly involve some particularities of the learning algorithm A (and then \mathcal{H}). This allows obtaining bounds that stand for a single hypothesis $h_{A(\mathcal{S})}$, the one learned with A from the learning sample \mathcal{S} . The form of such bounds is

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m} \left[|R_{\mathcal{D}}(h_{A(\mathcal{S})}) - R_{\mathcal{S}}(h_{A(\mathcal{S})})| \leq \epsilon \left(\frac{1}{\delta}, \frac{1}{m}, A \right) \right] \geq 1 - \delta.$$

For example, this approach has been used by Hardt et al. (2016) to derive generalization bounds for hypotheses learned by stochastic gradient descent.

PAC-Bayesian Bound. This paper leverages on PAC-Bayes bounds that stand in the PAC framework but borrow inspiration from the Bayesian probabilistic point of view that deals with randomness and uncertainty in machine learning (McAllester, 1998). That being said, considering the set $\mathcal{M}(\mathcal{H})$ of probability measures on \mathcal{H} , a PAC-Bayesian generalization bound provides a bound in expectation/average over the hypothesis set \mathcal{H} and involves a *prior* distribution $\mathcal{P} \in \mathcal{M}(\mathcal{H})$ on \mathcal{H} and a *posterior* $\mathcal{Q} \in \mathcal{M}(\mathcal{H})$ on \mathcal{H} learned after the observation of \mathcal{S} . The form of such a bound is

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m} \left[\forall \mathcal{Q}, \mathbb{E}_{h \sim \mathcal{Q}} |R_{\mathcal{D}}(h) - R_{\mathcal{S}}(h)| \leq \epsilon \left(\frac{1}{\delta}, \frac{1}{m}, \mathcal{Q} \right) \right] \geq 1 - \delta.$$

We recall below the classical PAC-Bayesian theorem in a slightly different form from the usual one: from Theorem 1, $\phi(h, \mathcal{S}) = \exp(\phi(h, \mathcal{S}))$ gives the usual form of the bounds.

Theorem 1 (General PAC-Bayesian Bounds). *For any data distribution \mathcal{D} on \mathcal{Z} , for any set \mathcal{H} of hypotheses $\mathcal{X} \rightarrow \mathcal{Y}$, for any prior distribution \mathcal{P} on \mathcal{H} , for any function $\phi : \mathcal{H} \times \mathcal{Z}^m \rightarrow \mathbb{R}^+$, we have*

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m} \left[\forall \mathcal{Q}, \mathbb{E}_{h \sim \mathcal{Q}} \ln(\phi(h, \mathcal{S})) \leq \right. \quad \text{(Germain et al., 2009)}$$

$$\left. \text{KL}(\mathcal{Q} \parallel \mathcal{P}) + \ln \left[\frac{1}{\delta} \mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathbb{E}_{h \sim \mathcal{P}} \phi(h, \mathcal{S}) \right] \right] \geq 1 - \delta, \quad (2.1)$$

$$\text{and } \mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m} \left[\forall \mathcal{Q}, \frac{\alpha}{\alpha-1} \ln \left(\mathbb{E}_{h \sim \mathcal{Q}} \phi(h, \mathcal{S}) \right) \leq \right. \quad \text{(Bégin et al., 2016)}$$

$$\left. D_{\alpha}(\mathcal{Q} \parallel \mathcal{P}) + \ln \left[\frac{1}{\delta} \mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathbb{E}_{h \sim \mathcal{P}} \phi(h, \mathcal{S})^{\frac{\alpha}{\alpha-1}} \right] \right] \geq 1 - \delta, \quad (2.2)$$

where $\text{KL}(\mathcal{Q} \parallel \mathcal{P}) \triangleq \mathbb{E}_{h \sim \mathcal{Q}} \ln \frac{\mathcal{Q}(h)}{\mathcal{P}(h)}$ is the *KL-divergence*, $D_{\alpha}(\mathcal{Q} \parallel \mathcal{P}) \triangleq \frac{1}{\alpha-1} \ln \left(\mathbb{E}_{h \sim \mathcal{P}} \left[\frac{\mathcal{Q}(h)}{\mathcal{P}(h)} \right]^{\alpha} \right)$ is the *Rényi divergence* (with $\alpha > 1$).

Note that Equation (2.2) is more general than Equation (2.1) since the Rényi divergence can be seen as a generalization of the KL-divergence: when α tends to 1, then $D_\alpha(\mathcal{Q}||\mathcal{P})$ tends to $\text{KL}(\mathcal{Q}||\mathcal{P})$. The advantage of these bounds is that they are rather general since they can be seen as the starting point of the derivation of different forms of bound by, firstly, instantiating the function ϕ to capture a deviation between the true risk and its empirical counterpart, secondly, upper-bounding the term $\mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathbb{E}_{h \sim \mathcal{P}} \phi(h, \mathcal{S})$. For instance, with $\phi(h, \mathcal{S}) = e^{m2[R_{\mathcal{D}}(h) - R_{\mathcal{S}}(h)]^2}$ we retrieve from Equation (2.1) the form proposed by McAllester (1998):

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m} \left[\forall \mathcal{Q}, \mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{D}}(h) \leq \mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{S}}(h) + \sqrt{\frac{1}{2m} \left[\text{KL}(\mathcal{Q}||\mathcal{P}) + \ln \frac{2\sqrt{m}}{\delta} \right]} \right] \geq 1 - \delta. \quad (2.3)$$

This bound has the advantage to easily illustrate the trade-off between the average empirical risk and the bound. Another example, that leads to a slightly tighter bound but less interpretable, is the Seeger (2002)'s bound that we can retrieve from Equation (2.1) with $\phi(h, \mathcal{S}) = e^{m \text{kl}[R_{\mathcal{S}}(h)||R_{\mathcal{D}}(h)]}$ where $\text{kl}[q||p] = q \ln \frac{q}{p} + (1-q) \ln \frac{1-q}{1-p}$:

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m} \left[\forall \mathcal{Q}, \mathbb{E}_{h \sim \mathcal{Q}} \text{kl}(R_{\mathcal{S}}(h)||R_{\mathcal{D}}(h)) \leq \frac{1}{m} \left[\text{KL}(\mathcal{Q}||\mathcal{P}) + \ln \frac{2\sqrt{m}}{\delta} \right] \right] \geq 1 - \delta. \quad (2.4)$$

Such PAC-Bayesian bounds are known to be tight, but they stand for a randomized classifier by nature (due to the expectation over \mathcal{H}). A key issue for usual machine learning tasks is then the derandomization of the bounds to obtain a guarantee for a deterministic classifier. In some cases, this derandomization is a result of the structure of the hypotheses such as for randomized linear classifier that can be directly expressed as one deterministic linear classifier (Germain et al., 2009). However, in other cases the derandomization is much more complex and specific to a class of hypothesis, such as in the case of neural networks; see, e.g., Neyshabur et al. (2018) or Nagarajan and Kolter (2019b, Appendix J).

In the next section, we present our contribution: a derandomization of the general PAC-Bayesian bounds based on a pointwise analysis to obtain bounds that stands for a single hypothesis from \mathcal{H} . Our approach provides a general framework not specific to particular families of hypotheses.

3 Three Pointwise PAC-Bayesian Theorems

We introduce a new kind of generalization bounds: the pointwise PAC-Bayesian bound that allows one to derandomize

the usual PAC-Bayesian bounds presented above. The form of such bounds can be expressed as follows:

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m, h \sim \mathcal{Q}_{\mathcal{S}}} \left[|R_{\mathcal{D}}(h) - R_{\mathcal{S}}(h)| \leq \epsilon \left(\frac{1}{\delta}, \frac{1}{m}, \mathcal{Q}_{\mathcal{S}} \right) \right] \geq 1 - \delta.$$

Here, we consider that the posterior distribution $\mathcal{Q}_{\mathcal{S}}$ is obtained through a deterministic algorithm taking the learning sample \mathcal{S} as input while the classic PAC-Bayesian bound holds for all posterior distribution over \mathcal{H} . More formally, we consider a *deterministic* algorithm $A: \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$ chosen *a priori* which (i) takes a learning sample $\mathcal{S} \in \mathcal{Z}^m$ as input and (ii) outputs a *data-dependent* distribution $\mathcal{Q}_{\mathcal{S}} \triangleq A(\mathcal{S})$ from the set $\mathcal{M}(\mathcal{H})$ of all the possible probability measures on \mathcal{H} . The above bound stands then for a unique hypothesis $h \sim \mathcal{Q}_{\mathcal{S}}$ instead of a randomized classifier: the individual risks are no longer averaged with respect to $\mathcal{Q}_{\mathcal{S}}$; this is our *pointwise-based derandomization*. The dependence in probability on $\mathcal{Q}_{\mathcal{S}}$ means that the bound is valid for any hypothesis drawn from $\mathcal{Q}_{\mathcal{S}}$. In practice, after learning the distribution $\mathcal{Q}_{\mathcal{S}}$ with an algorithm, then the unique hypothesis h is sampled from $\mathcal{Q}_{\mathcal{S}}$. In some sense this type of bounds is at the crossroads between the algorithmic-dependent bound and the PAC-Bayesian one recalled in Section 2. Under this principle, we present in the following three general pointwise PAC-Bayesian bounds (Theorem 2, and 3, and 6).

3.1 Pointwise Bounds Involving the Rényi Divergence

In the same spirit as Equation (2.2) our first result stated below can be seen as a general bound depending on the Rényi divergence $D_\alpha(\mathcal{Q}_{\mathcal{S}}||\mathcal{P})$ of order $\alpha > 1$.

Theorem 2 (Pointwise PAC-Bayesian Bound). *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any prior distribution \mathcal{P} on \mathcal{H} , for any function $\phi: \mathcal{H} \times \mathcal{Z}^m \rightarrow \mathbb{R}^+$, for any $\alpha > 1$, for any deterministic algorithm $A: \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, we have*

$$\mathbb{P}_{\substack{\mathcal{S} \sim \mathcal{D}^m \\ h \sim \mathcal{Q}_{\mathcal{S}}}} \left[\frac{\alpha}{\alpha-1} \ln(\phi(h, \mathcal{S})) \leq D_\alpha(\mathcal{Q}_{\mathcal{S}}||\mathcal{P}) + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}+1}} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha}{\alpha-1}} \right) \right] \geq 1 - 2\delta,$$

where $\mathcal{Q}_{\mathcal{S}}$ is output by A , i.e., $\mathcal{Q}_{\mathcal{S}} \triangleq A(\mathcal{S})$.

Proof Sketch (see Appendix for details). First, note that we obtain $\mathcal{Q}_{\mathcal{S}}$ after having seen the sample \mathcal{S} with algorithm A . Then, applying Markov inequality on $\phi(h, \mathcal{S})$ with the r.v. h and using Hölder inequality to introduce $D_\alpha(\mathcal{Q}_{\mathcal{S}}||\mathcal{P})$, we have with prob. at least $1 - \delta$ on $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_{\mathcal{S}}$

$$\begin{aligned} \frac{\alpha}{\alpha-1} \ln(\phi(h, \mathcal{S})) &\leq \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) \right) \\ &\leq D_\alpha(\mathcal{Q}_{\mathcal{S}}||\mathcal{P}) + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha}{\alpha-1}} \right). \end{aligned}$$

By applying again Markov inequality on $\phi(h, \mathcal{S})$ with the random variable \mathcal{S} , we have with *prob.* at least $1-\delta$ on $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_\mathcal{S}$

$$\begin{aligned} & \mathbb{P}_{\substack{\mathcal{S} \sim \mathcal{D}^m \\ h \sim \mathcal{Q}_\mathcal{S}}} \left[\ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha}{\alpha-1}} \right) \right] \\ & \leq \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1} + 1}} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha}{\alpha-1}} \right). \end{aligned}$$

By combining the two bounds with a union bound argument, we obtain the desired result. \square

Compared to Equation (2.2), our bound requires an additional term $-\ln \delta^{\frac{\alpha}{\alpha-1}}$. However, by taking $\phi(h, \mathcal{S}) = m \text{kl}(R_\mathcal{S}(h) \| R_\mathcal{D}(h))$ and $\alpha = 2$, the term $\ln \frac{1}{\delta^2}$ will be multiplied by $\frac{1}{m}$, allowing us to obtain a reasonable overhead to “derandomize” a bound into a pointwise one. For instance, if $m=5,000$ (reasonable sample size) and $2\delta = .05$, we have $\frac{1}{m} \ln \frac{1}{\delta^2} \approx .001$.

Note that [Rivasplata et al. \(2020, Th. 1\(ii\)\)](#) introduced the following bound that holds with probability at least $1 - \delta$ over the random choice of $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_\mathcal{S}$

$$\ln(\phi(h, \mathcal{S})) \leq \ln \left[\frac{\mathcal{Q}_\mathcal{S}(h)}{\mathcal{P}(h)} \right] + \ln \left(\frac{1}{\delta} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}') \right). \quad (3.1)$$

Even if Equation (3.1) has a better dependence in δ , the advantage of our bound is that it involves a divergence (*i.e.*, the Rényi divergence $D_\alpha(\mathcal{Q}_\mathcal{S} \| \mathcal{P})$). Indeed, their term $\ln \frac{\mathcal{Q}_\mathcal{S}(h)}{\mathcal{P}(h)}$ depends only the sampled hypothesis $h \sim \mathcal{Q}_\mathcal{S}$. Hence, depending on the hypothesis $h \sim \mathcal{Q}_\mathcal{S}$ and $\alpha > 1$, their term can be larger (or smaller) than $D_\alpha(\mathcal{Q}_\mathcal{S} \| \mathcal{P})$ which remains constant.

In the PAC-Bayes literature, parametrized bounds have been introduced (*e.g.*, [Catoni \(2007\)](#); [Thiemann et al. \(2017\)](#)) to control the trade-off between the empirical risk and the divergence along with the additional term (which can be useful from an algorithmic point of view). We now follow a similar approach to introduce a version of a pointwise Rényi divergence-based bound that has the advantage to be parametrizable.

Theorem 3 (Parametrizable Pointwise PAC-Bayesian Bound). *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any prior distribution \mathcal{P} on \mathcal{H} , for any function $\phi : \mathcal{H} \times \mathcal{Z}^m \rightarrow \mathbb{R}^+$, for any deterministic algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, we have*

$$\begin{aligned} & \mathbb{P}_{\substack{\mathcal{S} \sim \mathcal{D}^m \\ h \sim \mathcal{Q}_\mathcal{S}}} \left[\forall \lambda > 0, \ln(\phi(h, \mathcal{S})) \leq \ln \left(\frac{1}{2} \left[\lambda \exp[D_2(\mathcal{Q}_\mathcal{S} \| \mathcal{P})] \right. \right. \right. \\ & \quad \left. \left. \left. + \frac{1}{\lambda \delta^3} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^2 \right] \right) \right] \geq 1 - 2\delta, \end{aligned}$$

where $\mathcal{Q}_\mathcal{S} \triangleq A(\mathcal{S})$.

Proof Sketch (see Appendix for details). Applying Markov’s inequality on $\phi(h, \mathcal{S})$, and introducing the Rényi divergence ($\alpha=2$) with the fact that $\forall \lambda > 0, \sqrt{ab} \leq \frac{1}{2}(\lambda a + \frac{b}{\lambda})$, we have with *prob.* at least $1-\delta$ on $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_\mathcal{S} \triangleq A(\mathcal{S})$,

$$\begin{aligned} \forall \lambda > 0, \ln[\phi(h, \mathcal{S})] & \leq \ln \left[\frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_\mathcal{S}} \phi(h', \mathcal{S}) \right] \\ & \leq \ln \left[\frac{\lambda}{2} \exp[D_2(\mathcal{Q}_\mathcal{S} \| \mathcal{P})] + \mathbb{E}_{h' \sim \mathcal{P}} \frac{\phi(h', \mathcal{S})^2}{2\lambda\delta^2} \right]. \end{aligned}$$

From applying again Markov’s inequality on $\phi(h, \mathcal{S})$ with the random variable $\mathcal{S} \sim \mathcal{D}^m$, we have with *prob.* at least $1-\delta$ on $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_\mathcal{S}$,

$$\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}) \leq \frac{1}{\delta} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')$$

which is equivalent to

$$\begin{aligned} \forall \lambda > 0, \ln \left[\frac{\lambda}{2} \exp[D_2(\mathcal{Q}_\mathcal{S} \| \mathcal{P})] + \mathbb{E}_{h' \sim \mathcal{P}} \frac{\phi(h', \mathcal{S})^2}{2\lambda\delta^2} \right] \\ \leq \ln \left[\frac{\lambda}{2} \exp[D_2(\mathcal{Q}_\mathcal{S} \| \mathcal{P})] + \mathbb{E}_{\substack{\mathcal{S}' \sim \mathcal{D}^m \\ h' \sim \mathcal{P}}} \frac{\phi(h', \mathcal{S}')^2}{2\lambda\delta^3} \right]. \end{aligned}$$

The result is obtained by combining the two bounds with a union bound argument. \square

Note that the term $\exp[D_2(\mathcal{Q}_\mathcal{S} \| \mathcal{P})]$ is closely related to the χ -squared divergence since

$$\chi^2(\mathcal{Q}_\mathcal{S} \| \mathcal{P}) = \mathbb{E}_{h \sim \mathcal{P}} \left[\left(\frac{\mathcal{Q}_\mathcal{S}(h)}{\mathcal{P}(h)} \right)^2 - 1 \right] = \exp[D_2(\mathcal{Q}_\mathcal{S} \| \mathcal{P})] - 1.$$

An asset of Theorem 3 is the parameter λ that controls the trade-off between the exponentiated Rényi divergence $\exp[D_2(\mathcal{Q}_\mathcal{S} \| \mathcal{P})]$ and the term $\frac{1}{\delta^3} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^2$. A key point is that our bound is valid for all $\lambda > 0$, meaning that from a practical point of view one can learn the parameter to minimize the bound (and control the possible numerical instability due to $\exp[D_2(\mathcal{Q}_\mathcal{S} \| \mathcal{P})]$). Despite the algorithmic advantage of this bound, for a given \mathcal{P} and $\mathcal{Q}_\mathcal{S}$, we show in Proposition 4 that the optimal bound of Theorem 3 corresponds to the bound of Theorem 2.

Proposition 4. *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any prior distribution \mathcal{P} on \mathcal{H} and for any deterministic algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, let*

$$\begin{aligned} \lambda^* = \operatorname{argmin}_{\lambda > 0} \ln \left(\frac{\lambda}{2} \exp[D_2(\mathcal{Q}_\mathcal{S} \| \mathcal{P})] \right. \\ \left. + \frac{1}{2\lambda\delta^3} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^2 \right), \end{aligned}$$

$$\begin{aligned}
 \text{then } & \overbrace{2 \ln \left[\frac{\lambda^*}{2} \exp(D_2(Q_S \| \mathcal{P})) + \frac{\mathbb{E}_{\substack{S' \sim \mathcal{D}^m \\ h' \sim \mathcal{P}}} \phi(h', S')^2}{2\lambda^* \delta^3} \right]}^{\text{Theorem 3}} \\
 & = \underbrace{D_2(Q_S \| \mathcal{P}) + \ln \left[\frac{1}{\delta^3} \mathbb{E}_{\substack{S' \sim \mathcal{D}^m \\ h' \sim \mathcal{P}}} \phi(h', S')^2 \right]}_{\text{Theorem 2 with } \alpha = 2},
 \end{aligned}$$

$$\text{where } \lambda^* = \sqrt{\frac{\mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^2}{\delta^3 \exp(D_2(Q_S \| \mathcal{P}))}}.$$

Proof. The proof is deferred in Appendix. \square

In other words, Proposition 4 tells us that the optimal λ^* gives the same bound for Theorem 2 and Theorem 3. In the next subsection, we provide another result giving a new derandomization perspective.

3.2 Bound with Sibson's Mutual Information

The Rényi divergence between \mathcal{P} and \mathcal{Q} is sensitive to the choice of the learning sample \mathcal{S} : when the posterior \mathcal{Q} —learned from \mathcal{S} —differs greatly from the prior \mathcal{P} the divergence will be high. To avoid such behavior, we derive below a bound that depends on the Sibson's mutual information that involves an expectation over all the learning samples of a given size m , and measures the dependence between the random variables $\mathcal{S} \in \mathcal{Z}^m$ and $h \in \mathcal{H}$.

Definition 5 (Sibson's Mutual Information (Verdú, 2015)). *Given $\alpha > 1$, the Sibson's mutual information between the random variable $\mathcal{S} \in \mathcal{Z}^m$ and $h \in \mathcal{H}$ is*

$$I_\alpha(h; \mathcal{S}) \triangleq \min_{\mathcal{P} \in \mathcal{M}(\mathcal{H})} \frac{1}{\alpha-1} \ln \left[\mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathbb{E}_{h \sim \mathcal{P}} \left[\frac{Q_S(h)}{\mathcal{P}(h)} \right]^\alpha \right].$$

The higher $I_\alpha(h; \mathcal{S})$, the higher the correlation is, meaning that the sampling of h is highly dependent on the choice of the learning sample. Note that two interesting properties of the Sibson's mutual information lie in that it generalizes the mutual information (Verdú, 2015) and that it can be related to the Rényi divergence. Indeed, let $\rho(h, \mathcal{S}) = Q_S(h) \times \mathcal{D}^m(\mathcal{S})$ be the probability of sampling both the learning sample $\mathcal{S} \sim \mathcal{D}^m$ and the classifier $h \sim Q_S$, and the prior distribution $\pi(h, \mathcal{S}) = \mathcal{P}(h) \times \mathcal{D}^m(\mathcal{S})$ be the probability to sample both $\mathcal{S} \sim \mathcal{D}^m$ and the hypothesis $h \sim \mathcal{P}$. Then we can write:

$$\begin{aligned}
 I_\alpha(h; \mathcal{S}) &= \min_{\mathcal{P} \in \mathcal{M}(\mathcal{H})} \frac{1}{\alpha-1} \ln \left[\mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathbb{E}_{h \sim \mathcal{P}} \left[\frac{Q_S(h)}{\mathcal{P}(h)} \right]^\alpha \right] \\
 &= \min_{\mathcal{P} \in \mathcal{M}(\mathcal{H})} \frac{1}{\alpha-1} \ln \left[\mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathbb{E}_{h \sim \mathcal{P}} \left[\frac{Q_S(h) \mathcal{D}^m(\mathcal{S})}{\mathcal{P}(h) \mathcal{D}^m(\mathcal{S})} \right]^\alpha \right] \\
 &= \min_{\mathcal{P} \in \mathcal{M}(\mathcal{H})} D_\alpha(\rho \| \pi). \tag{3.2}
 \end{aligned}$$

From Verdú (2015) the optimal prior \mathcal{P}^* which minimizes Equation (3.2) is a *distribution-dependent* prior defined as

$$\mathcal{P}^*(h) = \frac{[\mathbb{E}_{S' \sim \mathcal{D}^m} Q_{S'}(h)^\alpha]^\frac{1}{\alpha}}{\mathbb{E}_{h' \sim \mathcal{P}} \frac{1}{\mathcal{P}(h')}} [\mathbb{E}_{S' \sim \mathcal{D}^m} Q_{S'}(h')^\alpha]^\frac{1}{\alpha}}.$$

This dependence's measure leads to a so-called *Information-Theoretic generalization bound* Xu and Raginsky (2017).

Theorem 6 (Pointwise Information-Theoretic Bound). *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any function $\phi : \mathcal{H} \times \mathcal{Z}^m$, for any $\alpha > 1$, for any deterministic algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, we have*

$$\begin{aligned}
 \mathbb{P}_{\substack{\mathcal{S} \sim \mathcal{D}^m \\ h \sim Q_S}} \left[\frac{\alpha}{\alpha-1} \ln(\phi(h, \mathcal{S})) \leq I_\alpha(h'; S') \right. \\
 \left. + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^{\frac{\alpha}{\alpha-1}} \right) \right] \geq 1 - \delta,
 \end{aligned}$$

where $Q_S \triangleq A(\mathcal{S})$.

Proof. The proof is deferred in Appendix. \square

From a theoretical viewpoint, Theorem 6 brings a different philosophy than the bounds of Section 3.1. Indeed, in Theorems 2 and 3, given a learning sample \mathcal{S} , the Rényi divergence term $D_\alpha(Q_S \| \mathcal{P})$ suggests that the learned distribution Q_S should be close enough to the prior \mathcal{P} in order to obtain a low bound. While in Theorem 6, the Sibson's Mutual Information $I_\alpha(h'; S')$ suggests that the random variable h has to be “not too much correlated” to the learning sample on average. However, the bound of Theorem 6 is not computable in practice due notably to the presence of the sample expectation over the unknown distribution \mathcal{D} in I_α . We find this result nevertheless interesting by the different interpretation it provides.

Esposito et al. (2020, Corollary 4) also introduced a bound involving the Sibson's mutual information. Their bound (with our notations) holds with probability at least $1 - \delta$:

$$2(R_S(h) - R_{\mathcal{D}}(h))^2 \leq \frac{1}{m} \left[I_\alpha(h'; S') + \ln \frac{2}{\delta^{\frac{\alpha}{\alpha-1}}} \right]. \tag{3.3}$$

For the sake of comparison, we introduce the following corollary of Theorem 6.

Corollary 7. *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any deterministic algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, with probability at least $1 - \delta$ over the random choice of $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim Q_S$, we have*

$$2(R_S(h) - R_{\mathcal{D}}(h))^2 \leq \frac{1}{m} \left[I_\alpha(h'; S') + \ln \frac{2\sqrt{m}}{\delta^{\frac{\alpha}{\alpha-1}}} \right], \tag{3.4}$$

$$\text{and } \text{kl}(R_S(h) \| R_{\mathcal{D}}(h)) \leq \frac{1}{m} \left[I_\alpha(h'; S') + \ln \frac{2\sqrt{m}}{\delta^{\frac{\alpha}{\alpha-1}}} \right]. \tag{3.5}$$

Equation (3.4) is a slightly looser version of Equation (3.3) since it involves an extra term of $\frac{\ln \sqrt{m}}{m}$. However, Equation (3.5) is tighter than Equation (3.3) when $\text{kl}(R_S(h)\|R_D(h)) - 2(R_S(h) - R_D(h))^2 \geq \frac{\ln \sqrt{m}}{m}$ (which becomes more frequent as m grows). Other bounds involving the mutual information instead of the Sibson’s mutual information have been proposed in the literature (see, e.g., Alabdulmohsin, 2017; Bassily et al., 2018).

4 The Derandomization in Action

So far we have introduced theoretical results allowing to derandomize PAC-Bayesian bounds. These bounds are rather general and can be instantiated to different settings. Among them, we propose to illustrate the usefulness of Theorem 2 on neural networks.

4.1 Specialization to Neural Network Classifiers

In this section, we consider neural networks parametrized by a weight vector $\mathbf{w} \in \mathbb{R}^d$ and overparametrized, i.e., $d \gg m$. We aim at learning the weights of the network that leads to the lowest true risk. To do so, practitioners usually proceed by epochs¹ and obtain one “intermediate” network after each epoch. Then, they select the best “intermediate” network associated with the lowest validation risk. We translate this practice into our PAC-Bayesian setting by considering one prior per epoch. Given T epochs, we hence have T priors $\mathbf{P} = \{\mathcal{P}_t\}_{t=1}^T$, where $\forall t \in \{1, \dots, T\}$, $\mathcal{P}_t = \mathcal{N}(\mathbf{v}_t, \sigma^2 \mathbf{I}_d)$ is a Gaussian distribution centered at \mathbf{v}_t the weight vector associated with the t -th “intermediate” network, with a covariance matrix of $\sigma^2 \mathbf{I}_d$ (where \mathbf{I}_d is the $d \times d$ -dimensional identity matrix). Assuming the T priors are learned from a set $\mathcal{S}_{\text{prior}}$ such that $\mathcal{S}_{\text{prior}} \cap \mathcal{S} = \emptyset$, then Corollaries 8 and 9 will guide us to learn a posterior $\mathcal{Q}_S = \mathcal{N}(\mathbf{w}, \sigma^2 \mathbf{I}_d)$ from the best prior on \mathcal{S} (we give more details on the procedure after the forthcoming corollaries).

Corollary 8 below is obtained by instantiating Theorem 2 to this setting. For the sake of comparison, we also instantiate Equation (3.1) (Rivasplata et al., 2020, Th. 1(i)) in Corollary 9. Each corollary states two bounds: a Seeger-like bound and a bound that depends on a parameter $C > 0$; the latter will serve for optimization purposes in Section 4.3, the former one being in general tighter than the latter one but harder to optimize. Indeed, from Germain et al. (2009, Prop. 2.1) and Letarte et al. (2019, Th. 3), we know that such parametrized bounds lead to the same bound values as Seeger-like bounds when C is optimal.

Corollary 8. *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any priors set $\mathbf{P} = \{\mathcal{P}_t\}_{t=1}^T$ of T prior \mathcal{P} on \mathcal{H} , for any algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, for any loss*

¹One epoch corresponds to one forward and backward pass of the entire learning set in the network.

$\ell : \mathcal{H} \times \mathcal{Z} \rightarrow [0, 1]$, with proba. at least $1 - 2\delta$ on the random choice of $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_S$, we have for all prior $\mathcal{P}_t \in \mathbf{P}$

$$\text{kl}(R_S(h)\|R_D(h)) \leq \frac{1}{m} \left[\frac{\|\mathbf{w} - \mathbf{v}_t\|_2^2}{\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta^3} \right], \quad (4.1)$$

and $\forall C > 0$,

$$R_D(h) \leq \frac{1 - e^{-CR_S(h) - \frac{1}{m} \left[\frac{\|\mathbf{w} - \mathbf{v}_t\|_2^2}{\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta^3} \right]}}{1 - e^{-C}}. \quad (4.2)$$

Corollary 9. *Under the same assumptions as in Corollary 8, with proba. at least $1 - \delta$ over the random choice of $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_S$, we have for all prior $\mathcal{P}_t \in \mathbf{P}$*

$$\text{kl}(R_S(h)\|R_D(h)) \leq \frac{1}{m} \left[\frac{\|\mathbf{w} + \boldsymbol{\epsilon} - \mathbf{v}_t\|_2^2 - \|\boldsymbol{\epsilon}\|_2^2}{2\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta} \right], \quad (4.3)$$

and $\forall C > 0$,

$$R_D(h) \leq \frac{1 - e^{-CR_S(h) - \frac{1}{m} \left[\frac{\|\mathbf{w} + \boldsymbol{\epsilon} - \mathbf{v}_t\|_2^2 - \|\boldsymbol{\epsilon}\|_2^2}{2\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta} \right]}}{1 - e^{-C}} \quad (4.4)$$

where $\boldsymbol{\epsilon} \sim \mathcal{N}(\mathbf{0}, \sigma^2 \mathbf{I}_d)$ is a Gaussian noise such that $\mathbf{w} + \boldsymbol{\epsilon}$ are the weights of $h \sim \mathcal{Q}_S$ with $\mathcal{Q}_S = \mathcal{N}(\mathbf{w}, \sigma^2 \mathbf{I}_d)$.

Proof of Corollaries 8 and 9. Deferred in Appendix. \square

The experiments of Section 4.3, involve the direct minimization of Equations (4.2) and (4.4). In order to obtain tight bounds, the divergence between one prior $\mathcal{P}_t \in \mathbf{P}$ and \mathcal{Q}_S must be low, i.e., the term $\|\mathbf{w} - \mathbf{v}_t\|_2^2$ (or $\|\mathbf{w} + \boldsymbol{\epsilon} - \mathbf{v}_t\|_2^2 - \|\boldsymbol{\epsilon}\|_2^2$) has to be small. However, this can be challenging in the case of neural networks since $d \gg m$. One solution to this issue is to split the learning sample into two non-overlapping subsets $\mathcal{S}_{\text{prior}}$ and \mathcal{S} , where $\mathcal{S}_{\text{prior}}$ is used to learn the prior, while \mathcal{S} is used both to learn the posterior and compute the bound. Hence, if we “pre-learn” a good enough prior $\mathcal{P}_t \in \mathbf{P}$ from $\mathcal{S}_{\text{prior}}$, then we can expect to have a small value for $\|\mathbf{w} - \mathbf{v}_t\|_2$.

Training Method

The initial training set is split in 2 distinct subsets: $\mathcal{S}_{\text{prior}}$ and \mathcal{S} . The training has two phases.

- (i) The prior distribution \mathcal{P} is “pre-learned” with $\mathcal{S}_{\text{prior}}$ and selected by early stopping the algorithm with \mathcal{S} as validation set (from an arbitrary learning algorithm A_{prior}).
- (ii) Given \mathcal{S} and \mathcal{P} , we learn the posterior \mathcal{Q}_S with the algorithm A (defined a priori and used by our bounds).

At first sight, the selection of the prior weights with \mathcal{S} by early stopping may appear to be “cheating”. However, this procedure can be seen as: (i) first constructing \mathbf{P} from the

T “intermediate” networks learned after each epoch from $\mathcal{S}_{\text{prior}}$, (ii) optimizing the bound with the prior that leads to the best risk on \mathcal{S} . This gives a statistically valid result, since Corollaries 8 and 9 are valid for every $\mathcal{P} \in \mathbf{P}$. Usually, practitioners consider the “best” prior as the final network. In our case, the advantage is that we refine this “best” prior on \mathcal{S} to learn the posterior $\mathcal{Q}_{\mathcal{S}}$.

Note that Pérez-Ortiz et al. (2020) have already introduced tight generalization bounds with data-dependent priors for—non derandomized—stochastic neural networks. However, our training method to learn the prior differs greatly since (i) we learn T networks (i.e., T priors) instead of one, (ii) we also fix the variance of the Gaussian in the posterior distribution $\mathcal{Q}_{\mathcal{S}}$. Hence, to the best of our knowledge, our training method for the prior is new.

4.2 A Note About Stochastic Neural Networks

Due to its stochastic nature, PAC-Bayes has been explored to study stochastic neural networks (e.g. Dziugaite and Roy, 2017; 2018; Zhou et al., 2019; Pérez-Ortiz et al., 2020). In Corollary 10 below, we instantiate the PAC-Bayes bound of Equation (2.1) for stochastic networks to empirically compare in Section 4.3 the stochastic and the deterministic networks associated to prior(s) and posterior distributions.

Recall that, in the current paper, a deterministic neural network is a *single* h sampled from the posterior distribution $\mathcal{Q}_{\mathcal{S}} = \mathcal{N}(\mathbf{w}, \sigma^2 \mathbf{I}_d)$ —output by the algorithm A —, meaning that for each example, the label prediction is performed by the same deterministic network: the one parametrized by the weights $\mathbf{w} \in \mathbb{R}^d$. Conversely, the stochastic neural network associated to a posterior distribution $\mathcal{Q} = \mathcal{N}(\mathbf{w}', \sigma^2 \mathbf{I}_d)$ predicts the label of a given example by (i) first sampling h according to \mathcal{Q} (i.e., the network parametrized by the weight vector $\mathbf{w}' \in \mathbb{R}^d$), (ii) then returning the label predicted by h .² Then, the risk of the stochastic network corresponds to the expected risk value $\mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{D}}(h)$, where the expectation is taken over *all* h sampled from \mathcal{Q} . We compute the empirical risk of the stochastic network from Monte-Carlo approximation: (i) we sample n weight vectors, and (ii) we average the risk over the n associated networks; we denote by \mathcal{Q}^n the distribution of such n -sample. In this context, we propose the following randomized PAC-Bayes bound.

Corollary 10. *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any priors’ set $\mathbf{P} = \{\mathcal{P}_t\}_{t=1}^T$ of T priors \mathcal{P} on \mathcal{H} , for any loss $\ell: \mathcal{H} \times \mathcal{Z} \rightarrow \{0, 1\}$ we have, with probability at least $1 - 2\delta$ over the random choice of $\mathcal{S} \sim \mathcal{D}^m$ and $\{h_1, \dots, h_n\} \sim \mathcal{Q}^n$, we have simultaneously*

$$\text{kl} \left(\frac{1}{n} \sum_{i=1}^n R_{\mathcal{S}}(h_i) \parallel \mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{S}}(h) \right) \leq \frac{1}{n} \ln \frac{2}{\delta}, \quad (4.5)$$

²In the PAC-Bayesian literature, such stochastic classifier is referred as the Gibbs classifier.

and, for all prior $\mathcal{P}_t \in \mathbf{P}$, for all posterior \mathcal{Q}

$$\text{kl} \left(\mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{S}}(h) \parallel \mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{D}}(h) \right) \leq \frac{1}{m} \left[\frac{\|\mathbf{w}' - \mathbf{v}_t\|_2^2}{2\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta} \right]. \quad (4.6)$$

Proof. The proof is deferred in Appendix. \square

This corollary shows two key features that allow considering it as an adapted baseline for a *fair* comparison between pointwise and classical PAC-Bayesian bounds, thus between deterministic and stochastic networks in the next subsection. On the one hand, it involves the same terms as Corollary 8. On the other hand, it is close to the bound of Pérez-Ortiz et al. (2020, Sec. 6.2), since (i) we adapt the KL divergence to our setting (i.e., $\text{KL}(\mathcal{Q} \parallel \mathcal{P}) = \frac{1}{2\sigma^2} \|\mathbf{w}' - \mathbf{v}_t\|_2^2$), (ii) the bound holds for T priors thanks to a union bound argument.

4.3 Experiments

In this section, we perform two experiments (EXP (1) and EXP (2)); we do not seek for state-of-the-art performance, in fact, we have a twofold objective. In EXP (1), we compare our pointwise bound (Corollary 8, Equation (4.2)) and the Rivasplata et al.-based bound that we instantiate (Corollary 9, Equation (4.3)). In EXP (2), we highlight that our pointwise bound (Corollary 8, Equation (4.1)) associated to the deterministic network is tighter than the randomized bound associated to the stochastic network (Corollary 10).

Training Method. To compare the bounds, we follow our Training Method (Section 4.1) in which we integrate the direct minimization of the bounds. First of all, we need to replace the non-differentiable 0-1 loss by a surrogate for which the bounds still hold: the bounded cross entropy loss (Dziugaite and Roy, 2018). This latter is defined in a multiclass setting with $y \in \{1, 2, \dots\}$ by $\ell(h, (\mathbf{x}, y)) = -\frac{1}{Z} \ln(\Phi(h[y])) \in [0, 1]$ where $\forall p \in [0, 1], \Phi(p) = e^{-Z} + (1 - 2e^{-Z})p$, where $h[y]$ denotes the y -th output of the network (we set $Z=4$). That being said, to learn the best prior $\mathcal{P} \in \mathbf{P}$ and the posterior $\mathcal{Q}_{\mathcal{S}}$, we run our Training Method with two gradient descent-based algorithms A_{prior} and A . In phase (i) algorithm A_{prior} consists in learning from $\mathcal{S}_{\text{prior}}$ the T priors $\mathcal{P}_1, \dots, \mathcal{P}_T \in \mathbf{P}$ (i.e., during T epochs) by minimizing the bounded cross entropy loss. Then, the best prior $\mathcal{P} \in \mathbf{P}$ is selected by early stopping on \mathcal{S} . In phase (ii), given \mathcal{S} and \mathcal{P} , algorithm A integrates the direct optimization of Equation (4.2) or (4.4) with the bounded cross entropy loss.

Optimization procedure. Let ω be the mean vector of a Gaussian distribution used as network weights that we are optimizing. In phases (i) and (ii), at each iteration of the optimizer³, we sample a noise $\epsilon \sim \mathcal{N}(0, \sigma^2 \mathbf{I}_d)$. Then,

³Our optimization is done with Adam optimizer.



EXP (1): Pointwise PAC-Bayesian bounds comparison

EXP (2): Pointwise versus randomized


FIGURE 4.1: **EXP (1)** shows the value of optimized pointwise bound and the test risk for Corollary 8 (“Ours”) and Corollary 9 (“Rivasplata”) in two different settings, *i.e.*, with a learning rate of 10^{-6} and 10^{-4} . **EXP (2)** illustrates the PAC-Bayes bound of Corollary 10 compared to our pointwise bound with a learning rate of 10^{-4} . The y-axis shows the values of the bounds and the test risks.

we forward the examples of the mini-batch in the network parametrized by the weight vector $\omega + \epsilon$ and we update ω according to the bounded cross entropy loss.

Datasets and Experimental Setting.⁴ We perform our experimental study on 3 datasets. As a sanity check, we use the MNIST dataset (LeCun et al., 1998) and a MNIST-like dataset: Fashion-MNIST (Xiao et al., 2017). We also consider a more complex datasets: CIFAR-10 (Krizhevsky, 2009). For the MNIST(-like) datasets we train a variant of the All Convolutional Network (Springenberg et al., 2015), and for CIFAR we train a ResNet network (He et al., 2016). We divide each original train set into 2 independent subsets $\mathcal{S}_{\text{prior}}$ and \mathcal{S} (with a split of 50%/50%). In **EXP (1)**, we measure the averaged test risks and bounds values over 400 deterministic networks sampled from $\mathcal{Q}_{\mathcal{S}}$ for two learning rates (10^{-4} and 10^{-6} ; we set the other optimizer parameters to their default values). In **EXP (2)**, for the learning rate of 10^{-4} , we additionally report the PAC-Bayes bound of Corollary 10 with $n=400$ (with the same 400 hypotheses as in **EXP (1)**); meaning that the test risk reported also corresponds to the risk of the stochastic network approximated with these 400 hypotheses. The results, computed on the original test sets, are reported on Figure 4.1; note that the variance of the results is in order of 10^{-3} , then we do not report it.

Analysis of EXP (1). The two learning rates lead to similar test risks (of the deterministic network) and similar values to our bound of Equation (4.2) (close to the risk), whereas the values of the bound of Rivasplata et al. are different: especially with the highest learning rate of 10^{-4} the value is clearly less tight than ours. This behavior comes to the fact that their bound takes into account the noise ϵ through the term $\|\mathbf{w} + \epsilon - \mathbf{v}_t\|_2^2 - \|\epsilon\|_2^2$ (ours simply computes $\|\mathbf{w} - \mathbf{v}_t\|_2^2$). Thus it seems that our pointwise bound leads to more precise and stable values, justifying its use for optimization purposes to learn a deterministic network.

⁴The complete description of our overparametrized networks and the hyper-parameters are provided in Appendix.

Analysis of EXP (2). We observe that the pointwise PAC-Bayesian bound leads to more precise bounds than the randomized PAC-Bayesian one. This imprecision is due to the non-avoidable sampling according to \mathcal{Q} done in Equation (4.5) (the higher n , the tighter is the bound). Thus, using a pointwise PAC-Bayesian bound avoids sampling a large number of networks to obtain a low risk.

5 Conclusion

We have provided three new and general PAC-Bayesian pointwise bounds providing some novel perspectives in the formalization of derandomized PAC-Bayesian bounds, allowing some parametrization and offering nice opportunities for the analysis and optimization of deterministic classifiers. We have notably illustrated the interest of our theoretical result on neural networks. Future work includes specializations to specific architectures and other models. Another direction is to study domain adaptation (DA) by relating single source DA to the single pointwise bound and multi-source DA to the multiple pointwise bound (the priors being related to the source and the posterior to the target).

References

- Ibrahim M. Alabdulmohsin. An Information-Theoretic Route from Generalization in Expectation to Generalization in Probability. In *AISTATS*, 2017.
- Amiran Ambroladze, Emilio Parrado-Hernández, and John Shawe-Taylor. Tighter PAC-Bayes Bounds. In *NIPS*, pages 9–16, 2006.
- Raef Bassily, Shay Moran, Ido Nachum, Jonathan Shafer, and Amir Yehudayoff. Learners that Use Little Information. In *ALT*, 2018.
- Luc Bégin, Pascal Germain, François Laviolette, and Jean-François Roy. PAC-Bayesian Bounds based on the Rényi Divergence. In *AISTATS*, pages 435–444, 2016.

- Gilles Blanchard and François Fleuret. Occam’s Hammer. In *COLT*, pages 112–126, 2007.
- Olivier Bousquet and André Elisseeff. Stability and Generalization. *JMLR*, 2:499–526, 2002.
- Olivier Catoni. PAC-Bayesian Supervised Classification: The Thermodynamics of Statistical Learning. *CoRR*, abs/0712.0248, 2007.
- Gintare Karolina Dziugaite and Daniel M. Roy. Computing Nonvacuous Generalization Bounds for Deep (Stochastic) Neural Networks with Many More Parameters than Training Data. In *UAI*, 2017.
- Gintare Karolina Dziugaite and Daniel M. Roy. Data-dependent PAC-Bayes priors via differential privacy. In *NeurIPS*, pages 8440–8450, 2018.
- Amedeo Roberto Esposito, Michael Gastpar, and Ibrahim Issa. Robust Generalization via α -Mutual Information. In *International Zurich Seminar on Information and Communication*, 2020.
- Pascal Germain, Alexandre Lacasse, François Laviolette, and Mario Marchand. PAC-Bayesian Learning of Linear Classifiers. In *ICML*, pages 353–360, 2009.
- Moritz Hardt, Ben Recht, and Yoram Singer. Train faster, generalize better: Stability of stochastic gradient descent. In *ICML*, pages 1225–1234, 2016.
- Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Deep Residual Learning for Image Recognition. In *CVPR*, pages 770–778, 2016.
- Alex Krizhevsky. Learning Multiple Layers of Features from Tiny Images. Master’s thesis, University of Toronto, 2009.
- John Langford. Tutorial on Practical Prediction Theory for Classification. *JMLR*, 6:273–306, 2005.
- John Langford and John Shawe-Taylor. Pac-bayes & margins. In *NIPS*, 2002.
- Yann LeCun, Corinna Cortes, and Christopher J.C. Burges. THE MNIST DATASET of handwritten digits, 1998. URL <http://yann.lecun.com/exdb/mnist/>.
- Gaël Letarte, Pascal Germain, Benjamin Guedj, and François Laviolette. Dichotomize and Generalize: PAC-Bayesian Binary Activated Deep Neural Networks. In *NeurIPS*, pages 6869–6879, 2019.
- Andreas Maurer. A Note on the PAC Bayesian Theorem. *CoRR*, cs.LG/0411099, 2004.
- David A. McAllester. Some PAC-Bayesian Theorems. In *COLT*, pages 230–234, 1998.
- Vaishnavh Nagarajan and J. Zico Kolter. Deterministic PAC-Bayesian generalization bounds for deep networks via generalizing noise-resilience. In *ICLR*, 2019a.
- Vaishnavh Nagarajan and J. Zico Kolter. Uniform convergence may be unable to explain generalization in deep learning. In *NeurIPS*, pages 11611–11622, 2019b.
- Behnam Neyshabur, Srinadh Bhojanapalli, and Nathan Srebro. A PAC-Bayesian Approach to Spectrally-Normalized Margin Bounds for Neural Networks. In *ICLR*, 2018.
- María Pérez-Ortiz, Omar Rivasplata, John Shawe-Taylor, and Csaba Szepesvári. Tighter risk certificates for neural networks. *CoRR*, 2020.
- Omar Rivasplata, Ilja Kuzborskij, Csaba Szepesvári, and John Shawe-Taylor. PAC-Bayes Analysis Beyond the Usual Bounds. In *NeurIPS*, 2020.
- Matthias Seeger. PAC-Bayesian generalisation error bounds for gaussian process classification. *Journal of machine learning research*, 3(Oct):233–269, 2002.
- John Shawe-Taylor and Robert C. Williamson. A PAC Analysis of a Bayesian Estimator. In *COLT*, pages 2–9, 1997.
- Jost Tobias Springenberg, Alexey Dosovitskiy, Thomas Brox, and Martin A. Riedmiller. Striving for Simplicity: The All Convolutional Net. In *ICLR*, 2015.
- Niklas Thiemann, Christian Igel, Olivier Wintenberger, and Yevgeny Seldin. A Strongly Quasiconvex PAC-Bayesian Bound. In *ALT*, pages 466–492, 2017.
- Leslie G. Valiant. A Theory of the Learnable. *Commun. ACM*, 27(11):1134–1142, 1984.
- Vladimir Vapnik. *The Nature of Statistical Learning Theory*. Springer, 2000.
- Sergio Verdú. α -mutual information. In *Information Theory and Applications Workshop*, 2015.
- Han Xiao, Kashif Rasul, and Roland Vollgraf. Fashion-MNIST: a Novel Image Dataset for Benchmarking Machine Learning Algorithms. *CoRR*, abs/1708.07747, 2017.
- Aolin Xu and Maxim Raginsky. Information-theoretic analysis of generalization capability of learning algorithms. In *NIPS*, 2017.
- Huan Xu and Shie Mannor. Robustness and generalization. *Mach. Learn.*, 86(3):391–423, 2012.

A GENERAL FRAMEWORK FOR THE DERANDOMIZATION OF PAC-BAYESIAN BOUNDS

Wenda Zhou, Victor Veitch, Morgane Austern, Ryan P. Adams, and Peter Orbanz. Non-vacuous Generalization Bounds at the ImageNet Scale: a PAC-Bayesian Compression Approach. In *ICLR*, 2019.

A GENERAL FRAMEWORK FOR THE DERANDOMIZATION OF PAC-BAYESIAN BOUNDS
SUPPLEMENTARY MATERIAL

This supplementary material is structured as follows. In Section 6, we give the proof of our Theorem 2. Section 7 presents the proof of our Theorem 3. The proof of Proposition 4 is provided in Section 8. Section 9 is devoted to the proofs of Theorem 6 and Corollary 7. The proofs of Corollaries 8 and 9 are introduced in Section 10. In Section 11, we give the proof of Corollary 10. Section 12 provides more details on our experimental setup.

6 Proof of Theorem 2

Theorem 2. *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any prior distribution \mathcal{P} on \mathcal{H} , for any function $\phi : \mathcal{H} \times \mathcal{Z}^m \rightarrow \mathbb{R}^+$, for any $\alpha > 1$, for any deterministic algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, we have*

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m, h \sim \mathcal{Q}_{\mathcal{S}}} \left[\frac{\alpha}{\alpha-1} \ln(\phi(h, \mathcal{S})) \leq D_{\alpha}(\mathcal{Q}_{\mathcal{S}} \| \mathcal{P}) + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1} + 1}} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha}{\alpha-1}} \right) \right] \geq 1 - 2\delta,$$

where $\mathcal{Q}_{\mathcal{S}}$ is output by A , i.e., $\mathcal{Q}_{\mathcal{S}} \triangleq A(\mathcal{S})$.

Proof. For any sample \mathcal{S} and deterministic algorithm A fixed which allow us to obtain the distribution $\mathcal{Q}_{\mathcal{S}}$, note that $\phi(h, \mathcal{S})$ is a non-negative random variable. Hence, from Markov's inequality we have

$$\mathbb{P}_{h \sim \mathcal{Q}_{\mathcal{S}}} \left[\phi(h, \mathcal{S}) \leq \frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) \right] \geq 1 - \delta \iff \mathbb{E}_{h \sim \mathcal{Q}_{\mathcal{S}}} \mathbf{I} \left[\phi(h, \mathcal{S}) \leq \frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) \right] \geq 1 - \delta.$$

Taking the expectation over $\mathcal{S} \sim \mathcal{D}^m$ to both sides of the inequality gives

$$\begin{aligned} \mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathbb{E}_{h \sim \mathcal{Q}_{\mathcal{S}}} \mathbf{I} \left[\phi(h, \mathcal{S}) \leq \frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) \right] &\geq 1 - \delta \\ \iff \mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m, h \sim \mathcal{Q}_{\mathcal{S}}} \left[\phi(h, \mathcal{S}) \leq \frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) \right] &\geq 1 - \delta. \end{aligned}$$

Taking the logarithm to both sides of the equality and multiplying by $\frac{\alpha}{\alpha-1} > 0$, we obtain

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m, h \sim \mathcal{Q}_{\mathcal{S}}} \left[\frac{\alpha}{\alpha-1} \ln(\phi(h, \mathcal{S})) \leq \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) \right) \right] \geq 1 - \delta.$$

We develop the right side of the inequality in the indicator function and make the expectation of the hypothesis over the prior distribution \mathcal{P} appears. Indeed, we have

$$\begin{aligned} \forall \mathcal{P} \in \mathcal{M}(\mathcal{H}), \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) \right) &= \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \frac{\mathcal{Q}_{\mathcal{S}}(h') \mathcal{P}(h')}{\mathcal{P}(h') \mathcal{Q}_{\mathcal{S}}(h')} \phi(h', \mathcal{S}) \right) \\ &= \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{P}} \frac{\mathcal{Q}_{\mathcal{S}}(h')}{\mathcal{P}(h')} \phi(h', \mathcal{S}) \right). \end{aligned}$$

Remark that $\frac{1}{r} + \frac{1}{s} = 1$ with $r = \alpha$ and $s = \frac{\alpha}{\alpha-1}$. Hence, we can apply Hölder's inequality:

$$\mathbb{E}_{h' \sim \mathcal{P}} \frac{\mathcal{Q}_{\mathcal{S}}(h')}{\mathcal{P}(h')} \phi(h', \mathcal{S}) \leq \left[\mathbb{E}_{h' \sim \mathcal{P}} \left(\frac{\mathcal{Q}_{\mathcal{S}}(h')}{\mathcal{P}(h')} \right)^{\alpha} \right]^{\frac{1}{\alpha}} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha}{\alpha-1}} \right]^{\frac{\alpha-1}{\alpha}}.$$

Then, by taking the logarithm; adding $\ln(\frac{1}{\delta})$ and multiplying by $\frac{\alpha}{\alpha-1} > 0$ to both sides of the inequality, we obtain

$$\begin{aligned} \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{P}} \frac{\mathcal{Q}_S(h')}{\mathcal{P}(h')} \phi(h', \mathcal{S}) \right) &\leq \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \left[\mathbb{E}_{h' \sim \mathcal{P}} \left(\frac{\mathcal{Q}_S(h')}{\mathcal{P}(h')} \right)^\alpha \right]^{\frac{1}{\alpha}} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha-1}{\alpha}} \right]^{\frac{\alpha-1}{\alpha}} \right) \\ &= \frac{1}{\alpha-1} \ln \left(\mathbb{E}_{h' \sim \mathcal{P}} \left(\frac{\mathcal{Q}_S(h')}{\mathcal{P}(h')} \right)^\alpha \right) + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha-1}{\alpha}} \right) \\ &= D_\alpha(\mathcal{Q}_S \| \mathcal{P}) + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha-1}{\alpha}} \right). \end{aligned}$$

From this inequality, we can deduce that

$$\mathbb{P}_{S \sim \mathcal{D}^m, h \sim \mathcal{Q}_S} \left[\forall \mathcal{P} \in \mathcal{M}(\mathcal{H}), \frac{\alpha}{\alpha-1} \ln(\phi(h, \mathcal{S})) \leq D_\alpha(\mathcal{Q}_S \| \mathcal{P}) + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha-1}{\alpha}} \right) \right] \geq 1 - \delta. \quad (6.1)$$

Note that $\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha-1}{\alpha}}$ is a non-negative random variable, hence, we apply Markov's inequality to have

$$\mathbb{P}_{S \sim \mathcal{D}^m} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha-1}{\alpha}} \leq \frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha-1}{\alpha}} \right] \geq 1 - \delta.$$

Since the inequality does not depend on the random variable $h \sim \mathcal{Q}_S$, we have

$$\begin{aligned} &\mathbb{P}_{S \sim \mathcal{D}^m} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha-1}{\alpha}} \leq \frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha-1}{\alpha}} \right] \\ &= \mathbb{E}_{S \sim \mathcal{D}^m} \mathbf{I} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha-1}{\alpha}} \leq \frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha-1}{\alpha}} \right] \\ &= \mathbb{E}_{S \sim \mathcal{D}^m} \mathbb{E}_{h \sim \mathcal{Q}_S} \mathbf{I} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha-1}{\alpha}} \leq \frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha-1}{\alpha}} \right] \\ &= \mathbb{P}_{S \sim \mathcal{D}^m, h \sim \mathcal{Q}_S} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha-1}{\alpha}} \leq \frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha-1}{\alpha}} \right]. \end{aligned}$$

Multiplying by $\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} > 0$ and taking the logarithm to both sides of the inequality give us

$$\begin{aligned} &\mathbb{P}_{S \sim \mathcal{D}^m, h \sim \mathcal{Q}_S} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha-1}{\alpha}} \leq \frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha-1}{\alpha}} \right] \geq 1 - \delta \\ \iff &\mathbb{P}_{S \sim \mathcal{D}^m, h \sim \mathcal{Q}_S} \left[\ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha-1}{\alpha}} \right) \leq \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}+1}} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha-1}{\alpha}} \right) \right] \geq 1 - \delta. \quad (6.2) \end{aligned}$$

Combining Equation (6.1) and Equation (6.2) with a union bound gives us the desired result. \square

7 Proof of Theorem 3

For the sake of completeness, we first prove an upper bound on \sqrt{ab} (see *e.g.*, [Thiemann et al., 2017](#)).

Lemma 11. *For any $a > 0, b > 0$, we have*

$$\sqrt{\frac{a}{b}} = \operatorname{arginf}_{\lambda > 0} \left(\frac{a}{\lambda} + \lambda b \right), \quad 2\sqrt{ab} = \inf_{\lambda > 0} \left(\frac{a}{\lambda} + \lambda b \right) \quad \text{and} \quad \forall \lambda > 0, \sqrt{ab} \leq \frac{1}{2} \left(\frac{a}{\lambda} + \lambda b \right).$$

Proof. Let $f(\lambda) = \left(\frac{a}{\lambda} + \lambda b \right)$. The first and second derivatives of f w.r.t. λ is

$$\frac{\partial f}{\partial \lambda}(\lambda) = \left(b - \frac{a}{\lambda^2} \right) \quad \text{and} \quad \frac{\partial f}{\partial \lambda^2}(\lambda) = \frac{2a}{\lambda^3}.$$

Hence, for $\lambda > 0$, we have $\frac{\partial f}{\partial \lambda^2}(\lambda) > 0$; f is strictly convex and admit a unique minimum. Solving $\frac{\partial f}{\partial \lambda}(\lambda) = 0$ we have $\lambda^* = \sqrt{\frac{a}{b}}$. Additionally, $f(\lambda^*) = 2\sqrt{ab}$ which proves the claim. \square

We can now prove Theorem 3 with Lemma 11.

Theorem 3. *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any prior distribution \mathcal{P} on \mathcal{H} , for any function $\phi : \mathcal{H} \times \mathcal{Z}^m \rightarrow \mathbb{R}^+$, for any deterministic algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, we have*

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m, h \sim \mathcal{Q}_{\mathcal{S}}} \left[\forall \lambda > 0, \ln(\phi(h, \mathcal{S})) \leq \ln \left(\frac{1}{2} \left[\lambda \exp[D_2(\mathcal{Q}_{\mathcal{S}} \parallel \mathcal{P})] + \frac{1}{\lambda \delta^3} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^2 \right] \right) \right] \geq 1 - 2\delta,$$

where $\mathcal{Q}_{\mathcal{S}} \triangleq A(\mathcal{S})$.

Proof. The proof is similar to the one of Theorem 2. Given a sample \mathcal{S} and a deterministic algorithm A (which allow us to obtain the distribution $\mathcal{Q}_{\mathcal{S}}$), the value $\phi(h, \mathcal{S})$ is a non-negative random variable. Hence, from Markov's inequality we have

$$\mathbb{P}_{h \sim \mathcal{Q}_{\mathcal{S}}} \left[\phi(h, \mathcal{S}) \leq \frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) \right] \geq 1 - \delta \iff \mathbb{E}_{h \sim \mathcal{Q}_{\mathcal{S}}} \mathbf{I} \left[\phi(h, \mathcal{S}) \leq \frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) \right] \geq 1 - \delta$$

Taking the expectation over $\mathcal{S} \sim \mathcal{D}^m$ to both sides of the inequality gives

$$\mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathbb{E}_{h \sim \mathcal{Q}_{\mathcal{S}}} \mathbf{I} \left[\phi(h, \mathcal{S}) \leq \frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) \right] \geq 1 - \delta \iff \mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m, h \sim \mathcal{Q}_{\mathcal{S}}} \left[\phi(h, \mathcal{S}) \leq \frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) \right] \geq 1 - \delta.$$

Using the fact that for all $\lambda > 0$, $\sqrt{ab} \leq \frac{1}{2} \left(\frac{a}{\lambda} + b\lambda \right)$ (see Lemma 11) with $a = \frac{1}{\delta^2} \phi(h', \mathcal{S})^2$ and $b = \frac{\mathcal{Q}_{\mathcal{S}}(h')^2}{\mathcal{P}(h')^2}$, we have

$$\forall \mathcal{P} \in \mathcal{M}(\mathcal{H}), \forall \lambda > 0, \frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) = \mathbb{E}_{h' \sim \mathcal{P}} \sqrt{\frac{\mathcal{Q}_{\mathcal{S}}(h')^2}{\mathcal{P}(h')^2} \frac{1}{\delta^2} \phi(h', \mathcal{S})^2} \leq \frac{1}{2} \left[\lambda \mathbb{E}_{h' \sim \mathcal{P}} \left(\frac{\mathcal{Q}_{\mathcal{S}}(h')}{\mathcal{P}(h')} \right)^2 + \frac{1}{\lambda \delta^2} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \right].$$

Then, taking the logarithm to both sides of the inequality, we obtain

$$\begin{aligned} \forall \lambda > 0, \ln \left(\frac{1}{\delta} \mathbb{E}_{h' \sim \mathcal{Q}_{\mathcal{S}}} \phi(h', \mathcal{S}) \right) &\leq \ln \left(\frac{1}{2} \left[\lambda \mathbb{E}_{h' \sim \mathcal{P}} \left(\frac{\mathcal{Q}_{\mathcal{S}}(h')}{\mathcal{P}(h')} \right)^2 + \frac{1}{\lambda \delta^2} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \right] \right) \\ &= \ln \left(\frac{1}{2} \left[\lambda \exp(D_2(\mathcal{Q}_{\mathcal{S}} \parallel \mathcal{P})) + \frac{1}{\lambda \delta^2} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \right] \right). \end{aligned}$$

Hence, we can deduce that

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m, h \sim \mathcal{Q}_{\mathcal{S}}} \left[\forall \mathcal{P} \in \mathcal{M}(\mathcal{H}), \forall \lambda > 0, \ln(\phi(h, \mathcal{S})) \leq \ln \left(\frac{1}{2} \left[\lambda \exp(D_2(\mathcal{Q}_{\mathcal{S}} \parallel \mathcal{P})) + \frac{1}{\lambda \delta^2} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \right] \right) \right] \geq 1 - \delta. \quad (7.1)$$

Note that $\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2$ is a non-negative random variable, hence, we apply Markov's inequality:

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \leq \frac{1}{\delta} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^2 \right] \geq 1 - \delta.$$

Since the inequality does not depend on the random variable $h \sim \mathcal{Q}_{\mathcal{S}}$, we have

$$\begin{aligned} &\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \leq \frac{1}{\delta} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^2 \right] = \mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathbf{I} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \leq \frac{1}{\delta} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^2 \right] \\ &= \mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathbb{E}_{h \sim \mathcal{Q}_{\mathcal{S}}} \mathbf{I} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \leq \frac{1}{\delta} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^2 \right] = \mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m, h \sim \mathcal{Q}_{\mathcal{S}}} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \leq \frac{1}{\delta} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^2 \right]. \end{aligned}$$

Additionally, note that multiplying by $\frac{1}{2\lambda\delta^2} > 0$, adding $\frac{\lambda}{2} \exp(D_2(\mathcal{Q}_{\mathcal{S}} \parallel \mathcal{P}))$, and taking the logarithm to both sides of the inequality results in the same indicator function. Indeed,

$$\begin{aligned}
& \mathbf{I} \left[\mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \leq \frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^2 \right] \\
&= \mathbf{I} \left[\forall \lambda > 0, \frac{1}{2\lambda\delta^2} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \leq \frac{1}{2\lambda\delta^3} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^2 \right] \\
&= \mathbf{I} \left[\forall \lambda > 0, \frac{\lambda}{2} \exp(D_2(\mathcal{Q}_S \| \mathcal{P})) + \frac{1}{2\lambda\delta^2} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \leq \frac{\lambda}{2} \exp(D_2(\mathcal{Q}_S \| \mathcal{P})) + \frac{1}{2\lambda\delta^3} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^2 \right] \\
&= \mathbf{I} \left[\forall \lambda > 0, \ln \left(\frac{\lambda}{2} \exp(D_2(\mathcal{Q}_S \| \mathcal{P})) + \frac{1}{2\lambda\delta^2} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \right) \leq \ln \left(\frac{\lambda}{2} \exp(D_2(\mathcal{Q}_S \| \mathcal{P})) + \frac{1}{2\lambda\delta^3} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^2 \right) \right].
\end{aligned}$$

Hence, we can deduce that

$$\begin{aligned}
& \mathbb{P}_{\substack{S \sim \mathcal{D}^m \\ h \sim \mathcal{Q}_S}} \left[\forall \lambda > 0, \ln \left(\frac{1}{2} \left[\lambda \exp(D_2(\mathcal{Q}_S \| \mathcal{P})) + \frac{1}{\lambda\delta^2} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^2 \right] \right) \right. \\
& \quad \left. \leq \ln \left(\frac{1}{2} \left[\lambda \exp(D_2(\mathcal{Q}_S \| \mathcal{P})) + \frac{1}{\lambda\delta^3} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^2 \right] \right) \right] \geq 1 - \delta. \tag{7.2}
\end{aligned}$$

Combining Equation (7.1) and Equation (7.2) with a union bound gives us the desired result. \square

8 Proof of Proposition 4

Proposition 4. For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any prior distribution \mathcal{P} on \mathcal{H} and for any deterministic algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, let

$$\lambda^* = \operatorname{argmin}_{\lambda > 0} \ln \left(\frac{\lambda}{2} \exp(D_2(\mathcal{Q}_S \| \mathcal{P})) + \frac{1}{2\lambda\delta^3} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^2 \right),$$

$$\text{then } \underbrace{2 \ln \left[\frac{\lambda^*}{2} \exp(D_2(\mathcal{Q}_S \| \mathcal{P})) + \frac{\mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^2}{2\lambda^*\delta^3} \right]}_{\text{Theorem 3}} = D_2(\mathcal{Q}_S \| \mathcal{P}) + \underbrace{\ln \left[\frac{1}{\delta^3} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^2 \right]}_{\text{Theorem 2 with } \alpha = 2},$$

$$\text{where } \lambda^* = \sqrt{\frac{\mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^2}{\delta^3 \exp(D_2(\mathcal{Q}_S \| \mathcal{P}))}}.$$

Proof. We first prove the closed form of λ^* , then, we compute the difference between the two bounds. For the sake of clarity, let $d = \exp(D_2(\mathcal{Q}_S \| \mathcal{P}))$, and $p = \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^2$ and $f(\lambda) = \ln \left(\frac{\lambda}{2} d + \frac{p}{2\lambda\delta^3} \right)$. We can find the first and second derivatives of f w.r.t. λ , indeed, we have

$$\frac{\partial f}{\partial \lambda}(\lambda) = \frac{\frac{d}{2} - \frac{p}{2\delta^3\lambda^2}}{\frac{d}{2} + \frac{p}{2\delta^3\lambda^2}} \quad \text{and} \quad \frac{\partial f}{\partial \lambda^2}(\lambda) = \frac{2p(\delta^3\lambda^2 d + p) - (\delta^3\lambda^2 d - p)^2}{\lambda^2(\delta^3\lambda^2 d + p)^2}.$$

Furthermore, when $\frac{\partial f}{\partial \lambda}(\lambda) = 0$, we have $\lambda = \sqrt{\frac{p}{\delta^3 d}}$, and we can prove that the associated minimum is unique (i.e., $\lambda = \lambda^*$) by studying the function.

Actually, we have $\frac{\partial f}{\partial \lambda^2} > 0 \iff p^2 + 4px - x^2 > 0$ where $x = \delta^3\lambda^2 d$. Solving this polynomial of order 2 ($ax^2 + bx + c > 0$ with $a = -1, b = 4p, c = p^2$), we have that

$$\frac{\partial f}{\partial \lambda^2}(\lambda) > 0 \iff 0 < \delta^3\lambda^2 d < p(2 + \sqrt{5}) \iff 0 < \lambda < \sqrt{\frac{p}{\delta^3 d}} \sqrt{2 + \sqrt{5}}.$$

Hence, for $\lambda \in]0, \sqrt{\frac{p}{\delta^3 d}} \sqrt{2 + \sqrt{5}}[$, f is strictly convex and λ^* is the unique minimizer on that interval. Furthermore, from the first derivative, we know that

$$\frac{\partial f}{\partial \lambda}(\lambda) > 0 \iff \lambda > \sqrt{\frac{p}{d\delta^3}}.$$

Hence, f is strictly increasing on $]\sqrt{\frac{p}{d\delta^3}}, +\infty[$, hence, $\lambda^* = \sqrt{\frac{p}{d\delta^3}}$ is the unique minimizer on $]0, +\infty[$.

Lastly, substituting λ by λ^* in the right term of the inequality in Theorem 3 gives the equality. \square

9 Proof of Theorem 6 and Corollary 7

Before proving Theorem 6, we prove the following lemma.

Lemma 12. *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any prior distribution \mathcal{P} on \mathcal{H} , for any function $\phi : \mathcal{H} \times \mathcal{Z}^m$, for any $\alpha > 1$, for any deterministic algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, we have*

$$\mathbb{P}_{S \sim \mathcal{D}^m, h \sim \mathcal{Q}_S} \left[\forall \mathcal{P} \in \mathcal{M}(\mathcal{H}), \frac{\alpha}{\alpha-1} \ln(\phi(h, S)) \leq D_\alpha(\rho \parallel \pi) + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^{\frac{\alpha}{\alpha-1}} \right) \right] \geq 1 - \delta.$$

where $\rho(h, S) = \mathcal{Q}_S(h) \times \mathcal{D}^m(S)$; $\pi(h, S) = \mathcal{P}(h) \times \mathcal{D}^m(S)$ and $\mathcal{Q}_S \triangleq A(S)$.

Proof. For any deterministic algorithm A fixed which allow us to obtain the distribution \mathcal{Q}_S from a sample S , note that $\phi(h, S)$ is a non-negative random variable. Hence, from Markov's inequality, we have

$$\mathbb{P}_{S \sim \mathcal{D}^m, h \sim \mathcal{Q}_S} \left[\phi(h, S) \leq \frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{Q}_{S'}} \phi(h', S') \right] \geq 1 - \delta.$$

Taking the logarithm to both sides of the equality and multiplying by $\frac{\alpha}{\alpha-1} > 0$, we have

$$\mathbb{P}_{S \sim \mathcal{D}^m, h \sim \mathcal{Q}_S} \left[\frac{\alpha}{\alpha-1} \ln(\phi(h, S)) \leq \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{Q}_{S'}} \phi(h', S') \right) \right] \geq 1 - \delta.$$

We develop the right side of the inequality in the indicator function and make the expectation of the hypothesis over the distribution \mathcal{P} appears:

$$\begin{aligned} \forall \mathcal{P} \in \mathcal{M}(\mathcal{H}), \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{Q}_{S'}} \phi(h', S') \right) &= \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{Q}_{S'}} \frac{\mathcal{Q}_{S'}(h')}{\mathcal{P}(h')} \frac{\mathcal{P}(h')}{\mathcal{Q}_{S'}(h')} \phi(h', S') \right) \\ &= \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \frac{\mathcal{Q}_{S'}(h')}{\mathcal{P}(h')} \phi(h', S') \right). \end{aligned}$$

Then, remark that $\frac{1}{r} + \frac{1}{s} = 1$ where $r = \alpha$ and $s = \frac{\alpha}{\alpha-1}$. Hence, Hölder's inequality gives

$$\mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{Q}_{S'}} \phi(h', S') \leq \left[\mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \left(\frac{\mathcal{Q}_{S'}(h')}{\mathcal{P}(h')} \right)^\alpha \right]^{\frac{1}{\alpha}} \left[\mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^{\frac{\alpha}{\alpha-1}} \right]^{\frac{\alpha-1}{\alpha}}.$$

Then, taking the logarithm, adding $\ln(\frac{1}{\delta})$, and multiplying by $\frac{\alpha}{\alpha-1} > 0$ to both sides of the inequality, we have

$$\begin{aligned} \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{Q}_{S'}} \phi(h', S') \right) &\leq \frac{\alpha}{\alpha-1} \ln \left(\frac{1}{\delta} \left[\mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \left(\frac{\mathcal{Q}_{S'}(h')}{\mathcal{P}(h')} \right)^\alpha \right]^{\frac{1}{\alpha}} \left[\mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^{\frac{\alpha}{\alpha-1}} \right]^{\frac{\alpha-1}{\alpha}} \right) \\ &= \frac{1}{\alpha-1} \ln \left(\mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \left(\frac{\mathcal{Q}_{S'}(h')}{\mathcal{P}(h')} \right)^\alpha \right) + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^{\frac{\alpha}{\alpha-1}} \right). \end{aligned}$$

Hence, we can deduce that

$$\begin{aligned} \mathbb{P}_{S \sim \mathcal{D}^m, h \sim \mathcal{Q}_S} \left[\forall \mathcal{P} \in \mathcal{M}(\mathcal{H}), \frac{\alpha}{\alpha-1} \ln(\phi(h, S)) \leq \frac{1}{\alpha-1} \ln \left(\mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \left[\frac{\mathcal{Q}_{S'}(h')}{\mathcal{P}(h')} \right]^\alpha \right) \right. \\ \left. + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', S')^{\frac{\alpha}{\alpha-1}} \right) \right] \geq 1 - \delta, \end{aligned}$$

where by definition we have $D_\alpha(\rho \parallel \pi) = \frac{1}{\alpha-1} \ln \left(\mathbb{E}_{S' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \left[\frac{\mathcal{Q}_{S'}(h')}{\mathcal{P}(h')} \right]^\alpha \right)$. \square

From Lemma 12, we prove Theorem 6.

Theorem 6. *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any function $\phi : \mathcal{H} \times \mathcal{Z}^m$, for any $\alpha > 1$, for any deterministic algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, we have*

$$\mathbb{P}_{\substack{\mathcal{S} \sim \mathcal{D}^m \\ h \sim \mathcal{Q}_{\mathcal{S}}}} \left[\frac{\alpha}{\alpha-1} \ln(\phi(h, \mathcal{S})) \leq I_{\alpha}(h'; \mathcal{S}') + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha}{\alpha-1}} \right) \right] \geq 1 - \delta,$$

where $\mathcal{Q}_{\mathcal{S}} \triangleq A(\mathcal{S})$.

Proof. Note that Sibson's mutual information is defined as $I_{\alpha}(h; \mathcal{S}) = \min_{\mathcal{P} \in \mathcal{M}(\mathcal{H})} D_{\alpha}(\rho \| \pi)$. Hence, in order to prove Theorem 6, we have to instantiate Lemma 12 with the optimal prior, *i.e.*, the prior \mathcal{P} which minimizes $D_{\alpha}(\rho \| \pi)$. Actually, this optimal prior has a closed-form solution (Verdú, 2015). For the sake of completeness, we derive it. First, we have

$$\begin{aligned} D_{\alpha}(\rho \| \pi) &= \frac{1}{\alpha-1} \ln \left(\mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathbb{E}_{h \sim \mathcal{P}} \left[\frac{\mathcal{Q}_{\mathcal{S}}(h)}{\mathcal{P}(h)} \right]^{\alpha} \right) \\ &= \frac{1}{\alpha-1} \ln \left(\mathbb{E}_{h \sim \mathcal{P}} \left[\mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathcal{Q}_{\mathcal{S}}(h)^{\alpha} \right] \mathcal{P}(h)^{-\alpha} \right) \\ &= \frac{1}{\alpha-1} \ln \left(\mathbb{E}_{h \sim \mathcal{P}} \left[\mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathcal{Q}_{\mathcal{S}}(h)^{\alpha} \right] \mathcal{P}(h)^{-\alpha} \left[\frac{\mathbb{E}_{h' \sim \mathcal{P}} \frac{1}{\mathcal{P}(h')}}{\mathbb{E}_{h' \sim \mathcal{P}} \frac{1}{\mathcal{P}(h')}} \left[\mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathcal{Q}_{\mathcal{S}'}(h')^{\alpha} \right]^{\frac{1}{\alpha}} \right]^{\alpha} \right) \\ &= \frac{\alpha}{\alpha-1} \ln \left(\mathbb{E}_{h' \sim \mathcal{P}} \frac{1}{\mathcal{P}(h')} \left[\mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathcal{Q}_{\mathcal{S}'}(h')^{\alpha} \right]^{\frac{1}{\alpha}} \right) + \frac{1}{\alpha-1} \ln \left(\mathbb{E}_{h \sim \mathcal{P}} \frac{1}{\mathcal{P}(h)^{\alpha}} \left[\frac{\mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathcal{Q}_{\mathcal{S}}(h)^{\alpha} \right]^{\frac{1}{\alpha}} \right)^{\alpha} \right) \\ &= \frac{\alpha}{\alpha-1} \ln \left(\mathbb{E}_{h' \sim \mathcal{P}} \frac{1}{\mathcal{P}(h')} \left[\mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathcal{Q}_{\mathcal{S}'}(h')^{\alpha} \right]^{\frac{1}{\alpha}} \right) + D_{\alpha}(\mathcal{P}^* \| \mathcal{P}), \end{aligned}$$

$$\text{where } \mathcal{P}^*(h) = \left[\frac{\mathbb{E}_{\mathcal{S} \sim \mathcal{D}^m} \mathcal{Q}_{\mathcal{S}}(h)^{\alpha} \right]^{\frac{1}{\alpha}} \left[\mathbb{E}_{h' \sim \mathcal{P}} \frac{1}{\mathcal{P}(h')} \left[\mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathcal{Q}_{\mathcal{S}'}(h')^{\alpha} \right]^{\frac{1}{\alpha}} \right].$$

From these equalities and using the fact that $D_{\alpha}(\mathcal{P}^* \| \mathcal{P})$ is minimal (*i.e.*, equal to zero) when $\mathcal{P}^* = \mathcal{P}$, we can deduce that

$$\operatorname{argmin}_{\mathcal{P} \in \mathcal{M}(\mathcal{H})} D_{\alpha}(\rho \| \pi) = \operatorname{argmin}_{\mathcal{P} \in \mathcal{M}(\mathcal{H})} \left[\frac{\alpha}{\alpha-1} \ln \left(\mathbb{E}_{h' \sim \mathcal{P}} \frac{1}{\mathcal{P}(h')} \left[\mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathcal{Q}_{\mathcal{S}'}(h')^{\alpha} \right]^{\frac{1}{\alpha}} \right) + D_{\alpha}(\mathcal{P}^* \| \mathcal{P}) \right] = \operatorname{argmin}_{\mathcal{P} \in \mathcal{M}(\mathcal{H})} D_{\alpha}(\mathcal{P}^* \| \mathcal{P}) = \mathcal{P}^*.$$

Note that \mathcal{P}^* is defined from the data distribution \mathcal{D} , hence, \mathcal{P}^* is a valid prior when instantiating Lemma 12 with \mathcal{P}^* . Then, we have with *prob.* at least $1 - \delta$ over the random choice of $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_{\mathcal{S}}$

$$\begin{aligned} \frac{\alpha}{\alpha-1} \ln(\phi(h, \mathcal{S})) &\leq D_{\alpha}(\rho \| \pi^*) + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha}{\alpha-1}} \right) \\ &= I_{\alpha}(h'; \mathcal{S}') + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S}')^{\frac{\alpha}{\alpha-1}} \right). \end{aligned}$$

where $\pi^*(h, \mathcal{S}) = \mathcal{P}^*(h) \times \mathcal{D}^m(\mathcal{S})$. □

From Theorem 6, we derive the following Corollary 7.

Corollary 7. *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any deterministic algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, with probability at least $1 - \delta$ over the random choice of $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_{\mathcal{S}}$, we have*

$$\underbrace{2(R_{\mathcal{S}}(h) - R_{\mathcal{D}}(h))^2 \leq \frac{1}{m} \left[I_{\alpha}(h'; \mathcal{S}') + \ln \frac{2\sqrt{m}}{\delta^{\frac{\alpha}{\alpha-1}}} \right]}_{\text{Equation (3.4)}} \quad \text{and} \quad \underbrace{\text{kl}(R_{\mathcal{S}}(h) \| R_{\mathcal{D}}(h)) \leq \frac{1}{m} \left[I_{\alpha}(h'; \mathcal{S}') + \ln \frac{2\sqrt{m}}{\delta^{\frac{\alpha}{\alpha-1}}} \right]}_{\text{Equation (3.5)}}.$$

Proof. First, we instantiate Theorem 6 with $\phi(h, \mathcal{S}) = \exp\left[\frac{\alpha-1}{\alpha} m \text{kl}(R_{\mathcal{S}}(h) \| R_{\mathcal{D}}(h))\right]$, we have (by rearranging the terms)

$$\text{kl}(R_{\mathcal{S}}(h) \| R_{\mathcal{D}}(h)) \leq \frac{1}{m} \left[I_{\alpha}(h'; \mathcal{S}') + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} e^{m \text{kl}(R_{\mathcal{S}'}(h') \| R_{\mathcal{D}}(h'))} \right) \right].$$

Then, from Maurer (2004), we upper-bound $\mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}} e^{m \text{kl}(R_{\mathcal{S}'}(h') \| R_{\mathcal{D}}(h'))}$ by $2\sqrt{m}$ to obtain Equation (3.4). Finally, to obtain Equation (3.4), we apply Pinsker's inequality, i.e., $2(R_{\mathcal{S}}(h) - R_{\mathcal{D}}(h))^2 \leq \text{kl}(R_{\mathcal{S}}(h) \| R_{\mathcal{D}}(h))$ on Equation (3.5). \square

10 Proof of Corollaries 8 and 9

For the sake of completeness, we this section includes a proof of Proposition 13 (Letarte et al., 2019, Th. 3). The latter is then used to prove Corollaries 8 and 9. Furthermore, we introduce Theorem 2' which takes into account a set of priors \mathbf{P} while Theorem 2 handles a unique prior \mathcal{P} .

Proposition 13. *For any hypothesis set \mathcal{H} , for any learning sample $\mathcal{S} \in \mathcal{Z}^m$, for any $h \in \mathcal{H}$, for any loss $\ell: \mathcal{H} \times \mathcal{Z} \rightarrow [0, 1]$, for any $\kappa > 0$, we have*

$$\sup_{R_{\mathcal{S}}(h) \leq r \leq 1} \left\{ r \mid \text{kl}(R_{\mathcal{S}}(h) \| r) \leq \kappa \right\} = \inf_{C > 0} \left[\frac{1 - \exp[-C R_{\mathcal{S}}(h) - \kappa]}{1 - \exp(-C)} \right].$$

The proposition proof builds on the following Lemma 14.

Lemma 14 (Proposition 2.1 of Germain et al. (2009)). *For any $p, q \in (0, 1)$, if $q \leq p$, we have*

$$\text{kl}(q \| p) = \max_{C \geq 0} \Delta(C, q, p), \quad \text{where} \quad \Delta(C, q, p) = -\ln(1 - p(1 - e^{-C})) - Cq.$$

Proof. First, we compute the first and second derivatives of $\Delta(C, q, p)$ with respect to C , we have

$$\frac{\partial \Delta(C, q, p)}{\partial C} = \frac{pe^{-C}}{pe^{-C} + (1-p)} - q = \frac{p}{p(1 - e^C) + e^C} - q, \quad \text{and} \quad \frac{\partial \Delta(C, q, p)}{\partial C^2} = \frac{p(p-1)e^C}{(p(1 - e^C) + e^C)^2} \leq 0.$$

Since the second derivative $\frac{\partial \Delta(C, q, p)}{\partial C^2}$ is negative, we can deduce that $\Delta(C, q, p)$ is concave. Furthermore, the maximum is attained when

$$\frac{\partial \Delta(C, q, p)}{\partial C} = 0 \iff C = \ln \left(\frac{p(q-1)}{q(p-1)} \right) = \ln \left(\frac{p(1-q)}{q(1-p)} \right) = C^*.$$

Hence, we can deduce that $C^* \geq 0$ when $q \leq p$. Finally, by plugging C^* into the function Δ , i.e., we have

$$\Delta(C^*, q, p) = \text{kl}(q \| p).$$

\square

We now can prove Proposition 13.

Proof. Let $r^* = \sup_{R_{\mathcal{S}}(h) \leq r \leq 1} \left\{ r \mid \text{kl}(R_{\mathcal{S}}(h) \| r) \leq \kappa \right\}$; We consider 3 cases.

(1) For $R_{\mathcal{S}}(h) \in (0, 1)$. Let $C^* = \text{argsup}_{C > 0} \Delta(C, R_{\mathcal{S}}(h), r^*)$. From Lemma 14 and since the KL divergence is strictly increasing for all $R_{\mathcal{S}}(h) \leq r \leq 1$, we have

$$\text{kl}(R_{\mathcal{S}}(h) \| r^*) = \Delta(C^*, R_{\mathcal{S}}(h), r^*) = \kappa, \quad \text{and} \quad \forall C > 0, \quad \Delta(C, R_{\mathcal{S}}(h), r^*) \leq \kappa.$$

Note that $C \neq 0$ since $r^* \neq R_{\mathcal{S}}(h)$ with $\kappa > 0$. When isolating r^* in the two equations, we obtain

$$r^* = \frac{1}{1 - e^{-C^*}} [1 - \exp(-C^* R_{\mathcal{S}}(h) - \kappa)], \quad \text{and} \quad \forall C > 0, \quad r^* \leq \frac{1}{1 - e^{-C}} [1 - \exp(-C R_{\mathcal{S}}(h) - \kappa)].$$

Thus, we have $r^* = \inf_{C>0} \left[\frac{1}{1 - e^{-C}} [1 - \exp(-C R_S(h) - \kappa)] \right]$ for $R_S(h) \in (0, 1)$.

(2) For $R_S(h) = 0$. We have $r^* = 1 - \exp(-\kappa)$ since $\text{kl}(0 \| r^*) = \text{kl}(0 \| 1 - \exp(-\kappa)) = \kappa$;
for all $C > 0$, we have $\frac{1 - e^{-\kappa}}{1 - e^{-C}} \geq 1 - e^{-\kappa}$ and $\lim_{C \rightarrow +\infty} \left[\frac{1 - e^{-\kappa}}{1 - e^{-C}} \right] = 1 - e^{-\kappa}$, hence, we have

$$r^* = \inf_{C>0} \left[\frac{1}{1 - e^{-C}} [1 - \exp(-C R_S(h) - \kappa)] \right] \text{ for } R_S(h) = 0.$$

(3) For $R_S(h) = 1$. We have $r^* = 1$ and $\text{kl}(1 \| 1) = 0$;

for all $C > 0$, we have $\frac{1 - e^{-C-\kappa}}{1 - e^{-\alpha}} \geq 1$ and $\lim_{C \rightarrow +\infty} \left[\frac{1 - e^{-C-\kappa}}{1 - e^{-C}} \right] = 1$, hence, we have

$$r^* = \inf_{C>0} \left[\frac{1}{1 - e^{-C}} [1 - \exp(-C R_S(h) - \kappa)] \right] \text{ for } R_S(h) = 1.$$

□

We now introduce Theorem 2' which takes into account a set of priors \mathbf{P} instead of a prior \mathcal{P} .

Theorem 2'. For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any priors set $\mathbf{P} = \{\mathcal{P}_t\}_{t=1}^T$ of T prior \mathcal{P} on \mathcal{H} , for any function $\phi : \mathcal{H} \times \mathcal{Z}^m \rightarrow \mathbb{R}^+$, for any $\alpha > 1$, for any deterministic algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, we have

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m, h \sim \mathcal{Q}_S} \left[\forall \mathcal{P}_t \in \mathbf{P}, \frac{\alpha}{\alpha-1} \ln(\phi(h, \mathcal{S})) \leq D_\alpha(\mathcal{Q}_S \| \mathcal{P}_t) + \ln \left(\frac{T}{\delta^{\frac{\alpha}{\alpha-1}+1}} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}_t} \phi(h', \mathcal{S}')^{\frac{\alpha}{\alpha-1}} \right) \right] \geq 1 - 2\delta,$$

Proof. The proof is essentially the same as Theorem 2. Indeed, we first derive the same equation as Equation (6.1), we have

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m, h \sim \mathcal{Q}_S} \left[\forall \mathcal{P} \in \mathcal{M}(\mathcal{H}), \frac{\alpha}{\alpha-1} \ln(\phi(h, \mathcal{S})) \leq D_\alpha(\mathcal{Q}_S \| \mathcal{P}) + \ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{h' \sim \mathcal{P}} \phi(h', \mathcal{S})^{\frac{\alpha}{\alpha-1}} \right) \right] \geq 1 - \delta.$$

Then, we apply Markov inequality (as in Theorem 2) T times with the T priors \mathcal{P}_t belonging to \mathbf{P} , however, we set the confidence to δ/T instead of δ , we have

$$\mathbb{P}_{\mathcal{S} \sim \mathcal{D}^m, h \sim \mathcal{Q}_S} \left[\ln \left(\frac{1}{\delta^{\frac{\alpha}{\alpha-1}}} \mathbb{E}_{h' \sim \mathcal{P}_t} \phi(h', \mathcal{S})^{\frac{\alpha}{\alpha-1}} \right) \leq \ln \left(\frac{T}{\delta^{\frac{\alpha}{\alpha-1}+1}} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}_t} \phi(h', \mathcal{S}')^{\frac{\alpha}{\alpha-1}} \right) \right] \geq 1 - \frac{\delta}{T}.$$

Finally, combining the $T + 1$ bounds with a union bound give us the desired result. □

We now prove Corollaries 8 and 9 from Proposition 13 and Theorem 2'.

Corollary 8. For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any priors set $\mathbf{P} = \{\mathcal{P}_t\}_{t=1}^T$ of T prior \mathcal{P} on \mathcal{H} , for any algorithm $A : \mathcal{Z}^m \rightarrow \mathcal{M}(\mathcal{H})$, for any loss $\ell : \mathcal{H} \times \mathcal{Z} \rightarrow [0, 1]$, with proba. at least $1 - 2\delta$ on the random choice of $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_S$, we have for all prior $\mathcal{P}_t \in \mathbf{P}$

$$\underbrace{\text{kl}(R_S(h) \| R_{\mathcal{D}}(h)) \leq \frac{1}{m} \left[\frac{\|\mathbf{w} - \mathbf{v}_t\|_2^2}{\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta^3} \right]}_{\text{Equation (4.1)}} \quad \text{and} \quad \forall C > 0, R_{\mathcal{D}}(h) \leq \underbrace{\frac{1 - e^{-C R_S(h) - \frac{1}{m} \left[\frac{\|\mathbf{w} - \mathbf{v}_t\|_2^2}{\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta^3} \right]}}{1 - e^{-C}}}_{\text{Equation (4.2)}}.$$

Proof. To prove Equation (4.1) we instantiate Theorem 2' with $\phi(h, \mathcal{S}) = \exp \left[\frac{\alpha-1}{\alpha} m \text{kl}(R_S(h) \| R_{\mathcal{D}}(h)) \right]$ and $\alpha = 2$: we have with probability at least $1 - 2\delta$ over the random choice of $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_S$, for all prior $\mathcal{P}_t \in \mathbf{P}$

$$\text{kl}(R_S(h) \| R_{\mathcal{D}}(h)) \leq \frac{1}{m} \left[D_2(\mathcal{Q}_S \| \mathcal{P}_t) + \ln \left(\frac{T}{\delta^3} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}_t} e^{m \text{kl}(R_{\mathcal{S}'}(h') \| R_{\mathcal{D}}(h'))} \right) \right].$$

From Maurer (2004) we upper-bound $\mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}_t} e^{m \text{kl}(R_{\mathcal{S}'}(h') \| R_{\mathcal{D}}(h'))}$ by $2\sqrt{m}$ for each prior \mathcal{P}_t . Hence, we have

$$\text{kl}(R_S(h) \| R_{\mathcal{D}}(h)) \leq \frac{1}{m} \left[D_2(\mathcal{Q}_S \| \mathcal{P}_t) + \ln \left(\frac{2T\sqrt{m}}{\delta^3} \right) \right].$$

Additionally, the Rényi divergence $D_2(\mathcal{Q}_S \| \mathcal{P}_t)$ between two multivariate Gaussians $\mathcal{Q}_S = \mathcal{N}(\mathbf{w}, \sigma^2 \mathbf{I}_d)$ and $\mathcal{P}_t = \mathcal{N}(\mathbf{v}_t, \sigma^2 \mathbf{I}_d)$ is well known: its closed-form solution is $D_2(\mathcal{Q}_S \| \mathcal{P}_t) = \frac{\|\mathbf{w} - \mathbf{v}_t\|_2^2}{\sigma^2}$. This leads to the given Equation (4.1).

By rewriting Equation (4.1) and with Proposition 13 where $\kappa = \frac{1}{m} \left[\frac{\|\mathbf{w} - \mathbf{v}_t\|_2^2}{\sigma^2} + \ln \left(\frac{2T\sqrt{m}}{\delta^3} \right) \right]$, we have

$$\begin{aligned} R_{\mathcal{D}}(h) &\leq \sup_{R_{\mathcal{S}}(h) \leq r \leq 1} \left\{ r \mid \text{kl}(R_{\mathcal{S}}(h) \| r) \leq \frac{1}{m} \left[\frac{\|\mathbf{w} - \mathbf{v}_t\|_2^2}{\sigma^2} + \ln \left(\frac{2T\sqrt{m}}{\delta^3} \right) \right] \right\} \\ &= \inf_{C > 0} \left[\frac{1 - \exp \left[-C R_{\mathcal{S}}(h) - \frac{1}{m} \left[\frac{\|\mathbf{w} - \mathbf{v}_t\|_2^2}{\sigma^2} + \ln \left(\frac{2T\sqrt{m}}{\delta^3} \right) \right] \right]}{1 - \exp(-C)} \right]. \end{aligned}$$

Hence, for all $C > 0$, Equation (4.2) holds. \square

Corollary 9 *Under the same assumptions as in Corollary 8, with proba. at least $1 - \delta$ over the random choice of $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_S$, we have for all prior $\mathcal{P}_t \in \mathbf{P}$*

$$\underbrace{\text{kl}(R_{\mathcal{S}}(h) \| R_{\mathcal{D}}(h)) \leq \frac{1}{m} \left[\frac{\|\mathbf{w} + \boldsymbol{\epsilon} - \mathbf{v}_t\|_2^2 - \|\boldsymbol{\epsilon}\|_2^2}{2\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta} \right]}_{\text{Equation (4.3)}} \text{ and } \forall C > 0, \underbrace{R_{\mathcal{D}}(h) \leq \frac{1 - e^{-C R_{\mathcal{S}}(h) - \frac{1}{m} \left[\frac{\|\mathbf{w} + \boldsymbol{\epsilon} - \mathbf{v}_t\|_2^2 - \|\boldsymbol{\epsilon}\|_2^2}{2\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta} \right]}}{1 - e^{-C}}}_{\text{Equation (4.4)},}$$

where $\boldsymbol{\epsilon} \sim \mathcal{N}(\mathbf{0}, \sigma^2 \mathbf{I}_d)$ is a Gaussian noise such that $\mathbf{w} + \boldsymbol{\epsilon}$ are the weights of $h \sim \mathcal{Q}_S$ with $\mathcal{Q}_S = \mathcal{N}(\mathbf{w}, \sigma^2 \mathbf{I}_d)$.

Proof. We instantiate Theorem 1(i) of [Rivasplata et al. \(2020\)](#) with $\phi(h, \mathcal{S}) = \exp[m \text{kl}(R_{\mathcal{S}}(h) \| R_{\mathcal{D}}(h))]$, however, we apply the theorem T times for each prior $\mathcal{P}_t \in \mathbf{P}$ (with a confidence δ/T instead of δ). Hence, for each prior $\mathcal{P}_t \in \mathbf{P}$, we have with probability at least $1 - \delta/T$ over the random choice of $\mathcal{S} \sim \mathcal{D}^m$ and $h \sim \mathcal{Q}_S$

$$\text{kl}(R_{\mathcal{S}}(h) \| R_{\mathcal{D}}(h)) \leq \frac{1}{m} \left[\ln \left[\frac{\mathcal{Q}_{\mathcal{S}}(h)}{\mathcal{P}_t(h)} \right] + \ln \left(\frac{T}{\delta} \mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}_t} e^{m \text{kl}(R_{\mathcal{S}'}(h') \| R_{\mathcal{D}}(h'))} \right) \right].$$

From [Maurer \(2004\)](#), we upper-bound $\mathbb{E}_{\mathcal{S}' \sim \mathcal{D}^m} \mathbb{E}_{h' \sim \mathcal{P}_t} e^{m \text{kl}(R_{\mathcal{S}'}(h') \| R_{\mathcal{D}}(h'))}$ by $2\sqrt{m}$. For each prior \mathcal{P}_t , we have

$$\text{kl}(R_{\mathcal{S}}(h) \| R_{\mathcal{D}}(h)) \leq \frac{1}{m} \left[\ln \left[\frac{\mathcal{Q}_{\mathcal{S}}(h)}{\mathcal{P}_t(h)} \right] + \ln \left(\frac{2T\sqrt{m}}{\delta} \right) \right].$$

Given the fact that the probability density functions of \mathcal{Q}_S and \mathcal{P}_t for $h \sim \mathcal{Q}_S$ (with the weights $\mathbf{w} + \boldsymbol{\epsilon}$) can be written as

$$\mathcal{Q}_S(h) = \left[\frac{1}{\sigma\sqrt{2\pi}} \right]^d \exp \left(-\frac{1}{2\sigma^2} \|\boldsymbol{\epsilon}\|_2^2 \right) \text{ and } \mathcal{P}_t(h) = \left[\frac{1}{\sigma\sqrt{2\pi}} \right]^d \exp \left(-\frac{1}{2\sigma^2} \|\mathbf{w} + \boldsymbol{\epsilon} - \mathbf{v}_t\|_2^2 \right),$$

we can derive a closed-form expression of $\ln \left[\frac{\mathcal{Q}_{\mathcal{S}}(h)}{\mathcal{P}_t(h)} \right]$. Indeed, we have

$$\begin{aligned} \ln \left[\frac{\mathcal{Q}_{\mathcal{S}}(h)}{\mathcal{P}_t(h)} \right] &= \ln [\mathcal{Q}_{\mathcal{S}}(h)] - \ln [\mathcal{P}_t(h)] \\ &= \ln \left(\left[\frac{1}{\sigma\sqrt{2\pi}} \right]^d \exp \left(-\frac{1}{2\sigma^2} \|\boldsymbol{\epsilon}\|_2^2 \right) \right) - \ln \left(\left[\frac{1}{\sigma\sqrt{2\pi}} \right]^d \exp \left(-\frac{1}{2\sigma^2} \|\mathbf{w} + \boldsymbol{\epsilon} - \mathbf{v}_t\|_2^2 \right) \right) \\ &= -\frac{1}{2\sigma^2} \|\boldsymbol{\epsilon}\|_2^2 + \frac{1}{2\sigma^2} \|\mathbf{w} + \boldsymbol{\epsilon} - \mathbf{v}_t\|_2^2 = \frac{1}{2\sigma^2} \left[\|\mathbf{w} + \boldsymbol{\epsilon} - \mathbf{v}_t\|_2^2 - \|\boldsymbol{\epsilon}\|_2^2 \right]. \end{aligned}$$

Finally, a union bound argument on the T bounds gives us the claimed Equation (4.3). Secondly, we obtain Equation (4.4) by rewriting Equation (4.3) and with Proposition 13 where $\kappa = \frac{1}{m} \left[\frac{\|\mathbf{w} + \boldsymbol{\epsilon} - \mathbf{v}_t\|_2^2 - \|\boldsymbol{\epsilon}\|_2^2}{2\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta} \right]$, we have

$$\begin{aligned} R_{\mathcal{D}}(h) &\leq \sup_{R_{\mathcal{S}}(h) \leq r \leq 1} \left\{ r \mid \text{kl}(R_{\mathcal{S}}(h) \| r) \leq \frac{1}{m} \left[\frac{\|\mathbf{w} + \boldsymbol{\epsilon} - \mathbf{v}_t\|_2^2 - \|\boldsymbol{\epsilon}\|_2^2}{2\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta} \right] \right\} \\ &= \inf_{C > 0} \left[\frac{1 - \exp \left[-C R_{\mathcal{S}}(h) - \frac{1}{m} \left[\frac{\|\mathbf{w} + \boldsymbol{\epsilon} - \mathbf{v}_t\|_2^2 - \|\boldsymbol{\epsilon}\|_2^2}{2\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta} \right] \right]}{1 - \exp(-C)} \right]. \end{aligned}$$

Hence, for all $C > 0$, Equation (4.4) holds. □

11 Proof of Corollary 10

Corollary 10. *For any distribution \mathcal{D} on \mathcal{Z} , for any hypothesis set \mathcal{H} , for any priors' set $\mathbf{P} = \{\mathcal{P}_t\}_{t=1}^T$ of T priors \mathcal{P} on \mathcal{H} , for any loss $\ell : \mathcal{H} \times \mathcal{Z} \rightarrow \{0, 1\}$ we have, with probability at least $1 - 2\delta$ over the random choice of $\mathcal{S} \sim \mathcal{D}^m$ and $\{h_1, \dots, h_n\} \sim \mathcal{Q}^n$, we have simultaneously*

$$\text{kl} \left(\frac{1}{n} \sum_{i=1}^n R_{\mathcal{S}}(h_i) \parallel_{h \sim \mathcal{Q}} \mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{S}}(h) \right) \leq \frac{1}{n} \ln \frac{2}{\delta}, \quad (\text{Equation (4.5)})$$

and, for all prior $\mathcal{P}_t \in \mathbf{P}$, for all posterior \mathcal{Q} ,

$$\text{kl} \left(\mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{S}}(h) \parallel_{h \sim \mathcal{Q}} \mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{D}}(h) \right) \leq \frac{1}{m} \left[\frac{\|\mathbf{w}' - \mathbf{v}_t\|_2^2}{2\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta} \right]. \quad (\text{Equation (4.6)})$$

Proof. We instantiate Equation (2.4) for each prior \mathcal{P}_t with $\mathcal{Q} = \mathcal{N}(\mathbf{w}', \sigma^2 \mathbf{I}_d)$ and $\mathcal{P}_t = \mathcal{N}(\mathbf{v}_t, \sigma^2 \mathbf{I}_d)$ with a confidence δ/T instead of δ . Indeed, for each prior \mathcal{P}_t , with probability at least $1 - \frac{\delta}{T}$ over the random choice of $\mathcal{S} \sim \mathcal{D}^m$, we have for all posterior \mathcal{Q} on \mathcal{H} ,

$$\text{kl} \left(\mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{S}}(h) \parallel_{h \sim \mathcal{Q}} \mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{D}}(h) \right) \leq \frac{1}{m} \left[\text{KL}(\mathcal{Q} \parallel \mathcal{P}_t) + \ln \frac{2T\sqrt{m}}{\delta} \right].$$

Note that the closed form solution of the KL divergence between the Gaussian distributions \mathcal{Q} and \mathcal{P}_t is well known, we have $\text{KL}(\mathcal{Q} \parallel \mathcal{P}_t) = \frac{\|\mathbf{w}' - \mathbf{v}_t\|_2^2}{2\sigma^2}$. Then, by applying a union bound argument over the T bounds obtained with the T priors \mathcal{P}_t , we have with probability at least $1 - \delta$ over the random choice of $\mathcal{S} \sim \mathcal{D}^m$, for all prior $\mathcal{P}_t \in \mathbf{P}$, for all posterior \mathcal{Q}

$$\text{kl} \left(\mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{S}}(h) \parallel_{h \sim \mathcal{Q}} \mathbb{E}_{h \sim \mathcal{Q}} R_{\mathcal{D}}(h) \right) \leq \frac{1}{m} \left[\frac{\|\mathbf{w}' - \mathbf{v}_t\|_2^2}{2\sigma^2} + \ln \frac{2T\sqrt{m}}{\delta} \right]. \quad (\text{Equation (4.6)})$$

Additionally, we obtained Equation (4.5) by a direct application the Theorem 2.2 of [Dziugaite and Roy \(2017\)](#). Finally, from a union bound of the two bounds in Equations (4.5) and (4.6) gives the claimed result. □

12 Experiments

We give more details on the architectures and the hyperparameters that we consider in Section 4.3.

Training the MNIST-like datasets. We train a variant of the *All Convolutional Network* of [Springenberg et al. \(2015\)](#). The model is a 3-hidden layers convolutional network with 96 channels. We use 5×5 convolutions with a padding of size 1, and a stride of size 1 everywhere except on the second convolution where we use a stride of size 2. We adopt the Leaky ReLU activation functions after each convolution. Lastly, we use a global average pooling of size 8×8 in order to obtain the desired output size. Furthermore, the weights are initialized with Xavier Normal initializer while the biases are left initialized with the default initializer. We learn the prior weights by using Adam optimizer for 10 epochs with a learning rate of 0.001 and a batch size of 32. Finally, we fix the variance of the prior and the posterior to $\sigma^2 = 0.0001$.

Training the CIFAR datasets. We train ResNet-20, *i.e.*, a ResNet network from [He et al. \(2016\)](#) with 20 layers. The weights are initialized with Kaiming Normal initializer and the initialization of the bias is the default one. The prior weights are learned for 100 epochs and the posterior weights for 10 epochs with a batch size of 32 by using Adam optimizer as well. Additionally, the learning rate to learn the prior for CIFAR-10 is 0.01. Finally, similarly to the training of the MNIST-like datasets, we fix the variance to $\sigma^2 = 0.0001$.