



## Domestication of South American Camelids: new Archaeological investigations

Manon Le Neün, Elise Dufour, Nicolas Goepfert, Thomas Cucchi

### ► To cite this version:

Manon Le Neün, Elise Dufour, Nicolas Goepfert, Thomas Cucchi. Domestication of South American Camelids: new Archaeological investigations: From hunters to herders in the central Andes. the World Meeting on Heritage, Sciences and Technologies and The Scientific Symposium Frontiers in Heritage Science, Feb 2019, Paris, France. hal-03146224

**HAL Id: hal-03146224**

**<https://hal.science/hal-03146224>**

Submitted on 18 Feb 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Domestication of South American Camelids: new Archaeological investigations

## From hunters to herders in the central Andes



Manon LE NEÜN<sup>1@</sup>, Elise DUFOUR<sup>1</sup>, Nicolas GOEPFERT<sup>2</sup> and Thomas CUCCHI<sup>1</sup>

### Context

Archaeological evidence suggests that South American Camelids (SAC), llamas (*Lama glama*) and alpacas (*Vicugna pacos*), were domesticated from potential wild ancestors, guanacos (*Lama guanicoe*) and vicunas (*Vicugna vicugna*), around 6000 years ago in the Andes. However, their domestication process is still poorly understood. Their taxonomic identification in the archaeological records remains challenging for zooarchaeologists due to their morphological similarity. A clear distinction between the four species is highly required to further document their domestication process.

### First results

We first investigated how much tooth morphology could contribute to identify the taxonomic diversity of SAC using 2D Geometric morphometrics (GMM) of teeth. This approach proved unable to distinguish current wild and domestic taxa. However, we found a great divergence between the archaeological and modern variation of SAC, suggesting that the camelids Pre-hispanic diversity could not be explored using modern SAC as comparatives.



### New investigations

To bring new insights into the SAC domestication process, my PhD research project will focus on the unique site of Telarmachay (Peru) where evidence of a domestication process, 2000 years before the rest of Andean area, has been proposed. We will use a multidisciplinary approach combining 3D geometrics morphometrics on post-cranial bones and isotopics geochemistry analyses to unveiling the domestication process.

