



HAL
open science

Late Neanderthal short-term and specialized occupations at the Abri du Maras (South-East France, level 4.1, MIS 3)

Marie-Helene Moncel, M. Gema Chacón, Delphine Vettese, Marie-Agnès Courty, Aleix Eixea, Camille Daujeard, Paul Fernandes, Ethel Allué, Bruce Hardy, Rosalia Gallotti, et al.

► To cite this version:

Marie-Helene Moncel, M. Gema Chacón, Delphine Vettese, Marie-Agnès Courty, Aleix Eixea, et al.. Late Neanderthal short-term and specialized occupations at the Abri du Maras (South-East France, level 4.1, MIS 3). *Archaeological and Anthropological Sciences*, 2021, 13 (3), pp.45. 10.1007/s12520-021-01285-5 . hal-03146408

HAL Id: hal-03146408

<https://hal.science/hal-03146408>

Submitted on 31 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Late Neanderthal short-term and specialized occupations at the Abri du Maras 2 (South-East France, level 4.1, MIS 3)

3
4 Marie-Hélène Moncel¹(*), M. Gema Chacón¹⁻²⁻³, Delphine Vettese¹⁻⁴⁻⁵, Marie-Agnès Courty⁶,
5 Camille Daujeard¹, Aleix Eixea⁷⁻⁸, Paul Fernandes⁹, Ethel Allué²⁻³, Bruce Hardy¹⁰, Florent
6 Rivals²⁻³⁻¹⁴, Philippe Béarez¹¹, Rosalia Gallotti¹²⁻¹³, Simon Puaud¹

7
8 ¹ UMR7194 – HNHP (CNRS – MNHN –UPVD – Sorbonne Universités), 1 rue René Panhard, 75013 Paris,
9 France (* Correspondant author : marie-helene.moncel@mnhn.fr)

10 ² Institut Català de Paleoecologia Humana i Evolució Social (IPHES), Zona Educacional 4, Campus Sescelades
11 URV (Edifici W3), 43007 Tarragona, Spain

12 ³ Universitat Rovira i Virgili (URV), Departament d'Història i Història de l'Art, Av. Catalunya 35, 43002
13 Tarragona, Spain

14 ⁴ Università di Ferrara, Dipartimento di Studi Umanistici, Sezione di Scienze Preistoriche e Antropologiche,
15 Corso Ercole I D'Este 32, Ferrara, 44100, Italy

16 ⁵ Instituto Internacional de Investigaciones Prehistóricas de Cantabria, (Universidad de Cantabria, Santander,
17 Gobierno de Cantabria), Santander, 39005, Spain

18 ⁶ CNRS-UPR 8521 PROMES. Procédés et Matériaux Solaires. Rambla de la Thermodynamique. Tecnosud,
19 Perpignan, FR 66100, France

20 ⁷ Universitat de València, Departament de Prehistòria, Arqueologia i Història Antiga, Av. Blasco Ibañez 28,
21 46010 Valencia, Spain

22 ⁸ Grupo de Investigación Prehistoria del Mediterráneo Occidental (PREMEDOC), Universitat de València,
23 Spain

24 ⁹ Paleotime, Villars-de-Lens, France

25 ¹⁰ Dept. of Anthropology, Kenyon College, Gambier, Ohio, USA

26 ¹¹ Archéozoologie, archéobotanique : sociétés, pratiques et environnements (UMR 7209, AASPE), MNHN,
27 CNRS, 55 rue Buffon, CP 56, 75005 Paris, France

28 ¹² Université Paul Valéry Montpellier 3, CNRS, UMR 5140 Archéologie des sociétés méditerranéennes,
29 Campus Saint Charles, 34199 Montpellier (France)

30 ¹³ Université Bordeaux 1 UMR 5199 PACEA-PPP, Bâtiment B18 allée Geoffroy Saint-Hilaire CS 50023 F -
31 33615 Pessac Cedex (France)

32 ¹⁴ ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain

33
34
35

This article is part of the Topical Collection on *Settlement Patterns Dynamics of the Middle Paleolithic and Middle Stone Age*

36 37 38 **Abstract**

39
40 Level 4.1 from the Abri du Maras (Ardèche, France) is chronologically attributed to the beginning of MIS 3
41 and is one example of late Neanderthal occupations in the southeast of France. Previous work on the faunal
42 and lithic remains suggest that this level records short-term hunting episodes of reindeer associated with
43 fragmented lithic reduction sequences. During fieldwork, the high density of the material did not allow
44 identification of clear spatial patterning of these activities. In order to try to decipher the palimpsest of
45 these short-term occupations, we combined contextual micro-stratigraphic analysis with interdisciplinary
46 and methodological approaches to obtain high-resolution intra-site spatial data. The former was performed
47 by studying microfacies variability of occupation layers at meso to microscales. A combination of spatial
48 techniques based on GIS and kernel density analysis, along with faunal and lithic refitting was used and
49 focused on the horizontal distribution of the whole archaeological assemblage.

50 The results demonstrate that quantitative approaches, associated with the interdisciplinary empirical
51 processing of data, are suitable and adequate methods for describing the spatio-temporal formation of the
52 archaeological assemblage. This integrated approach allowed us to identify a temporal succession of
53 occupational events marked by distinctive anthropic imprints in the host matrix in well-preserved activity

54 areas. The analysis of their spatial patterns reveals differential treatment of lithic and faunal remains. We
55 describe the possible organization of the settlement patterns dynamics of these specialized short-term
56 occupations.

57

58 **Keywords:** Middle Palaeolithic, late Neanderthal, Abri du Maras, France, fauna and lithic refits, spatial
59 pattern analysis, short-term occupations

60

61 1. Introduction

62

63 The interpretation of lithic and faunal assemblages in terms of human settlement patterns is one of the
64 main issues in Prehistoric Archaeology for reconstructing behavioural strategies. Most Middle Palaeolithic
65 assemblages are viewed as palimpsests formed from successive episodes of occupation that cannot be
66 individually identified (Bailey 2007). They originate from the convergence into a single space of
67 anthropogenic events with depositional, erosional and post-depositional processes (Lucas 2005, 2012)
68 which is exclusively defined by stratigraphic criteria (Vaquero 2008). Even in archaeological contexts of high
69 temporal resolution, each stratigraphically-expressed palimpsest is acknowledged to integrate successive
70 occupation events (i.e. Julien et al. 1992; Vaquero 2008, Malinsky-Buller et al. 2011; Henry 2012; Vaquero et
71 al. 2012a; Chacón et al. 2015; Machado et al. 2013, 2019; Mallol et al. 2019; Real et al. 2019).

72

73 A variety of methods have been used to attempt to identify behaviourally significant spatial patterns in
74 terms of activities performed by Neanderthals (i.e. Adler and Conard 2005; Henry 2012; Bargalló et al. 2016;
75 2020; Chacón et al. 2015; Dibble et al. 2018; Ekshtain et al. 2019; Marín et al. 2019). The critical issue of the
76 palimpsest relevance has been more specifically approached by refitting analysis of lithics and faunal
77 remains and spatial analysis (Henry 2012) in order to reconstruct the archaeological assemblage formation
78 from a spatio-temporal perspective. Quantitative approaches, particularly using geostatistical tools, have
79 recently provided a more refined analysis of human occupational patterns (Romagnoli and Vaquero 2016;
80 Gabucio et al. 2018; Marín et al. 2019)

81

82 Previous interdisciplinary studies on the Maras 4.1 archaeological assemblage have provided solid data to
83 recognize distinctive events of short-term and specialized occupation (Moncel et al. 2010, 2014; Hardy et al.
84 2013, 2020; Vettese 2014, Richard et al. 2015; Vettese et al. 2017; Daujeard et al. 2019a). The combination
85 of zooarchaeological, cementochronological and tooth microwear analyses reveals a single-species
86 dominated spectrum, with catastrophic mortality and repeated autumnal deaths during a cold and humid
87 phase of MIS 3. This integrated approach points to short-term hunting episodes of reindeer herds. The
88 technological analysis strategies indicate the fragmentation of the reduction processes in a local and semi-
89 local perimeter around the cave. The lithic procurement shows a flexible Neanderthal strategy combining
90 forethought and planning. Results on micro-wear traces and residues point to mainly non-specialized
91 activities. Considering the abundance of flint sources in the area, Neanderthal mobility appears to have
92 been organized around the procurement of other important resources, including fauna and plants for food
93 or tools as well.

94

95 Our objective here is to apply an integrated approach to the lithic and faunal materials based on refitting
96 and spatial distribution in order to more fully understand the behaviours, activities and duration of the
97 occupations of late Neanderthals in south-eastern France. We intend to interrogate our previously
98 proposed circulating model of regional mobility that strongly depended upon a detailed knowledge of the
99 regional micro-topography and the seasonally available resources (Daujeard and Moncel 2010; Moncel and
100 Daujeard 2012; Daujeard et al. 2012; Daujeard et al. 2016). We also aim to challenge our initial hypothesis
101 of reindeer mass hunting strategies (Daujeard et al. 2019a), in order to refine our interpretation of the site
102 function for the level 4.1 occupation phase.

103 This should also help us to better understand the topographical constraints of this widely opened shelter
104 adjacent to a small, narrow valley on the types of occupation, their spatial pattern and their duration in
105 comparison to other examples of short-term Middle Paleolithic occupation (i.e. Vallverdú et al. 2005;
106 Costamagno et al 2006; Hovers et al. 2011; Sánchez-Hernández et al. 2014; Bargalló et al. 2016; 2020; Castel
107 et al. 2017; Villaverde et al. 2017; Mallol et al. 2019; Real et al. 2019; Mora Torcal et al. 2020; Picin et al.
108 2020; Bargalló et al. 2020).

109 Moreover, the chronological attribution of the level 4.1 at the beginning of the MIS 3 between 46 ± 3 ka and
110 40 ± 3 ka (Richard et al. 2015) places the occupation among the most recent Neanderthal occupations in
111 Western Europe (Higham et al. 2014). The Abri du Maras is not isolated in the area and it is close to other
112 late Middle Palaeolithic occupations, such as Le Figuier, Mandrin Cave (Vandeveldt et al. 2017), Baume
113 Néron (Defleur et al. 1994), Saint Marcel (Moncel et al. 2008; Szmids et al., 2010) and Abri des Pêcheurs.

114 Likewise, the nearby Chauvet cave with its Aurignacian painting attests to the presence of Modern Humans
115 in the region at 32-36 ka (Nomade et al. 2016). It is now well established that the replacement of
116 Neanderthal by Modern Humans between 50 and 40 ka was characterized by a regionalization of lithic
117 traditions, as shown for instance by various “transitional industries”. Technological strategies and land use
118 patterns varied greatly in different regions (Quina/Levallois technologies for instance in southwestern
119 France (Delagnes and Rendu 2011, Rendu et al. 2012), or “bout coupé” in England (Ruebens and Wragg
120 Sykes 2016). Therefore, the level 4.1 from the Abri du Maras site could provide one of the latest traces of
121 Neanderthals in the area.

122
123

124 **2. The Abri du Maras site and the level 4.1**

125

126 **2.1 The site**

127 The site of Abri du Maras (170 m a.s.l.) is located on the right bank of the Middle Rhône Valley (**Fig. 1**), in a
128 small dry valley of the Ardèche River, at the outer edge of the gorges. This rockshelter opens to the south-
129 east and is the relict of a large cave at the end of the small valley (Debard 1988; Moncel et al. 2010, 2015).

130

131 The first archaeological investigations at Abri du Maras revealed a stratigraphic sequence of about 3 m
132 including eight levels with Middle Palaeolithic deposits and a Levallois laminar debitage (upper part of the
133 sequence, level 1) (Gilles 1950; Combiér 1967; Debard 1988; Moncel et al. 1994). The new excavations that
134 started in 2006 have since provided a wide exposure of the succession of occupation layers. The excavations
135 were conducted with classical methodologies, using a square meter grid and plotting each object with a
136 total station. A GIS database recorded the spatial distribution (vertical and horizontal) information.
137 Sediments were regularly dry and wet screened for microvertebrates, micro-flakes and shells.

138

139 The new excavations include the middle and lower parts of the sequence partially excavated during past
140 fieldwork (previously assigned levels 4 to 8). Six large units or layers have been identified but their
141 relationship with previously assigned levels is unclear due to loss of overlying infill. Stratigraphic layers 6
142 and 5 are at present the oldest units currently known. They lie on the limestone substratum, visible on
143 different parts of the excavated areas far from the actual shelter overhang. The overlying layer 4 is a 0.5 to 1
144 m thick, silty to sandy-silty loessic deposit, mostly of aeolian origin. The coarse components come from frost
145 shattering and gravitational fall of the limestone walls of the rock shelter. The loessic fine fraction originates
146 primarily from the wind erosion of the fluvio-glacial terraces of the Middle Rhône valley during the cold and
147 dry conditions of MIS 3 (Puaud et al. 2015). The initial loessic deposits have been weakly affected by gentle
148 runoff and pedogenic transformations that are essentially expressed by an intense earthworm bioturbation
149 and moderate redistribution of calcium carbonate along root channels (Puaud et al. 2015). The excavation
150 has revealed the occurrence of two archaeological levels, 4.2 and 4.1, clearly separated by a sterile loessic
151 layer, which have been exposed over 50 m² in the front of the remaining shelter (Moncel et al. 2014). The
152 two distinctive occupation phases both have a high density of artefacts, the lack of *in situ* combustion
153 evidence, the occurrence of patchy greyish to reddish-brown microaggregate lenses that have been
154 suspected to be hearth remains during the excavation. Rare artefacts have been encountered in the
155 overlying layers (3, 2, and 1). U/Th dating applied to bones from the top of unit 5/bottom of layer 4 yielded
156 ages of 72 ± 3 ka, 87 ± 5 ka, 89 ± 4 ka, and 91 ± 4 ka (Moncel et al. 1994; Moncel and Michel 2000). New
157 preliminary ESR-U/Th dating of ungulate teeth confirmed the chronology of layer 5 (90 ± 9 ka) but gave
158 more recent ages for layer 4, attributed at the most to MIS 3. At present, the upper part of layer 4 defined
159 as level 4.1 can be framed between 46 ± 3 ka (n=2) and 40 ± 3 ka, while 4.2 level would range from 55 ± 2 ka
160 (n=3) to 42 ± 3 ka, therefore both belonging to the MIS 3 (Richard et al. 2015). This chronological
161 attribution is consistent with Heinrich events H5 and H4, correlated to the D-O #12 event, a record in the
162 Villars speleothems (Genty et al. 2010) of a woody steppe with a relatively warm and humid climate in
163 contrast to the earlier periods.

164

165 **2.2 Layer 4 and level 4.1**

166

167 At the upper part of the large layer 4, the excavation of sub-level 4.1 (34 m² surface) (**Fig. 2, 3**) exposed a
168 dense accumulation of archaeological remains throughout the 20 cm thick deposit. Its meticulous
169 excavation has not revealed the occurrence of any significant change in colour, structure or texture that
170 would be closely associated with the occurrence of the archaeological items, emphasizing the homogeneity
171 of the clayey-sandy silt sedimentary matrix (see **Fig. S1**). The accumulation of the archaeological materials
172 was observed to correlate with increased amount of coarse components issued from the pedogenic
173 weathering of the limestone cliff together with the presence of collapsed limestone slabs for levels 4.1 and
174 4.2. This repeated joint occurrence has allowed us to associate each phase of human presence with specific
175 environmental conditions that were favourable to vegetation colonization of the limestone cliff and soil
176 development on the uppermost limestone plateau.

177
178 In contrast, the fine silt texture typical of intact aeolian dust, the weak effect of pedogenesis and the lack of
179 coarse limestone fraction in the sterile deposit that accumulated between 4.1 and 4.2 levels indicates that
180 the site was not occupied when the climate was colder and drier. Microvertebrate remains, and pollen
181 grains are very rare at the site due to poor preservation, whereas charcoals are commonly encountered
182 throughout the two occupation layers showing in general a good preservation independently of a highly
183 variable abundance. The number of charcoal fragments in level 4.1 is low with a dominance of *Pinus*
184 *sylvestris* type and *Betula* sp. (Daujeard et al. 2019a). A lack of preferential orientation of the archaeological
185 materials, confirmed by statistical tests on the trends in orientation of the materials shows the minimal
186 effect of post-depositional processes (**Fig. S2**). Moreover, the occurrence of some herbivore bones found in
187 close anatomical association, with no evidence of post-fragmentation and broken bones *in situ* by post-
188 depositional factors, along with the abundance of small lithic and faunal microremains (≤ 1 cm) confirm the
189 very low impact of post-depositional processes (**Fig. 3**).

190
191

192 **2.2.1 The faunal assemblage of level 4.1**

193

194 The small and large mammal assemblage that is dominated by reindeer (88% of the NISP, representing 16
195 individuals) with a minor occurrence of horse, bison, giant deer, red deer, ibex and lagomorphs, highlights
196 open landscapes under cold and arid climate. There are no carnivore remains (Daujeard et al. 2019a).

197 For reindeer, long bones largely dominate the accumulation and, among them, the richest elements in meat
198 and marrow, indicate primary carcass processing on the kill site with subsequent selective transport of the
199 carcass elements. The scarcity of the axial skeleton and of the limb extremities is interpreted as
200 abandonment at the kill site and/or an *in situ* differential and/or spatial preservation or treatment of spongy
201 bones (bone grease extraction). Presence of cut and percussion marks on the remains of all types of species
202 indicates that all the steps of the carcass treatment were processed *in situ*: skinning, dismembering and
203 removing tendons, periosteum, flesh, marrow and possibly bone grease extraction.

204 The high number of reindeer individuals, catastrophic mortality profile, restricted season of death and
205 intensive butchery may correspond with a planned mass hunting strategy of a particular species, with
206 repeated autumnal deaths. The zooarchaeological data highlight diverse subsistence activities managed at
207 the site, including the use of bones for fuel, with the high rates of carbonised bone fragments (majority less
208 than 25 mm long) and the *in situ* consumption of meat and marrow. The presence of non-specialised
209 activities differs from other Neanderthal reindeer-dominated assemblages, often connected with
210 specialised sites such as hunting camps, with evidence of food storage and no or scant use of fire, i.e.
211 Salzgitter-Lebenstedt (Gaudzinski and Roebroeks 2000; Gaudzinski-Windheuser, 2006), Pech de l'Azé IV
212 (Niven 2013), Jonzac (Niven et al. 2012), Roc de Marsal (Castel et al. 2017) and Les Pradelles (Costamagno et
213 al. 2006). Here, the data rather suggest the recurrent use of the site as a dwelling camp.

214

215 **2.2.2 The lithic assemblage of level 4.1**

216

217 The lithic material in level 4.1 totals 2041 items (excavations from 2006-2018 seasons) (**Table 1, Tables S1-**
218 **S4, Fig. S2**). The applied technical strategies indicate the fragmentation of the "chaîne opératoire" into local
219 and semi-local perimeters around the site (*i.e.* Moncel et al. 2014). Flint is the main raw material followed
220 by quartz and occasionally basalt, limestone, granite, sandstone, quartzite and schist. The assemblage is

221 mainly composed of flakes and fragments of flakes, including 4.3% Levallois flakes (n=88). Laminar products
222 (blades and bladelets) total 208 pieces (10.2%), including 58 Levallois blades. Points total 81 pieces (3.9%),
223 including some clear Levallois points (n=44). There are 51 cores (2.5%), Levallois, discoidal-type, on flake
224 and others. Only 22 cores can be related to the Levallois concept (preferential flake, unipolar convergent
225 removals for points and centripetal). The other cores are unifacial unipolar, discoid, Kombewa, trifacial,
226 orthogonal or crude. Most of them are not exhausted. The original form is possibly a flake for some of
227 them, and for the others a nodule, a pebble, a fragment of slab or undetermined.
228 Pebble tools are rare (one in quartz), while we total 32 entire or broken pebbles in various stones. Flake
229 tools are rare as well (n = 46, 1.1%). They are mainly scrapers and denticulates, including one Quina scraper
230 made on a thick and large flint flake. Levallois products are not retouched.
231 The presence of cortical products (both backed and unbacked) suggests that a part of the debitage has been
232 performed on the site with an introduction of already prepared cores. This is also the case for the blades
233 with both cortical and non-cortical blades and points (**Fig. 4**). There are also backed blades and bladelets. If
234 we compare the ratio of blades and points, there is an over-representation of these products compared to
235 cores. Cores are mainly smaller and few exhausted on nodules or flakes except the Levallois and some
236 centripetal cores made on flakes. A deep reduction would be necessary to be flaked again. The largest
237 products could not be produced on the site and any large by-products are present.
238 The largest flint flakes, Levallois blades (both cortical and non-cortical, mainly with unipolar schemes) and
239 points were produced elsewhere, to the north and south of the site (up to 20-30 km and then brought to the
240 shelter (**Fig. 4**). We total 55 pieces between 60 and 105 mm long (25 blades, 20 flakes and 10 points).
241 According to geological analysis applied to a sample of more than 200 pieces, 13 raw material types were
242 identified, with several types still unknown. They may have been collected from the Rhône Valley coming
243 from the Alps Mountains. The main flint types so far identified attest to: (1) flint procurement on the
244 plateau to the north of the site at a distance of 3 to 30 km (local to semi-local Barremian formations) from
245 secondary flint outcrops and conglomerates, (2) secondary collection to the south, on a plateau located
246 between 2 and 20 km from the site, involving crossing the Ardèche River (Ludian and Lutecian formations).
247 The diversity of types is due to frequent gathering from conglomerates and indicates the indiscriminate use
248 of all kinds of existing flint in each outcrop. Neanderthals collected flint while performing other activities, in
249 a perimeter of 30 km around the site as an anticipation of future domestic needs.
250 In terms of tool use, evidence of impact fractures suggests that some points, blades and flakes, could have
251 been used as projectile tips, possibly as part of a complex projectile system (Hardy et al. 2013).
252 Neanderthals exploited a wide range of resources including large mammals, fish, birds (possibly waterfowl
253 and raptors), rabbits, plants, wood, and possibly mushrooms. Twisted fibres on stone tools provide evidence
254 of making string or cordage (Hardy et al. 2020). Most of the stone tools were used for cutting, whatever
255 their size and shape (including the Quina scraper). Imported artefacts and artefacts made on the site were
256 used for the same range of activities and materials (butchery, plant and woodworking - Moncel et al. 2014).
257

258 **3. Materials and methods**

259 **3.1 Contextual stratigraphic analysis**

260 This methodology focuses on the attributes of the sedimentary matrix immediately associated with the
261 artefacts in order to examine the three-dimensional spatial relationships among objects and deposits
262 (Goldberg and Berna 2009). The aim is to identify at the finest scales the stratigraphic reality of human
263 occupation episodes of level 4.1 with their distinctive microfacies assemblage resulting from the
264 interference of depositional events and anthropic activities (Courty 2001).
265

266 **

267 As previously debated (Daujeard et al. 2019a), the contrast between the homogeneity at macroscale of the
268 level 4.1 and the dense accumulations of archaeological remains makes interpreting the imprints of human
269 occupation on the sedimentary matrix difficult. Also, the common occurrence of burnt bone fragments and
270 the abundance of dispersed charcoals but no evidence of *in situ* heating makes a link occupation with fire-
271 related activities unclear. The occurrence of greyish to reddish-brown thin lenses contrasting from the
272 embedding loessic sediment by their granular structure were suspected to be possible remains of
273

274 combustion deposits (Daujeard et al. 2019a). This suggested that the level 4.1 homogeneity could either
275 express weak impact of human activities on the host sediment during short occupation events or complete
276 alteration of anthropic attributes by post-depositional processes.
277 In the absence of a distinctive microstratification throughout level 4.1 or of clear living-floors, contextual
278 stratigraphic analysis was performed in the field by testing the link between any noticeable changes of
279 microfacies and the occurrence of human artefacts. All subtle variations of microfacies identified by
280 contrasting structure, texture or colour from the homogeneous host sediment have been exhaustively
281 sampled to perform a systematic analysis of the microresidues from the bulk sediment samples. A sample of
282 bulk matrix has also been systematically collected with each burnt bone or charcoal fragment that was
283 exposed during the excavation. The same strategy was adopted for the polymer compounds that are similar
284 to the previously identified ones in similar Neanderthal contexts (Courty et al. 2012). The polymer films,
285 filaments and aggregates were identified with the naked eye during the excavation with their host matrix.
286 . The exposure in square I6 of a 5 cm thick microstratified sublayer showing a succession of three greyish
287 brown microaggregate lenses interwoven with abundant humified plant fragments and layered faunal
288 remains with anatomical connections has provided a unique chance to perform a high resolution
289 microfacies analysis during the excavation (**Fig. 5**).
290 At an early stage of the excavation, a continuous column of undisturbed sediment blocks was sampled in a
291 benchmark section for the fabrication of thin sections for micromorphological study. The column sampling
292 was performed in an area with low density of human artifacts.
293 All the bulk samples collected along the course of the excavation were water-sieved to obtain the total
294 dispersion of the weakly resistant aggregates, clean grain surfaces and separation by sized-fractions (>2 mm,
295 2-1 mm, 1mm-500 μ m, 500-250 μ m, 250-100 μ m) after separation of light components by flotation. A
296 morphological and mineralogical classification of the water-sieved residues was performed under the
297 binocular microscope in order to perform a microfacies analysis of layer 4.1 and to identify unusual
298 elements which would contrast from the bulk components. A selection of these unusual elements was
299 subjected to scanning electron microscope and microprobe analyses (SEM-EDS).

300

301 **3.2 Spatial pattern analysis**

302

303 Level 4.1 is possibly a palimpsest of several occupations. The density and thickness of the material (Moncel
304 and Daujeard 2012; Daujeard et al. 2019a). makes it difficult to distinguish all occupational events and the
305 intensity or duration of each one (i.e. Mora Torcal et al. 2020). Based on previous results of archaeological
306 assemblage studies, we decided to test spatial patterning focused on the horizontal distribution of all the
307 remains from level 4.1 (Moncel et al., 2014; Daujeard et al. 2019a). The spatial patterning of the lithic and
308 faunal remains is then compared with the data from lithic and faunal refitting and RMU lithic analysis
309 (Bargalló et al. 2016; 2020; Romagnoli and Vaquero 2019; Fernández-Laso et al. 2020; Spagnolo et al. 2020a,
310 2020b). These analyses provide insight into some spatial behaviours of Neanderthal groups occupying the
311 rock shelter. Moreover, they provide a key to improving our knowledge of their cultural capabilities
312 (Vaquero and Pastó 2001). However, given the limited extent of currently excavated surface, the analyses do
313 not consider the entire area occupied by Neanderthals at the site. GIS analysis was carried out using ArcGIS
314 10 software (ArcMap 10.4) (ESRI 2014). Nearest Neighbour Analysis and Kernel density analyses were used
315 to identify the distribution pattern statically and visually. The research radius in the Kernel density analysis
316 was defined as half standard deviation of our data.

317

318 **3.3 Faunal refits**

319

320 Bone fragments are less often included in refitting program than lithic materials. However, to discuss both
321 the synchronic and diachronic components of transport and carcass processing events, bone refitting is
322 essential (Rosell et al. 2012, 2019; Bargalló et al. 2016; Romagnoli and Vaquero 2019). Faunal remain
323 refitting is one of the last steps of the zooarchaeological analyses, after taxonomic and anatomical
324 determinations and taphonomic studies and can aid in the understanding of the taphonomic dynamics of
325 the assemblage. Bone refitting can be complex because of species diversity within faunal assemblages and
326 the quantity of bone fragments. Furthermore, material weakness increased the post-depositional
327 fragmentation which may add to the perimortem fragmentation (Poplin 1976; Soulier 2013).

328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381

All the three-dimensionally recorded archaeological remains (n= 1946) and the $\leq 1-1.5$ cm material non-recorded collected by sieving (n= 242) are taken into account.

In our work, mechanical and anatomical refitting (articulation and bilateral pairs) focused mostly on long bones and teeth of large- and medium-size ungulates (Morín et al. 2005; Lyman 2008; Enloe 2010; Fernández-Laso 2010). Spongy undetermined fragments were excluded in systematic refitting but undetermined shaft fragments (≥ 2 cm) were included. Green bone or dry bone fracture edges help determine the homogeneity of assemblages through the identification of the *in situ* fragment movements (vertically and horizontally, intra- and inter-levels) (Enloe and David 1989; Villa and Mahieu 1991). Some refits were found during zooarchaeological analyses while other were notices during excavation. However, due to the scarcity of the long bone epiphyses and other elements of axial skeleton and carpals and tarsal, the anatomical refitting yielded few results compared to mechanical refitting, especially the bilateral pairs, which is quite impossible to identify in archaeological context. The identification of refits and anatomical connections during the excavation of level 4.1 revealed a weak pre- and post-depositional spatial disturbance of the archaeological material and offered us the possibility of distinguishing possible areas of activity (Rosell et al. 2012, 2019; Vettese 2014). Due to these numerous refits and the good spatial conservation of the elements, we established a systematic refit protocol for the long bones of Level 4.1 (Vettese 2014, Vettese et al., 2017; Daujeard et al. 2019a):

- (1) Selection and sorting of the identified remains by anatomical parts (ex. humerus, radio-ulna...taking account all the remains ≥ 2 cm);
- (2) Sorting by portions and sides (anterior, medial, posterior and lateral) (Fig. 2, Vettese et al. 2017);
- (3) Classification of all the remains by lateral side (right, left or undetermined);
- (4) Comparison of all the remains between right and left side and also with the undetermined ones, square by square and between squares;
- (5) Spatial distribution analysis of the bone refits and connections to identify the fauna-linked activity areas;
- (6) Cross-comparisons with the lithic refits and other disciplines in order to reach a comprehensive view on the type of occupation based on multiple resources.

3.4 Lithic Raw Material Units (RMU) and refit lithic analysis

The Raw Material Units (RMU) and the refit analysis allowed the reconstruction of the temporal formation of the lithic assemblage and identification of specific events of technical activity. This is defined as an activity episode related to the manipulation of lithic resources that is performed continuously, without location changes or temporal interruption (Roebroeks 1988; Vaquero 2008).

Refits are the most direct evidence to identify pieces belonging to the same RMU. However, other kinds of data must be considered. First, the macroscopic characteristics of the raw materials, including colour, grain, size, texture, inclusions and type of cortex. The technical categories, the size of the pieces, together with their spatial pattern are also used to integrate different artefacts in the same RMU (Machado et al. 2011, 2013; Vaquero et al. 2012a, 2012b; Chacón et al. 2015). In addition, RMU have been characterized by considering two main features: introducing events defined by the mode of introduction and knapping events referring to the intentional modification performed inside the site. We have also used petrographic analysis to group some RMUs on patinated flint due to the very high percentage of patinated flint material (around 58 %) in level 4.1 (Moncel et al. 2014).

For the refit analysis, we used the methods and terminology proposed by Czesla (1990) and Sisk and Shea (2008). This methodology has been already used in other Middle Paleolithic sites with very good results to reconstruct the formation and composition of the lithic assemblages (among others: El Salt, Abric del Pastor, Abric Romaní, Teixoneres cave - Roebroeks 1988; Vaquero 2008; Chacón et al. 2015; Machado et al. 2011, 2013, 2019; Vaquero et al. 2012a, 2012b, 2019; Picin et al. 2020).

Three kinds of connections are distinguished: production-sequence, breakage and retouch refits. For production and retouch refits, connection lines have been defined according to the chronological order of

382 detachments, starting with the first refitted flake to the core or tool. For the refits of broken artefacts, the
383 connection lines have been established by considering the contacting surface (Vaquero et al. 2012b). The
384 refitting procedure allows us to distinguish two forming processes: (1) sets of artefacts splitting from each
385 other by controlled conchoidal fracture (e.g., a flake whose ventral side joins to the surface of a core or the
386 dorsal surface of another flake); and (2) conjoins: fragments of artefacts broken by forces other than
387 conchoidal fractures (e.g., natural flaws, bending fractures, burning, etc.) (Sisk and Shea 2008).
388

389 A total of 2050 lithic items recorded for level 4.1 were considered for this study, discarding the badly
390 preserved pieces (with patina or thermal alteration) and pieces ≤ 1 cm long. The amount is not the same
391 that in **Table 1** (n=2041) because some microflakes from the sieving materials were identified as parts of
392 RMUs due to their macroscopic attributes.
393

394 As a final step for the faunal and lithic refits, their spatial distribution together with the one of the RMU
395 groups has been analysed. The results obtained are interpreted by integrating the multidisciplinary analysis
396 of the different resources in order to identify the activity events (technical or related to the exploitation of
397 carcasses) and their relevance in terms of occupation type for level 4.1.
398
399

400 **4. Results and interpretation**

401 **4.1 Contextual stratigraphic analysis**

402
403
404
405 The micromorphological study performed on the benchmark section covering the entire level 4.1 confirms
406 its homogeneity throughout the 20 cm thick deposit with a well-developed biogenic microstructure showing
407 dense packing of earthworm pellets. The absence of distinctive vertical channels and the uniform
408 composition of the microaggregated fine mass indicate an accumulation of biogenic reworked soil-materials
409 rather than post-depositional effects of bioturbation. This extensive deposition of previously bioturbated
410 soil-materials suggests that the limestone cliff was extensively colonized by woody vegetation with dense
411 roots that trapped the loessic sediment. Incipient *in situ* redistribution of secondary carbonates along the
412 packing porosity of the biogenic microaggregates has acted as a weak cementing agent of the deposits, thus
413 favouring their intact preservation following their accumulation at the surface. In addition to the major silt
414 component of aeolian origin, level 4.1 also displays an abundant subrounded calcareous coarse fraction (20
415 to 40 %) of local origin and a minor population of angular quartz fine sands. The surface corrosion of the
416 local coarse components that is typical of the microbial weathering occurring in root-colonized soil materials
417 contrasts with the fresh angular sands, thus suggesting a possible exogenous origin. The random
418 distribution of the limestone components and the lack of any freeze-thaw pedofeatures in the fine mass
419 indicate that level 4.1 did not suffer from significant effects of frost during its accumulation.

420 These microstructural characteristics are coherent with the close association of the archaeological materials
421 with the large slabs that collapsed from the limestone walls and partially fragmented *in situ* due to
422 weathering by root colonization, as noticed during the excavation. The formation of level 4.1 can thus be
423 concluded to have occurred under a mild temperate climate that allowed the formation of a dense
424 vegetation cover protecting the limestone cliff from the direct effect of rainfall thus offering a dry shelter to
425 the occupants. In contrast, the fine silt texture, the weak effect of pedogenesis and the lack of coarse
426 limestone fraction in the sterile loessic deposits that bracket level 4.1 indicates that the site was not
427 occupied when the limestone cliff was no longer protected by a dense vegetation cover and therefore
428 directly exposed to winds and rainfall.
429

430
431 A few diffuse, cm-thick discontinuities that are marked by a more open micro-aggregation have been
432 identified in the thin sections. These have sub-horizontal biochannels, well-preserved humidified organic-
433 fragments, slight depletion of the calcitic fine mass and of the coarse fraction, and higher abundance of the
434 angular quartz sands. This particular microfacies is typical of soil surface that was covered for a while by

435 biogenic crusts most probably due to moisture increase, thus inducing microbial activity (Courty et al.
436 1989). However, the scarcity of microartefacts (bone and charred residues) and the absence of a distinctive
437 trampled-linked microstructure do not allow us to consider these subtle discontinuities as residual living
438 floors (Gé et al. 1993).

439 These discontinuities differ from the greyish to reddish-brown microaggregate lenses by their diffuse
440 characteristics and the lack of distinctive attributes for relating them to human activities.

441 In contrast, the greyish to reddish-brown microaggregate lenses have been clearly identified in the field
442 along the course of excavation by their sharp contact with the host homogeneous sediment, their loose
443 microstructure and the common occurrence of microartefacts (**Fig.5a, b**).

444
445 Their microfacies analysis reveals a dense packing of coarser aggregates (5 mm in average) that have a high
446 structural stability and a marked hydrophobicity as compared to the homogeneous level 4.1. A great
447 diversity of abundant millimetre-sized microresidues with clean, hydrophobic surfaces have been retrieved
448 from the fine mass (**Fig.6a**). They comprise a high amount of the angular quartz sands that are similar to the
449 ones in the weak discontinuities, along with abundant polymer components, siliceous sandstones, glassy
450 breccia, metal films, fine charcoal, vitreous chars, shell fragments, flint micro-flakes, humified plant
451 fragments, whitish ash grains, burnt, calcined or pyrolyzed bone fragments and elements of flying insects.

452
453 The analysis of the water-sieved residues from the matrix associated with charcoals and chars that were
454 collected throughout level 4.1 revealed similar translucent polymer films and coloured filaments, often in
455 the form of twisted bundles.

456
457 Within these diverse microresidues, the SEM-EDS analyses showed a common occurrence of
458 nanostructured deposits of native metals which are either filling micro-cavities, forming aligned droplets, or
459 being attached as twisted films to vesicles of the microdebris (**Fig.6**). All these exotic components have been
460 previously identified in Neanderthal hearths at the Abric Romani (Courty et al. 2012) and have been since
461 shown to be traces of flash-pyrolysis by-products that are issued from lightning-triggered aerosol
462 transformation during periods of enhanced atmospheric electrification (Courty 2017). The occurrence of
463 these flash pyrolysis by-products in the chars and charcoals of level 4.1 indicates the use of wood fuel that
464 accumulated lightning-formed refractory components of enhanced conductivity in the plant tissues during
465 tree growth (Courty et al. 2020).

466
467 The microfacies of the microaggregate lenses with their heteroclite microresidues does not match any type
468 of firing by-products which have been encountered so far in present or past fires of human or natural origin
469 (Courty et al. 1989; Courty 2012; Macphail and Goldberg 2018). Furthermore, only part of the micro-
470 residues displays burning traces whereas other are unburnt, although showing metal encrustations. In
471 addition, the sharp contact of these distinctive microfacies with the surrounding and underlying sediment
472 and the lack of large earthworm channels indicate that the microaggregation was not formed by *in situ*
473 reworking of combustion deposits. In contrast to the adjacent bioturbated matrix of level 4.1, the
474 abundance of human-related microresidues refutes an origin of soil materials falling from the limestone cliff.
475 Therefore, the microaggregate microfacies must be an earth-made anthropogenic material mixed with fire
476 residues and lightning-formed organo-mineral components. The observation during the laboratory
477 treatment by moderate heating at 80°C of a new generation of sticky polymer films bounding the water-
478 sieved residues showed that active catalyzers are still present in the microaggregates and are still reactive.
479 Similar kinds of highly reactive catalyzers have been observed in recent situations of lightning impact at the
480 ground along to highly stable polymers and metal pulverisation (Courty 2017; Courty et al. 2020). Compared
481 to the reference data base, their great abundance in the microaggregate lenses of level 4.1 at the Abri du
482 Maras helps to establish their anthropogenic deposition from use of lightning-impacted organic materials.
483 This correlation is confirmed by the good match between the vertical distribution of the archaeological
484 objects and the depth of the microaggregate lenses shown by the archeostratigraphic profiles.

485
486 The contextual stratigraphic analysis of level 4.1 provides solid support to establish the multiplicity of
487 occupation episodes for a short duration, which coincides with periods of enhanced lightning in response to
488 atmospheric increase of ionized aerosols (Courty 2017; Courty et al. 2020). The absence of a continuous

489 particle-size range between the coarse and fine components of the artefacts clearly shows the lack of
490 mechanical fragmentation which would have been caused by intense trampling of long-used surfaces. This
491 moderate mechanical effects of human activity is confirmed by the microstructural properties of the
492 successive level 4.1 occupation phases. The weak anthropogenic impact associated with the abundance of
493 lightning-related microresidues suggests that the successive short-time occupation episodes share in
494 common a set of activities linked to the use of lightning products.

495
496 The highest density of lithic remains in the main part of the excavated area indicates that the occupations
497 were preferentially located at a certain distance from the shelter wall, probably at the edge of the shelter
498 roof (**Fig. 7**). As suggested by Combier (1967) and Debard (1988), at the time of level 4.1 deposition, the
499 shelter was already largely collapsed and therefore limited to the right side of the small valley. There is no
500 clear concentration of artefacts within the site, except a higher density of lithic remains close to the areas
501 with the microaggregate greyish to brown-reddish lenses at the Northeast part and at the centre of the
502 excavated area.

503 504 **4.2 Spatial patterning**

505
506 Level 4.1 has been excavated over a surface of 34 m² and a total thickness of 20 cm. Due to the dense
507 accumulation of archaeological remains over the entire surface, it was presumed that level 4.1 did not
508 present well preserved living floors (Daujeard et al. 2019a, b). After the study of the total archaeological
509 assemblage (fauna, lithics and others; **Fig. S4-8**) from different methodological approaches, we decided to
510 apply some spatial geostatistics in order to identify, from a quantitative point of view, the possible existence
511 of accumulation areas that could correspond with domestic activity areas (following the definition of
512 Stevenson 1991; Vaquero and Pastó 2001).

513
514 The Nearest Neighbour Analyses of all the remains, including faunal and lithic materials, show that the
515 spatial distribution is significantly clustered (all: score z: -55.38, p-value<0.001; bone: z-score: -40.33, p-
516 value<0.001; lithic: z-score: -31.04, p-value<0.001). To visualize the clusters, we applied a Kernel-density
517 analysis and the results show some clear accumulations of remains. These clusters are associated with the
518 higher density of archaeological remains (fauna and lithics) and the microaggregate greyish to brown-
519 reddish lenses (**Fig. 8**). The general pattern of level 4.1 shows four main accumulations areas (A to D) and
520 three secondary ones (E to G), whereas the other remains are dispersed throughout the entire occupied
521 surface. Some remains dispersed within the first interval (0–7%, lower concentration of remains) were
522 located on the areas with few artefacts present, possibly in sectors where the level is residual.

523
524 The spatial patterning of the faunal (**Fig. 9**) and lithic remains (**Fig. 10 and 11**) shows significant differences
525 despite a broadly similar general distribution. The lithics show a larger dispersion of remains (from lines 5 to
526 8) that the faunal ones (from lines 5 to 7). Thus, a greater quantity of artefacts was produced by knapping
527 activities than the exploitation of carcasses and their normal dispersion is usually ≤ 1.5-2 m around the
528 knapping locations (see refits section). In addition, the lithics present a higher number of clusters than for
529 the faunal remains, especially for the main largest/dense accumulations areas (**Fig. 8**).

530
531
532 Lithic refits show pieces that connect two accumulations (n=7 refits) whereas faunal refits show only one
533 (see section 4.3 and 4.4.2 – **Fig. 9 and 11**).

534
535 Considering the general distribution of the remains and the morphology of the shelter over time, the main
536 activities areas that are now identified are located just at the centre towards the outside part of the site.
537 The geological study and the previous excavations showed that the slanted topography of the limestone
538 substratum with a series of steps encouraged the occupants to settle a bit far from the shelter wall
539 (Combier 1967). Therefore, the successive occupations followed the shortening of the roof in order to
540 always settle at the best place, some meters away from the shelter wall. This open, well-lit area was also
541 conducive to subsistence activities. This pattern is similar to the ones found in other Middle Paleolithic sites
542 and can also be found in numerous ethnographic accounts (Binford 1978, 1996; Henry et al. 1996, 2004;

543 O'Connell 1987; Stevenson 1991; De la Torre et al. 2012; Modolo et al. 2015; Ekshtain et al. 2019;
544 Fernández-Laso et al. 2020; Spagnolo et al. 2020a, 2020b).

545 546 **4.3 Bone refits**

547
548 Based on 2188 remains analysed, we identified 222 remains (10%), split into 88 bone refits in level 4.1.
549 Mechanical refits represent the most frequent type, with 46 green bone fracture refits and 21 dry bone
550 fracture refits, whereas there were only 21 anatomical refits. The most numerous refits are those
551 composed of 2 remains (n=64 refits), followed by 3 remains (n=13 refits). Refit groups with more remains
552 (between 3 and 8) are less frequent (n=11 refits) (**Table 2**). The longest refit is anatomical and connects
553 eight elements (including seven complete ones).

554
555 Regarding all refits, 134 connection lines were identified (anatomical refits (n=47), green bone fracture
556 refits (n=62), dry bone fracture refits (n=25)). The longest distance recorded was 2.51 meters for a green
557 bone fracture connecting two bones (**Table 3**). However, the distance of most of the remains refitted was
558 equal to 0 (52 % of the refits).

559
560
561 At taxonomic level, *Rangifer tarandus* showed the largest number of refits (61 refits) incorporating 547
562 remains (**Table 4**), reflecting its dominance at the site. *Equus ferus* was the next most common with a
563 majority of anatomical refits (3/6). We also found refits of non-taxonomically identified elements (20.5 %).
564 Most of the bone refits were the partial reconstruction of long bone remains, mostly forelimbs (**Table 5**). All
565 the bone refit of the head were anatomical refits.

566
567 In horizontal distribution, the bone material followed the pattern observed with the clusters resulted of the
568 analysis of all the remains. However, areas C and D are grouped together. Most of the refits are in the main
569 cluster areas defined from the Kernel density (**Fig. 8 and 9**). These zones are not connected, except for one
570 mechanical refit, a green bone fracture, connecting the main area D with the secondary area E.

571 572 573 **4.4 RMU and lithic refits**

574
575 As we previously mentioned (see materials and method section), the percentage of patinated flint material
576 for level 4.1 is high (around 58 %). A total of 2050 lithic items were considered for this study. Patinated,
577 thermally altered, and pieces ≤ 1 cm long were excluded from this study. Nevertheless 52 RMU and 22
578 refitting groups were recognized.

579 580 **4.4.1 RMU analysis**

581
582 The 52 RMU identified total 593 pieces (26.3 % of the lithic assemblage, **Table 6**), mainly in flint and quartz.
583 Each RMU corresponds to individual technical events and they were categorized as representing either
584 introduction to the site (introducing events - IE) or modification performed on the site (knapping events -
585 KE).

586
587 Some RMUs indicate several complete or almost complete knapping sequences produced on-site. All the
588 technical categories are observed in the same RMU, including the production of the first cortical flakes and
589 the different phases of reduction sequences and cores. Others show the fragmentation of the reduction
590 sequences. The smallest RMUs are composed of a single object (isolated artefacts). These pieces were
591 introduced to the site already knapped and are interpreted as part of tool kits transported by the human
592 groups while moving throughout their territory. These tools demonstrate the fragmentation of some
593 reduction processes and the existence of planned behaviour strategies in relation to (1) raw material
594 constraints and, (2) specific needs in relation to the activities performed within and outside the site. **Table 7**
595 synthesizes and describes each identified RMU.

596

597 The IE are mainly knapping products (flakes, blades and points) of different sizes, produced by both
598 Levallois and discoidal strategies (e.g. Isol-Fl 1 to 11, Isol-Qtzite 3, Fl-1 – **Table 7 and Fig. 10b-d**). Some cores
599 are introduced in an advanced state of reduction and are abandoned at the site after yielding a small
600 number of flakes (around 2-3) (e.g. RMU Fl-16b, **Table 6**).

601
602 The KEs demonstrate the co-existence of different core technologies, with a predominance of centripetal
603 bifacial strategies (e.g. RMUs Fl-5b, Fl-13, Fl-16d to f, Qtz-1). Levallois technology only employs flint and
604 limestone with unipolar recurrent and recurrent centripetal methods most common. For other stone types
605 (i.e. quartz) the most frequent methods are orthogonal and discoidal. Nodules, angular fragments and large
606 flakes form the basis of cores.

607
608 The high degree of reduction often produces cores that are over-exhausted, making it difficult to identify
609 the method, but the knapping products present the typical attributes necessary to describe the core
610 technologies.

611
612 The technological analysis of the whole assemblage (Moncel et al. 2014), as well as the RMUs analysis,
613 shows a high degree of fragmentation of the lithic reduction processes (i.e. Vaquero et al. 2012b; Turq et al.
614 2013; Vaquero et al. 2012b). Some isolated pieces are introduced to the site already knapped while some
615 RMUs show that the main technical processes were realized on the site. However, some pieces are missing,
616 mainly predetermined flakes and cores (e.g. RMUs Fl-12, Qtz-5). Knapping products and flake tools may
617 therefore be mobile or curated pieces (i.e. Vaquero et al. 2012a, Vaquero et al. 2012b; Moncel et al., 2014;
618 Spagnolo et al. 2020b; Mayor et al. 2020; Picin et al. 2020). By contrast, some RMUs show the entire
619 reduction but not the core, indicating that a core could have been transported away from the site as a new
620 mobile tool kit.

621
622 There are many isolated artefacts produced in flint (**Table 6 and 7**). These pieces are all produced with
623 Levallois technology and are mainly flakes, blades and points. There are some in quartz, quartzite and
624 basalt, with one quartzite piece that looks like a Levallois product. The other artefacts are unretouched
625 flakes or fragments coming from discoidal and orthogonal technologies.

626
627 The spatial distribution of the RMUs does not show any special spatial pattern either by RMU, by type of
628 stone or technological composition. Their density and spatial distribution correspond to the pattern
629 observed in the spatial Kernel analysis of the lithic and faunal assemblage (**Fig. 8 and 10**). Some of them
630 present a normal dispersion of a knapping area (e.g. RMU Fl-12, Fl-15) (Newcomer and Sieveking 1980;
631 Böeda and Pelegrin 1985; Czesla, 1990 Vaquero et al. 2019), while others are spread over the whole
632 occupation area (e.g. RMU Fl-13, Fl-16). A clear example is the spatial patterning of the RMU Lim-1 (unique
633 RMU in limestone and formed only by Levallois flakes - **Table 7 and Fig. 11d**) that corresponds with the
634 main and largest accumulation identified by the Kernel analysis as well as for the lithics than the bones.
635 Consequently, this RMU may relate to one specific occupation event. The differences and overlap in RMU
636 spatial patterning support the hypothesis of repeated short-term occupations. (Vaquero et al. 2012;
637 Spagnolo et al. 2016; 2020b).

638 639 **4.4.2 Lithic refits analysis**

640
641 After the analysis of the RMU, some pieces < 1 cm long recovered from the sieving material were also
642 taking in account for the refitting.

643 A total number of 22 refits groups comprising 49 pieces have been identified. The lithic refit rate of the
644 whole lithic assemblage (n=2050) is 2.4% with 8.2% coming only from the identified RMU groups (n=593).
645 Flint and quartz present the highest number of refits. Most of them are composed of two elements (82%)
646 and with the largest comprising 4 pieces (**Table 8**).

647 The majority are knapping products, mainly flakes and flake fragments. There are only 3 cores and 1 flake-
648 tool (**Table 9**). These products come mainly from Levallois and discoidal strategies (**Fig. 11b, 11d-f**).

649 The refits give 27 connections lines (see **Table 10**). Most of the lines correspond to knapping connections
650 (55.5%), knapping fractures and conjoins (22.2 % each). Due to the difficulties in refitting, small retouch

651 flakes as well as the high number of isolated artefacts carried out to the site already knapped, it was
652 difficult to identify refits related to retouch (**Table 10, Fig. 4 and Fig 10c, d**). Where refits were possible,
653 knapping connection lines are dominant no matter the stone type. All of them following the same pattern,
654 even if), core reduction activities were carried out *in situ* (e.g. limestone).
655 The distances of the connection lines vary from several centimeters to almost 4 meters (**Table 11**). They are
656 within the normal dispersion range observed by experimental archaeology for on-site knapping sequences
657 that are normally less than 1 m in diameter and rarely exceeding 2 m (Newcomer and Sieveking 1980;
658 Böeda and Pelegrin 1985; Czesla, 1990; Vaquero et al. 2019). Most connection lines are shorter than 1 m
659 (n=17-63%), six refits (22.2%) are between 1 and 2 m, and only 4 (15%) between 2 and 4 m. The shortest
660 connection lines belong mainly to the conjoins (post-depositional breakages produce by trampling). The
661 longest connections lines are all related to knapping activities with only one refit of 3 pieces (belong to a
662 conjoin) presenting a distance for more than 1.5 m (**Fig. 11**). Since *in situ* preservation of anatomical bone
663 connections and broken artefacts suggests that natural processes have had minimal impact on the site,
664 these longer connections lines suggest intentional anthropogenic movement of some pieces during daily
665 activities,. These pieces were knapped in one area and moved to other part of the settlement and/or
666 abandoned at the place of their last use. These pieces with intra-site mobility are all products of on-site
667 knapping and are not distinguished by a particular raw material type.
668 The spatial patterning of the lithic refits reveals two clear patterns. First, the refits are clearly located in the
669 main accumulation areas identified by the Kernel analysis (**Fig. 11**). Second, there is an important number of
670 refits that connect some of the identified clusters (e.g. A with D, E with D, B with G and one between E, F
671 and B). Consequently, the lithic refits show a higher degree of articulation of the occupied surface than the
672 one observed by the faunal refits (**Fig. 9**). However, we cannot confirm that these connections between
673 areas are showing their temporal synchronicity, because (1) the unidirectionality patterns of pieces moved
674 and (2) any recycling pattern have been observed in any of these pieces (Vaquero et al. 2015; Romagnoli
675 and Vaquero 2019).

676
677

678 **5. Discussion and conclusion**

679
680 The results of this work demonstrate that quantitative approaches, associated with the interdisciplinary
681 empirical processing of data, are suitable and adequate methods for describing the spatio-temporal
682 formation of the archaeological assemblages and can be used to reconstruct human occupation patterns.
683 We identified seven different accumulations areas, all related to subsistence activities and associated with
684 remains related to the use of fire by-products (domestic areas *sensu* Stevenson 1991; Vaquero and Pastó
685 2001, **Fig. 8**), although *in situ* combustion deposits and structured hearths were not identified. This is also
686 confirmed by the results of the faunal and lithic refits and the RMU analysis (**Fig. 10 and 11**). Despite the 34
687 m² excavated surface we know from the shape of the substratum observed in previous excavations, that the
688 successive occupations followed the shelter ceiling and floor. The oldest occupations are farther from the
689 shelter wall and are located on a lower limestone step.
690 Level 4.1 of the Abri du Maras provides detailed evidence on the types of occupations under an open
691 shelter during MIS 3. This site reveals some unusual patterns for Middle Palaeolithic short-term occupations
692 (i.e. Sánchez-Hernández et al. 2014; Gabucio et al. 2018; Moncel et al. 2018, 2019; Bargalló et al. 2020;
693 Cascalheira and Picin 2020; Picin et al. 2020). Previous studies have highlighted various types of
694 Neanderthal cave and shelter occupations with a farsighted circulating model and recurrent base camps of
695 various duration, from seasonal camps to bivouacs (Daujeard and Moncel, 2010; Daujeard et al. 2016).
696 Instead, we have intensive *in situ* carcass processing, non-specialized dwelling activities with evidence of fire
697 use, and no traces of carnivore activity (Hardy et al. 2013; Daujeard et al. 2019a). Furthermore, this study
698 highlights low material displacement, more faunal refits than lithic ones, and a relatively low ratio of flake-
699 tools. This pattern is uncommon compared to the other examples of short-term occupations both in the
700 region and in Europe in general.

701
702 Short-term occupations are characterized here by the introduction of large with secondary flaking *in situ*,
703 raw material collection within 30 km of the, and behaviours organized around the exploitation of the

704 surroundings during short-term autumnal reindeer hunting episodes (can be seen in other sites *i.e.*
705 Costamagno et al. 2006; Richards et al. 2008; Gaudzinski et al. 2009; Daujeard and Moncel 2010; Britton et
706 al. 2011, 2012; Niven et al. 2012; Rendu et al. 2012; Daujeard et al. 2016, 2019b; Discamps and Royer 2016;
707 Discamps and Faivre 2017; Cascalheira and Picin 2020).

708
709 Zooarchaeological and taphonomic studies, combined with spatial analyses applied to faunal material
710 (including bone refits), is a useful tool for identifying areas of activity (Vaquero et al. 2012a, 2012b; Rosell et
711 al. 2012, 2019; Spagnolo et al. 2019). Faunal refitting allows us to test the time between carcass processing
712 events. According to our analysis, we observe minor or no displacement of bone remains, with most of the
713 refits having a distance of zero. Anatomical refits demonstrate a high fidelity of spatial distribution. Among
714 the high number of bone refits, most are green bone fractures (Daujeard et al. 2019a). Root-etching
715 alterations are the principal modifying agents of bone surfaces with half of the assemblage presenting more
716 than one-third dissolved surface. However, this did not prevent the identification of numerous refits. The
717 one dry bone fracture with a zero distance could be the consequence of sedimentary compaction without
718 visible hiatus. Together, this shows a high preservation of the spatial integrity of the occupation level with a
719 vertical distribution related to sediment compaction. It highlights the high probability that the level is a
720 palimpsest of several occupations. Almost all the bones (in particular the ones involved in refits) have no
721 traces of weathering, thus suggesting rapid burial. Moreover, we observed very few trampling marks and no
722 carnivore disturbance. This indicated also that the rapid burial of the bone remains. Overall, this suggests
723 that the excavated area has a high temporal resolution.

724
725 Most of the faunal refits have a small or a zero distance. Moreover, most of the activity areas identified are
726 not connected by bone refits, except for one green bone fracture refit involving a displacement before the
727 bone dried and before burial. Butchery activity distributions seem to be clustered indicating different
728 carcass processing episodes. All the activities of the butchering “chaîne opératoire” were carried out *in situ*
729 (Vettese et al 2017; Daujeard et al 2019a). However, the absence of hearth does not allow specifying the
730 hypothesis on the functionality of these areas (toss or drop zones, or activity areas) based on ethnographic
731 observations (Binford 1978, Rosell et al. 2012). Clusters of bone and lithics are quite similar. We also
732 observed some patterning relating possible butchery activities with the microaggregate lenses. Moreover,
733 we noted a scarcity of bone remains around the microaggregate lenses.

734
735 The implication of monospecific faunal spectra in late Neanderthal occupation contexts reflect specific
736 behaviors which particularly raises questions regarding food meat and marrow abundance in the case of this
737 specialized hunting strategies (Gaudzinski and Roebroeks 2000; Costamagno et al. 2006; Niven et al. 2012;
738 Niven 2013; Castel et al. 2017). The high frequency of reindeer in level 4.1 shows that Neanderthals used
739 the site in the autumn specifically for this task, targeting the most common species in the area (Gaudzinski
740 and Roebroeks 2000; Costamagno et al. 2006; Niven et al. 2012; Niven 2013; Castel et al. 2017). Moreover,
741 the selective transport of the most nutritionally valuable elements within the rock shelter shows planning
742 and forethought (Vettese et al. 2017; Daujeard et al. 2019a). Intact metapodials with no evidence of
743 butchery suggests a relative abundance of food and relatively short occupations of the site. This is further
744 supported by extensive long bone breakage with little evidence of exploitation of phalanges or mandibles
745 for marrow. The emphasis on bones with large amounts of marrow suggests a period of abundance of meat
746 and marrow. Bone retouchers are absent indicating that bone was not reused after butchery. All this leads
747 to evidence of several rather short occupations that are repeated over time and mostly centered on the
748 reindeer carcass processing.

749
750 The technological attributes observed on the RMUs and the refits confirm the preliminary results of the
751 technological analysis of the whole lithic assemblage (Moncel et al. 2014) and emphasize several main
752 features. First, the high degree of fragmentation of the reduction sequences and the high number of
753 isolated artefacts brought to the site already knapped are evidence of planning ahead based on anticipated
754 raw materla constraints and subsistence activities (Kuhn 1992, 1995). Second, the evidence of partial
755 reduction sequences could be interpreted as a feature of repeated short-term occupations in relation to
756 the hunting of reindeer (Turq et al. 2013; Mayor et al. 2020; Picin et al 2020). Moreover, the variability of

757 RMUs spatial distribution (see **Fig. 10, Table 7**) with some more clustered and some more scattered,
758 reflects a series of short term occupations.
759

760 The spatial patterning of the archaeological remains and the morphology of the shelter over time shows
761 that the main activities areas that have been so far identified are located in the centre of the excavated
762 area, about 4 meters from the shelter wall. The shelter substratum, with its successive steps, appears to
763 have strongly influenced the location of occupation. The occupations are located in the best area under the
764 roof edge to develop the subsistence activities with more light. This pattern is similar to other Middle
765 Paleolithic sites and ethnographical data. The direct contact of the latest occupation with the limestone
766 substratum at the foot of the shelter wall suggests that part of the sequence has been possibly eroded. The
767 occurrence of the large limestone slabs that collapsed at some distance from the shelter wall appears to
768 have greatly aided in the preservation of the level 4.1 occupational sequence.
769

770

771 The good preservation of anthropogenic microfacies and scattered charcoal fragments throughout level
772 4.1, together with the lack of evidence for *in situ* bioturbation, show that the lack of hearth structures or of
773 distinctive firing deposits is not a bias caused by post-depositional alteration. The profusion of scattered
774 highly fired bones and vitreous chars in an unburnt host matrix suggests that the fire-related residues have
775 been displaced from the original activity area where they were produced and intentionally used for such as
776 other activities, as suggested by the scattered charcoals. The excavated area may not correspond to the
777 particular location devoted to fire maintenance and more likely represents an open well-lit area devoted to
778 subsistence strategies since it is far from the shelter wall (Kedar et al. 2020). Wood was probably the main
779 fuel used (see charcoals of trees collected on the site) and was probably collected from the nearby areas
780 based on the gathering of fallen branches. The selection was based on branch sizes and condition of the
781 wood as has been suggested for other sites (Uzquiano et al. 2012; Solé et al. 2013; Vidal-Matutano et al.
782 2015; Allué et al. 2017; Vidal-Matutano 2017).
783

784 The spatial distribution of the lithic refits reveals two patterns. First, the refits are clearly located to the
785 main accumulation areas identified by the Kernel analysis (**Fig. 8**). Second, there are a significant number of
786 refits connecting some of the identified clusters (e.g. A with D, E with D, B with G and one between E, F and
787 B). All the faunal refits are well clustered on the accumulation area and only one connect two of them. This
788 faunal refit with a long connection line could suggest possible food sharing between the group's members
789 or different areas of treatment of the carcasses (Gabucio et al. 2018; Marín et al. 2019) (butchery
790 specialized areas). By contrast, the lithic refits show a higher degree of articulation and intra-site mobility in
791 the occupied area (n=7 refits) compared to the faunal refits (n=1 refit) (**Fig. 9 and 11**), even if the
792 synchronicity cannot be demonstrated. This feature is also observed at other Palaeolithic sites (*i.e.* Vaquero
793 et al. 2012a; Bargalló et al. 2016; 2020). Carcasses were exploited in the same areas and only humans and
794 artefacts moved between the different subsistence activities areas.
795

796 To sum up, level 4.1 of the Abri du Maras represents recurrent short-term occupations, and not a mass
797 hunting of reindeers during one main occupation, marked by faint living floors and distinctive
798 anthropogenic facies. Neanderthals brought materials from a collecting perimeter of 30 km and introduced
799 some already knapped artefacts, especially during autumn for hunting reindeers and/or processing
800 carcasses of reindeers. Some types of flint show that Neanderthal had to cross the Ardèche River to reach
801 the shelter. Carcass processing indicates a low level of fracturing implying the recovery of meat and
802 consumption on the spot of some marrow. Micro-wear traces and residues support that both butchering
803 and domestic activities were performed at the site. Four pieces that refit show traces related with cutting
804 plants and whittling wood (Hardy et al. 2013). There are not clear areas of specialized activities but there is
805 intentional movement of some artefacts. At this juncture, it is not possible to estimate the size of the group
806 in question but it may have been a few individuals or a family group. The absence of carnivore tooth marks
807 on the bones abandoned on the site could be possibly explained by rapid burial, or repeated human visits to
808 the site despite the indications of primarily autumnal occupations.
809

810 The results of the level 4.1 of the Abri du Maras raise questions about the definition of mobile groups and

811 ephemeral occupations for Neanderthals. (Gabucio et al. 2018; Marín et al. 2019). At the nearby site of
812 Payre, during MIS 7, we observed short-term occupations with a diversity of flint, mainly local (less than 30
813 km), the differences being in the ratio of alluvial or colluvial flint collected (Moncel et al. 2018). Most of the
814 types of flints Payre were similar to those of the Abri du Maras indicating that, over time, Neanderthals in
815 the region used a similar land use strategy for flint collecting (for instance, for good quality Barremian
816 Bedoulian flint - F14 and F34). For MIS 4-3, the Abri des Pêcheurs, located to the west in the Chassezac
817 Valley, and the Barasses II cave (Balazuc) along the Ardèche River, the records suggest brief and recurrent
818 occupations (bivouacs) (Moncel et al. 2008, 2018, 2019; Daujeard et al., 2019b). These sites only contain
819 flakes and a few cores in flint with limited use of local quartz or basalt pebbles. They also have a wider
820 diversity of flint types from alluvial deposits and colluviums from a large perimeter extending in several
821 directions, from 10 to 30 km. This suggests that toolkits may have circulated from site to site.

822
823 Our combined results provide new data on late Neanderthal short-term occupations regarding level 4.1 of
824 the Abri du Maras. The use of toolkits (points, flakes, blades and some cores brought to the site), specialized
825 seasonal hunting of reindeer, carcass processing *in situ* and repeat disposal of lightning-related materials at
826 the same place indicate specialized, short occupations of the site by Neanderthal groups. The lack of a
827 formal hearth may not mean lack of fire in general (contra Dibble et al. 2018) but in contrast a long-
828 maintained knowledge of suitable fuel issued from lightning-struck wood. Indeed, distinct activity areas
829 which vary in time and space suggest repeated occupation by one or several groups, probably centered on
830 this specialized and seasonal reindeer hunt. Moreover, this evidence could represent shared common
831 traditions regarding lithic, hunting and butchery practices. Comparison with Neanderthal MIS 3 sites in
832 Europe, such as Abric Romani, Teixoneres, El Salt, El Pastor or La Quebrada, indicates a high diversity of
833 modes of occupation and solutions for exploitation of the local territory. The picture that emerges from Abri
834 du Maras and other MIS 3 sites in Europe shows that late Neanderthals found multiple solutions to the
835 exploitation of local conditions across southern Europe (Delagnes and Rendu 2011; Di Modica 2011;
836 Sánchez-Hernández et al. 2014; Vidal-Matutano et al. 2015; Villaverde et al. 2017; Mallol et al. 2019; Mayor
837 et al. 2020; Picin et al. 2020), emphasizing their behavioural adaptability.

838 839 **Acknowledgements**

840 We thank the editors of this special issue of the Topical Collection on Settlement Patterns Dynamics of the
841 Middle Paleolithic and Middle Stone Age to include this manuscript.

842 We would like to thank Dr Vincenzo Spagnolo and the two anonymous reviewers for their useful comments
843 that improved the papers.

844 845 **Funding information**

846 The excavations at the Abri du Maras are supported by the French Ministry of Culture, Service Régional de
847 l'Archéologie Rhône-Alpes-Auvergne under the supervision of M-H. Moncel. MG Chacón, F. Rivals and E.
848 Allué research is funded by the CERCA Programme/Generalitat de Catalunya, the Spanish Ministry of
849 Science, Innovation and Universities (MICINN) projects HAR2016-76760-C3-1-P and PID2019-103987GB-C31
850 and the Catalan Government Research Group 2017 SGR 836.

851 852 **Compliance with ethical standards**

853 **Conflict of interests:** The authors declare that they have no conflict of interests.

854 855 **Authors Contributions**

856 M.-H.M., M.G.CH., D.V. and M-A.C. designed and conceptualized the paper, wrote the initial draft and
857 prepared the figures, tables and supplementary information. Spatial analyses were made by M.G.CH. and
858
859
860
861
862
863

864 D.V., for the lithic and faunal remains respectively. D.V. performed the faunal refits and M.G.CH. and A.E. the
865 lithic ones. Zooarchaeology and taphonomy analysis of large mammals was carried out by C.D and D.V.
866 Ungulate dental micro-wear was done by F.R. Fish remains were studied by P.B. Microstratigraphic analysis
867 was made by M.-A.C. Geology was done by S.P. Lithics were studied by M.-H.M., M.G.CH. and A.E. B.H. did
868 the micro-wear and residues analysis on lithics. Petrographic analysis of flint was done by P.F. E.A. made the
869 charcoal analysis. R.G. performed the databases. Excavations were conducted by M.-H.M. All the authors
870 have interpreted and discussed the data obtained and commented on the manuscript.

871

872

873 **References**

874

875 Adler DS, Conard NJ (2005) Tracking Hominins during the Last Interglacial complex in the Rhineland. In:
876 Gamble C, Porr M (eds) *The Hominid Individual in context. Archaeological Investigations of Lower and Middle*
877 *Palaeolithic Landscapes, Locales and Artefacts*. Routledge. London and New York, pp 133-153

878

879 Allué E, Solé A, Burguet-Coca A (2017). Fuel exploitation among Neanderthals based on the anthracological
880 record from Abric Romaní (Capellades, NE Spain). *Quat. Int.*, 431:6-15

881

882 Bargalló A, Gabucio MJ, Rivals F (2016). Puzzling out a palimpsest: testing an interdisciplinary study in level
883 O of Abric Romaní. *Quat Inter* 417: 51-65.

884

885 Bargalló A, Gabucio MJ, Gómez de Soler B, et al (2020) A Snapshot of a Short Occupation in the Abric
886 Romaní Rock Shelter: Archaeo-Level Oa. In: Cascalheira J, Picin A (eds) *In Short-Term Occupations in*
887 *Paleolithic Archaeology*. Springer International Publishing, Cham, pp 217–235

888

889 Bailey G (2007) Time perspectives, palimpsests and the archaeology of time. *J Anthropol Archaeol* 26:198–
890 223

891

892 Binford LR (1978) Dimensional analysis of behavior and site structure: learning from an Eskimo hunting
893 stand. *Am Antiq* 43(3):330–361

894

895 Binford LR (1996) Hearth and home: the spatial analysis of ethnographically documented rock shelter
896 occupations as a template for distinguishing between human and hominid use of sheltered space. In
897 *Proceedings of the XIII UISPP Congress, ABACO, Forli* pp 229-239

898

899 Böeda E, Pelegrin J (1985) Approche expérimentale des amas de Marsagny. *Archéologie Expérimentale*
900 1:19–36

901

902 Britton K, V Grimes, L Niven, T Steele, S McPherron, M Soressi, TE Kelly, J Jaubert, JJ Hublin, MP Richards
903 (2011) Strontium isotope evidence for migration in late Pleistocene Rangifer: Implications for Neanderthal
904 hunting strategies at the Middle Palaeolithic site of Jonzac, France. *J Hum Evol* 61:176-185

905

906 Britton K, Gaudzinski-Windheuser S, Roebroeks W, Kindler L, Richards MP (2012) Stable isotope analysis of
907 well-preserved 120,000-year-old herbivore bone collagen from the Middle Palaeolithic site of Neumark-
908 Nord 2, Germany reveals niche separation between bovids and equids. *Palaeogeogr Palaeoclimatol*
909 *Palaeoecol* 333-334:168-177

910

911 Cascalheira J, Picin A (2020) *Short-Term Occupations in Paleolithic Archaeology: Definition and*
912 *Interpretation. Interdisciplinary Contributions to Archaeology*. Springer edition. USA

913

914 Castel JC, Discamps E, Soulier MC, et al (2017) Neanderthal subsistence strategies during the Quina
915 Mousterian at Roc de Marsal (France). *Quat Int* 433:140–156

916

917
918 Chacón MG, Bargalló A, Gabucio MJ, Rivals F, Vaquero M (2015) Neanderthal Behaviors from a Spatio-
919 Temporal Perspective: An Interdisciplinary Approach to Interpret Archaeological Assemblages. In: Conard
920 NJ, Delagnes A (eds) Settlement Dynamics of the Middle Paleolithic and Middle Stone Age, Vol. IV Kerns
921 Verlag, Tübingen, pp 253-294
922
923 Combiér J (1967) Le Paléolithique de l'Ardèche dans son cadre bioclimatique. Delmas, Bordeaux
924
925 Costamagno S, Meignen L, Beauval C, Vandermeersch B, Maureille B (2006) Les Pradelles (Marillac-le-Franc,
926 France): A mousterian reindeer hunting camp? *J Anthropol Archaeol* 25:466-484
927
928 Courty MA (2001) Microfacies analysis assisting archaeological stratigraphy. In: Goldberg P, Holliday VT,
929 Ferring CR (eds) *Earth Sciences and Archaeology*. Plenum-Kluwer, New York, pp 205–239
930
931 Courty MA (2012) Transformation pédologique des produits de combustion : approche expérimentale.
932 *Étude et Gestion des Sols*, 19(1) : 41-45.
933
934 Courty MA (2017) Fuel origin and firing product preservation in archaeological occupation contexts. *Quat*
935 *Int* 431:116-130
936
937 Courty MA, Goldberg P, Macphail R. (1989). *Soils and micromorphology in archaeology*. Cambridge:
938 Cambridge.
939
940 Courty MA, Carbonell E, Poch JV, Banerjee R (2012). Microstratigraphic and multi-analytical evidence for
941 advanced Neanderthal pyrotechnology at Abri Romani (Capellades, Spain). *Quaternary International*, 247:
942 294-312.
943
944 Courty MA, Allué E, Henry A (2020). Forming mechanisms of vitrified charcoals in archaeological firing-
945 assemblages. *Journal of Archaeological Science: Reports*, 30, 102215
946
947 Czesla E (1990) On refitting of stone artifacts. In: Czesla E, Eickhoff S, Arts N, Winter D (eds) *The big puzzle*.
948 *International symposium on refitting stone artefacts. Monrepos 1987, Studies in modern archaeology*, vol
949 1. Holos, Bonn, pp 9–44
950
951 Daujeard C, Moncel MH (2010) On Neanderthal subsistence strategies and land use: a regional focus on the
952 Rhône Valley area in southeastern France. *J Anthropol Archaeol* 29(3):368-391
953
954 Daujeard C, Fernandes P, Guadelli JL, Moncel MH, Santagata C, Raynal JP (2012) Neanderthal subsistence
955 strategies in Southeastern France between the plains of the Rhône Valley and the mid-mountains of the
956 Massif Central (MIS 7 to MIS 3). *Quat Int* 252:32-47
957
958 Daujeard C, Abrams G, Germonpré M, Le Pape JM, Wampach A, Di Modica K, Moncel MH (2016)
959 Neanderthal and animal karstic occupations from southern Belgium and south-eastern France: Regional or
960 common features? *Quat Int* 411:179-197
961
962 Daujeard C, Vettese D, Britton K, Béarez P, Boulbes N, Crégut-Bonnoure E, Desclaux E, Lateur N, Pike-Tay A,
963 Rivals F, Allue E, Chacón MG, Courty MA, Gallotti R, Hardy B, Puaud S, Moncel MH (2019a) Neanderthal
964 selective hunting of reindeer in the Upper Pleistocene of Western Europe: the example of Abri du Maras
965 (Southeastern France, Level 4.1, MIS 3). *Archaeol Anthropol Sci* 11:985–1011
966
967 Daujeard C., Brugal J.-P., Moncel M.-H., Fernandes P., Delvigne V., Lafarge A., Le Pape J.-M., Raynal J.-P.
968 (2019b) - Neanderthals and Caprines. Two Upper Pleistocene Caves of Southeastern France. In: Gourichon L,
969 Daujeard C, Brugal JP (eds) *Rencontres internationales d'Antibes 2018. Hommes et Caprinés. De la*
970 *montagne à la steppe, de la chasse à l'élevage*. COLANTIB XXXIX, pp 77-99

- 971
972 Debard E (1988) Le Quaternaire du Bas-Vivarais d'après l'étude des remplissages d'avens, de grottes et
973 d'abris sous roche. Dynamique sédimentaire, paléoclimatologie et chronologie. Doc Lab Géologie Lyon 103
974
975 Defleur A, Bez JF, Crégut-Bonnoure E, Desclaux E, Fontugne M, Jeannet M, Magnon F, Talon B, Thinon M,
976 Comnier J (1994) Industries, biostratigraphie, restes humains et datation du gisement moustérien de la
977 Baume Néron (Soyons, Ardèche). CR Acad Sci Paris 318(II):1409-1414
978
979 Delagnes A, Rendu W (2011) Shifts in Neanderthal mobility, technology and subsistence strategies in
980 western France. J Archaeol Sci 38:1771-1783
981
982 De la Torre I, Martínez-Moreno J, Mora R (2012) When Bones are Not Enough: Lithic Refits and Occupation
983 Dynamics in the Middle Palaeolithic Level 10 of Roca dels Bous (Catalonia, Spain). Bones tools - tools bones
984 interplay between objects Object 1349:13–23
985
986 Dibble, Harold L., Sandgathe, Dennis, Goldberg, Paul, et al (2018) Were Western European Neanderthals
987 able to make fire? J Paleo Archaeol 1(1):54-79
988
989 Di Modica K (2011) Les productions lithiques du Paléolithique moyen de Belgique : variabilité des systèmes
990 d'acquisition et des technologies en réponse à une mosaïque d'environnements contrastés. Dissertation,
991 Muséum National d'histoire Naturelle, Paris
992
993 Discamps E, Royer A (2017) Reconstructing palaeoenvironmental conditions faced by Mousterian hunters
994 during MIS 5 to 3 in southwestern France: A multi-scale approach using data from large and small mammal
995 communities. Quatern Int 433:64-87
996
997 Discamps E, Faivre JP (2017) Substantial biases affecting Combe-Grenal faunal record cast doubts on
998 previous models of Neanderthal subsistence and environmental context. J Archaeol Sci 36:128-132
999
1000 Ekshtain R, Malinsky-Buller A, Greenbaum N, Mitki N, Stahlschmidt MC, Shahack-Gross R, et al. (2019)
1001 Persistent Neanderthal occupation of the open-air site of 'Ein Qashish, Israel. PLoS ONE 14(6):e0215668
1002
1003 Enloe JG (2010) Refitting bones: negative evidence, site structure and social organisation. Lithic Technol
1004 35(1):63-71
1005
1006 Enloe J, David F (1989) Le remontage des os par individus : le partage du renne chez les Magdaléniens de
1007 Pincevent (La Grande Paroisse, Seine-et-Marne). Bull Soc Préhist Fr 86(9) :275-281
1008
1009 ESRI (2014) ArcGIS for Desktop. Environmental Systems Research Institute, Redlands, 10.2.2.
1010 <http://www.esri.com>
1011
1012 Fernández-Laso MC (2010) *Remontajes de restos faunísticos y relaciones entre áreas domésticas en los*
1013 *niveles K, L y M del Abric Romaní (Capellades, Barcelona, España)*. Dissertation, Univeritat Rovira i Virgili,
1014 Tarragona
1015
1016 Fernández-Laso MC, Rosell J, Blasco R, Vaquero M (2020) Refitting bones: Spatial relationships between
1017 activity areas at the Abric Romaní Level M (Barcelona, Spain). J Archaeol Sci Reports 29:102188
1018
1019 Gabucio MJ, Cáceres I, Rivals F, Bargalló A, Rosell J, Saladié P, Vallverdú J, Vaquero M, Carbonell E. 2018.
1020 Unraveling a Neanderthal palimpsest from a zooarcheological and taphonomic perspective. Archaeo
1021 Anthropol Sci. 10:197–222
1022
1023 Gaudzinski-Windheuser S (2006) Monospecific or species-dominated faunal assemblages during the Middle
1024 Paleolithic in Europe. In: Hovers E, Kuhn SL (eds) Transitions before the Transition. Springer, New York, pp

1025 137-147
1026
1027 Gaudzinski S, Roebroeks W (2000) Adults only. Reindeer hunting at the Middle Palaeolithic site Salzgitter
1028 Lebenstedt, Northern Germany. *J Hum Evol* 38: 497-521.
1029
1030 Gaudzinski-Windheuser S, Niven L (2009) Hominin subsistence patterns during the Middle and Late
1031 Paleolithic in Northwestern Europe. In: Hublin JJ, Richards MP (eds) *The Evolution of Hominid Diets: Integrating Approaches to the Study of Palaeolithic Subsistence*. Springer, Dordrecht, pp 97-109.
1032
1033
1034 Ge, T., Courty, M.A., Watzel, J. and Matthews, W. (1993). Sedimentary formation processes of occupation
1035 surfaces. In « Formation processes in archaeological context », P. Goldberg, M. Petraglia and D.T. Nash
1036 (eds.). *Monographs in World Archaeology* 17: 149-163. Madison, Wisconsin: Prehistory Press
1037
1038 Genty D, Combourieu-Nebout N, Peyron O, Blamart D, Wainer K, Mansuri F, Ghaleb B, Isabello L, Dormoy I,
1039 von Grafenstein U, Bonelli S, Landais A, Brauer A (2010) Isotopic characterization of rapid climatic events
1040 during OIS3 and OIS4 in Villars Cave stalagmites (SW-France) and correlation with Atlantic and
1041 Mediterranean pollen records. *Quat Sci Rev* 29:2799-2820
1042
1043 Gilles R (1950) Présentation des silex taillés d'un abri-sous-roche du département de l'Ardèche. *Bull Soc*
1044 *Préhist Fr* 47:202
1045
1046 Goldberg, P., and Berna, F. (2010). Micromorphology and context. *Quaternary International* 214.1-2 : 56-62.
1047
1048 Hardy BL, Moncel MH, Daujeard C, Fernandes P, Béarez P, Desclaux E, Chacon Navarro MG, Puaud S, Gallotti
1049 R (2013) Impossible Neanderthals? Making string, throwing projectiles and catching small game during
1050 Marine Isotope Stage 4 (Abri du Maras, France). *Quat Sci Rev* 82:23-40.
1051
1052 Hardy B. L., Moncel M-H., Kerfant C., Lebon M., Bellot-Gurlet L., Mélard N. (2020) Direct evidence of
1053 Neanderthal fibre technology and its cognitive and behavioral implications. *Nature. Scientific Report* 10,
1054 4889. <https://doi.org/10.1038/s41598-020-61839-w>
1055
1056 Henry DO (2012) the Palimpsest Problem, Hearth Pattern Analysis, and Middle Paleolithic Site Structure.
1057 *Quatern Intern* 247:246-266
1058
1059 Henry DO, Hall SA, Hietala HJ, Demidenko YE, Usik VI, Rosen AM, Thomas PA (1996) Middle Paleolithic
1060 behavioral organization: 1993 excavation of Tor Faraj, southern Jordan. *J Field Archaeol* 23:31-53
1061
1062 Henry DO, Hietala HJ, Rosen AM, Demidenko YE, Usik VI, Armagan TL (2004) Human behavioural
1063 organization in the Middle Paleolithic: were Neanderthals different? *Amer Anthropol* 106:17-31
1064
1065 Higham T, Douka K, Wood R, Bronk Ramsey C, Brock F, Basell L, Camps M, Arrizabalaga A, Baena J, Barroso-
1066 Ruíz C, Bergman C, Boitard C, Boscato P, Caparrós M, Conard NJ, Draily C, Froment A, Galván B, Gambassini
1067 P, Garcia-Moreno A, Grimaldi S, Haesaerts P, Holt B, Iriarte-Chiapusso MJ, Jelinek A, Jordá Pardo JF, Maíllo-
1068 Fernández JM, Marom A, Maroto J, Menéndez M, Metz L, Morin E, Moroni A, Negrino F, Panagopoulou E,
1069 Peresani M, Pirson S, de la Rasilla M, Riel-Salvatore J, Ronchitelli A, Santamaria D, Semal P, Slimak L, Soler J,
1070 Soler N, Villaluenga A, Pinhasi R, Jacobi R (2014) The timing and spatiotemporal patterning of Neanderthal
1071 disappearance. *Nature* 512:306-309.
1072
1073 Hovers E, Malinsky-Buller a, Goder-Goldberger M, Ekshtain R (2011) Capturing a moment: Identifying short-
1074 lived activity locations in Amud Cave, Israel. In: Le Tensorer JM, Jagher R, Otte M (eds) *The Lower and Middle Paleolithic in the Middle East and Neighbouring Regions*. Basel Symposium (May 8-10 2008) ERAUL
1075 126, pp 101-114
1076
1077
1078 Julien M, Karlin C, Valentin B (1992) Déchets de silex, déchets de pierres chauffés. De l'intérêt des

- 1079 remontages à Pincevent (France). In: Hofman JL, Enloe JG (eds.) *Piecing Together the Past: Applications of*
1080 *Refitting Studies in Archaeology*. BAR International Series, vol. 578, Oxford, pp 287–295
- 1081
- 1082 Kedar Y, Kedar G, Barkai R (2020) Setting fire in a Paleolithic Cave: The influence of cave dimensions on
1083 smoke dispersal. *J Archaeol Sci Rep* 29:102112
- 1084
- 1085 Kuhn SL (1992) On planning and curated technologies in the Middle Palaeolithic. *J Anthropol Res* 48:185214
- 1086
- 1087 Kuhn SL (1995) *Mousterian Lithic Technology. An Ecological Perspective*. Princeton. University Press, New
1088 Jersey
- 1089
- 1090 Lucas G (2005) *The Archaeology of time*. Routledge, London
- 1091
- 1092 Lucas G (2012) *Understanding the Archaeological record*. Cambridge university Press
- 1093
- 1094 Lyman RL (2008) (Zoo) Archaeological Refitting: A Consideration of Methods and Analytical. *J Anthropol Res*
1095 64(2):229-248
- 1096
- 1097 Macphail RI, Goldberg P. (2018) *Applied soils and micromorphology in archaeology*. Cambridge University
1098 Press.
- 1099
- 1100 Machado J, Hernández CM, Galván B (2011) Contribución teórico-metodológica al análisis histórico de
1101 palimpsestos arqueológicos a partir de la producción lítica. Un ejemplo de aplicación para el Paleolítico
1102 medio en el yacimiento de El Salt (Alcoy, Alicante). *Recerques Museo d'Alcoi* 20:33-46
- 1103
- 1104 Machado J, Hernández CM, Mallol C, Galván B (2013) Lithic production, site formation and Middle
1105 Palaeolithic palimpsest analysis: in search of human occupation episodes at Abric del Pastor Stratigraphic
1106 Unit IV (Alicante, Spain). *J Archaeol Sci* 40(5):2254-2273
- 1107
- 1108 Machado J, Mayor A, Hernández CM, Galván B (2019) Lithic refitting and the analysis of Middle Palaeolithic
1109 settlement dynamics: a hightemporal resolution example from El Pastor rock shelter (Eastern Iberia).
1110 *Archaeol Anthropol Sci* 11:4539-4554
- 1111
- 1112 Malinsky-Buller A, Hovers E, Marder O (2011) Making time: 'living floors', 'palimpsests' and site formation
1113 processes—a perspective from the open-air Lower Paleolithic site of Revadim Quarry, Israel. *J Anthropol*
1114 *Archaeol* 30(2):89–101
- 1115
- 1116 Mallol C, Hernández C, Mercier N, Falguères C, Carrancho Á, Cabanes D. et al (2019) Fire and brief human
1117 occupations in Iberia during MIS 4: Evidence from Abric del Pastor (Alcoy, Spain). *Sci Rep* 9(1):1-11
- 1118
- 1119 Marín J, Rodríguez-Hidalgo A, Vallverdú J, Gómez de Soler B, Rivals F, Rabuñal JR, Pineda A, Chacón MG,
1120 Carbonell E, Saladié P (2019) Neanderthal logistic mobility from a taphonomic perspective of Abric Romaní
1121 (Capellades, Spain). *Quat Sci Rev* 225:106033
- 1122
- 1123 Mayor A, Hernández, CM, Machado J, Mallol C, Galván B (2020) On identifying Palaeolithic single occupation
1124 episodes: archaeostratigraphic and technological approaches to the Neanderthal lithic record of
1125 stratigraphic unit Xa of El Salt (Alcoy, eastern Iberia) *Archaeol Anthropol Sci* 12:84.
- 1126
- 1127 Modolo M, Rosell Ardèvol J, Thun Hohenstein U (2015) L'impiego dei rimontaggi ossei in due realtà del
1128 Paleolitico Medio: Abric Romani e Riparo Tagliente. In: Thun Hohenstein U, Cangemi M, Fiore I, De Grossi
1129 Mazzorin J (eds) *Atti del 7° Convegno Nazionale di Archeozoologia*. Ferrara, pp 13–20
- 1130
- 1131 Moncel MH, Gaillard C, Patou-Mathis M (1994) L'Abri du Maras (Ardèche) : une nouvelle campagne de
1132 fouilles dans un site paléolithique moyen. *Bull Soc Préhist Fr* 91(6):363-368

- 1133
1134 Moncel MH, Michel V (2000) Première tentative de datation par U-Th du site paléolithique moyen de l'abri
1135 du Maras. *Bull Soc Préhist Fr* 93(3):371-375
1136
1137 Moncel MH, Brugal JP, Prucca A, Lhomme G (2008) Mixed occupation during the Middle Palaeolithic: case
1138 study of a small pit-cave-site of Les Pêcheurs (Ardèche, south-eastern France). *J Anthropol Archaeol* 27(3):
1139 382-398
1140
1141 Moncel MH, Daujeard C, Cregut-Bonnoure E, Boulbes N, Puaud S, Debard E, Bailon S, Desclaux E, Escude E,
1142 Roger T, Dubar M (2010) Nouvelles données sur les occupations humaines du début du Pléistocène
1143 supérieur de la moyenne vallée du Rhône (France). Les sites de l'Abri des Pêcheurs, de la Baume Flandin, de
1144 l'Abri du Maras et de la Grotte du Figuier (Ardèche). *Quatern* 21(4):389-415
1145
1146 Moncel MH, Daujeard C (2012) The variability of the Middle Palaeolithic on the right bank of the Middle
1147 Rhône Valley (southeast France): Technical traditions or functional choices? *Quatern Int* 247:103-124
1148
1149 Moncel MH, Chacón MG, La Porta A, Fernandes P, Hardy B, Gallotti R (2014) Fragmented reduction
1150 processes: Middle Palaeolithic technical behaviour in the Abri du Maras shelter, southeastern France.
1151 *Quatern Int* 350:180-204
1152
1153 Moncel MH, Allué E, Bailon S, Barshay-Szmidt C, Béarez P, Crégut-Bonnoure E, Daujeard C, Desclaux E,
1154 Debard E, Lartigot-Campin AS, Puaud S, Roger T (2015) Evaluating the integrity of palaeoenvironmental and
1155 archaeological records in MIS 5 to 3 karst sequences from southeastern France. *Quatern Int* 378:22-39
1156
1157 Moncel MH, Raynal JP, Fernandes P, Delvigne V, Hardy B (2018) Les différentes phases d'occupation de la
1158 cavité : fonction(s) du site. Le matériel lithique. In : Daujeard C (dir.) *La Grotte des Barasses II (Balazuc) :*
1159 *entre Néandertaliens, bouquetins et carnivores. Des occupations du Pleistocène supérieur en moyenne*
1160 *vallée de l'Ardèche DARA 46*, pp 109-128
1161
1162 Moncel MH, Fernandes P, Willmes M, James H, Grün R (2019) Rocks, Teeth, and Tools: New insights into
1163 early Neanderthal subsistence strategies in South-Eastern France from lithic reconstructions and strontium
1164 isotope analysis, *PLoS ONE* 14(4): e0214925
1165
1166 Mora R, Roy M, Martínez-Moreno J, Benito-Calvo A, Samper S (2020) Inside the Palimpsest: Identifying
1167 Short Occupations in the 497D Level of Cova Gran (Iberia). In: Cascalheira J, Picin A (eds) *Short-Term*
1168 *Occupations in Paleolithic Archaeology, Interdisciplinary Contributions to Archaeology*. Springer
1169 International Publishing, Cham, pp 39-69
1170
1171 Morin E, Tsanova T, Sirakov N, Rendu W, Mallye JB, Lévêque F (2005) Bone refits in stratified deposits:
1172 Testing the chronological grain at Saint-Césaire. *J Archaeol Sci* 32(7):1083-1098
1173
1174 Newcomer MH, Sieveking G d G (1980) Experimental flake scatter-patterns: a new interpretative technique.
1175 *J Field Archaeol* 7:345-352
1176
1177 Niven L, Steele TE, Rendu W, Mallye JB, McPherron SP, Soressi M, Jaubert J, Hublin JJ (2012) Neanderthal
1178 mobility and large-game hunting: The exploitation of reindeer during the Quina Mousterian at Chez-Pinaud
1179 Jonzac (Charente-Maritime, France). *J Hum Evol* 63:624-635
1180
1181 Niven L (2013) A diachronic evaluation of Neanderthal cervid exploitation and site use at Pech de l'Azé IV,
1182 France. In: Clark JL, Speth JD (eds.) *Zooarchaeology and Modern Human Origins: Human Hunting Behavior*
1183 *during the Later Pleistocene, Vertebrate Paleobiology and Paleoanthropology*. Springer, Dordrecht, pp 151-
1184 161.
1185
1186 Nomade, S., Genty, D., Sasco, R., Scao, V., Féruglio, V., Baffier, D., Guillou H, Bourdier C, Valladas H, Reigner

1187 E, Debard E, Pastre J-F, Geneste J-M (2016) A 36,000-year-old volcanic eruption depicted in the Chauvet-
1188 Pont d'Arc Cave (Ardèche, France)?. *PloS one* 11(1).
1189
1190 O'Connell JF (1987) Alyawara site structure and its archaeological implications. *Am Antiq* 52(1):74–108
1191
1192 Picin A, Chacón MG, Gómez de Soler B, Blasco R, Rivals F, Rosell J (2020) Neanderthals mobile toolkit in
1193 short-term occupations at Teixoneres Cave (Moia, Spain). *J Archaeol Sci Rep* 29: 102165
1194
1195 Poplin F (1976) Remarques théoriques et pratiques sur les unités utilisées dans les études d'ostéologie
1196 quantitative, particulièrement en archéologie préhistorique. *Thèmes spécialisés, IXe Congrès UISPP, Nice,*
1197 *124-140.*
1198
1199 Puaud S, Nowak M, Pont S, Moncel MH (2015) Minéraux volcaniques et alpins à l'abri du Maras (Ardèche,
1200 France) : témoins de vents catabatiques dans la vallée du Rhône au Pléistocène supérieur. *C R Palevol*
1201 *14:331-341*
1202
1203 Real C, Sanchis A, Eixea A, Villaverde V (2019) Neanderthal subsistence and short-term human occupation
1204 patterns during MIS 5. New data from Abrigo de la Quebrada (Valencia, Spain). *J Archeol Sci Rep* 28:102056
1205
1206 Rendu W, Costamagno S, Meignen L, Soulier MC (2012) Monospecific faunal spectra in Mousterian contexts:
1207 Implications for social behavior. *Quatern Int* 247:50-58
1208
1209 Richard M, Falguères C, Pons-Branchu E, Bahain JJ, Voinchet P, Lebon M, Valladas H, Dolo JM, Puaud S, Rué
1210 M, Daujeard C, Moncel MH, Raynal JP (2015) Contribution of ESR/U-series dating to the chronology of late
1211 Middle Palaeolithic sites in the middle Rhône valley, southeastern France. *Quat Geochronol* 30(B):529-534
1212
1213 Richards MP, Taylor G, Steele T, McPherron SP, Soressi M, Jaubert J, Orschiedt J, Mallye JB, Rendu W, Hublin
1214 JJ (2008) Isotopic dietary analysis of a Neanderthal and associated fauna from the site of Jonzac (Charente-
1215 Maritime), France. *J Hum Evol* 55(1):179-185
1216
1217 Roebroeks W (1988) From find scatters to Early hominid behaviour. A study of Middle Palaeolithic riverside
1218 settlements at Maastricht-Belvédère (The Netherlands). *Analecta Praehistorica Leidensia* 21, Leiden
1219
1220 Romagnoli F, Vaquero M (2016) Quantitative stone tools intra-site point and orientation patterns of a
1221 Middle Palaeolithic living floor: A GIS multi-scalar spatial and temporal approach. *Quartär* 63, 47-60
1222
1223 Romagnoli F, Vaquero M (2019) The challenges of applying refitting analysis in the Palaeolithic archaeology
1224 of the twenty-first century: an actualised overview and future perspectives. *Archaeol Anthropol Sci*
1225 *11:4387–4396*
1226
1227 Rosell J, Blasco R, Fernández-Laso MC, Vaquero M, Carbonell E (2012) Connecting areas: Faunal refits as a
1228 diagnostic element to identify synchronicity in the Abric Romaní archaeological assemblages, *Quatern Int*
1229 *252:56-67*
1230
1231 Rosell J, Modesto-Mata M, Fernández-Laso C, Modolo M, Blasco R (2019) Refitting bones to reconstruct the
1232 diversity in Middle Palaeolithic human occupations: the case of the Abric Romaní site (Capellades,
1233 Barcelona, Spain). *Archaeol Anthropol Sci* 1: 4601-4619.
1234
1235 Ruebens K, Wragg Sykes RM (2016) Spatio-temporal variation in late Middle Palaeolithic Neanderthal
1236 behaviour: British bout coupé handaxes as a case study. *Quatern Int* 411:305-326
1237
1238 Sánchez-Hernández C, Rivals F, Blasco R, Rosell J (2014) Short, but repeated Neanderthal visits to Teixoneres
1239 Cave (MIS 3, Barcelona, Spain): a combined analysis of tooth microwear patterns and seasonality. *J Archaeol*
1240 *Science* 49:317-325

1241
1242 Sisk ML, Shea JJ (2008) Intrasite spatial variation of the Omo Kibish middle stone age assemblages: artifact
1243 refitting and distribution Patterns. *J Hum Evol* 55(3):486-500
1244
1245 Solé A, Allué E, Carbonell E (2013) Hearth-related wood remains from Abric Romaní layer M (Capellades,
1246 Spain). *J. Anthropol. Res* 69:535-559
1247
1248 Szmidski CC, Moncel M-H, Daujeard C (2010) New data on the late Mousterian in Mediterranean France: first
1249 radiocarbon (AMS) dates at Saint-Marcel Cave (Ardèche). *C. R. Palevol* 9(4): 185-199
1250
1251 Soulier MC (2013) Entre alimentaire et technique : l'exploitation animale aux débuts du Paléolithique
1252 supérieur. Stratégies de subsistance et chaînes opératoires de traitement du gibier à Isturitz, La Quina aval,
1253 Roc-de-Combe et Les Abeilles. Dissertation. Université de Toulouse 2 Le Mirail
1254
1255 Spagnolo V, Marciani G, Aureli D, Berna F, Boscato P, Ranaldo F, Ronchitelli A (2016) Between hearths and
1256 volcanic ash: the SU 13 palimpsest of the Oscurusciuto rock shelter (Ginosa - southern Italy): analytical and
1257 interpretative questions. *Quatern Int* 417:105-121.
1258
1259 Spagnolo V, Marciani G, Aureli D, Berna F, Toniello, G, Astudillo F, Boschini F, Boscato P, Ronchitelli A (2019)
1260 Neanderthal activity and resting areas from stratigraphic unit 13 at the Middle Palaeolithic site of
1261 Oscurusciuto (Ginosa - Taranto, Southern Italy). *Quat Sci Rev* 217:169–193.
1262
1263 Spagnolo V, Marciani G, Aureli D, et al (2020a) Climbing the time to see Neanderthal behaviour's continuity
1264 and discontinuity: SU 11 of the Oscurusciuto Rockshelter (Ginosa, Southern Italy). *Archaeol Anthropol Sci*
1265 12:54.
1266
1267 Spagnolo V, Aureli D, Martini I, Ekberg I, Boschini F, Crezzini J, Poggi G, Boscato P, Ronchitelli A (2020b) Short
1268 and close in time: overlapped occupation from the layer 56 of the Molare Rock shelter (Southern Italy).
1269 *Archaeol Anthropol Sci* 12: 1-35.
1270
1271 Stevenson MG (1991) Beyond the formation of hearth-associated artifact assemblages. In: Kroll EM, Price
1272 TD (eds.) *The Interpretation of Archaeological Spatial Patterning*. New York and London, Plenum Press, pp
1273 269-299
1274
1275 Turq A, Roebroeks W, Bourguignon L, Faivre JP (2013) The fragmented character of Middle Palaeolithic
1276 stone tool technology. *J Hum Evol* 65(5):641–655
1277
1278 Uzquiano P, Yravedra J, Zapata BR, García MJG, Sesé C, Baena J (2012) Human behaviour and adaptations to
1279 MIS 3 environmental trends (> 53–30 ka BP) at Esquilleu cave (Cantabria, northern Spain). *Quat. Int.* 252:82-
1280 89
1281
1282 Vallverdú J, Allué E, Bischoff JL, Cáceres I, Carbonell E, Cebrià A, García-Antón D, Huguet R, Ibáñez N,
1283 Martínez K, Pastó I, Rosell J, Saladié P, Vaquero M (2005) Short human occupations in the Middle
1284 Palaeolithic level i of the Abric Romaní rock-shelter (Capellades, Barcelona, Spain). *J Hum Evol* 48(2):157–
1285 174
1286
1287 Vandevelde S, Brochier JÉ, Petit C, Slimak L (2017) Establishment of occupation chronicles in Grotte Mandrin
1288 using sooted concretions: Rethinking the middle to upper paleolithic transition. *J Hum Evol* 112:70-78
1289
1290 Vaquero M (2008) The History Of Stones: Behavioural Inferences And Temporal Resolution Of An
1291 Archaeological Assemblage From The Middle Palaeolithic. *J Archaeol Sci* 35(12):3178-3185
1292
1293 Vaquero M, Pastó I (2001) The definition of spatial units in Middle Palaeolithic sites. The hearth-related
1294 assemblages. *J Archaeol Sci* 28:1209-1220

1295
1296 Vaquero M, Chacón MG, García-Antón MD, Gómez de Soler B, Martínez K, Cuartero F (2012a) Time and
1297 space in the formation of lithic assemblages: the example of Abric Romaní level J *Quat Int* 247:162–181
1298
1299 Vaquero M, Chacón MG, Cuartero F, García-Antón MD, Gómez de Soler B, Martínez K (2012b) The lithic
1300 assemblage of level J. In: Carbonell E (ed) *High Resolution Archaeology and Neanderthal Behavior: Time and*
1301 *Space In Level J of Abric Romaní (Capellades, Spain)*. *Vertebr Paleobiol Paleoanthropol Book Series*. Springer,
1302 Dordrecht, pp 189-311
1303
1304 Vaquero M, Bargalló A, Chacón MG, Romagnoli F, Sañudo, P (2015) Lithic recycling in a Middle Paleolithic
1305 expedient context: Evidence from the Abric Romaní (Capellades, Spain), *Quatern Inter* 361:212-228
1306
1307 Vaquero M, Romagnoli F, Bargalló A, Chacón MG, Gómez de Soler B, Picin A, Carbonell E (2019) Lithic
1308 refitting and intrasite artifact transport: a view from the Middle Paleolithic. *Archaeol Anthropol Sci*
1309 11(9):4491-4513
1310
1311 Vettese D (2014) *Le traitement des carcasses d'ongulés chez les Néanderthaliens de l'Abri du Maras*
1312 *(Ardèche, MIS4) / individualisme ou poids des traditions?*. Master Dissertation. Muséum national d'Histoire
1313 naturelle, Paris
1314
1315 Vettese D, Daujeard C, Blasco R, Borel A, Caceres I, Moncel MH (2017) Neanderthal long bone breakage
1316 process: Standardized or random patterns? The example of Abri du Maras (Southeastern France, MIS 3). *J*
1317 *Archaeol Sci Rep* 13:151-163
1318
1319 Vidal-Matutano P (2017) Firewood and hearths: Middle Palaeolithic woody taxa distribution from El Salt,
1320 stratigraphic unit Xb (Eastern Iberia). *Quatern Intern* 457:74-84
1321
1322 Vidal-Matutano P, Hernández CM, Galván B, Mallol C (2015) Neanderthal firewood management: evidence
1323 from Stratigraphic Unit IV of Abric del Pastor (Eastern Iberia). *Quat. Sci. Rev.* 111:81-93
1324
1325 Villa P, Mahieu E (1991) Breakage patterns of human long bones. *J Hum Evol* 21(1):27-48
1326
1327 Villaverde V, Eixea A, Zilhão J, Sanchis A, Real C, Bergadà M (2017). Diachronic variation in the Middle
1328 Paleolithic settlement of Abrigo de la Quebrada (Chelva, Spain). *Quat Int* 435:164-179.
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344

1345
1346
1347

Table 1 Categories of artefacts and raw materials of the level 4.1

| | Basalt | Limestone | Siliceous limestone | Granit | Sandstone | Quartz | Quartzite | Schist | Flint | Ind | Total |
|------------------------|-----------|-----------|---------------------|-----------|-----------|------------|-----------|----------|-------------|----------|--------------|
| Debris | | 1 | 1 | 1 | 1 | 36 | | | 137 | 3 | 180 |
| Flakes | 6 | 2 | 3 | 3 | 1 | 45 | | 1 | 646 | 1 | 708 |
| Fragments of flake | 1 | | 2 | 2 | | 8 | | | 327 | | 340 |
| Entire-broken pebbles | 5 | 1 | | 8 | 2 | 9 | 2 | 1 | 2 | 2 | 32 |
| Blades-bladelets | 1 | | 2 | | | 2 | 1 | | 202 | | 208 10.2% |
| Micro-debris | | | | | | 3 | | | 164 | | 167 |
| Cores | | | | | | | | | 51 | | 51 2.5% |
| Pebble-tools | | | | | | 1 | | | | | 1 |
| Small flakes (< 15 mm) | | | 1 | | | 4 | | | 268 | | 273 |
| Points | | 1 | 1 | | | | | | 79 | | 81 3.9% |
| Total | 13 | 5 | 10 | 14 | 4 | 108 | 3 | 2 | 1876 | 6 | 2041 |

1348
1349
1350
1351
1352
1353
1354

Table 2 Distribution of refitting groups according to the number of remains and the type of refit: mechanical refit (green bone fracture refit [Gbf Refit] and dry bone fracture refit [Dbf Refit]) and anatomical refit (Anat. Refit) values between brackets are %, NSP (Number of Specimen)

| Nb of remains | Anat. Refits | Gbf Refits | Dbf Refits | Total refits | Total NR |
|---------------|------------------|------------------|------------------|--------------|------------|
| 2 | 11 (12.5) | 35 (39.8) | 18 (20.5) | 64 | 128 |
| 3 | 4 (4.5) | 7 (8.0) | 2 (2.3) | 13 | 39 |
| 4 | 2 (2.3) | 3 (3.4) | 1 (1.1) | 6 | 24 |
| 5 | 1 (1.1) | 1 (1.1) | | 2 | 10 |
| 6 | 1 (1.1) | | | 1 | 6 |
| 7 | 1 (1.1) | | | 1 | 7 |
| 8 | 1 (1.1) | | | 1 | 8 |
| Total | 21 (23.9) | 46 (52.3) | 21 (23.9) | 88 | 222 |

1355
1356
1357
1358
1359

Table 3 Distances of the faunal refits connections lines (cm): average, Max. (Maximum), Min (Minimum), Sd (standard deviation)

| | All refits | Gbf refits | Dbf refits | Anat. refits |
|------------------------|------------|------------|------------|--------------|
| Max | 251.31 | 251.31 | 114.11 | 96.03 |
| Min | 0 | 0 | 0 | 0 |
| Average | 14.07 | 15.40 | 9.07 | 16.19 |
| Sd | 34.18 | 41.25 | 24.95 | 24.37 |
| Nb. Refit=0 | 46 | 28 | 13 | 5 |
| Nb total Refits | 88 | 46 | 21 | 21 |

1360
1361

1362
1363
1364
1365

Table 4 Bone refits according the species. Nb refit (refit Number), NSP, values between brackets are %

| | Nb. refits | NR refits |
|------------------------------|-----------------|------------------|
| <i>Rangifer tarandus</i> | 61 (69.3) | 152 (68.5) |
| <i>Equus ferus</i> | 6 (6.8) | 20 (9) |
| <i>Megaloceros giganteus</i> | 3 (3.4) | 9 (4.1) |
| Und. | 18 (20.5) | 41 (18.5) |
| Total | 88 (100) | 222 (100) |

1366
1367
1368
1369
1370
1371

Table 5 Bone refits according to anatomical part. Nb refits (Refit number), NR (Number of remains), Gbf refit (green bone fracture refit), Dbf refit (dry bone fracture refit), Anat. Refit (Anatomical refit); of total NR and total Nb refit. Values between brackets are %

| | Anat. Refits | Dry b.fract | Green b.fract | Total |
|----------------|------------------|------------------|-------------------|------------------|
| Ant. Autopode | 7 (3.2) | 4 (1.8) | 4 (1.8) | 15 (6.8) |
| Axial | 2 (.9) | | | 2 (.9) |
| Forlimb | 2 (.9) | 5 (2.3) | 43 (19.4) | 50 (22.5) |
| Head | 26 (11.7) | 2 (.9) | | 28 (12.6) |
| Hindlimb | 6 (2.7) | 6 (2.7) | 17 (7.7) | 29 (13.1) |
| Phal. Und. | 6 (2.7) | | | 6 (2.7) |
| Post. Autopode | 19 (8.6) | 13 (5.9) | 28 (12.6) | 60 (27) |
| Und | | 16 (7.2) | 16 (7.2) | 32 (14.4) |
| Total | 68 (30.6) | 46 (20.7) | 108 (48.6) | 222 (100) |

1372
1373
1374
1375

Table 6 Total of pieces by type of stone and general type of RMU

| | Basalt | Quartzite | Quartz | Limestone | Flint | Total |
|--|----------|-----------|-----------|-----------|------------|------------|
| Isolated pieces | - | 3 | 6 | - | 11 | 20 |
| Groups | 2 | - | 7 | 1 | 22 | 32 |
| RMUs Total nb. pieces | 6 | 3 | 95 | 19 | 459 | 593 |

1376
1377
1378
1379
1380
1381

Table 7 Detailed descriptions of each RMU identified in level 4.1 by stone type, total of pieces, technical categories, number of refits and technological consideration (see also Fig. 10 and 11)

| Stone type | RMU Number (n°) | Total n° pieces | Technical categories | N° of refits by RMU | General description |
|------------|-----------------|-----------------|-----------------------------------|---------------------|--|
| Quartzite | Isol-Qtzite 1 | 1 | 1 big fragment of pebble | - | Rectangular quartzite pebble fractured intentionally in the middle. Presents unifacial retouch on distal part. There are percussion and macro-traces of use on the distal edge (unknown material use). |
| | Isol-Qtzite 2 | 1 | 1 broken flake | - | Broken Levallois quartzite flake (proximal zone conserved) retouched in the proximal part with a simple notch. Fine brownish grain good for knapping. Flake-tool introduced into the site. |
| | Isol_Qtzite 3 | 1 | 1 flake | - | Levallois lengthen flake slightly <i>déjàtê</i> knapped on a fine grain quartzite. Use wear macro traces on lateral edge. |
| Basalt | Bas-1 | 5 | 1 core 2 fragments 3 flakes | 1 (n=2) | Medium size pebble introduced in the site where is used to obtain two or three flakes and then transform in a massive tool with a thick edge (<i>robot</i>). |

| | | | | | |
|-----------|-------------------|----|---|------------|---|
| | Bas-2 | 1 | 1 fragment | | Fragment of hammerstone |
| Limestone | Lim-1 | 19 | 13 flakes 6 fragments of flakes | 2 (n=4) | Medium size flakes and fragments of flakes. The presence of cortical products and flakes from reduction sequences (even without cores) shows that the knapping strategies were realized in the site following recurrent centripetal Levallois strategies. This could be corroborated by the 2 identified knapping refits (4 pieces), one of them with the longest connection line (371 cm). |
| Quartz | Isol_Qtz (1 to 6) | 6 | 3 fragments of flakes 2 flakes 1 flake-tool | - | Isolated products of different varieties of quartz that cannot be possible associate to the other RMU. All are products coming from bifacial bipolar strategies (discoïdal type) |
| | Qtz-1 | 30 | 1 core 1 flake-tool 7 flakes 19 fragments of flakes 2 fragmented flakes | 4 (n=9) | The 30 pieces concerning all phases of the reduction sequence (cortical flakes, <i>débordant</i> elements, small fragments, tools, etc.). It seems that there are 2 different pebbles that are introduced to be knapped in the site. The tools of large formats are especially notches and denticulates. The first flakes of cortical removal of the nodule confirm the complete lithic sequences. Flakes are normally dissymmetric and with thick formats (four refits). |
| | Qtz-2 | 4 | 3 flakes 1 fragment of flake | - | 4 isolated cortical pieces (more than 90 % of the surface) and 1 without that are shows macro-traces. Reduction sequence very fragmented. Possible introduction of one big cortical flake. |
| | Qtz-3 | 2 | 2 fragments of flakes | - | Both are completely cortical. Flakes of hammerstone produced by the impact? |
| | Qtz-4 | 3 | 3 fragments of flakes | - | There are no more pieces of this RMU, only fragments of flakes and some completely cortical. |
| | Qtz-5 | 40 | 2 flake-tools 10 flakes 17 fragments of flakes 11 fragmented flakes | 3 (n=7) | Pebble introduced and knapped in the site, all the stages of <i>chaîne opératoire</i> were identified. The knapping products show that some left, perhaps a part of them should be transported after leaving the site as tool kit. realized. Some big flakes were transformed by retouch in sidescraper and denticulate. Presence of Siret fractures in various cases (e.g.1 refit). |
| | Qtz-6 | 5 | 4 fragments of flake 1 fragmented flake | - | 2 cortical flakes and some small fragments of flakes. Very fragmented reduction sequence, introduction of 1 big cortical flake. |
| | Qtz-7 | 5 | 1 flake 1 flake-tool 3 fragments of flakes | - | 2 isolated pieces, without cortex, retouched as a notch and denticulate. Some fragments of broken flakes (3). Fragmented reduction sequence and only presence of the final stages. |
| Flint | Isol-Fl (1 to 11) | 11 | 8 flakes 2 fragments of flake 1 flake-tool | - | These isolated pieces belong all of them to bifacial centripetal strategies, mainly Levallois but some of them come from discoïdal ones. We should point out two typical Levallois points and one big Quina scraper. All these pieces are pieces introduced already knapped to the site (tool kit) |
| | Fl-1 | 2 | 2 flakes | - | 2 elongated Levallois flakes introduced into the site (tool kit). |
| | Fl-2 | 12 | 1 core 7 flakes 2 fragmented flakes 1 fragment of flake | 2 (n=4) | Complete reduction sequence: cortical flakes (refit) peeling the nodule and recurrent centripetal Levallois exploitation (refit) obtaining large flakes. The core is abandoned although it is not exhausted. |
| | Fl-3 | 6 | 5 flakes 1 fragmented flake | - | Advanced fragmented reduction sequence with just few flakes and small <i>débitage</i> (no refits). |
| | Fl-4 | 3 | 2 flakes 1 fragment of flake | - | One elongated Levallois flake and two small broken flakes. Fragmented reduction sequence that implies pieces introduced already knapped. |
| | Fl-5a | 17 | 11 flakes 6 fragments of flakes 1 fragmented flake | - | Advanced and fragmented reduction sequence with no cortical elements. The knapping products are varied (configuration, <i>réaffûtage</i> , etc.) and belongs to Levallois strategies. |
| | Fl-5b | 11 | 11 flakes | 1 (n=4) | Reduction sequence of flakes by recurrent centripetal Levallois strategies. |
| | Fl-6 | 4 | 4 flakes | - | Initial phase of the reduction sequence with the presence of 4 cortical flakes. There are no refits but RMU are located in |

| | | | | | |
|--------|-----|---|-------------|--|---|
| | | | | | the same area of the site. |
| FI-7 | 5 | 1 flake 3 fragments of flake 1 fragmented flake | - | | A set of 1 flake and 4 broken flakes (fragmented reduction sequence). |
| FI-8 | 2 | 2 fragmented flakes | - | | 1 isolated and a broken flake (no cortex). Pieces introduced into the site already knapped (tool kit). |
| FI-9 | 5 | 3 flakes 2 fragments of flake | 1 (n=2) | | Advanced reduction sequence. These knapping products belong to centripetal recurrent Levallois strategies. |
| FI-10 | 9 | 3 fragments of flake 3 fragmented flakes 3 flakes (1 point) | - | | Set of 7 Levallois flakes introduced to the site already knapped. All come from Levallois strategies. |
| FI-11 | 6 | 4 flakes 1 fragment of flake 1 fragmented flake | - | | Isolated pieces related with an advanced production phase. All come from Levallois strategies. |
| FI-12 | 34 | 1 flake-tool 14 flakes 7 fragmented flakes 12 fragments of flakes | - | | Complete reduction sequence where all the technological products are represented: cortical flakes, backed elements, advanced flakes and tools, but no cores. Two possibilities (1) the core was transported after the occupation, or (2) some of the cores very patinated belongs to this RMU. |
| FI-13 | 231 | 129 flakes 7 flake-tools 1 core 24 fragmented flakes 70 fragments of flakes | 6 (n=13) | | Complete reductions sequences. It is probably that exist two or three pebbles which are knapped at the site. They were differentiated by the type of cortex and some macroscopic characteristics. They come from the same geological formation (same type of flint). |
| FI-14 | 4 | 2 flakes 1 fragment of flake 1 fragmented flake | - | | 3 semi-cortical flakes (around 25-50 %), two with macro traces of use. Reduction sequence is partial and fragmented, no cores associated. |
| FI-15 | 4 | 3 fragments of flake 1 flake-tool | 1 (n=2) | | Isolated pieces, some cortical. |
| FI-16a | 2 | 1 flake 1 fragmented flake | - | | Isolated cortical pieces. |
| FI-16b | 5 | 1 core 1 flake 1 fragment of flake 1 fragmented flake | - | | Isolated pieces of different phases of the reduction sequence, so very fragmented. They belong to Levallois strategies. |
| FI-16c | 3 | 3 flakes | - | | Isolated cortical pieces coming from the first phase of the reduction sequence. |
| FI-16d | 32 | 2 cores 24 flakes 4 fragments of flake 2 fragmented flakes | 1 (n=2) | | Complete reductions sequences knapped at the site. All of them are related with different modalities of Levallois strategies. |
| FI-16e | 32 | 3 cores 20 flakes 1 fragmented flake 6 fragments of flakes | - | | It is probably that in some cases exist two or three pebbles which were differentiated by the type of cortex and some macroscopic characteristics. They come from the same geological formation (same type of flint). Only one refit had been found, perhaps some pieces left and where transported after leaving the site (fragmented reduction sequence). |
| FI-16f | 30 | 3 cores 17 flakes 3 fragmented flakes 9 fragments of flakes | - | | |

1383
1384
1385

Table 8 Lithic refits by type of stone and number of pieces/refits from Level 4.1

| | Flint | Limestone | Basalt | Quartz | Total |
|------------------------------|-----------|-----------|----------|----------|-----------|
| Refits of 2 pieces | 10 | 2 | 1 | 5 | 18 |
| Refits of 3 pieces | 1 | - | - | 2 | 3 |
| Refits of 4 pieces | 1 | - | - | - | 1 |
| Total Nb of refitting | 12 | 2 | 1 | 7 | 22 |

1386
1387
1388
1389
1390

Table 9 Technological categories by type of stone of the refitted pieces of the Level 4.1

| | | Technological categories | | | | | Total | |
|---------------|-----------|--------------------------|-----------|--------------------|---------------|-------------|----------|-----------|
| | | Cores | Flakes | Fragments of flake | Broken flakes | Flake-tools | | Debris |
| Raw materials | Basalt | | | | | | 2 | 2 |
| | Quartz | 2 | 4 | 5 | 5 | | | 16 |
| | Limestone | | 4 | | | | | 4 |
| | Flint | 1 | 12 | 10 | 3 | 1 | | 27 |
| Total | | 3 | 20 | 15 | 8 | 1 | 2 | 49 |

1391
1392
1393
1394
1395
1396

Table 10 Type of connection lines by type of stone for Level 4.1

| | flint | limestone | basalt | quartz | Total |
|--|-----------|-----------|----------|----------|-----------|
| Knapping connection lines | 8 | 2 | 1 | 4 | 15 |
| Knapping fracture connection lines | 1 | - | - | 5 | 6 |
| Conjoin (post-depositional Fractures) connection lines | 6 | - | - | - | 6 |
| Total | 15 | 2 | 1 | 9 | 27 |


1397
1398
1399
1400
1401
1402

Table 11 Distances of the fauna refits connections lines (cm): average, Max. (Maximum), Min (Minimum), Sd (standard deviation)

| | | All refits |
|------------------------|------------------------------|------------|
| Distance | Max | 370.91 |
| | Min | 0 |
| | Average | 60.41 |
| | Sd | 101.30 |
| | Nb. Refit=0 | 1 |
| | Nb total of connection lines | 27 |
| Nb total refits | | 22 |

1403
1404

1405
1406
1407




1408
1409
1410
1411

Fig. 1 Location of the Abri du Maras. Site situation on the edges of the Rhône valley and detailed view location near the Saint Martin d'Ardèche village




A



B



C



D

1412
 1413 **Fig. 2** A excavated area of level 4.1; B Vertical projection of the archaeological remains by levels (projection
 1414 band 6 green level 4.1, red 4.2 and yellow); C Map of the excavated area for the level 4.1; D Vertical view of
 1415 the site before the beginning of the excavation of 2009. The upper limit of the photo corresponds to the
 1416 limit of the shelter wall (see the limit on map C).
 1417
 1418
 1419



Fig. 3 Example of square J6, level 4.1 and density of bone remains and artefacts

1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436




Fig. 4 Examples of elongated pieces and points brought to the site already worked

1437
1438
1439
1440
1441
1442
1443




Fig. 1/MAC

Fig. 5 **a** Field view of the greyish microaggregate lens in I6 (260-268 cm); **b** Field view of the greyish microaggregated lens in L6 C4.1 L6 (283 cm); **c** Field view of a microaggregate wrapped in a polymer film from a diffuse greyish microaggregate lens in K6 (271 cm).

1444
1445
1446
1447
1448





Fig. 2/MAC


1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462

Fig. 6 **a** Stereomicroscope view of the sand-sized fraction extracted by water-sieving of a greyish microaggregate lens in I5 (258-262 cm): abundant white and coloured polymer films, filaments, aggregates, humified organic fragments, metal-splashed angular grains; **b** Back-scattered (BSE) SEM view of the greyish lens in I6 showing the imbrication of polymerized plant fibres (in black) with the silty clay microaggregates and the native metal inclusions (bright spots); **c** BSE-SEM view of the connected fish-scales and bones lying above the lens viewed in (b), also with abundant native metal inclusions (bright spots); **d to k** typical components of the microaggregate lenses in level 4.1; **d** nanostructured twisted polymer films with metal inclusions; **e** nanostructured composite filament showing lightning-formed quartz with Fe-Cr-Ni metal splash (in bright); **f** flash-pyrolyzed partly vitrified charcoal with metal inclusions (in bright); **g** flash-pyrolyzed bone with metal inclusions (in bright); **h** exogenous angular quartz showing lightning-formed Fe-Cr-Ni metal splash on the angular edges; **i** flash-burnt shell fragment showing in **j** degassed vesicles and metal inclusions; **k** aligned droplets of Fe-Cr-Ni metal splash on a flint-flake.




1463
 1464
 1465
 1466

Fig. 7 Density maps and visual maps of all the archaeological remains (Kernel density analysis) with the localisation of all type of microaggregate lenses from Maras level 4.1.




1467
 1468 **Fig. 8** Density maps and visual maps of all the remains from Maras level 4.1 (Kernel density analysis). Red
 1469 circles show the accumulations areas with higher density of remains and the blue ones those with less
 1470 remains
 1471




1472
 1473
 1474
 1475
 1476

Fig. 9 A Density maps and visual the faunal assemblage, the faunal refits and their connection lines maps from Maras level 4.1, (Kernel density analysis); (B) Example of anatomical refit; (C) Example of green bone fracture refit and (D) Example of dry bone refit. (Photos D. Vettesse & C. Daujeard)



1477
 1478
 1479
 1480
 1481
 1482

Fig. 10 A Spatial patterning of the lithic assemblage and RMU identified from Maras level 4.1 (Kernel density analysis); B Example of RMU in limestone; C Example of isolated pieces (tool kits) in quartzite; D Example of isolated pieces (tool kits) in flint; E Example of some of the identified RMUs in flint (Photos M.G. Chacón & A. Eixea)



1483
 1484
 1485
 1486
 1487

Fig. 11 A Density maps and visual maps of the lithic assemblage and refits identified with their connection lines from Maras level 4.1, (Kernel density analysis); B, E, F Example of refits in flint; C Example of refit in basalt; D Example of refit in limestone; G Example refit in quartz (Photos M.G. Chacón & A. Eixea)