



HAL
open science

“I sing of arms and of a man...”: medial epicondylosis and the sexual division of labour in prehistoric Europe

Sébastien Villotte, Christopher J. Knüsel

► To cite this version:

Sébastien Villotte, Christopher J. Knüsel. “I sing of arms and of a man...”: medial epicondylosis and the sexual division of labour in prehistoric Europe. *Journal of Archaeological Science*, 2014, 43, pp.168-174. 10.1016/j.jas.2013.12.009 . hal-03147057

HAL Id: hal-03147057


<https://hal.science/hal-03147057v1>

Submitted on 22 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

AUTHOR QUERY FORM

 ELSEVIER	Journal: YJASC Article Number: 3917	Please e-mail or fax your responses and any corrections to: E-mail: corrections.esch@elsevier.tnq.co.in Fax: +31 2048 52789
--	--	---

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof.

Location in article	Query / Remark: Click on the Q link to find the query's location in text Please insert your reply or correction at the corresponding line in the proof
Q1	Please check the article footnote, and correct if necessary.
Q2	References "Chapman, 1997; O'Shea and Zvelebil, 1984; Rega, 1997; Stig-Sorensen, 1997; Treherne, 1995; Joyce 2004; Keeley 1986; Jobe and Ciccotti, 1984; Cain et al., 2003; Villotte et al., 2010" are cited in the text but not provided in the reference list. Please provide them in the reference list or delete these citations from the text.
Q3	Uncited references: This section comprises references that occur in the reference list but not in the body of the text. Please cite each reference in the text or, alternatively, delete it. Any reference not dealt with will be retained in this section.
Q4	Please confirm that given names and surnames have been identified correctly. <div data-bbox="304 1410 895 1587" style="border: 1px solid black; padding: 5px; margin-top: 10px;"> <p style="color: red;">Please check this box or indicate your approval if you have no corrections to make to the PDF file</p> <div style="text-align: right; margin-right: 20px;"> <input data-bbox="791 1470 876 1551" type="checkbox"/> </div> </div>

Thank you for your assistance.

Contents lists available at [ScienceDirect](#)

Journal of Archaeological Science

journal homepage: <http://www.elsevier.com/locate/jas>

Highlights

- Thrower's elbow (medial epicondylitis) is a rare occupational syndrome today.
- The lateral to medial (*L/M*) epicondylar ratio is greater than 1 in living populations.
- The *L/M* ratio permits comparisons in past populations.
- European prehistoric males display a specific pattern of lesions for the right elbow.
- Right upper limb use in throwing motions is a long-term status-linked male proclivity.

UNCORRECTED PROOF



Contents lists available at ScienceDirect

Journal of Archaeological Science

journal homepage: <http://www.elsevier.com/locate/jas>

“I sing of arms and of a man...”: medial epicondylitis and the sexual division of labour in prehistoric Europe[☆]

Q4 Sébastien Villotte^{a,*}, Christopher J. Knüsel^b

^a University of Bordeaux, CNRS, PACEA, UMR 5199, F-33405 Talence, France

^b Department of Archaeology, University of Exeter, Exeter, UK

ARTICLE INFO

Article history:

Received 19 October 2013

Received in revised form

10 December 2013

Accepted 16 December 2013

Keywords:

Sexual division of labour

Activity markers

Entheseal changes

Throwing motion

ABSTRACT

Sexual division of labour in European prehistory is usually inferred by indirect means: ethnographic analogy, pictorial representation, or from grave inclusions. The study of skeletal activity-related morphology seems the most direct means by which to interrogate the question of sexual division of labour in past societies. In this paper we present the results of an analysis of enthesopathies (i.e. lesions of the tendon attachments) of the elbow in three time-successive population samples spanning the prehistoric, pre-industrial historic, and modern European eras. We employ an innovative analytical procedure, the lateral to medial epicondylar ratio (*L/M* ratio) to assess limb use. Results indicate a tendency for lateral epicondylitis in all samples, except for prehistoric males, who possess medial epicondylitis more frequently, and for the right side only. The increased prevalence of pathological changes of the right medial epicondyle suggests lateralized limb use that corresponds with “thrower’s elbow”. This indicates that males, but not females, preferentially employed movements involving throwing motions in these hunter-gatherer and early farming groups. Based on this evidence we postulate the existence of a persistent sexual division of labour in these prehistoric European populations involving one or several strenuous activities linked to unilateral limb use.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction

In prehistory sexual division of labour or gendered activity, a part of gender relations that contributes to the roles that men and women play in society, is often inferred from analyses of grave goods, as a part of the social identity of the deceased (see, for example, Chapman, 1997; O’Shea and Zvelebil, 1984; Rega, 1997; Stig-Sorensen, 1997; Treherne, 1995), and artistic representations, both pictographs and figurines (Conkey, 1991; Duhard, 1993; Joyce 2004; Nelson, 1997). Interpretations depend on analogies drawn from cross-cultural studies that provide evidence for ostensibly universal sex differences in labour (Wood and Eagly, 2002). Within non-industrial societies, the majority of tasks are strongly gendered, but substantial diversity exists across societies with respect to which sex performs a particular task (Gilchrist, 2004; Moore, 1994; with regard to the effect of age and social change,

respectively). In almost all groups hunting large game and warfare appear to be largely, albeit not exclusively, men’s activities (Keeley 1986: 35; Knüsel and Smith, 2013; Murdock and Provost, 1973; Testart, 1986; Wood and Eagly, 2002). With regard to hunting, Testart (1986) points out that among modern hunter-gatherers, females can and do hunt, but almost never with weapons that penetrate the flesh. In a sample of 50 worldwide traditional societies from the Human Relations Area Files probability sample, men as opposed to women overwhelmingly engage in games and sports employing projectiles, such as spears, sticks, javelins, darts, and stones (Deaner and Smith, 2012). As a whole, men perform movements involving percussive force, as in the use of an adze or an axe (Testart, 1986); these require similar movements of the upper limb to those employed in throwing, the biomechanics and injurious effects of which has been studied in detail for some years (Bennett, 1941, 1947, 1959; Fleisig et al., 1996; Gainor et al., 1980; Jobe et al., 1983; McCue et al., 1985; Priest et al., 1974; Tullos and Bryan, 1985).

Throwing is among the most dynamic and demanding of physical activities, requiring complex sequential movements of the shoulder, arm, and forearm, as well as each segment of the body from the fingers to the toes in various combinations for the different forms of projectile use. The power of the throw is as much a product of the trunk, hips and lower limbs as it is of the upper

[☆] From the opening lines of Virgil’s *Aeneid*.

* Corresponding author. Anthropologie des Populations Passées et Présentes, PACEA, UMR 5199, Université Bordeaux 1 – CNRS, Bat. B8, Avenue des Facultés, Université Bordeaux 1, 33405 Talence, France.

E-mail address: s.villotte@pacea.u-bordeaux1.fr (S. Villotte).

Table 1
Reported prevalence rates of epicondylitis in modern clinical groups performing various trades, professions, and activities. The higher lesion frequencies for lateral or medial epicondyles is indicated in bold.

Population	N	Lateral epicondyle	Medial epicondyle	L/M ratio	Reference
General practice population	8500	0.7%	0.1%	6.0	Hamilton, 1986
Nursery school cooks (women only)	209	20.1%	11.5%	1.7	Ono et al., 1998
Social welfare employees (other than cook, women only)	366	7.7%	3.0%	2.6	Ono et al., 1998
Automobile manufacturing workers	1198	3.3%	2.2%	1.5	Gold et al., 2009
Assembly-line packers (women only)	152	2.6%	3.3%	0.8	Luopajarvi et al., 1979
Shop assistants (women only)	133	2.3%	0.0%	/	Luopajarvi et al., 1979
General population	4783	1.3%	0.4%	3.3	Shiri et al., 2006
General population (men)	2696	1.3%	0.6%	2.2	Walker-Bone et al., 2004
General population (women)	3342	1.1%	1.1%	1.0	Walker-Bone et al., 2004
Blue collar workers (shoe factory)	253	2.0%	0.0%	/	Roquelaure et al., 2002
Workers in repetitive/constrained jobs (men)	393	1.5%	2.3%	0.7	Nordander et al., 2009
Workers in repetitive/constrained jobs (women)	1106	3.3%	1.6%	2.1	Nordander et al., 2009
Workers in varied/mobile jobs (men)	522	1.5%	0.6%	2.5	Nordander et al., 2009
Workers in varied/mobile jobs (women)	656	1.7%	0.5%	3.4	Nordander et al., 2009
Workers in strenuous jobs	Not clear	0.6%	0.2%	3.0	Viikari-Juntura et al., 1991
Workers in non-strenuous jobs	Not clear	0.5%	0.3%	1.7	Viikari-Juntura et al., 1991
Sewing machine operators (women only)	82	4.9%	0.0%	/	Anderson and Gaardboe, 1993
Auxiliary nurses and home helpers (women only)	25	0.0%	0.0%	/	Anderson and Gaardboe, 1993
Public gas- and waterworks employees, high exposure (right side)	46	10.9%	2.2%	5.0	Ritz, 1995
Public gas- and waterworks employees, high exposure (left side)	46	6.5%	6.5%	1.0	Ritz, 1995
Public gas- and waterworks employees, moderate exposure (right side)	52	7.7%	5.8%	1.3	Ritz, 1995
Public gas- and waterworks employees, moderate exposure (left side)	52	13.5%	1.9%	7.0	Ritz, 1995
Public gas- and waterworks employees, no exposure (right side)	82	7.3%	2.4%	3.0	Ritz, 1995
Public gas- and waterworks employees, no exposure (left side)	82	7.3%	2.4%	3.0	Ritz, 1995
Cases of work-related peri-articular injuries (France, 1998)	8972	16.1%	1.9%	8.6	Gary and Costa-Salute, 2001
workers exposed to repetitive movements	1759	12.3%	4.0%	3.1	Bourgeois et al., 2000
workers not exposed to repetitive movements	343	7.9%	3.5%	2.3	Bourgeois et al., 2000
Baseball pitchers (9–12 years old)	298	11.1%	21.8%	0.5	Lyman et al., 2001

extremities (Toyoshima et al., 1974; Pappas et al., 1985), but the shoulder and elbow joints and their associated soft tissue structures are those most often injured (Ciccotti et al., 2004; Jobe and Ciccotti, 1984).

The torques generated by dynamic movements (i.e. those in excess of the range of normal joint motions) produce injuries in the shoulder and elbow joints of the dominant limb (McCue et al., 1985; Tullos and Bryan, 1985). The overhead throwing motion places valgus stress on the elbow that is transmitted via the flexor-pronator muscles to the medial epicondyle of the humerus, producing “valgus extension overload syndrome”, a form of repetitive stress syndrome that is the most common cause of elbow injuries in the throwing athlete (Cain et al., 2003). Medial epicondylitis,¹ also called “thrower’s elbow” is a condition affecting the site of attachment of the medial collateral ligament and origin of the common flexor tendon at the medial epicondyle (Baccarani and Simonini, 1968; Bramhall et al., 1994; Ciccotti et al., 2004; Jobe and Ciccotti, 1994; Miller, 1960; Ouellette et al., 2008). It is a rare condition seen mainly in sports involving overhead throwing but also in occupational activities creating similar forces (as in hammering in carpentry) (Ciccotti et al., 2004; Jobe and Ciccotti, 1994). Lateral epicondylitis or “tennis elbow” is more common than medial epicondylitis and described in connection with numerous sports and occupational activities (Jobe and Ciccotti, 1994; Shiri and Viikari-Juntura, 2011). When medial and lateral epicondylitis are studied together in living populations the frequency of lateral epicondylitis divided by the frequency of medial epicondylitis (the L/M ratio) is almost always greater than 1 (Table 1). Of those groups studied, baseball pitchers are the only

group where the occurrence of medial epicondylitis is clearly greater than lateral epicondylitis (Table 1). Unilateral enthesopathy of the medial epicondyle has been recognized previously as a good skeletal marker of throwing and used to discuss the sexual division of labour in prehistoric groups (Dutour, 1986; Villotte et al., 2010a) but for small samples, and the L/M ratio has never been studied in detail. The goal of this article is to present the frequencies of lateral and medial epicondylitis in several prehistoric and historic samples and use the L/M ratio in order to discuss the sexual division of labour in activities associated with the throwing motion. Because these repeated movements affect the skeleton in a patterned way, their relative distribution in male and female skeletons provides an unrivalled opportunity to assess sexual division of labour directly from the remains of those who performed tasks.

2. Material and methods

The frequencies of lateral and medial epicondyle enthesopathies in several prehistoric and historic groups have been calculated, and the L/M ratio has been determined for three large samples of prehistoric, pre-industrial historic, and modern documented populations (Table 2). Two different analyses were performed: one involving the total number of observed right and left condyles, and a second of humeri with both condyles preserved and recorded. A single observer (S.V.) performed all entheses examinations employing the same methodological criteria (Villotte, 2006; Villotte et al., 2010b). These samples include adult specimens only. Age at death and sex are known for the identified skeletal collections (the modern documented populations) (see Villotte, 2009 for a description of these collections). Estimation of age-at-death and sex of the prehistoric and pre-industrial historic individuals was based on recently published methods with demonstrably good reliability (references in Villotte et al., 2010a).

¹ Contrary to “epicondylitis”, the more neutral term “epicondylitis” does not suggest inflammation and is thus more appropriate (even if less used). As in most of the cases of medial or lateral epicondyle enthesopathies there is no histopathological evidence of either acute or chronic inflammation (Khan et al., 1999).

Table 2
Frequencies of lateral and medial epicondylitis for each sub-sample. The higher lesion frequencies for lateral or medial epicondyles is indicated in bold.

Samples	Collections	Periods	Females						Males													
			Right epicondyles			Left epicondyles			Right epicondyles			Left epicondyles										
			N total and %	Lateral	Medial	N total and %	Lateral	Medial	N total and %	Lateral	Medial	N total and %	Lateral	Medial								
			L/M ratio			L/M ratio			L/M ratio			L/M ratio										
Modern Documented Sample (223 females and 430 males)	Spitalfields (England) Coimbra (Portugal) Sassari (Italy) Bologna (Italy) Total Modern Documented Sample	18–20th c. AD	67	68.7%	64	39.1%	1.8	60	70.0%	55	34.5%	2.0	53	39.6%	46	19.6%	2.0	49	36.7%	50	12.0%	3.1
			131	35.1%	118	16.1%	2.2	124	21.8%	111	14.4%	1.5	119	20.2%	121	6.6%	3.1	133	11.3%	112	7.1%	1.6
			/	/	/	/	/	/	/	/	/	/	/	64	39.1%	56	8.9%	4.4	65	29.2%	56	12.5%
Pre-industrial Historic Sample (211 females and 89 males)	Post-medieval (France) Medieval (England) Antiquity (France) Total Pre-industrial Historic Sample	16–18th c. AD 10–11th c. AD 2nd–3rd c. AD	106	59.4%	92	33.7%	1.8	96	53.1%	85	21.2%	2.5	11	72.7%	16	43.8%	1.7	16	56.3%	14	28.6%	2.0
			39	64.1%	40	45.0%	1.4	34	58.8%	41	31.7%	1.9	32	68.8%	36	33.3%	2.1	34	47.1%	39	20.5%	2.3
			4	50.0%	6	0.0%	/	7	28.6%	8	12.5%	2.3	20	25.0%	17	11.8%	2.1	16	18.8%	18	16.7%	1.1
Prehistoric Sample (169 females and 139males)	Middle Neolithic (Switzerland) Middle Neolithic (France) Early Neolithic (Germany) Late Mesolithic (Portugal) Early/Middle Mesolithic (Europe) Upper Palaeolithic (Europe) Total Prehistoric Sample	3.2–2700 BC 4.5–4000 BC 5000 BC 6.4–5300 BC 10–7000 BC 30–10,000 BC	21	42.9%	18	22.2%	1.9	20	25.0%	21	23.8%	1.1	15	26.7%	15	53.3%	0.5	9	22.2%	11	18.2%	1.2
			12	41.7%	16	12.5%	3.3	14	35.7%	15	13.3%	2.7	11	18.2%	11	27.3%	0.7	9	22.2%	10	0.0%	/
			25	20.0%	28	10.7%	1.9	24	8.3%	22	9.1%	0.9	35	28.6%	36	36.1%	0.8	29	24.1%	31	6.5%	3.7
			27	29.6%	42	11.9%	2.5	22	22.7%	21	9.5%	2.4	19	0.0%	24	12.5%	0.0	15	6.7%	15	6.7%	1.0
			14	21.4%	11	9.1%	2.4	12	8.3%	11	9.1%	0.9	11	27.3%	9	33.3%	0.8	10	20.0%	10	0.0%	/
			10	10.0%	10	0.0%	/	9	11.1%	10	0.0%	/	15	6.7%	16	25.0%	0.3	17	11.8%	17	0.0%	/
			109	28.4%	125	12.0%	2.4	101	18.8%	100	12.0%	1.6	106	18.9%	111	30.6%	0.6	89	18.0%	94	5.3%	3.4

3. Results

Fig. 1 displays the frequencies of medial and lateral epicondyle enthesopathies in the modern documented (i.e. identified) sample. Frequencies of enthesopathies increase with age, but for each age category and for both sides the *L/M* ratio falls into the range of variation seen for living samples (Table 1). For most of the age categories considered, enthesopathies of the lateral epicondyle are two or three times more common than they are for the medial epicondyle, as for living samples. Moreover, there is no apparent correlation between age-at-death and the ratio. For all comparisons except one, enthesopathies are more frequent for the lateral epicondyle. The ratio below 1 for the first age group for the left side, as well as the very high ratio for the 30–39 age-class, can be explained by the very low frequency of cases seen for the medial epicondyle (one in both cases).

Frequencies for the right and left sides for each collection are presented in Table 2. As there is no control for age, frequencies cannot be compared between groups, but this can be done within each sub-sample. In all modern documented and pre-industrial historic collections, enthesopathies are more common at the lateral epicondyle for both sides and both sexes. For the prehistoric groups, the situation is dramatically different. For females, medial epicondylitis is more frequent in two cases (from 12 comparisons): for the site of Stuttgart-Mühlhausen (Early Neolithic) and for the sample dating from the Early/Middle Mesolithic, a minor difference (less than 1%) is observed for the left side. For males and for each of the groups considered, the ratio clearly favours the medial epicondyle of the right side. When groups are pooled (Fig. 2) the *L/M* ratio is always positive and varies from 1.8 to 3.4, except in the right side of the male prehistoric group (0.6), indicating that, although the lateral epicondyle is affected, it is more often accompanied by medial epicondylar enthesopathies in these males than for their same sex counterparts from more recent times and for females. When only humeri with both medial and lateral epicondyles scored are taken into account and when only one enthesopathy is recorded, there is a significant tendency for the lesion to occur at the lateral epicondyle for all groups and both sides, except for prehistoric males for the right side (Table 3).

4. Discussion

Although enthesal changes are age-related (e.g. Villotte, 2009; Villotte et al., 2010b), for the *L/M* ratio there is no associated age correlation (Fig. 1). We did not attempt to control for age-at-death in the present study due the number of individuals for which it had not been possible to assess it, or for which the assessment provided a very broad age range. Thus, frequencies of medial or lateral epicondylitis cannot be compared across samples, but the *L/M* ratio can. Moreover, the present study clearly shows that this ratio is almost always greater than 1 in the archaeological and clinical samples. The *L/M* ratio for the right side in the prehistoric male sample, below the range of variation seen for historic samples presented here, as well as in clinical samples (Table 1), highlights a specific pattern for the right arm. This very low *L/M* ratio is seen today only in young baseball pitchers (Lyman et al., 2001). This provides strong evidence for the mechanical origin of these lesions, as the pattern would have been seen for both sides if they had resulted from hormonal or other intrinsic factors. Medial epicondyle enthesopathy frequently co-occurs with lateral epicondyle enthesopathy when it is unrelated to the throwing motion (Gold et al., 2009; Walker-Bone et al., 2004). When only humeri with both medial and lateral epicondyles scored are taken into account, and when only one enthesopathy (medial or lateral) is recorded, the right medial epicondyle tends to be more often injured in




Fig. 1. Modern documented sample (four collections of individuals of known sex and age at death). Frequencies of enthesopathies and *L/M* ratios (triangles) for each age group, Top: right side, down: left side.




Fig. 2. *L/M* ratios for the prehistoric, pre-industrial historic, and modern documented samples. Full line and dashed lines: 25, 50th, and 75th percentiles calculated from *L/M* ratios from clinical studies (Table 1), excepted the group of pitchers (for which the ratio is presented by a dotted line).

prehistoric males (Table 3). Considering the high specificity of medial epicondyle injuries today, we suggest that the unilateral pattern of medial epicondylosis indicates that most of these lesions were likely due to the throwing motion. Cross-culturally, humans employ a single hand for tasks demanding skilled use of implements. This is most often the right hand (Coren and Porac, 1977; Calvin, 1983; Faurie and Raymond, 2004; Steele, 2002), and our study reveals that this was the same in the past. Today, ballistic skills, as opposed those requiring fine motor control, disproportionately employ the right hand (Annett, 1970). Within each prehistoric sample, the frequency of medial versus lateral epicondylar enthesopathies is higher in males. The consistency across these groups of hunter-gatherers and farmers provides strong support for a universal biocultural phenomenon linked to throwing, rather than being an artefact of sampling bias.

Hunting was essential during the Upper Palaeolithic and Mesolithic, but also appears to have played a considerable role in European Neolithic groups, providing dietary supplements and as a prestige-enhancing means of social distinction (Hachem, 2001, 2011; Jeunesse, 2010; Uerpmann, 2001). The use of complex projectile weapons, like the bow and arrow, spear-thrower, and dart, which do not seem to especially stress the medial epicondyle (see Villotte et al., 2010 and references therein), is attested in Europe for the latter part of the Upper Palaeolithic, Mesolithic and Neolithic

Table 3

Humeri with both lateral and medial epicondyles scored. The *p*-value (McNemar's test) indicates a significant tendency for increased numbers of cases of lateral epicondylitis in all samples, except for prehistoric males, where individuals with medial epicondylitis alone are more prevalent than those with the lateral form, and on the right side only.

		Right				Left			
		N	Lateral epicondylitis only	Medial epicondylitis only	<i>P</i>	N	Lateral epicondylitis only	Medial epicondylitis only	<i>P</i>
Modern documented sample	Females	169	36	2	<0.001	149	29	3	<0.001
	Males	293	53	8	<0.001	304	39	9	<0.001
Pre-industrial historic sample	Females	115	32	4	<0.001	106	32	0	<0.001
	Males	56	20	2	<0.001	58	12	0	0.002
Prehistoric sample	Females	89	20	6	0.011	74	10	0	0.004
	Males	92	4	15	0.022	78	11	3	0.003

(Cattelain, 2006; Cattelain and Bellier, 2002; Clark, 1963; Rozoy, 1999; Shea, 2006). However, this does not exclude the use of more simple technology simultaneously: barbed points, javelins, killing sticks and boomerangs are known for the same period (Bahn, 1987; Burov, 1998; Cattelain and Bellier, 2002; Valde-Nowak et al., 1987). Even thrown stones are highly effective in hunting and warfare (Isaac, 1987). As documented in ethnographic literature (see Churchill, 1993 for a review), hand-thrown projectiles likely formed a frequent complement to the use of more complex weaponry in the prehistoric groups studied here. Thus, it appears that most cases of medial epicondylitis observed for these individuals are related to the use of overhead throwing during games, warfare, and hunting. Overhead throwing with force and accuracy is a skill uniquely developed in our species that played an important role in human evolution (Calvin, 1983; Darlington, 1975; Darwin, 1871; Isaac, 1987; Knüsel, 1992). Effective throwing behaviour (probably of stones initially) may date as far back as 4 million years ago (Calvin, 1983; Darlington, 1975; Isaac, 1987). Today, males perform significantly better than females in spatial ability tests, as in gauging movements across the visual field (Jardine and Martin, 1983; Voyer et al., 1995), as well as in both throwing velocity and accuracy exercises (Jardine and Martin, 1983; Tillaar and Ettema, 2004; Whittaker and Kamp, 2006). Significant correlations have been found between throwing accuracy and spatial ability, and these sex differences may reflect the evolutionary adaptations required of male hunters and later as armed warriors (Jardine and Martin, 1983; Kolakowski and Malina, 1974). Because this proclivity appears to form a fundamental part of masculine identity, prolonged training and socialization to achieve proficiency may be implicated in the occurrence of observed medial epicondyle enthesopathies.

Other activities may be associated with some of the medial epicondylitis seen in prehistoric males. The sling, an ancient implement used by herders to tend flocks, farmers to protect crops from scavengers – and as a weapon especially well known in the classical ancient world (Paunov and Dimitrov, 2000; Dohrenwend, 2002) – suggests one such activity, with evidence for what may be sling stones from prehistoric sites worldwide (Finney, 2006; Rosenberg, 2009; Vega and Craig, 2009), including the European Neolithic (Paunov and Dimitrov, 2000). Interestingly, as for other throwing activities men have a greater range in sling use (Vega and Craig, 2009). The intensive use of axes and adzes, objects which are well known from the Neolithic and throughout the Mesolithic (e.g. Valdeyron, 2011), may also have produced valgus stress in the medial compartment of the humerus similar to that seen in overhead throwing. Unilateral wood-working is a masculine activity in non-industrial societies (the average percentage of male participation calculated from 185 societies is 98.8%; Murdock and Provost, 1973), and it is noteworthy that thick-bladed adzes are found almost exclusively in male burials from the Early Neolithic

Linearbandkeramik (Bakels, 1987), perhaps as a means of signalling masculine identity.

5. Conclusion

Sexual division in the use of throwing weapons is one of the most striking universal occurrences noted in cross-cultural studies (Murdock and Provost, 1973; Testart, 1986; Wood and Eagly, 2002). Our results indicate that sex differences indicative of this proclivity date back at least 30,000 years. Moreover, this sexual division of labour appears to have been a persistent feature of human groups through millennia and across societies with radically different subsistence economies and living a diverse range of environments. An ability to throw with accuracy and at increasing distance may represent one of the fundamental structuring factors in past human societies, with both biological and social implications.

Uncited references

Churchill and Rhodes, 2009; Dunsworth et al., 2003; Knüsel, 2000; Rhodes and Churchill, 2009.

Acknowledgements

The authors thank the two reviewers for their useful comments. S.V. thanks the many curators across Europe who allowed him access to the material.

Grant sponsors: Maison des Sciences de l'Homme d'Aquitaine; European Union (Synthesys GB-TAF-636); CNRS; Ministère de l'Enseignement Supérieur et de la Recherche (PhD research grant, and "Aires Culturelles"); Fondation Fyssen (Post doctoral research fellowship), British Academy (Small Research Grant).

References

- Anderson, J., Gaardboe, O., 1993. Musculoskeletal disorders of the neck and upper limb among sewing machine operators: a clinical investigation. *Am. J. Ind. Med.* 24 (6), 689–700.
- Annett, M., 1970. A classification of hand preference by association analysis. *Br. J. Psychol.* 61 (3), 303–321.
- Baccarani, G., Simonini, D., 1968. Su due casi di epitrocleite in lanciatori di giavellotto. *Osp. d'Italia Chir.* 19 (4–5), 409–413.
- Bahn, P.G., 1987. Return of the Euro-boomerang. *Nature* 329 (6138), 388.
- Bakels, C.C., 1987. On the adzes of the Northwestern Linearbandkeramik. *Analecta Praehist. Leidensia* 20, 53–85.
- Bennett, G.E., 1941. Shoulder and elbow lesions of the professional baseball pitcher. *J. Am. Med. Assoc.* 117, 510–514.
- Bennett, G.E., 1947. Shoulder and elbow lesions distinctive of baseball players. *Ann. Surg.* 26, 107–110.
- Bennett, G.E., 1959. Elbow and shoulder lesions of baseball players. *Am. J. Surg.* 98, 484–492.
- Bourgeois, F., Lemarchand, C., Hubault, F., Brun, C., Polin, A., Fauchoux, J., 2000. Troubles musculosquelettiques et travail. Quand la santé interroge

- l'organisation. Editions de l'ANACT (Agence nationale pour l'Amélioration des Conditions de Travail), Lyon.
- Bramhall, J.P., Scarpinato, D.F., Andrews, J.R., 1994. Injuries in throwing sports. In: Renström, P. (Ed.), *Clinical Practice of Sports Injury Prevention and Care*, The Encyclopaedia of Sports Medicine, vol. 5. Blackwell Scientific Publications, Oxford, pp. 446–449.
- Burov, G.M., 1998. The use of vegetable materials in the Mesolithic of Northeast Europe. In: Zvelebil, M., Domańska, L., Dennell, R. (Eds.), *Harvesting the Sea, Farming the Forest: the Emergence of Neolithic Societies in the Baltic Region*. Sheffield Academic Press, Sheffield (UK), pp. 53–64.
- Calvin, W.H., 1983. A stone's throw and its launch window: timing precision and its implications for language and hominid brains. *J. Theor. Biol.* 104, 121–135.
- Cattelain, P., 2006. Apparition et évolution de l'arc et des pointes de flèches dans la préhistoire européenne (Paléo-, Mésol-, Néolithique). In: Bellintani, P. (Ed.), *Catene Operative Dell'arco Preistorico*. Giunta della Provincia Autonoma di Trento, Trento, pp. 45–66.
- Cattelain, P., Bellier, C., 2002. La chasse dans la Préhistoire du Paléolithique au Néolithique en Europe... et ailleurs. Treignes. Éditions du CEDARC.
- Churchill, S.E., 1993. Weapon technology, prey size selection, and hunting methods in modern hunter-gatherers: implications for hunting in the Palaeolithic and Mesolithic. *Archeol. Pap. Am. Anthropol. Assoc.* 4 (1), 11–24.
- Churchill, S.E., Rhodes, J.A., 2009. The evolution of the human capacity for "killing at a distance": the human fossil evidence for the evolution of projectile weaponry. In: Hublin, J.-J., Richards, M.P. (Eds.), *The Evolution of Hominin Diets: Integrating Approaches to the Study of Palaeolithic Subsistence*. Springer, Dordrecht, pp. 201–210.
- Ciccotti, M.C., Schwartz, M.A., Ciccotti, M.G., 2004. Diagnosis and treatment of medial epicondylitis of the elbow. *Clin. Sports Med.* 23 (4), 693–705.
- Clark, J.G.D., 1963. Neolithic bows from Somerset, England, and the prehistory of archery in North-Western Europe. *Proc. Prehist. Soc.* 29 (3), 50–98.
- Conkey, M.W., 1991. Contexts of action, contexts for power: material culture and gender in the Magdalenian. In: Gero, J., Conkey, M.W. (Eds.), *Engendering Archaeology: Women and Prehistory*. Wiley-Blackwell, New York, pp. 57–92.
- Coren, S., Porac, C., 1977. Fifty centuries of right-handedness: the historical record. *Science* 198, 631–632.
- Darlington, P.J., 1975. Group selection, altruism, reinforcement, and throwing in human evolution. *Proc. Natl. Acad. Sci.* 72 (9), 3748–3752.
- Darwin, C.R., 1871. *The Descent of Man, and Selection in Relation to Sex*, first ed., vol. 1. John Murray, London.
- Deaner, R.O., Smith, B.A., 2012. Sex differences in sports across 50 societies. *Cross-Cult. Res. J. Comp. Soc. Sci.* Available at: SSRN: <http://ssrn.com/abstract=2009329>.
- Dohrenwend, R.E., 2002. The sling: forgotten firepower of antiquity. *J. Asian Martial Arts* 11 (2), 29–49.
- Duhard, J.-P., 1993. Upper Palaeolithic figures as a reflection of human morphology and social organization. *Antiquity* 67, 83–91.
- Dunsword, H., Challis, J., Walke, A., 2003. Throwing and bipedalism: a new look at an old idea. In: Franzen, J., Köhler, M., Moyà-Solà, S. (Eds.), *Walking Upright*. Senckenberg Institute, Frankfurt, pp. 105–110.
- Dutour, O., 1986. Enthesopathies (lesions of muscular insertions) as indicators of the activities of Neolithic Saharan populations. *Am. J. Phys. Anthropol.* 71 (2), 221–224.
- Faurie, C., Raymond, M., 2004. Handedness frequency over more than ten thousand years. *Proc. R. Soc. Lond. B* 271 (Suppl.), S43–S45.
- Finney, J.B., 2006. Middle Iron Age Warfare of the Hillfort Dominated Zone c. 400 B.C. to c. 150 B.C. In: *British Archaeological Reports (British Series) 423*. Archaeopress, Oxford.
- Fleisig, G.S., Barrentine, S.W., Escamilla, R.F., Andrews, J.R., 1996. Biomechanics of overhead throwing with implications for injuries. *Sports Med.* 21, 421–437.
- Gainor, B.J., Piotrowski, G., Puhl, J., Allen, W.C., Hagen, R., 1980. The throw: biomechanics and injury. *Am. J. Sports Med.* 8 (2), 114–118.
- Gary, Y., Costa-Salute, C., 2001. La CRAM et les troubles musculo-squelettiques du membre supérieur. De l'analyse épidémiologique à la prévention. In: Hérisson, C., Fouquet, B., Codine, P. (Eds.), *Membre supérieur et pathologie professionnelle*. Masson, Paris, pp. 6–16.
- Gilchrist, R., 2004. Archaeology and the life course: a time and age for gender. In: Meskell, L., Preucel, R.W. (Eds.), *A Companion to Social Archaeology*. Blackwell Publishing, Malden (MA), pp. 142–160.
- Gold, J.E., d'Errico, A., Katz, J.N., Gore, R., Punnett, L., 2009. Specific and non-specific upper extremity musculoskeletal disorder syndromes in automobile manufacturing workers. *Am. J. Ind. Med.* 52 (2), 124–132.
- Hachem, L., 2001. La conception du monde animal sauvage chez les éleveurs du Rubané. In: Arbogast, R.-M., Jeunesse, C., Schibler, J. (Eds.), *Rôle et statut de la chasse dans le Néolithique ancien danubien (5500–4900 av J-C)*, Premières rencontres danubiennes, Actes de la table ronde de Strasbourg 20 et 21 novembre 1996. International Archäologie, Arbeitsgemeinschaft Symposium Tagung Kongress 1. Verlag Marie Leidorf GmbH, Rahden/Westfalen, pp. 91–111.
- Hachem, L., 2011. Les faunes du Néolithique moyen dans le nord de la France: bilan et pistes de recherches. In: Bostyn, F., Martial, E., Praud, I. (Eds.), *Le Néolithique du Nord de la France dans son contexte européen: habitat et économie aux 4^e et 3^e millénaires avant notre ère*, Actes du 29^e colloque international sur le Néolithique Revue Archéologique de Picardie, n° spécial 28, pp. 313–329.
- Hamilton, P.G., 1986. The prevalence of humeral epicondylitis: a survey in general practice. *J. R. Coll. Gen. Pract.* 36 (291), 464–465.
- Isaac, B., 1987. Throwing and human evolution. *Afr. Archaeol. Rev.* 5, 3–17.
- Jardine, R., Martin, N.G., 1983. Spatial ability and throwing accuracy. *Behav. Genet.* 13 (4), 331–340.
- Jeunesse, C., 2010. Changements sociaux et signification de la chasse dans les sociétés du Néolithique circumalpin. Une hypothèse alternative au déterminisme écologique. In: Matuschik, I., Strahm, C., Eberschweiler, B., Fingerlin, G., Hafner, A., Kinsky, M., Mainberger, M., Schobel, G. (Eds.), *Vernetzungen - Aspekte siedlungsarchäologischer Forschung*, Festschrift für Helmut Schlichtherle zum 60 Geburtstag. Lavori Verlag, Freiburg im Breisgau, pp. 127–139.
- Jobe, F., Ciccotti, M., 1994. Lateral and medial epicondylitis of the elbow. *J. Am. Acad. Orthop. Surg.* 2 (1), 1–8.
- Jobe, F.W., Tibone, J.E., Perry, J., Moynes, D., 1983. An EMG analysis of the shoulder in throwing and pitching: a preliminary report. *Am. J. Sports Med.* 11 (1), 3–5.
- Khan, K.M., Cook, J.L., Bonar, F., Harcourt, P., Astrom, M., 1999. Histopathology of common tendinopathies. Update and implications for clinical management. *Sports Med.* 27 (6), 393–408.
- Knüsel, C.J., 1992. Throwing and hominid evolution. *Hum. Evol.* 7 (1), 1–7.
- Knüsel, C.J., 2000. Bone adaptation and its relationship to physical activity in the past. In: Cox, M., Mays, S. (Eds.), *Human Osteology in Archaeology and Forensic Science*. Greenwich Medical Media, London, pp. 381–402.
- Knüsel, C.J., Smith, M.J., 2013. The osteology of conflict: what does it all mean. In: Knüsel, C.J., Smith, M.J. (Eds.), *The Routledge Handbook of the Bioarchaeology of Human Conflict*. Routledge, London, pp. 656–694.
- Kolakowski, D., Malina, R.M., 1974. Spatial ability, throwing accuracy and man's hunting heritage. *Nature* 251 (5474), 410–412.
- Luopajarvi, T., Kuorinka, I., Virolainen, M., Holmberg, M., 1979. Prevalence of tenosynovitis and other injuries of the upper extremities in repetitive work. *Scand. J. Work Environ. Health* 5 (3), 48–55.
- Lyman, S., Fleisig, G.S., Waterbor, J.W., Funkhouser, E.M., Pulley, L., Andrews, J.R., Osinski, E.D., Roseman, J.M., 2001. Longitudinal study of elbow and shoulder pain in youth baseball pitchers. *Med. Sci. Sports Exerc.* 33 (11), 1803–1810.
- McCue III, F.C., Gieck, J.H., West, J.O., 1985. Throwing injuries to the shoulder. In: Zarins, B., Andrews, J.R., Carson, W.A. (Eds.), *Injuries to the Throwing Arm*. W.B. Saunders and Co, Philadelphia, pp. 95–111.
- Miller, J.E., 1960. Javelin thrower's elbow. *J. Bone Jt. Surg. Br.* 42 (4), 788–792.
- Moore, H., 1994. Understanding sex and gender. In: Ingold, T. (Ed.), *Companion Encyclopedia of Social Anthropology: Humanity, Culture, and Social Life*. Routledge, London and New York, pp. 813–830.
- Murdoch, G.P., Provost, C., 1973. Factors in the division of labor by sex: a cross-cultural analysis. *Ethnology* 12 (2), 203–225.
- Nelson, S.M., 1997. *Gender in Archaeology*. Altamira, Walnut Creek (CA).
- Nordander, C., Ohlsson, K., Akesson, I., Arvidsson, I., Balogh, I., Hansson, G., Strömberg, U., Rittner, R., Skerfving, S., 2009. Risk of musculoskeletal disorders among females and males in repetitive/constrained work. *Ergonomics* 52 (10), 1226–1239.
- Ono, Y., Nakamura, R., Shimaoka, M., Hiruta, S., Hattori, Y., Ichihara, G., Kamijima, M., Takeuchi, Y., 1998. Epicondylitis among cooks in nursery schools. *Occup. Environ. Med.* 55 (3), 172–179.
- Ouellette, H., Bredella, M., Labis, J., Palmer, W., Torriani, M., 2008. MR imaging of the elbow in baseball pitchers. *Skelet. Radiol.* 37 (2), 115–121.
- Pappas, A.M., Zawacki, R.M., Sullivan, T.J., 1985. Biomechanics of baseball pitching: a preliminary report. *Am. J. Sports Med.* 13 (4), 216–222.
- Paunov, E., Dimitrov, D.Y., 2000. New data on the use of war sling in Thrace (4th–1st century BC). *Archaeol. Bulg.* IV (3), 44–57.
- Priest, J.D., Jones, H.H., Nagel, D.A., 1974. Elbow injuries in highly skilled tennis players. *J. Sports Med.* 2 (3), 137–149.
- Rhodes, J.A., Churchill, S.E., 2009. Throwing in the Middle and Upper Paleolithic: inferences from an analysis of humeral retroversion. *J. Hum. Evol.* 56 (1), 1–10.
- Ritz, B.R., 1995. Humeral epicondylitis among gas- and waterworks employees. *Scand. J. Work Environ. Health* 21 (6), 478–486.
- Roquelaure, Y., Mariel, J., Fanello, S., Boissière, J.-C., Chiron, H., Dano, C., Bureau, D., Penneau-Fontbonne, D., 2002. Active epidemiological surveillance of musculoskeletal disorders in a shoe factory. *Occup. Environ. Med.* 59 (7), 452–458.
- Rosenberg, D., 2009. Flying stones – the slingstones of the Wadi Rabah Culture of the southern Levant. *Paléorient* 35 (2), 99–112.
- Rozoy, J.-G., 1999. Le mode de vie au Mésolithique. In: Thévenin, A., Bintz, P. (Eds.), *L'Europe des derniers chasseurs. Épipaléolithique et Mésolithique*, Actes du Colloque International U.I.S.P.P. (Grenoble septembre 1995). CTHS, Paris, pp. 39–50.
- Shea, J.J., 2006. The origins of lithic projectile point technology: evidence from Africa, the Levant and Europe. *J. Archaeol. Sci.* 33, 823–846.
- Shiri, R., Viikari-Juntura, E., 2011. Lateral and medial epicondylitis: role of occupational factors. *Best Pract. Res. Clin. Rheumatol.* 25 (1), 43–57.
- Shiri, R., Viikari-Juntura, E., Varonen, H., Heliövaara, M., 2006. Prevalence and determinants of lateral and medial epicondylitis: a population study. *Am. J. Epidemiol.* 164 (11), 1065–1074.
- Steele, J., 2002. When did directional asymmetry enter the record? In: Crow, T.J. (Ed.), *The Speciation of Modern Homo Sapiens Proceedings of the British Academy vol. 106*, 153–168.
- Testart, A., 1986. *Essai sur les fondements de la division sexuelle du travail chez les chasseurs-cueilleurs*. EHESS, Paris.
- Tillaar, R., Ettema, G., 2004. Effect of body size and gender in overarm throwing performance. *Eur. J. Appl. Physiol.* 91 (4), 413–418.
- Toyoshima, S., Hoshikawa, T., Miyashita, M., Oguri, T., 1974. Contribution of body parts to throwing performance. In: Melson, R.C., Morehouse, C.A. (Eds.), *Biomechanics IV*. University Park Press, Baltimore (MD), pp. 169–174.

- Tullos, H.S., Bryan, W.J., 1985. Examination of the throwing elbow. In: Zarins, B., Andrews, J.R., Carson, W.A. (Eds.), *Injuries to the Throwing Arm*. W.B. Saunders and Co, Philadelphia, pp. 191–200.
- Uerpman, M., 2001. Animaux sauvages et domestiques du Rubané “le plus ancien” (LBK 1) en Allemagne. In: Arbogast, R.-M., Jeunesse, C., Schibler, J. (Eds.), *Rôle et statut de la chasse dans le Néolithique ancien danubien (5500–4900 av J-C)*, Premières rencontres danubiennes. Actes de la table ronde de Strasbourg 20 et 21 novembre 1996. Rahden/Westfahlen: International Archäologie, Arbeitsgemeinschaft Symposium Tagung Kongress 1. Verlag Marie Leidorf GmbH, pp. 57–75.
- Valde-Nowak, P., Nadachowski, A., Wolsan, M., 1987. Upper Palaeolithic boomerang made of a mammoth tusk in south Poland. *Nature* 329 (6138), 436–438.
- Valdeyron, N., 2011. Derniers chasseurs et premiers bûcherons? La question des haches et des herminettes dans le Mésolithique européen. In: Servelle, C. (Ed.), *Haches de pierre au néolithique, les premiers paysans du Tarn*. Comité départemental d'archéologie du Tarn, Castres, pp. 419–434.
- Vega, M.B., Craig, N., 2009. New experimental data on the distance of sling projectiles. *J. Archaeol. Sci.* 36, 1264–1268.
- Viiikari-Juntura, E., Kurppa, K., Kuosma, E., Huuskonen, M., Kuorinka, I., Ketola, R., Konni, U., 1991. Prevalence of epicondylitis and elbow pain in the meat-processing industry. *Scand. J. Work Environ. Health* 17 (1), 38–45.
- Villotte, S., 2006. Connaissances médicales actuelles, cotation des enthésopathies: nouvelle méthode. *Bull. Mém. Soc. d'Anthropol. Paris* 18 (1–2), 65–85.
- Villotte, S., 2009. Enthésopathies et activités des hommes préhistoriques - Recherche méthodologique et application aux fossiles européens du Paléolithique supérieur et du Mésolithique. In: *British Archaeological Reports (International Series) 1992*. Archaeopress, Oxford.
- Villotte, S., Churchill, S.E., Dutour, O., Henry-Gambier, D., 2010a. Subsistence activities and the sexual division of labor in the European Upper Paleolithic and Mesolithic: evidence from upper limb enthesopathies. *J. Hum. Evol.* 59 (1), 35–43.
- Villotte, S., Castex, D., Couallier, V., Dutour, O., Knüsel, C.J., Henry-Gambier, D., 2010b. Enthesopathies as occupational stress markers: evidence from the upper limb. *Am. J. Phys. Anthropol.* 142 (2), 224–234.
- Voyer, D., Voyer, S., Bryden, M.P., 1995. Magnitude of sex differences in spatial abilities: a meta-analysis and consideration of critical variables. *Psychol. Bull.* 117 (2), 250–270.
- Walker-Bone, K., Palmer, K., Reading, I., Coggon, D., Cooper, C., 2004. Prevalence and impact of musculoskeletal disorders of the upper limb in the general population. *Arthritis Rheum.* 51 (4), 642–651.
- Whittaker, J.C., Kamp, K.A., 2006. Primitive weapons and modern sport: atlatl capabilities, learning, gender, and age. *Plains Anthropol.* 51 (198), 213–221.
- Wood, W., Eagly, A.H., 2002. A cross-cultural analysis of the behavior of women and men: implications for the origins of sex differences. *Psychol. Bull.* 128 (5), 699–727.