



**HAL**  
open science

## Contribution of *Posidonia oceanica* meadows in the context of climate change mitigation in the Mediterranean Sea

Christine Pergent-Martini, Gérard Pergent, Briac Monnier, Charles-François Boudouresque, Christophe Mori, Audrey Valette-Sansevin

### ► To cite this version:

Christine Pergent-Martini, Gérard Pergent, Briac Monnier, Charles-François Boudouresque, Christophe Mori, et al.. Contribution of *Posidonia oceanica* meadows in the context of climate change mitigation in the Mediterranean Sea. *Marine Environmental Research*, 2020, 10.1016/j.marenvres.2020.105236 . hal-03150734

**HAL Id: hal-03150734**

**<https://hal.science/hal-03150734v1>**

Submitted on 2 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

# 1 Contribution of *Posidonia oceanica* meadows in the context of climate change 2 mitigation in the Mediterranean Sea

3

4 Christine Pergent-Martini<sup>(1)</sup>, Gérard Pergent<sup>(1)</sup>, Briac Monnier<sup>(1)</sup>, Charles-François  
5 Boudouresque<sup>(2)</sup>, Christophe Mori<sup>(1)</sup>, Audrey Valette<sup>(1)</sup>

6

7 <sup>(1)</sup>FRES 3041 – UMR 6134, Coastal Ecosystems Team, University of Corsica, 20250 Corte, France

8 <sup>(2)</sup>Aix-Marseille University and University of Toulon, MIO (Mediterranean Institute of Oceanography),  
9 CNRS, IRD, Campus de Luminy, Marseille, France

10

11

## 12 Abstract

13 Coastal marine vegetation has been recently highlighted for its highly efficient carbon  
14 storage capacity. Among the sixty-four species of seagrass, *Posidonia oceanica*, a  
15 Mediterranean endemic species, appears to be the most effective in carbon fixation and  
16 storage. Based on new data from the study of one of the largest *P. oceanica* meadows in the  
17 Mediterranean Sea (100 km of coastline, 20 425 ha), and a synthesis of available data from  
18 the whole of the Mediterranean basin, the aim of this work is to evaluate the amount of  
19 carbon fixed each year by *P. oceanica* and sequestered in the mat, in relation with the  
20 mitigation of the impact of climate change (carbon sink).

21 The mean total carbon fixation (blades, sheaths and rhizomes) per year varies between 33.5  
22 and 426.6 g C.m<sup>-2</sup> and the mean carbon sequestration (long-term sink in the mat),  
23 corresponding to the sheath and rhizome tissues, varies between 7.7 and 84.4 g C.m<sup>-2</sup>, with  
24 a clear decreasing trend according to depth because of the meadow density decrease.

25 The synthesis of a hundred measurements made throughout the Mediterranean Sea and at  
26 depths between 0.5 and 32.0 m provides a basis for estimating the average annual carbon  
27 fixation and sequestration rate throughout the Mediterranean basin. The fixation of the  
28 blades is estimated at 1 024 t C.ha<sup>-1</sup>.yr<sup>-1</sup>, that of the sheaths at 220 t C ha<sup>-1</sup>.yr<sup>-1</sup> and that of  
29 the rhizomes at 58 t C ha<sup>-1</sup>.yr<sup>-1</sup>; *i.e.* a total fixation rate of 1 302 t C ha<sup>-1</sup>.yr<sup>-1</sup> and  
30 sequestration rate (dead sheaths and rhizomes) of 278 t C ha<sup>-1</sup>.yr<sup>-1</sup>. This annual carbon  
31 fixation represents only 0.61% on average of CO<sub>2</sub> emissions/releases for all Mediterranean  
32 countries but in the large Mediterranean islands this fixation is on average 3.1% and can  
33 reach almost 14.4% for Corsica. Moreover, the major advantage of the *P. oceanica* meadow  
34 lies in its capacity to store carbon from annual carbon sequestration for centuries to  
35 millennia and can be compared to several terrestrial ecosystems considered to be efficient in  
36 carbon storage (peatlands).

37

38 **Key words:** Seagrass; *Posidonia oceanica*; Carbon Fixation; Carbon Sequestration;  
39 Mediterranean Sea; Corsica.

40

41

## 42 Introduction

43 At the 21st meeting of the Conference of the Parties (COP 21) of the Climate Change  
44 Convention, all participants adopted the Paris Agreement. This agreement stipulates in  
45 Article 5 that '*Parties should take action to conserve and enhance, as appropriate, sinks and*  
46 *reservoirs of greenhouse gases*' (FCCC, 2015). On the other hand, while previous conferences  
47 had focused on forest-based sinks (Kyoto Protocol), for the first time COP 21 noted '*the*

48 *importance of ensuring the integrity of all ecosystems, including oceans'*. Major carbon sinks  
49 such as wetlands, peatlands, coastal vegetation and phytoplankton were, therefore, for the  
50 first time taken into account.

51  
52 The concept of carbon sink covers several very different compartments, which are often  
53 erroneously confused in the literature. (i) Long-term sinks correspond to carbon which is not  
54 re-mineralized after the death of the primary producer and therefore sequestered for  
55 thousand or million years. This sink corresponds to coal, peat and modern deposits. (ii)  
56 Short-term sinks correspond to the amount of carbon which is incorporated into living  
57 matter, for a few days to millennia, depending upon the life span of the organisms: days for  
58 phytoplankton, months for leaves, centuries to millennia for tree trunks. In fact, this  
59 corresponds to the biomass. Part of the so-called carbon sequestration by the European  
60 forests is just due the expansion of forests, following agricultural decline, and to the ageing  
61 of forests, which accumulate carbon in the wood; this carbon is more or less slowly  
62 mineralized after the death of the leaves or of the whole plant. (iii) A number of plants and  
63 habitats, such as *Posidonia oceanica* (Linnaeus) Delile (see below), may fall in between, due  
64 to the very low rate of re-mineralization of some tissues (rhizomes and roots), accumulated  
65 in the soil, able to persist during several millennia.

66  
67 Despite a very small surface area (less than 0.5% of the oceans), coastal vegetation plays a  
68 major role in the fixation and sequestration of Blue Carbon (Nellemann *et al.*, 2009;  
69 Fourqurean *et al.*, 2012). In fact, these ecosystems, such as seagrass beds, kelp forests,  
70 mangroves, salt marshes and wetlands, are major carbon sinks (Chmura *et al.*, 2003; Duarte  
71 *et al.*, 2005; Bouillon *et al.*, 2008; Reed & Brzezinski, 2009).

72  
73 Carbon sequestration by seagrass is estimated at 15% of the blue carbon while covering only  
74 17.7 to 61.0 million hectares at the biosphere scale (Duarte & Chiscano, 1999; Spalding *et al.*,  
75 2003; Kennedy & Björk, 2009 ; UNEP-WCMC, 2018). Among the sixty-four species of  
76 seagrass (Guiry & Guiry, 2020), *P. oceanica*, a Mediterranean endemic species, appears to be  
77 the most effective in carbon storage (Pergent *et al.*, 1994; Mateo *et al.*, 2006; Kennedy &  
78 Björk, 2009; Pergent *et al.*, 2014; Jamaludin, 2015. The *P. oceanica* meadow is the only  
79 ecosystem able to 'compete' with peatlands (IPS, 2008) and mangroves (Bouillon *et al.*,  
80 2008) because it builds a unique structure: the matte (Boudouresque *et al.*, 2012). Made up  
81 of rhizomes and roots with sediment that fills the interstices, this little putrescible structure  
82 can reach several meters in height, and the organic and inorganic carbon inside can persist  
83 for millennia (Romero *et al.*, 1994; Mateo *et al.*, 1997; Lo Iacono *et al.*, 2008; Tomasello *et al.*,  
84 2009; Boudouresque *et al.*, 2016; Monnier *et al.*, *in press*).

85  
86 The surface area covered by *P. oceanica* meadows in the Mediterranean Sea has been the  
87 subject of several estimates (Pasqualini *et al.*, 1998, UNEP-MAP-RAC/SPA, 2009; Giakoumi *et al.*,  
88 2013; Telesca *et al.*, 2015). The scarcity of data for the eastern basin detracts from the  
89 accuracy of these estimates, although recent mapping works of Topouzelis *et al.* (2018) and  
90 Traganos *et al.* (2018), for the Greek coastline (32% of the Mediterranean coastline), Akçali  
91 *et al.* (2019) and Duman *et al.* (2019) for 3 000 km of Turkish Mediterranean coastline, have  
92 made it possible to improve the accuracy of the previous estimates.

93

94 On the other hand, the increase of the amount of data concerning primary production by *P.*  
95 *oceanica* at many sites including different locations, depths, meadow structure and human  
96 pressures, has made it possible to estimate a probable mean value of the amount of carbon  
97 fixed by *P. oceanica* at the scale of the Mediterranean basin (Buia *et al.*, 1992; Pergent-  
98 Martini *et al.*, 1994; Pergent *et al.*, 1997; Torricelli & Peirano, 1997; Zupo *et al.*, 1997;  
99 Guidetti *et al.*, 2000; Cebrian & Duarte, 2001; Vela *et al.*, 2006). Moreover, the use of  
100 lepidochronology associated with the analysis of the elementary composition of tissues  
101 allows determination of the part of this carbon sequestered within the matte mainly through  
102 rhizomes and dead sheaths (Pergent *et al.*, 1994; Pergent-Martini *et al.*, 1994; Cebrian &  
103 Duarte, 2001; Gobert *et al.*, 2005; Vela *et al.*, 2006; Scartazza *et al.*, 2017).

104

105 Based on new data concerning one of the largest *P. oceanica* meadows in the Mediterranean  
106 Sea (100 km of coastline, 20 425 ha; Valette *et al.*, 2019) and the synthesis of available data  
107 from the whole of the Mediterranean basin, the aim of this work is (i) to evaluate the  
108 amount of carbon fixed each year by *P. oceanica* and sequestered in the matte, and (ii) to  
109 better understand this ecosystem service in relation with the mitigation of the impact of  
110 climate change (carbon sink) for each Mediterranean country.

111

112

### 113 **Material and method**

114 The distribution of *P. oceanica* along the Corsican coast is conventionally subdivided into 14  
115 sectors with similar environmental conditions (Meinesz *et al.*, 1990; Pasqualini *et al.*, 1998;  
116 Valette *et al.*, 2019). The study area is located along the eastern coast of Corsica (Western  
117 Mediterranean Sea) characterized by the presence of an extensive *P. oceanica* meadow,  
118 designated as a NATURA 2000 site 'FR9402014 - Grand Herbier de la Côte Orientale' (Valette  
119 *et al.*, 2019; Fig. 1). Sampling stations are located at 5, 10, 15, 20, 25 and 30 m depth and  
120 distributed from north to south.

121




Figure 1: Left: Corsica island, with sectors 1 through 14. Middle: The studied sector (sector 2). Right: Location of the study sites and sampling stations of *P. oceanica* shoots.

The meadow density (number of shoots per square meter – shoot.m<sup>-2</sup>) was measured for each station (0.16 m<sup>2</sup> quadrat, 20 replicates) while 20 orthotropic shoots of *P. oceanica* were collected every three months, from July 2015 to July 2016.

A phenological analysis (Giraud, 1979), and a lepidochronological study (Pergent, 1990) were performed. The primary production was estimated according to the crossed lepidochronological method (Vela, 2006), derived from Pergent & Pergent-Martini (1990). A leaf is formed by a basal non-photosynthetic sheath and a blade. The blade primary production was estimated considering: the average number of annually produced leaves, the average length of the rank 3 adult blades (usually the longest leaf of the shoot), the average width of the rank 1 adult blades, the average density (ratio of leaf area to leaf dry mass) of the rank 1 adult blades (leaf growth completed). The sheath production was estimated considering: the average number of annually produced leaves, the average length, width and density of the rank 1 adult sheaths.

Rhizome production was assessed from the segments of the rhizome, with inserted roots, between each pair of dead sheaths of minimum thickness (corresponding to the tissue produced during a one-year period) cut and then dried for 48 h at 70 °C, until constant dry weight was achieved. Hereafter, 'rhizomes' always means 'rhizomes and roots' due to the fact that root tissues are very limited. The portions corresponding to the two most recent

146 years were not considered because their growth is not yet complete (Boudouresque *et al.*  
147 1984).

148 The carbon content (% C) of adult blades, sheaths and rhizomes, expressed in dry mass (DM)  
149 percentage, were determined for each sample using elementary analysis (Elementar Vario  
150 MICRO Cube®, Elementar Analysensysteme GmbH, Germany).


151  
152 A 95% confidence interval for each mean was computed. A Shapiro-Wilk test is applied to  
153 check the normality of the means; this is one of the most powerful normality tests (Ghasemi  
154 & Zahediasl, 2012). Student's t-test is used for comparing the means, while confidence  
155 interval (95%) is reported for non-linear regression (Minitab software™). Inter-relationships  
156 between variables were investigated by performing Pearson correlation test. The correlation  
157 coefficient was calculated together with p-values to determine the significance and strength  
158 of each relationship. A significant difference is considered as a p-value<0.05.

159  
160

## 161 Results

162 Meadow density decreases regularly with depth, from 550.5 shoots.m<sup>-2</sup> at -5 m to 106.1  
163 shoots.m<sup>-2</sup> at -30 m (Fig. 2). The number of leaves produced each year also exhibits a  
164 declining trend from shallow water to deep water (from 8.55 to 6.35 leaves.yr<sup>-1</sup>; Fig. 2).


165



166  
167 Figure 2: Meadow density (mean number of shoots.m<sup>-2</sup> and confidence level) and number of leaves  
168 produced annually (mean number per shoot and confidence level) according to depth. Density trend  
169 curve:  $y = -226.1 \times \ln(\text{depth}) + 914.8$  ( $R^2 = 0.91$ ); leaf number trend curve:  $y = -1.045 \times \ln(\text{depth}) +$   
170  $10.093$  ( $R^2 = 0.73$ ).

171  
172 Whatever the depth, the adult blade of rank 1 (external leaf) is on average shorter than the  
173 adult blade of rank 3 (t-test, p-value<0.0001) due to the higher percentage of leaves broken  
174 by herbivore grazing and water movement. The mean density of the blades of rank 1 and  
175 rank 3 is not significantly different ( $3.68 \pm 0.44$  and  $3.85 \pm 0.38$  mg DM.cm<sup>-2</sup>, respectively; t-

176 test, p-value=0.579). Conversely, the density of blades is always significantly smaller than  
 177 sheaths ( $9.11 \pm 1.27 \text{ mg DM.cm}^{-2}$ ; t-test, p-value<0.0001). Blade and sheath density decrease  
 178 significantly with depth (respectively  $R^2 = 0.98$ , p-value<0.0001 and  $R^2 = 0.89$ , p-value<0.005;  
 179 Fig. 3).  
 180  
 181



182  
 183 Figure 3: Blades and sheaths density (mean and confidence level 95%) according to depth. The high  
 184 values of the confidence interval are related to seasonal variations in leaf tissue density.  
 185

186 The mean carbon content in the leaves shows no significant difference between blades  
 187 ( $41.3 \pm 0.4 \%$ ) and sheaths ( $41.1 \pm 0.7 \%$ ) (t-test, p-value=0.663; Table I). However, the mean  
 188 carbon content in rhizomes is higher ( $42.2 \pm 0.7 \%$ ) and exhibits a significant difference with  
 189 blades (t-test, p-value=0.022) and sheaths (t-test, p-value=0.024).  
 190

191 Table I: Carbon content (in %; mean and confidence level 95%) in *P. oceanica* leaves according to  
 192 depth and sampling period (July 2015 to July 2016).

	-5 m		-10 m		-15 m	
	Blades	Sheaths	Blades	Sheaths	Blades	Sheaths
<b>July</b>	41.2 ± 1.9	43.0 ± 2.5	41.9 ± 1.2	43.0 ± 2.2	42.2 ± 0.6	42.6 ± 2.8
<b>October</b>	41.3 ± 1.1	40.2 ± 1.2	42.9 ± 0.3	41.7 ± 2.3	40.4 ± 1.1	41.0 ± 0.3
<b>January</b>	40.7 ± 0.2	37.7 ± 3.1	40.6 ± 0.4	38.2 ± 3.7	40.1 ± 0.2	37.6 ± 2.7
<b>April</b>	40.5 ± 4.2	40.0 ± 3.1	41.7 ± 2.6	40.4 ± 3.9	42.4 ± 4.8	41.6 ± 1.5
	-20 m		-25 m		-30 m	
	Blades	Sheaths	Blades	Sheaths	Blades	Sheaths
<b>July</b>	42.2 ± 0.9	43.2 ± 2.4	41.7 ± 1.1	41.7 ± 3.0	41.5 ± 1.6	39.8 ± 1.6
<b>October</b>	41.3 ± 0.5	41.8 ± 0.4	41.5 ± 0.8	41.3 ± 0.2	40.4 ± 8.0	45.3 ± 7.3
<b>January</b>	40.4 ± 0.2	39.0 ± 1.8	39.4 ± 0.2	38.6 ± 5.8	40.5 ± 0.3	40.2 ± 2.7
<b>April</b>	41.5 ± 1.2	40.0 ± 2.5	41.6 ± 2.7	40.1 ± 5.5	41.0 ± 1.4	40.6 ± 1.3

193

194 Blade and sheath production can be estimated for each depth (Table II). The entire leaf  
 195 production varies between 1 844.3 mg DM.shoot<sup>-1</sup>.yr<sup>-1</sup> at -5 m and 736.1 mg DM.shoot<sup>-1</sup>.yr<sup>-1</sup>  
 196 at -30 m (752.2 to 301.4 mg C.shoot<sup>-1</sup>.yr<sup>-1</sup>). Rhizome production varies between 59.2 and  
 197 34.8 mg DM.shoot<sup>-1</sup>.yr<sup>-1</sup> (24.9 to 14.7 mg C.shoot<sup>-1</sup>.year<sup>-1</sup>). Total primary production (blades,  
 198 sheaths and rhizomes) decreases with depth. The mean production, in carbon per shoot,  
 199 corresponds to 79 % for blades, 17 % for sheaths and 4 % for rhizomes.


200  
 201

Table II: Leaf primary production according to depth at the studied site.

Depth (m)	Mean leaf number.yr <sup>-1</sup>	Mean blade mass (mg DM)	Blade production (mg DM.shoot <sup>-1</sup> )	Carbon (%)	Blade production (mg C.shoot <sup>-1</sup> )
5	8.6	177.7	1 519.3	40.9	621.5
10	7.5	139.5	1 046.3	41.7	436.8
15	6.6	164.7	1 093.6	41.3	451.4
20	7.0	121.6	851.2	41.4	352.0
25	7.3	124.1	909.7	41.0	373.2
30	6.4	93.8	595.6	40.8	243.1
Depth (m)	Mean leaf number.yr <sup>-1</sup>	Mean sheath mass (mg DM)	Sheath production (mg DM.shoot <sup>-1</sup> )	Carbon (%)	Sheath production (mg C.shoot <sup>-1</sup> )
5	8.6	38.0	324.9	40.2	130.7
10	7.5	33.1	248.6	40.8	101.5
15	6.6	32.7	217.3	40.7	88.4
20	7.0	25.2	176.3	41.0	72.2
25	7.3	26.1	191.6	40.4	77.4
30	6.4	22.1	140.4	41.5	58.3

202  
 203  
 204  
 205

The mean total carbon fixation (blades, sheaths and rhizomes) varies between 426.6 and 33.5 g C.m<sup>-2</sup> with a clear decreasing trend according to depth (Fig. 4).




206



207 Figure 4: Annual mean carbon fixation (dotted line) and sequestration (solid line) by *P. oceanica*  
 208 meadow ( $\text{g C}\cdot\text{m}^{-2}$ ) according to depth; confidence interval (95%) is also reported. Fixation trend  
 209 curve:  $y = -202.5 \times \ln(\text{depth}) + 724.6$  ( $R^2 = 0.95$ ); sequestration trend curve:  $y = -40.5 \times \ln(\text{depth}) +$   
 210  $145.5$  ( $R^2 = 0.98$ ).  
 211

212 The mean carbon sequestration (long-term sink in the matte), corresponding to the dead  
 213 sheath and rhizome tissues, varies between  $84.4$  and  $7.7 \text{ g C}\cdot\text{m}^{-2}$ .  
 214

215 The integration of the carbon fixation and sequestration values over the entire bathymetric  
 216 range of the *P. oceanica* meadow at the study site makes it possible to estimate the total  
 217 fixation rate at  $22\,068 \text{ t C}\cdot\text{yr}^{-1}$ , *i.e.* an average value of  $108.0 \text{ g C}\cdot\text{m}^{-2}\cdot\text{yr}^{-1}$ , and the  
 218 sequestration rate at  $4\,564.7 \text{ t C}\cdot\text{yr}^{-1}$ , *i.e.* an average value of  $22.3 \text{ g C}\cdot\text{m}^{-2}\cdot\text{yr}^{-1}$ , corresponding  
 219 to an average depth of  $21 \text{ m}$  over the entire site (Fig. 5).  
 220



221  
 222 Figure 5: Annual carbon fixation (A) and sequestration (B) by *P. oceanica* meadow ( $\text{g C}\cdot\text{m}^{-2}$ ) according  
 223 to depth.  
 224  
 225

226 **Discussion**

227 Carbon fixation and sequestration by the *P. oceanica* meadow, measured in sector 2 (Fig. 1),  
 228 corresponds to the sector with the largest extension of this habitat (more than 38% of the  
 229 areas covered by *P. oceanica* along the Corsican coast; Valette *et al.*, 2019). Moreover,  
 230 recorded values are similar to those measured sporadically in the Bay of Calvi, on the  
 231 western coast of Corsica (Bay, 1984; Pergent-Martini *et al.*, 1994; Vela *et al.*, 2006).

232  
 233 Based on the values measured in sector 2, it is therefore possible to extrapolate the carbon  
 234 fixation and sequestration, by *P. oceanica* meadows, for the entire coast of Corsica (Fig. 4).  
 235 The areas covered in each sector as well as their average bathymetric extension provide a  
 236 basis for determining the carbon fixation and sequestration rate (Table III). Total fixation is  
 237 estimated at 69 275 t C.yr<sup>-1</sup>, an average of 1.29 t C.ha<sup>-1</sup>.yr<sup>-1</sup>, while sequestration rate in the  
 238 mat (dead sheaths and rhizomes) is estimated at 14 250 t C.yr<sup>-1</sup>, an average of 0.27 t C.ha<sup>-1</sup>.yr<sup>-1</sup>  
 239 (Table III).


240  
 241 Table III: *P. oceanica* meadow surface area, mean depth (from Valette *et al.*, 2019), carbon fixation (-  
 242 202.5 x ln(depth) + 724.56 and carbon sequestration (-40.46 x ln(depth) + 145.53) in the different  
 243 sectors. In grey, sector 2, corresponding to the studied site, \*= updated in this study; C = carbon.

	<i>P. oceanica</i> surface area (ha)	Mean depth (m)	Mean C fixation (t C.ha <sup>-1</sup> .yr <sup>-1</sup> )	Carbon fixation (t C.yr <sup>-1</sup> )	Mean C sequestration (t C.ha <sup>-1</sup> .yr <sup>-1</sup> )	Carbon sequestration (t C.yr <sup>-1</sup> )
Sector 1	4 137	20	1.21	5 005	0.25	1 032
Sector 2	20 425	21*	1.08	22 068	0.22	4 565
Sector 3	3 078	19	1.34	4 116	0.27	846
Sector 4	6 179	16	1.68	10 395	0.34	2 124
Sector 5	3 534	18	1.42	5 001	0.29	1 026
Sector 6	1 430	16	1.70	2 424	0.35	495
Sector 7	2 765	18	1.38	3 804	0.28	781
Sector 8	2 981	18	1.40	4 168	0.29	856
Sector 9	763	20	1.21	927	0.25	191
Sector 10	719	18	1.44	1 038	0.30	213
Sector 11	1 505	18	1.35	2 029	0.28	417
Sector 12	2 940	20	1.21	3 572	0.25	736
Sector 13	1 784	17	1.55	2 766	0.32	566
Sector 14	1 497	19	1.31	1 961	0.27	403

244  
 245 Although the *P. oceanica* meadow in sector 2 covers a very large area, it only allows an  
 246 average fixation rate of 1.08 t C.ha<sup>-1</sup>.yr<sup>-1</sup> and a sequestration rate of 0.22 t C.ha<sup>-1</sup>.yr<sup>-1</sup>, the  
 247 lowest values recorded along the coast of Corsica, because the average bathymetric range of  
 248 this meadow is the deepest one (-21 m; Table III). This is explained by its deep upper limit  
 249 due to significant hydrodynamic forces (exposed coast) and of areas unsuitable for *P.*  
 250 *oceanica*, near the mouth of coastal rivers (Fig. 1; Pasqualini *et al.*, 1998; Boudouresque *et*  
 251 *al.*, 2012). Conversely, for the sectors where the average bathymetric extension of the  
 252 meadow is shallower, the carbon fixation rate reaches 1.70 t C.ha<sup>-1</sup>.yr<sup>-1</sup> and the  
 253 sequestration rate 0.35 t C.ha<sup>-1</sup>.yr<sup>-1</sup> (sector 4 = East coast of the Natural Reserve of the Strait  
 254 of Bonifacio; sector 6 = Gulf of Valinco).


255

256 The primary production of the *P. oceanica* meadow has been the subject of numerous  
 257 studies, but they mainly concern the North-Western Mediterranean Sea (Bay, 1984; Buia *et al.*,  
 258 1992; Pergent-Martini *et al.*, 1994; Cebrian *et al.*, 1997; Torricelli & Peirano, 1997;  
 259 Guidetti & Fabiano 2000; Romero, 2004; Vela *et al.*, 2006), even if some studies are available  
 260 for other areas of the Mediterranean Sea; Algeria (Boumaza & Semroud, 2000), Croatia  
 261 (Bakran-Petricioli & Schultz, 2010), Sicily (La Loggia *et al.*, 2004), Turkey (Pergent-Martini *et al.*,  
 262 1994). These data confirm that, for a continuous meadow, primary production (mg  
 263 DM.shoot<sup>-1</sup>) decreases with depth (Fig. 6). However, for a given depth, the values can be  
 264 highly variable from one site to another (*e.g.* between 310 and 2 250 g DM.shoot<sup>-1</sup> at -2 m;  
 265 Boumaza & Semroud, 2000, Vela *et al.*, 2006).  
 266



267  
 268 Figure 6: Annual primary production of *P. oceanica* (mg DM.shoot<sup>-1</sup>) according to depth and site, R<sup>2</sup> =  
 269 determination coefficient is reported (from Pergent-Martini *et al.*, 1994; Bakran-Petricioli & Schultz,  
 270 2010 and this study).  
 271

272 The synthesis of a hundred measurements made throughout the Mediterranean Sea and at  
 273 depths between 0.5 and 32.0 m provides a basis for estimation of the average primary leaf  
 274 production (in g DM.m<sup>-2</sup>) according to the depth (Fig. 7). For a depth of 15 m, which  
 275 corresponds to the average depth of *P. oceanica* meadows in the Mediterranean Sea (mean  
 276 extension between -2 and -28 m, in UNEP/MAP-RAC/SPA, 2015), annual primary leaf  
 277 production (blades and sheaths) is estimated at 343 g DM.m<sup>-2</sup>.  
 278



279  
 280 Figure 7: Leaf annual primary production of *P. oceanica* meadow (g DM.m<sup>-2</sup>) according to depth in  
 281 the Mediterranean Sea (data from: Wittmann & Ott, 1982; Bedhomme *et al.*, 1983; Bay, 1984; Thelin  
 282 & Giorgi, 1984; Wittmann, 1984; Esteban 1989; Romero, 1989; Esteban *et al.*, 1990; Buia *et al.*, 1992;  
 283 Pergent-Martini *et al.*, 1994; Pergent *et al.*, 1997; Torricelli & Peirano, 1997; Zupo *et al.*, 1997;  
 284 Boumaza & Semroud, 2000; Guidetti, 2000; Guidetti *et al.*, 2000; Guidetti & Fabiano, 2000; La Loggia  
 285 *et al.*, 2004; Vela, 2006; Vela *et al.*, 2006; Bakran-Petricioli & Schultz, 2010; this study). Trend curve:  $y$   
 286  $= -227.2 \times \ln(\text{depth}) + 958.4$  ( $R^2 = 0.43$ ).  
 287

288 Carbon concentrations in the leaf tissue of *P. oceanica* show great variability depending on  
 289 the site (Table IV); thus, they are on average higher in Corsica (38%, Gobert *et al.*, 1995;  
 290 41.2%, this study), low at Alicante (27.9%, Romero *et al.*, 1998) and intermediate at Ischia  
 291 (30.6 %, Romero *et al.*, 1992; 31.3 %, Scartazza *et al.*, 2017; 34.9 %, Garrard, 2013).  
 292 However, the carbon concentration in the tissues does not seem to be significantly impacted  
 293 by the depth or the season (Garrard, 2013; Table I) although Alcovero *et al.* (2000) highlights  
 294 a slight variation between winter and spring, probably related to the age of the leaf tissue.  
 295 The average of the values available in the Mediterranean Sea, for all sites and depths  
 296 combined, is estimated at  $37.8 \pm 1.6\%$  for leaf tissue and  $42.0 \pm 1.1\%$  for rhizomes (Table IV).  
 297 These values are lower than those measured for terrestrial Magnoliophyta: 47.5% as a mean  
 298 (Dupouey *et al.*, 2000) and 44% in the case of mangrove trees (Bouillon *et al.*, 2008).  
 299

300 Table IV: Carbon concentration (%) in leaf tissue and in the rhizomes of *P. oceanica* in the  
 301 Mediterranean Sea.

Country / Location	Depth (m)	% C in leaves	% C in rhizome	References
Spain / Catalonia	5.0 – 13.0	34.1 - 35.3	35.6	Romero, 1989
Italy / Ischia	20.0	30.6	35.7	Romero <i>et al.</i> , 1992
Spain	0.5 - 15	38.1		Enriquez <i>et al.</i> , 1995
France / Corsica	10.0	30.5 - 45.5		Gobert <i>et al.</i> , 1995
Spain / Alicante	5.0 – 20.0	26.9 - 29.0	38.1 - 38.6	Romero <i>et al.</i> , 1998
Spain / Catalonia	5.0	30.5 - 35.0	39.0	Alcovero <i>et al.</i> , 2000
Spain / Murcia	7.5	40.0		Ruiz, 2000

France / Corsica	10.0	32.1		Gobert <i>et al.</i> , 2005
France / Corsica	11.0	35.6		Leoni <i>et al.</i> , 2007
France / Provence	2.0	38.2		Vela, 2006
France / Corsica	2.0	38.7		Vela, 2006
Italy / Sardinia	2.0	38.0		Vela, 2006
Tunisia / Cap Bon	2.0	37.7		Vela, 2006
Tunisia / South	2.0	37.4		Vela, 2006
Italy / Ischia	3.0	33.5 - 35.8		Garrard, 2013
Italy / Ischia	3.0	31.3	38.3	Scartazza <i>et al.</i> , 2017
France / Corsica	5.0 - 30.0	37.6 - 43.2	38.0 - 45.8	This study
France / Corsica	8.0 - 37.0	35.1 - 42.4		Gobert unpublished

302

303 The average annual carbon fixation and sequestration rate throughout the Mediterranean  
304 basin can be estimated from (i) the annual primary leaf production (blades and sheaths) for  
305 the integrating depth of -15 m (343 g DM.m<sup>-2</sup>), (ii) the average proportion of the different  
306 tissues produced estimated in this study (blades 79%, sheaths 17%, rhizomes 4%), and (iii)  
307 the average carbon concentration of these tissues (37.8 ± 1.6% for leaf tissue and 42.0 ±  
308 1.1% for rhizomes). Thus, the fixation rate of the blades is estimated at 102.4 g C.m<sup>-2</sup>.yr<sup>-1</sup>,  
309 that of the sheaths at 22.0 g C m<sup>-2</sup>.yr<sup>-1</sup> and that of the rhizomes at 5.8 g C m<sup>-2</sup>.yr<sup>-1</sup>; *i.e.* a total  
310 fixation rate of 130.2 g C m<sup>-2</sup>.yr<sup>-1</sup> and sequestration rate (dead sheaths and rhizomes) of  
311 27.8 g C m<sup>-2</sup>.yr<sup>-1</sup>.

312

313 Based on the available meadow areas for each country, to date (UNEP-MAP-SPA/RAC, 2019),  
314 it is possible to make a first evaluation of the amount of carbon fixed and sequestered by the  
315 *P. oceanica* meadow each year (Table V). The total amount of carbon fixed is estimated at 3  
316 638 909 tons per year and the amount sequestered at 776 971 tons per year, equivalent  
317 respectively to 13 343 878 and 2 849 154 tons per year of CO<sub>2</sub> emission/release equivalent.

318

319 Table V: Annual carbon fixation and sequestration by *P. oceanica* meadows in different  
320 Mediterranean countries. \* UNEP-MAP-SPA/RAC, 2019 ; \*\* Telesca *et al.*, 2015 ; \*\*\* Duman *et al.*,  
321 2019 ; \*\*\*\* Pergent-Martini *et al.*, 2013.

	Available <i>P. oceanica</i> surface area (ha)*	Surface mapped (%)**	Estimated <i>P. oceanica</i> surface (ha)	Estimated C fixation (t C.yr <sup>-1</sup> )	Estimated sequestration (t C.yr <sup>-1</sup> )
Spain	115 904	100 %	115 904	150 907	32 221
France	87 680	100 %	87 680	114 159	24 379
Monaco	14	100 %	14	18	4
Italy	337 611	100 %	337 611	439 570	93 856
Slovenia	1	100 %	1	1	0
Croatia	31 437	14 %	224 550	292 364	62 425
Bosnia Herzegovina	Not present	100 %*	0		
Montenegro	51	10 %*	512	667	142
Albania	4 803	100 %	4 803	6 254	1 335
Malta	5 860	100 %	5 860	7 630	1 629
Greece	251 010	100 %*	251 010	326 815	69 781
Turkey	39 983	58 %***	68 937	89 756	19 164
Cyprus	3 627	23 %****	15 770	20 533	4 384

Syria, Lebanon, Palestinian territories, Israel	Not present				
Egypt	No data		0		
Libya	1 235	11 %	11 227	14 618	3 121
Tunisia	1 332 815	81 %	1 645 451	2 142 377	457 435
Algeria	4 085	16 %	25 531	33 241	7 098
Morocco	Not present	100 %	0		
<b>Total</b>	<b>2 216 116</b>		<b>2 794 861</b>	<b>3 638 909</b>	<b>776 974</b>

322

323

324

325

326

327

328

329

330

331

332

333

334

Fixation and sequestration of carbon in *P. oceanica* meadows can present a significant part of the CO<sub>2</sub> emissions/releases of the various Mediterranean countries (Table VI). Indeed, even if this fixation represents only 0.61% on average for all Mediterranean countries, it must be borne in mind that (i) seven countries, some of which have a large population (*e.g.* Egypt, Morocco), have not identified any *P. oceanica* meadow, and (ii) that the length of the Mediterranean coast is sometimes very limited compared to the surface area of the country (*e.g.* Slovenia, Turkey, Morocco). However, this fixation may be high for certain countries (*e.g.* 24.6% for Tunisia, 5.6% for Croatia).

Table VI: Carbon fixation in the different Mediterranean countries. Meadow surface area is estimated from the UNEP-MAP-SPA/RAC (2019), 2018 population and national CO<sub>2</sub> emissions from the site <http://www.globalcarbonatlas.org/en/CO2-emissions>.

	Estimated C fixation (t C)	Estimated CO <sub>2</sub> release* (Mt yr <sup>-1</sup> )	Population (inhabitant)	Estimated CO <sub>2</sub> release (t CO <sub>2</sub> .inhab <sup>-1</sup> )	CO <sub>2</sub> fixation by <i>P. oceanica</i> (%)
Spain	150 907	268.0	46 692 858	5.74	0.21
France and Monaco	114 178	338.0	64 990 511	5.20	0.12
Italy	439 570	338.0	60 627 291	5.58	0.48
Slovenia	1.3	14.0	2 077 837	6.74	0.00
Croatia	292 364	19.0	4 156 405	4.57	5.64
Bosnia Herzegovina	0	22.0	3 323 925	6.62	0.00
Montenegro	667	2.0	627 809	3.19	0.12
Albania	6 254	4.6	2 882 740	1.60	0.50
Malte	7 630	1.6	439 248	3.64	1.75
Greece	326 815	74.0	10 522 246	7.03	1.62
Turkey	89 756	428.0	82 340 088	5.20	0.08
Cyprus	20 533	7.5	1 189 265	6.31	1.00
Syria	0	28.0	16 945 057	1.65	0.00
Lebanon	0	24.0	6 859 408	3.50	0.00
Ter. Palestinians	0	3.2	4 862 979	0.66	0.00
Israel	0	64.0	8 381 516	7.64	0.00
Egypt	0	239.0	98 423 598	2.43	0.00
Libya	14 618	54.0	6 678 559	8.09	0.10
Tunisia	2 142 377	32.0	11 565 201	2.77	24.55
Algeria	33 241	156.0	42 228 408	3.69	0.08
Morocco		66.0	36 029 093	1.83	0.00
<b>Total (mean)</b>	<b>3 638 909</b>	<b>2 182.9</b>	<b>511 844 042</b>	<b>4.26</b>	<b>0.61</b>

335  
 336  
 337  
 338  
 339  
 340  
 341  
 342  
 343  
 344  
 345

The large Mediterranean islands deserve special attention because the carbon fixation by *P. oceanica* is higher given the lower population density and the large areas of seagrass beds they shelter (Table VII). The annual carbon fixation is on average 3.1% of CO<sub>2</sub> emissions/releases but can reach almost 14.4% for Corsica. Moreover, the percentage of carbon fixation in comparison with carbon release/emission in the large Mediterranean islands appears higher in the Western basin.

Table VII: Carbon fixation in the large Mediterranean islands. Meadow surfaces are estimated from the report of the UNEP-MAP-SPA/RAC (2019) and from Valette *et al.* (2019); national CO<sub>2</sub> emissions from the site <http://www.globalcarbonatlas.org/en/CO2-emissions>; \* : This study.

	Surface <i>P. oceanica</i> (ha)	Carbon fixation (t C.yr <sup>-1</sup> )	Release (t CO <sub>2</sub> .yr <sup>-1</sup> )	Fixation by <i>P. oceanica</i> (%)
Balearic Islands	63 316	82 437	6 818 687	4.4
Corsica	53 736	69 275	1 763 983	14.4
Sardinia	153 382	199 703	9 143 077	8.0
Sicily	76 000	98 952	27 875 235	1.3
Cyprus	15 770	20 533	7 500 000	1.0
Crete	14 500	18 879	4 465 284	1.6
<b>Total</b>	<b>396 927</b>	<b>516 799</b>	<b>59 166 267</b>	<b>3.1</b>

346  
 347  
 348  
 349  
 350  
 351  
 352  
 353  
 354

Inter-relationships between variables were investigated by performing a Pearson correlation test (Table VIII); variables correspond to ‘**Po cover 0-40**’ (% *P. oceanica* surface covered/total surface available between 0 and -40 m in Valette *et al.*, 2019), ‘**C Fixation**’ (total island carbon fixation by *P. oceanica* in t C.yr<sup>-1</sup>), ‘**C Sequestration**’ (total island carbon sequestration by *P. oceanica* in t C.yr<sup>-1</sup>), ‘**% CO<sub>2</sub> Fixed**’ (% CO<sub>2</sub> fixed by *P. oceanica*/CO<sub>2</sub> released in t CO<sub>2</sub>.yr<sup>-1</sup>) and ‘**CO<sub>2</sub> Release Inhab**’ (total island CO<sub>2</sub> released per inhabitant in t CO<sub>2</sub>.yr<sup>-1</sup>.inhab<sup>-1</sup> according to <http://www.globalcarbonatlas.org/en/CO2-emissions>).

355  
 356  
 357  
 358  
 359  
 360  
 361  
 362  
 363  
 364

The areas covered by the *P. oceanica* meadows between 0 and -40 m are significantly greater for the islands located to the north ( $r = 0.941$ ;  $p$ -value<0.01) and to the west ( $r = 0.904$ ;  $p$ -value<0.05) of the Mediterranean Sea (*i.e.* Sardinia, Corsica, Balearic Islands). Conversely, CO<sub>2</sub> emissions per capita/inhabitant are negatively correlated with latitude ( $r = 0.863$ ;  $p$ -value<0.05) reflecting higher CO<sub>2</sub> emissions for the inhabitants of the islands in the south of the Mediterranean basin (*i.e.* Cyprus, Crete). Thus, the northwestern Mediterranean islands, characterized by lower CO<sub>2</sub> emissions per capita/inhabitant and greater coverage of *P. oceanica* meadows within the bathymetric range from 0 to -40 m, appear as sectors where the ratio CO<sub>2</sub> fixed / CO<sub>2</sub> released is the highest ( $r = 0.880$ ;  $p$ -value<0.05)

365  
 366

Finally, the carbon sequestered by *P. oceanica* meadows in the matte is significantly correlated with the fixation of carbon produced by the plant ( $r = 0.979$ ;  $p$ -value<0.001).

367  
 368  
 369  
 370

Table VIII: Pearson’s correlation matrix between geographical, environmental and carbon variables. Level of significance: \* $p$ ≤0.05, \*\* $p$ ≤0.01, \*\*\* $p$ ≤0.001; NS,  $p$ ≥0.05; Significant correlations in bold (r value).

Variables	Longitude	Latitude	Po cover 0-40	C Fixation	C Sequestration	% CO <sub>2</sub> Fixed	CO <sub>2</sub> Release Inhab
Longitude		<b>-0.854</b>	<b>-0.904</b>	-0.574	-0.621	-0.529	0.720

Latitude	*		<b>0.941</b>	0.544	0.547	<b>0.880</b>	<b>-0.863</b>
Po cover 0-40	*	**		0.473	0.511	0.794	-0.681
C Fixation	NS	NS	NS		<b>0.979</b>	0.316	-0.594
C Sequestration	NS	NS	NS	***		0.312	-0.523
% CO <sub>2</sub> Fixed	NS	*	NS	NS	NS		-0.673
CO <sub>2</sub> Release Inhab	NS	*	NS	NS	NS	NS	

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

Carbon fixation by *P. oceanica* is high with an average of 1 302 t C ha<sup>-1</sup>.yr<sup>-1</sup>. Even if the areas covered by *P. oceanica* meadows are smaller than those of other terrestrial ecosystems, this ecosystem represents around 4% of the areas covered by forest ecosystems in all Mediterranean countries (FAO & Plan Bleu, 2018) and more than 20% for the six large Mediterranean islands (FAO & Plan Bleu, 2018).

Moreover, the major advantage of the *P. oceanica* meadow lies in its capacity to store carbon in soil from centuries to millennia. The average carbon stock found in the mat, resulting from the accumulation of 21% of annual primary production, varies between 40 and 237 kg C.m<sup>-2</sup> (Romero *et al.* 1994, Mateo *et al.* 1997, Serrano *et al.*, 2012; 2014). These values are similar or higher than those measured in several terrestrial ecosystems considered to be efficient in carbon storage such as peatlands (120 kg C.m<sup>-2</sup>; Warner *et al.*, 1993), wetlands (13-73 kg C.m<sup>-2</sup>; Laffoley & Grimsditch 2009) and the boreal forests (9-34 kg C.m<sup>-2</sup>; Serrano *et al.* 2014) and Mediterranean forest (5.7 kg C.m<sup>-2</sup>; FAO & Plan Bleu, 2018).

### Acknowledgments

389

390

391

392

393

394

395

This work would not have been possible without the financial support of the Agence Française pour la Biodiversité, the Collectivité de Corse (PADDUC-CHANGE programme), the Uffiziu di Ambiente di a Corsica. A special thanks to Maria Cristina Buia (Stazione Zoologica Anton Dohrn, Ischia - Italy) and Sylvie Gobert (University of Liège) for providing raw data and Michael Paul for proof-reading the manuscript.

396

### References

397

398

399

400

401

402

403

404

405

406

407

408

409

410

- Akçalı B., Kaboğlu G., Güçlüsoy H., 2019. A review on *Posidonia oceanica* (Linnaeus) Delile coverage along the Turkish coasts until 2019. *J. Black Sea/Mediterranean Environment*, 25(1): 115-124.
- Alcoverro T., Manzanera M., Romero J., 2000. Nutrient mass balance of the seagrass *Posidonia oceanica*: the importance of nutrient retranslocation. *Marine Ecology Progress Serie*, 194: 13-21.
- Bakran-Petricioli T., Schultz S., 2010. Biometry and leaf lepidochronology of the seagrass *Posidonia oceanica* in the Croatian Adriatic. *Proceedings of the 4<sup>th</sup> Mediterranean Symposium on marine vegetation* (Yasmine-Hammamet, Tunisia), 2-4 December 2010, El Asmi S., Langar H., Belkacem W (eds.), RAC/SPA publ., Tunis: 28-31.
- Bay D., 1984. A field study of the growth dynamics and productivity of *Posidonia oceanica* (L.) Delile in Calvi Bay, Corsica. *Aquatic Botany*, 20: 43-64.
- Bédhomme A.L., Thelin I., Boudouresque C.F., 1983. Mesure de la production primaire des feuilles de *Posidonia oceanica*: modification de la méthode de Zieman. *Botanica*



411 *marina*, 26 : 35-43.

412 Boudouresque C.F., Jeudy de Grissac A., Meinesz A., 1984. Relation entre la sédimentation et  
413 l'allongement des rhizomes orthotropes de *Posidonia oceanica* dans la baie d'Elbu  
414 (Corse). International Workshop on *Posidonia oceanica* Beds. C.F. Boudouresque. A.  
415 Jeudy de Grissac and J. Olivier Eds. GIS Posidonie publ., Marseille: 185-191.

416 Boudouresque C.F., Bernard G., Bonhomme P., Charbonnel E., Diviacco G., Meinesz A.,  
417 Pergent G., Pergent-Martini C., Ruitton S., Tunesi L., 2012. Protection and  
418 conservation of *Posidonia oceanica* meadows. RAMOGE and RAC/SPA publ., Tunis.  
419 202pp.

420 Boudouresque C.F., Pergent G., Pergent-Martini C., Ruitton S., Thibaut T., Verlaque V., 2016.  
421 The necromass of the *Posidonia oceanica* seagrass meadow: Fate, role, ecosystem  
422 services and vulnerability. *Hydrobiologia*, 781: 25-42.

423 Bouillon S., Borges A.V., Castañeda-Moya E., Diele K., Dittmar T., Duke N.C., Kristensen E.,  
424 Lee S.Y., Marchand C., Middelburg J.J., Rivera-Monroy V.H., Smith III T.J., Twilley R.R.,  
425 2008. Mangrove production and carbon sinks: A revision of global budget estimates.  
426 *Global Biogeochemical Cycles*, 22: 1-12.

427 Boumaza S., Semroud S., 2000. Evaluation of past and present primary production in  
428 *Posidonia oceanica* beds from the cove of Kouali (Tipasa, Algeria) using the  
429 lepidochronological method. *Biologia marina mediterranea*, 7 (2): 26-30.

430 Buia M.C., Zupo V., Mazzella L., 1992. Primary production and growth dynamics in *Posidonia*  
431 *oceanica*. *PSZN Marine Ecology*, 13 (1): 2-16.

432 Cebrián J., Duarte C.M., 2001. Detrital stocks and dynamics of the seagrass *Posidonia*  
433 *oceanica* (L.) Delile in the Spanish Mediterranean. *Aquatic Botany*, 70: 295-309.

434 Cebrián J., Duarte C.M., Marbà N., Enríquez S., 1997. Magnitude and fate of the production  
435 of four co-occurring Western Mediterranean seagrass species. *Marine Ecology*  
436 *Progress Serie*, 155: 29-44.

437 Chmura G.L., Anisfeld S.C., Cahoon D.R., Lynch J.C., 2003. Global carbon sequestration in  
438 tidal, saline wetland soils. *Global Biogeochemical Cycles*, 17 (4).

439 Duarte C.M., Chiscano C.L., 1999. Seagrass biomass and production: a reassessment. *Aquatic*  
440 *Botany*, 65: 159–174.

441 Duarte C.M., Middelburg J.J., Caraco N., 2005. Major role of marine vegetation on the  
442 oceanic carbon cycle. *Biogeosciences*, 2: 1-8.

443 Duman M., Eronat A.H., Ilhan T., Talas E., Küçüksezgin F., 2019. Mapping *Posidonia oceanica*  
444 (Linnaeus) Meadows in the Eastern Aegean Sea Coastal Areas of Turkey: Evaluation of  
445 Habitat Maps Produced Using the Acoustic Ground-Discrimination Systems.  
446 *International Journal of Environment and Geoinformatics (IJEgeo)*, 6(1): 67-75. DOI:  
447 10.30897/ijegeo.

448 Dupouey J.L., Pignard G., Badeau V., Thimonier A., Dhote J.F., Nepveu G., Berges L., Augusto  
449 L., Belkacem S., Nys C., 2000. Carbon stocks and flux in French forests; Stocks et flux  
450 de carbone dans les forets francaises. *Revue Forestiere Francaise* : 139-154.

451 Enríquez S., Duarte C.M., Sand-Jensen K., 1995. Patterns in the photosynthetic metabolism of  
452 Mediterranean macrophytes. *Marine Ecology Progress Serie*, 199: 243-252.

453 Esteban J.L., 1989. Dinamica, ciclo de hojas y produccion foliar en praderas de *Posidonia*  
454 *oceanica* del litoral de la comunidad valenciana (Mediterraneo occidental). Tesis de  
455 Licenciatura, Universitat de Valencia: 1-142.

456 Esteban, J.L., Giner, I.M., Benedito, V., Torres, J., Capaccioni, R., Garcia-Carrascosa, A.M.,  
457 1990. Leaf biomass and production of *Posidonia oceanica* at Spanish Eastern Coast.

458 Rapp. P-V Reun. Comm. Int. Explor. Sci. Mer. Mediterr. Monaco, 32: 8.

459 FAO and Plan Bleu, 2018. State of Mediterranean Forests 2018. Food and Agriculture  
460 Organization of the United Nations, Rome and Plan Bleu, Marseille: 331pp.

461 FCCC, 2015. Conference of the Parties: Twenty-first session. United Nations - Framework  
462 Convention on Climate Change. Paris, 30 November to 11 December 2015, 32pp.

463 Fourqurean J.W., Duarte C.M., Kennedy H., Marbà N., Holmer M., Mateo M.A., Apostolaki  
464 E.T., Kendrick G.A., Krause-Jensen D., Mcglathery K.J., Serrano O., 2012. Seagrass  
465 ecosystems as a globally significant carbon stock. *Nature Geoscience*, 5: 505-509.

466 Garrard S.L., 2013. The effect of ocean acidification on plant animal interactions in a  
467 *Posidonia oceanica* meadow. Thesis of the Open University Milton Keynes (UK) and  
468 Stazione Zoologica Anton Dohrn: 1-318.

469 Ghasemi A., Zahediasl S., 2012. Normality Tests for Statistical Analysis: A Guide for Non-  
470 Statisticians. *International Journal Endocrinology Metabolism*, 10 (2): 486–489.

471 Giakoumi S., Sini M., Gerovasileiou V., Mazor T., Beher J., Possingham H.P., Abdulla A., Cinar  
472 M.E., Dendrinou P., Gucu A.C., Karamanlidis A.A., Rodic P., Panayotidis P., Taskin E.,  
473 Jaklin A., Voultziadou E., Webster C., Zenetos A., Katsanevakis S., 2013. Ecoregion-  
474 Based Conservation Planning in the Mediterranean: Dealing with Large-Scale  
475 Heterogeneity. *PLoS ONE*, 8 (10) : e76449. doi:10.1371/journal.pone.0076449.

476 Giraud G., 1979. Sur une méthode de mesure et de comptage des structures foliaires de  
477 *Posidonia oceanica* (Linnaeus) Delile. *Bulletin Museum Histoire Naturelle Marseille*,  
478 39 : 33-39.

479 Global Carbon Atlas. <http://www.globalcarbonatlas.org/en/CO2-emissions>. Accessed 28 May  
480 2020.

481 Gobert S., Belkhiria S., Dauby P., Havelange S., Soullard M., Bouquegneau J.M., 1995.  
482 Variations temporelles de la phénologie et de la composition biochimique de la  
483 phanérogame marine *Posidonia oceanica* en baie de Calvi. *Bulletin Société Royale  
484 Sciences Liège*, 64 (4-5) : 263-284.

485 Gobert S., Lejeune P., Lepoint G., Bouquegneau J.M., 2005. C, N, P concentrations and requi-  
486 rements of flowering *Posidonia oceanica* shoots. *Hydrobiologia*, 533: 253-259.

487 Guidetti P., 2000. Leaf primary production in *Posidonia oceanica*: two reconstructive aging  
488 techniques give similar results. *Aquatic Botany*, 68: 337–343.

489 Guidetti P., Buia M.C., Mazzella L., 2000. The Use of Lepidochronology as a Tool of Analysis  
490 of Dynamic Features in the Seagrass *Posidonia oceanica* of the Adriatic Sea. *Botanica  
491 Marina*, 43: 1-9.

492 Guidetti P., Fabiano M., 2000. The use of lepidochronology to assess the impact of  
493 terrigenous discharges on the primary leaf production of the Mediterranean seagrass  
494 *Posidonia oceanica*. *Marine Pollution Bulletin*, 40 (5): 449-453.

495 Guiry M.D., Guiry G.M., 2020. AlgaeBase. World-wide electronic publication. National  
496 University of Ireland. Galway. <https://www.algaebase.org>; searched on 09 April 2020.

497 IPS, 2008. Peatlands and Climate Change. Maria Strack, University of Calgary, Canada (Edit);  
498 International Peat Society (Pub.): 227pp.

499 Jamaludin M.R., 2015. Carbon storage and preservation in seagrass meadows. Thesis in  
500 Environmental Science, Edith Cowan University – Western Australia: 188pp.

501 Kennedy H., Björk M., 2009. Seagrass Meadows. In: Laffoley D., Grimsditch G. (Eds), The  
502 Management of Coastal Carbon Sinks. IUCN, Gland, Switzerland: pp.23-30.

503 La Loggia G., Calvo S., Ciruolo G., Mazzola A., Pirrotta M., Sara G., Tomasello A., Vizzini S.,  
504 2004. Influence of hydrodynamic conditions on the production and fate of *Posidonia*

505 *oceanica* in a semi-enclosed shallow basin (Stagnone di Marsala, Western Sicily).  
506 *Chemistry Ecology*, 20 (3): 183-201.

507 Laffoley D, Grimsditch G 2009. The Management of Natural Coastal Carbon Sinks. IUCN,  
508 Gland, Switzerland: 53pp.

509 Leoni V., Pasqualini V., Pergent-Martini C., Vela A., Pergent G., 2007. Physiological responses  
510 of *Posidonia oceanica* to experimental nutrient enrichment of the canopy water.  
511 *Journal of Experimental Marine Biology Ecology*, 349 (1): 73-83.

512 Lo Iacono C., Mateo M.A., Gracia E., Guasch L., Carbonell R., Serrano L., Serrano O.,  
513 Danõbeitia J., 2008. Very high-resolution seismo-acoustic imaging of seagrass  
514 meadows (Mediterranean Sea): Implications for carbon sink estimates. *Geophysical*  
515 *Research Letters*, 35 (18): 1-5.

516 Mateo M.A., Romero J., Pérez M., Littler M.M., Littler D.S., 1997. Dynamics of millenary  
517 organic deposits resulting from the growth of the Mediterranean seagrass *Posidonia*  
518 *oceanica*. *Estuarine, Coastal Shelf Science*, 44: 103-110.

519 Mateo M.A., Cebrián J., Dunton K., Mutchler T., 2006. Carbon flux in seagrass ecosystems. In  
520 *Seagrass: Biology, Ecology and Conservation*, Larkum AWD, Orth RJ, Duarte CM eds,  
521 Springer, New York: 159-192.

522 Meinesz A., Genot I., Hesse B., 1990. Données quantitatives sur les biocénoses littorales  
523 marines de la Corse et impact de l'aménagement du littoral. GIS Posidonie et DRAE  
524 Corse publ., Marseille : 1-22.

525 Monnier B., Pergent G., Valette-Sansevin A., Boudouresque C.F., Mateo M.A., Pergent-  
526 Martini C.; *In Press*. The *Posidonia oceanica* matte: A unique coastal carbon sink for  
527 climate change mitigation and implications for management. *Vie & Milieu*.

528 Nellemann C., Corcoran E., Duarte C.M., Valdés L., De Young C., Fonseca L., Grimsditch G.,  
529 2009. Blue carbon. The role of healthy oceans in binding carbon. A rapid response  
530 assessment. United Nations Environment Programme publ., GRID-Arendal, Norway:  
531 1-80.

532 Pasqualini V., Pergent-Martini C., Clabaut P., Pergent G., 1998. Mapping of *Posidonia*  
533 *oceanica* using aerial photographs and side-scan sonar: Application of the island of  
534 Corsica (France). *Estuarine Coastal Shelf Science*, 47: 359-367.

535 Pergent G., 1990. Lepidochronological analysis of the seagrass *Posidonia oceanica* (L.) Delile:  
536 a standardized approach. *Aquatic Botany*, 37: 39-54.

537 Pergent G., Bazairi H., Bianchi C.N., Boudouresque C.F., Buia M.C., Calvo S., Clabaut P.,  
538 Harmelin-Vivien M., Mateo M.A., Montefalcone M., Morri C., Orfanidis S., Pergent-  
539 Martini C., Semroud R., Serrano O., Thibaut T., Tomasello T., Verlaque M., 2014.  
540 Climate change and Mediterranean seagrass meadows: a synopsis for environmental  
541 managers. *Mediterranean Marine Science*. 15/2 Free Open Access.

542 Pergent G., Pergent-Martini C., 1990. Some Applications of Lepidochronological Analysis in  
543 the Seagrass *Posidonia oceanica*. *Botanica Marina*, 33: 299-310.

544 Pergent G., Rico-Raimondino V., Pergent-Martini C., 1997. Fate of primary production in  
545 *Posidonia oceanica* meadows of the Mediterranean. *Aquatic Botany*, 59: 307-321.

546 Pergent G., Romero J., Pergent-Martini C., Mateo M.A., Boudouresque C.F., 1994. Primary  
547 production stocks and fluxes in the Mediterranean seagrass *Posidonia oceanica*.  
548 *Marine Ecology Progress Series*, 106: 139-146.

549 Pergent-Martini C., Bonacorsi M., Bein A., Braschi J., Kremmer L., Pergent G., 2013. Mapping  
550 of coastlines of the Republic of Cyprus – Progress report, April 2013. Agreement

551 DFMR Republic of Cyprus, GIS Posidonie & AP Marine Environmental Consultancy Ltd,  
552 Ref n°1115, GIS Posidonie Publ., Corte: 1-38.

553 Pergent-Martini C., Rico-Raimondino V., Pergent G., 1994. Primary production of *Posidonia*  
554 *oceanica* in the Mediterranean basin. *Marine Biology*, 120: 9-15.

555 Reed D.C., Brzezinski M.A., 2009. Kelp forests in Laffoley D. d'A. and Grimsditch G. (eds). The  
556 management of natural coastal carbon sinks. IUCN. Gland. Switzerland: 31-37.

557 Romero J., 2004. Las praderas de Fanerógamas marinas. La producción primaria y su destino.  
558 Características de los restos de la planta. Luque A.A., Templado J. edit., *Praderas y*  
559 *bosques marinos de Andalucía*, Consejería de Medio Ambiente, Junta de Andalucía  
560 publ., Sevilla: 74-81.

561 Romero J., 1989. Primary production of *Posidonia oceanica* beds in the Medas Islands  
562 (Girona, NE Spain). *International Workshop on Posidonia beds*, Boudouresque C.F.,  
563 Meinesz A., Fresi E., Gravez V. edit., GIS Posidonie publ., Marseille, 2: 85-91.

564 Romero J., Pergent G., Pergent-Martini C., Mateo M.A., Regnier C., 1992. The detritic  
565 compartment in a *Posidonia oceanica* meadow: litter features, decomposition rates  
566 and mineral stocks. *PSZN Marine Ecology*, 13 (1): 69-83.

567 Romero J., Perez M., Mateo M.A., Sala E., 1994. The belowground organs of the  
568 Mediterranean seagrass *Posidonia oceanica* as a biogeochemical sink. *Aquatic*  
569 *Botany*, 47: 13-19.

570 Romero J., Perez M., Alcoverro T., Mateo M.A., Sanchez-Lizaso J.L., 1998. Production ecology  
571 of *Posidonia oceanica* (L.) Delile meadows in Nueva Tabarca marine Reserve: growth,  
572 biomass and nutrient stocks along a bathymetric gradient. *Oecologia aquatica*, 11:  
573 111-121.

574 Ruiz J.M., 2000. Respuesta de la fanerogama marina *Posidonia oceanica* (L.) Delile. A  
575 perturbaciones antropicas. Tesis Doctoral, Universidad de Murcia: 212pp.

576 Scartazza A., Moscatello S., Gavrichkova O., Buia M.C., Lauteri M., Batistelli A., Lorenti M.,  
577 Garrard S.L., Calfapietra C., Brugnoli E., 2017. Carbon and nitrogen allocation strategy  
578 in *Posidonia oceanica* is altered by seawater acidification. *Science of the Total*  
579 *Environment*, 607-608: 954-964.

580 Serrano O., Lavery PS, Rozaimi M, Mateo MA 2014. Influence of water depth on the carbon  
581 sequestration capacity of seagrass. *Global Biogeochem Cy* 28: 950-961.

582 Serrano O., Mateo M.A., Renom P., Julià R., 2012. Characterization of soils beneath a  
583 *Posidonia oceanica* meadow. *Geoderma*, 185: 26-36.

584 Spalding M., Taylor M., Ravilious C., Short F., Green E., 2003. The distribution and status of  
585 seagrasses. *World Atlas of Seagrasses*: 5-26.

586 Telesca L., Belluscio A., Criscoli A., Ardizzone G., Apostolaki E.T., Fraschetti S., Gristina M.,  
587 Knittweis L., Martin C.S., Pergent G., Alagna A., Badalamenti F., Garofalo G., Gerakaris  
588 V., Pace M.L., Pergent-Martini C., Salomidi M., 2015. Seagrass meadows (*Posidonia*  
589 *oceanica*) distribution and trajectories of change. *Scientific Reports*, 5: 1-14 (12505).

590 Tomasello A., Di Maida G., Calvo S., Pirrota M., Borra M., Procaccini G., 2009. Seagrass  
591 meadows at the extreme of environmental tolerance: the case of *Posidonia oceanica*  
592 in a semi-enclosed coastal lagoon. *Marine Ecology*, 30: 288-300.

593 Thelin I., Giorgi J., 1984. Production de feuilles dans un herbier superficiel à *Posidonia*  
594 *oceanica* évaluée par une méthode dérivée de la méthode de ZIEMAN. *International*  
595 *Workshop on Posidonia oceanica beds*, Boudouresque C.F., Jeudy De Grissac A.,  
596 Olivier J. edit., GIS Posidonie publ., Marseille, 1: 271-276.

597 Topouzelis K., Makri D., Stoupas N., Papakonstantinou A., Katsanevakis S., 2018. Seagrass  
598 mapping in Greek territorial waters using Landsat-8 satellite images. *International*  
599 *Journal Applied Earth Observation Geoinformation*, 67: 98-113.

600 Torricelli L., Peirano A., 1997. Primary leaf production in the *Posidonia oceanica* meadow of  
601 Monterosso al Mare (eastern Ligurian Sea): variations along a depth gradient (in  
602 Italian). *Biologia Marina Mediterranea*, 4 (1): 81-85.

603 Traganos D., Aggarwal B., Poursanidis D., Topouzelis K., Chrysoulakis N., Reinartz P., 2018.  
604 Towards global-scale seagrass mapping and monitoring using Sentinel-2 on Google  
605 Earth Engine: The case study of the Aegean and Ionian seas. *Remote Sensing*, 10 (8):  
606 1227.

607 UNEP-MAP-RAC/SPA. 2009. State of knowledge on the geographical distribution of marine  
608 magnoliophyta meadows in the Mediterranean. Information Document for the Ninth  
609 meeting of the Focal Points for SPAs. prepared by R. Leonardini. G. Pergent & C.F.  
610 Boudouresque. Floriana Malta. 3-6 June 2009. RAC/SPA Publ.. UNEP(DEPI)/MED WG  
611 331/Inf 5: 1-374.

612 UNEP-MAP-RAC/SPA, 2015. Guidelines for Standardization of Mapping and Monitoring  
613 Methods of Marine Magnoliophyta in the Mediterranean. Christine Pergent-Martini,  
614 Edits., RAC/SPA publ., Tunis: 48pp + Annexes.

615 UNEP-MAP-SPA/RAC, 2019. Updating of the Action Plan for the Conservation of Marine  
616 Vegetation in the Mediterranean Sea. Fourteenth Meeting of SPA/BD Thematic Focal  
617 Points. Portorož, Slovenia, 18-21 June 2019: 25pp + Annexes.

618 UNEP-WCMC, 2018. Global distribution of seagrasses (version 6.0). Sixth update to the data  
619 layer used in Green and Short (2003). Cambridge (UK): UN Environment World  
620 Conservation Monitoring Centre. URL: <http://data.unep-wcmc.org/datasets/7>.

621 Valette-Sansevin A., Pergent G., Buron K., Damier E., Pergent-Martini C., 2019. Continuous  
622 mapping of benthic habitats along the coast of Corsica: A tool for the inventory and  
623 monitoring of blue carbon ecosystems. *Mediterranean Marine Science*, 20 (3): 585-  
624 593.

625 Vela A., 2006. Fonctionnement et production primaire des herbiers à *Posidonia oceanica* (L.)  
626 Delile en Méditerranée. Thèse Doctorat Université de Corse : 1-155.

627 Vela A., Leoni V., Pergent G., Pergent-Martini C., 2006. Relevance of Leaf matter loss in the  
628 functioning of *Posidonia oceanica* system. *Biologia Marina Mediterranea*. 13(4): 102-  
629 106.

630 Warner B.G., Clymo R.S., Tolonen K., 1993. Implications of peat accumulation at Point  
631 Escuminac, New Brunswick. *Quaternary Res.*, 39: 245-248.

632 Wittmann K.J., 1984. Temporal and morphological variations of growth in a natural stand of  
633 *Posidonia oceanica*. *PSZN Marine Ecology*, 5 (4): 301-316.

634 Wittmann K.J., Ott J.A., 1982. Effects of cropping on growth in the Mediterranean seagrass  
635 *Posidonia oceanica* (L.) Delile. *PSZN Marine Ecology*, 3 (2): 151-159.

636 Zupo V., Buia M.C., Mazzella L., 1997. A production model for *Posidonia oceanica* based on  
637 temperature. *Estuarine Coastal Shelf Science*, 44: 485-492.

638