



HAL
open science

Upper bounds on the one-arm exponent for dependent percolation models

Vivek Dewan, Stephen Muirhead

► **To cite this version:**

Vivek Dewan, Stephen Muirhead. Upper bounds on the one-arm exponent for dependent percolation models. 2021. hal-03154641

HAL Id: hal-03154641

<https://hal.science/hal-03154641v1>

Preprint submitted on 1 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

UPPER BOUNDS ON THE ONE-ARM EXPONENT FOR DEPENDENT PERCOLATION MODELS

VIVEK DEWAN¹ AND STEPHEN MUIRHEAD²

ABSTRACT. We prove upper bounds on the one-arm exponent η_1 for dependent percolation models; while our main interest is level set percolation of smooth Gaussian fields, the arguments apply to other models in the Bernoulli percolation universality class, including Poisson-Voronoi and Poisson-Boolean percolation. More precisely, in dimension $d = 2$ we prove $\eta_1 \leq 1/3$ for Gaussian fields with rapid correlation decay (e.g. the Bargmann-Fock field), and in general dimensions we prove $\eta_1 \leq d/3$ for finite-range fields and $\eta_1 \leq d - 2$ for fields with rapid correlation decay. Although these results are classical for Bernoulli percolation (indeed they are best-known in general), existing proofs do not extend to dependent percolation models, and we develop a new approach based on exploration and relative entropy arguments. We also establish a new Russo-type inequality for smooth Gaussian fields which we use to prove the sharpness of the phase transition for finite-range fields.

1. INTRODUCTION

The critical phase of percolation models is believed (see, e.g., [23, Chapter 9]) to be described by *critical exponents* which govern the power-law behaviour of macroscopic observables at, or near, criticality. In this paper we consider the *one-arm exponent*; we introduce this in the classical setting of Bernoulli percolation, before generalising to a class of dependent percolation models induced by the excursion sets of smooth Gaussian fields (‘Gaussian percolation’).

Fix a dimension $d \geq 2$, consider the lattice $\mathbb{Z}^d = (\mathcal{V}, \mathcal{E})$, and declare each edge $e \in \mathcal{E}$ to be ‘open’ independently with probability $p \in [0, 1]$. The resulting law \mathbb{P}_p of the open subset of \mathcal{E} is known as *Bernoulli percolation on \mathbb{Z}^d with parameter p* . Defining the connection event

$$\{A \longleftrightarrow B\} := \{\text{there exists a path of open edges that intersects } A \text{ and } B\}$$

where $A, B \subset \mathcal{V}$,¹ and denoting by $\Lambda_R := [-R, R]^d \subset \mathcal{V}$ the box of size R , it is well known [23] that there exists $p_c = p_c(d) \in (0, 1)$, satisfying $p_c(2) = 1/2$ and $p_c(d) < 1/2$ for $d \geq 3$, such that

$$\theta(p) := \mathbb{P}_p[0 \longleftrightarrow \infty] := \lim_{R \rightarrow \infty} \mathbb{P}_p[0 \longleftrightarrow \partial\Lambda_R] = \begin{cases} 0 & \text{if } p < p_c, \\ > 0 & \text{if } p > p_c. \end{cases}$$

Although it is still open to prove $\theta(p_c) = 0$ for $d \geq 3$, it has been shown that [11, 2]

$$(1.1) \quad \theta(p) \geq c(p - p_c)$$

for a constant $c = c(d)$ and $p > p_c$ sufficiently close to p_c ; this is known as the *mean-field lower bound* and is expected to be tight for dimensions $d \geq d_c = 6$ in which critical exponents take their mean-field value.

At criticality $p = p_c$ it is believed that connection probabilities between scales obey a power law, in the sense that there exists $\eta_1 > 0$ such that, as $R \rightarrow \infty$ and for $r = o(R)$,

$$(1.2) \quad \mathbb{P}_{p_c}[\Lambda_r \longleftrightarrow \partial\Lambda_R] = (r/R)^{-\eta_1 + o(1)}.$$

¹INSTITUT FOURIER, UNIVERSITÉ GRENOBLE ALPES

²SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE

E-mail addresses: vivek.dewan@univ-grenoble-alpes.fr, smui@unimelb.edu.au.

Date: February 25, 2021.

2010 *Mathematics Subject Classification.* 60G60 (primary); 60F99 (secondary).

Key words and phrases. Percolation, critical exponents, Gaussian fields.

¹We allow the path to be empty, so that $\{A \longleftrightarrow B\}$ occurs if $A \cap B \neq \emptyset$.

While the existence of the *one-arm exponent* η_1 is not known rigorously, since we are interested in upper bounds we define

$$(1.3) \quad \eta_1 := \liminf_{R \rightarrow \infty} \frac{-\log \mathbb{P}_{p_c}[0 \longleftrightarrow \partial \Lambda_R]}{\log R}.$$

Clearly upper bounds on (1.3) imply upper bounds on the exponent in (1.2) assuming its existence. Note however that the choice of \liminf , rather than \limsup , in the definition of η_1 is deliberate and yields a priori weaker upper bounds (see however Remark 1.3).

The phenomenon of *universality* suggests that a wide class of dependent percolation models behave similarly to Bernoulli percolation at, or near, criticality, and in particular η_1 should be identical inside this class. In this paper we consider the following class of dependent models. Let f be a continuous stationary-ergodic centred Gaussian field on \mathbb{R}^d , and for $\ell \in \mathbb{R}$ write $\mathbb{P}_\ell[\cdot]$ to denote $\mathbb{P}[f + \ell \in \cdot]$ (abbreviating $\mathbb{P} = \mathbb{P}_0$). Then the excursion sets $\{f + \ell \geq 0\} := \{x \in \mathbb{R}^d : f(x) + \ell \geq 0\}$ induce a stationary ergodic percolation model on \mathbb{R}^d via the connectivity relation

$$\{A \longleftrightarrow B\} := \{\text{there exists a path in } \{f \geq 0\} \text{ that intersects } A \text{ and } B\}$$

for closed sets $A, B \subset \mathbb{R}^d$. Recalling the box $\Lambda_R := [-R, R]^d$ (now considered a subset of \mathbb{R}^d), by monotonicity there exists $\ell_c = \ell_c(f) \in [-\infty, \infty]$ such that

$$\theta(\ell) := \mathbb{P}_\ell[\Lambda_1 \longleftrightarrow \infty] := \lim_{R \rightarrow \infty} \mathbb{P}_\ell[\Lambda_1 \longleftrightarrow \partial \Lambda_R] = \begin{cases} 0 & \text{if } \ell < \ell_c, \\ > 0 & \text{if } \ell > \ell_c, \end{cases}$$

where the choice of $\{\Lambda_1 \longleftrightarrow \partial \Lambda_R\}$ rather than $\{0 \longleftrightarrow \partial \Lambda_R\}$ is to avoid the possibility of local obstructions (relevant only in the case that the FKG inequality is not available; see the comments after (POS')). Under general conditions it is known that $\ell_c = 0$ if $d = 2$ and $\ell_c \in (-\infty, 0]$ if $d \geq 3$ (see [44, 38, 22, 42, 37] and [35, 36] respectively for sufficient conditions, which are implied by the assumptions in this paper, namely Assumption 1.4 below). Similarly to for Bernoulli percolation, for this class of models we define

$$(1.4) \quad \eta_1 := \liminf_{R \rightarrow \infty} \frac{-\log \mathbb{P}_{\ell_c}[\Lambda_1 \longleftrightarrow \partial \Lambda_R]}{\log R}.$$

In this case the mean-field lower bound (1.1) has not yet been established; indeed in this paper we prove it for Gaussian fields with *finite-range dependence*.

1.1. Upper bounds on the one-arm exponent. We now present our main results, which are upper bounds on η_1 . We begin with Bernoulli percolation; although the results are not new in this case, they are illustrative for general models.

Theorem 1.1. *For Bernoulli percolation on \mathbb{Z}^d ,*

$$\eta_1 \leq \begin{cases} 1/3 & d = 2, \\ d/3 & d \geq 3. \end{cases}$$

Remark 1.2. If $d = 2$, the bound $\eta_1 \leq 1/3$ was first given in [29]² which established that $\mathbb{P}_{p_c}[0 \longleftrightarrow \partial \Lambda_R] \geq cR^{-1/3}$. Recently this was improved to $\geq cR^{-1/6}$ [14], giving $\eta_1 \leq 1/6$. It is believed that $\eta_1 = 5/48$, but this is known rigorously only for very specific models [47].

In general dimension the hyperscaling inequality $\eta_1 \leq d/(1 + \delta)$ has been established rigorously [9], where δ is the critical exponent governing the volume of critical clusters. In light of the mean-field bound $\delta \geq 2$ [2], this implies $\eta_1 \leq d/3$.³ In high dimension $d \geq 11$ it is known that η_1 takes its mean-field value $\eta_1 = 2$ [31, 20], see also Corollary 1.2 below. It is believed that

$$\eta_1 = \begin{cases} 0.48 \dots & d = 3, & 0.95 \dots & d = 4, \\ 1.5 \dots & d = 5, & 2 & d \geq d_c = 6. \end{cases}$$

²In the paper the argument is attributed to van den Berg.

³Although [9] assumes the existence of the exponent δ , one can extract the unconditional bound $\eta_1 \leq d/3$ from the proof.

In particular, the bound $\eta_1 \leq d/3$ is expected to be tight at the upper-critical dimension $d_c = 6$ (indeed, it implies $d_c \geq 6$ since it shows that $\eta_1 = 2$ cannot occur for $d \leq 5$).

As for lower bounds on η_1 , in $d = 2$ one can use RSW estimates to prove that $\eta_1 > \varepsilon$ (which can be quantified, but is small), however if $d \in \{3, 4, 5, 6\}$ it is still wide open to prove $\eta_1 > 0$ (there are partial results in intermediate dimensions, e.g. for spread-out models).

Remark 1.3. For the bound $\eta_1 \leq 1/3$ in $d = 2$, we could replace the \liminf in the definition of η_1 in (1.3) with \limsup , since the argument yields $\mathbb{P}_{p_c}[0 \longleftrightarrow \partial\Lambda_R] \geq cR^{-1/3}$, see (2.9). In fact, the argument gives the stronger bound $\mathbb{P}_{p_c}[A_2(R)] \geq cR^{-2/3}$, where $A_2(R)$ is the (*polychromatic*) *two-arm event*; see Section 2 for the definition and more details.

Similarly, if $d \geq 3$ one can modify our argument to give $\mathbb{P}_{p_c}[0 \longleftrightarrow \partial\Lambda_R] \geq cR^{-d/3}$ by working under an (unproven) assumption that critical ‘box-crossing’ probabilities do not converge to 1, which to our knowledge is a new inference; see Remark 2.6 for details. Note that this assumption is expected to be true if $d < d_c = 6$, but likely not if $d \geq 6$.

Previous proofs of Theorem 1.1 rely heavily on specific properties of Bernoulli percolation (such as the BK inequality, used to prove $\delta \geq 2$, and in the case of the stronger bound $\eta_1 \leq 1/6$ if $d = 2$, on the ‘parafermionic observable’), and hence do not extend easily to dependent percolation models. On the contrary, we give a new proof of Theorem 1.1 that extends naturally to a wide class of dependent models; our next result illustrates this for Gaussian percolation.

Let us begin by stating some assumptions. Recall that f is a continuous stationary-ergodic centred Gaussian field. We will always assume that f has a *spatial moving average* representation $f = q \star W$, where $q \in L^2(\mathbb{R}^d) \neq 0$ is Hermitian (i.e. $q(x) = q(-x)$), W is the white noise on \mathbb{R}^d , and \star denotes convolution; a sufficient condition is that the covariance kernel $K(\cdot) := \mathbb{E}[f(0)f(\cdot)] = (q \star q)(\cdot)$ is in $L^1(\mathbb{R}^d)$, since then we may define $q := \mathcal{F}[\sqrt{\rho}]$, where \mathcal{F} denotes the Fourier transform and $\rho = \mathcal{F}[K] \in C^0(\mathbb{R}^d)$ is the *spectral density* of the field.

For our main results we will further assume that q satisfies the following basic properties:

Assumption 1.4 (Basic assumptions on the Gaussian field).

- (a) (*Regularity*) q is three-times differentiable and each of these derivatives is in $L^2(\mathbb{R}^d)$.
 (b) (*Decay of correlations, with parameter $\beta > d$*) There exists a $c > 0$ such that, for all $x \in \mathbb{R}^d$,

$$\max\{|q(x)|, |\nabla q(x)|\} \leq c|x|^{-\beta}.$$

- (c) (*Symmetry*) q is symmetric under negation and permutation of the coordinate axes.

Let us explain some consequences of Assumption 1.4. The regularity condition implies that $K = q \star q \in C^6(\mathbb{R}^d)$, and hence f is C^2 -smooth almost surely (see [1, Theorem 1.4.1]). The decay condition implies that $q \in L^1(\mathbb{R}^d)$ and so also $K \in L^1(\mathbb{R}^d)$, which ensures that the spectral density is continuous and $(f, \nabla f, \nabla^2 f)$ is non-degenerate (i.e. its evaluation on a finite number of distinct points is a non-degenerate Gaussian vector, see [6, Lemma A.2]). The symmetry assumption is crucial in $d = 2$ (for instance, to prove RSW estimates), but it also simplifies some aspects of the proof in all dimensions. Finally, as we mentioned above, if $d = 2$ then Assumption 1.4 is sufficient to prove that $\ell_c = 0$ (see [37, Theorem 1.3] and Remark 1.9 therein).

For most of our results we also assume

$$(POS) \quad \int q := \int_{\mathbb{R}^d} q(x) dx > 0.$$

This is equivalent to the spectral density being positive at the origin, and is a natural assumption when studying how properties of a Gaussian field change with the level; see e.g. [38, 5].

For some of our results we further assume that f is *positively correlated*:

$$(POS') \quad K(x) = (q \star q)(x) \geq 0 \text{ for all } x \in \mathbb{R}^d.$$

This is equivalent to the *FKG inequality* holding for the field (i.e. the field is *positively associated*), so that events increasing with respect to the field are positively correlated [41].⁴ Note that

⁴Although in [41] this is proven only for finite Gaussian *vectors*, one can deduce positive associations for all increasing events considered in this paper via standard approximation arguments, see [43, Lemma A.12].

(POS') is stronger than (POS) (the former implies that the spectral density is positive definite, and so strictly positive at the origin unless $K = 0$).

Recall that the mean-field lower bound (1.1) is not known for Gaussian percolation. We introduce it as an assumption: There exists $c > 0$ such that, for $\ell > \ell_c$ sufficiently close to ℓ_c ,

$$(MFB) \quad \theta(\ell) := \mathbb{P}_\ell[\Lambda_1 \longleftrightarrow \infty] \geq c(\ell - \ell_c).$$

While we expect (MFB) to hold in great generality, in this paper we prove it only for finite-range dependent fields, see Theorem 1.14 below.

For Gaussian percolation we prove the following upper bounds on the one-arm exponent:

Theorem 1.5. *Suppose $f = q \star W$ satisfies Assumption 1.4 with parameter β and (POS).*

- (1) *If $d \geq 3$ and $\beta > 4d - 4$, then $\eta_1 \leq \min\{\frac{d-2+\alpha(3d-1)}{1+2\alpha}, d-1\}$ where $\alpha = \frac{3d-4}{2\beta-5d+4}$.*
- (2) *If $d \geq 2$, $\beta > 2d - 1$, and (MFB) holds, then $\eta_1 \leq \min\{\max\{\frac{d}{3} + \frac{\alpha(d-1)}{3}, \frac{\alpha(2d-1)}{3}\}, d-1\}$ where $\alpha = \frac{3d-2}{2\beta-d}$.*
- (3) *If $d = 2$, $\beta > \frac{8}{3}$, and (POS') holds, then $\eta_1 \leq \min\{\frac{1}{3} + \frac{5}{6(\beta-1)}, \frac{1}{2}\}$.*

To illustrate Theorem 1.5 consider the example of the *Bargmann-Fock* field with covariance kernel $K(x) = e^{-|x|^2/2}$ (see [4] for background and motivation), which is easily seen to satisfy Assumption 1.4 for every parameter β and also (POS'). According to the *Harris criterion* (see [51], or [7] for further discussion), it is expected that Gaussian percolation is in the Bernoulli percolation universality class if $K(x) \ll c|x|^{-2/\nu}$ where $\nu = \nu(d)$ is the correlation length exponent of Bernoulli percolation (see Section 1.2). In particular the Bargmann-Fock field is expected to belong to this class, and hence possess the same exponents as Bernoulli percolation.

Corollary 1.6. *Suppose $f = q \star W$ satisfies Assumption 1.4 for every parameter β and (POS') (e.g. the Bargmann-Fock field). Then*

$$\eta_1 \leq \begin{cases} 1/3 & d = 2, \\ d - 2 & d \geq 3. \end{cases}$$

Further, if (MFB) holds then $\eta_1 \leq d/3$.

Proof. Take $\beta \rightarrow \infty$ in Theorem 1.5. □

One can also consider the example of finite-range dependent fields, i.e. for which

$$(BOU) \quad q \text{ has bounded support,}$$

noting that this supersedes the second condition in Assumption 1.4.

Corollary 1.7. *Suppose $f = q \star W$ satisfies Assumption 1.4 and (POS)-(BOU). Then $\eta_1 \leq d/3$.*

Proof. Take $\beta \rightarrow \infty$ in the second statement of Theorem 1.5 ((MFB) holds by Theorem 1.14). □

Remark 1.8. Previously for Gaussian percolation it was known only that $\eta_1 \leq 1/2$ in $d = 2$, and $\eta_1 \leq d-1$ in $d \geq 3$ (the former is a consequence of RSW estimates [4, 43, 38], and see [12], or the proof of Theorem 1.5, for the latter); hence the bounds in Theorem 1.5 exceed what was known in all dimensions for large β . Notably, as for Bernoulli percolation, the bound $d/3$ is expected to be tight if $d = 6$. We emphasise that Corollary 1.7 does not assume positive correlations, and so applies to a class of models that lack positive associations.

Remark 1.9. As in Remark 1.3, if $d = 2$ we could replace the liminf in the definition of η_1 with limsup, since the proof yields polynomial lower bounds on $\mathbb{P}_{\ell_c}[\Lambda_1 \longleftrightarrow \partial\Lambda_R]$ (see (3.12)). Indeed the proof gives polynomial lower bounds on the *two-arm event*; for example for the Bargmann-Fock field we prove that, for every $\varepsilon > 0$ there is a $c > 0$ such that

$$\mathbb{P}_{\ell_c}[\{\text{there exists a path in } \{f = 0\} \text{ that intersects } \Lambda_1 \text{ and } \partial\Lambda_R\}] \geq cR^{-2/3-\varepsilon}.$$

1.2. Relations to other critical exponents. The methods used to prove the above results also give bounds on η_1 in terms of other critical exponents. For simplicity we state these only for Bernoulli percolation, but similar bounds can be proven for Gaussian percolation (which, under Assumption 1.4 and (POS')-(BOU), would match those in Theorem 1.10 below).

Let us introduce the relevant exponents. Recall the mean-field lower bound (1.1) on $\theta(p)$. It is expected that $\theta(p) \rightarrow 0$ as a power law as $p \downarrow p_c$; although this is not known rigorously (except in high dimension), we will assume that the corresponding exponent exists

$$\beta = \lim_{p \downarrow p_c} \frac{\log \theta(p)}{\log |p_c - p|} \in (0, 1].$$

Below criticality $p < p_c$, it is known that connection probabilities decay exponentially [34, 2, 18] and that the limit

$$\frac{1}{\xi(p)} := \lim_{R \rightarrow \infty} \frac{-\log \mathbb{P}_p[0 \longleftrightarrow \partial \Lambda_R]}{R} \in (0, \infty)$$

exists [23, Theorem 6.10]. The *correlation length* $\xi(p)$ is expected to diverge as a power law as $p \uparrow p_c$, and we will again assume that the corresponding exponent exists

$$\nu = \lim_{p \uparrow p_c} \frac{-\log \xi(p)}{\log |p_c - p|} \in (0, \infty).$$

Similarly, as $p \uparrow p_c$ the *susceptibility* $\chi(p) := \sum_{v \in \mathbb{Z}^d} \mathbb{P}_p[0 \longleftrightarrow v] < \infty$ is expected to diverge as a power law, and we will assume the existence of

$$\gamma = \lim_{p \uparrow p_c} \frac{-\log \chi(p)}{\log |p_c - p|} \in (0, \infty).$$

Finally we also assume that the critical *two-point function* decays as a power law with exponent

$$d - 2 + \eta := \lim_{|v|_\infty \rightarrow \infty} \frac{-\log \mathbb{P}_{p_c}[0 \longleftrightarrow v]}{\log |v|_\infty} \in (0, \infty),$$

where $|\cdot|_\infty$ denotes the sup-norm. It is well known that $\nu \geq 2/d$ [10], $\gamma \geq 1$ [3], and $\eta \leq 1$ [24].

Theorem 1.10. *For Bernoulli percolation on \mathbb{Z}^d , assuming the existence of β, ν, γ and η ,*

$$(1.5) \quad \frac{2 - \gamma}{\nu} \leq \eta_1 \leq \bar{\eta}_1 \leq \min \left\{ d - \frac{2}{\nu}, \frac{2 - \eta}{2/\beta - 1} \right\},$$

where $\bar{\eta}_1$ is defined as in (1.3) with *limsup* replacing *liminf*. Moreover

$$\eta_1 \leq \frac{d}{2/\beta + 1} \quad \text{and} \quad \bar{\eta}_1 \leq 1 - \frac{1}{\nu}, \quad \text{if } d = 2.$$

Remark 1.11. To our knowledge the bounds in (1.5) are new even for Bernoulli percolation, and $\eta_1 \geq \frac{2-\gamma}{\nu}$ may be of particular interest as a *lower* bound on η_1 . The bound $\eta_1 \leq \frac{d}{2/\beta+1}$ is implied by the hyperscaling inequality in [9], and for $\bar{\eta}_1 \leq 1 - \frac{1}{\nu}$ if $d = 2$ see [29, 50].

For Bernoulli percolation in sufficiently high dimension it is known that the exponents ν, γ and η exist and take their mean-field values $\nu = 1/2$ [26], $\gamma = 1$ [3], and $\eta = 0$ [27]. Hence Theorem 1.10 gives a new proof of the result of Kozma and Nachmias that $\eta_1 = 2$ in high dimension:

Corollary 1.12 ([31]). *For Bernoulli percolation on \mathbb{Z}^d , there exists $d_0 > 0$ such that, if $d \geq d_0$,*

$$\lim_{R \rightarrow \infty} \frac{-\log \mathbb{P}_{p_c}[0 \longleftrightarrow \partial \Lambda_R]}{\log R} = 2.$$

Remark 1.13. Our argument is significantly simpler than the one in [31], however it yields only

$$c_1 R^{-2} \leq \mathbb{P}_{p_c}[0 \longleftrightarrow \partial \Lambda_R] \leq c_2 R^{-2} (\log R)^4$$

whereas [31] proved that $\mathbb{P}_{p_c}[0 \longleftrightarrow \partial \Lambda_R] \asymp R^{-2}$ in the sense of bounded ratios (see Remark 2.7). Another difference is that we deduce $\eta_1 = 2$ in any dimension from the bounds $\nu \leq 1/2$ and $\eta \geq 0$ (see Remark 2.7, or recall the Fischer inequality $\gamma/\nu \leq 2 - \eta$), whereas the argument in [31] uses as input $d > 6$ and the two-sided bound $\eta = 0$ (or more precisely $\mathbb{P}_{p_c}[0 \longleftrightarrow v] \asymp |v|_\infty^{-d+2}$).

1.3. Sharpness of the phase transition for smooth Gaussian fields. As well as bounds on the one-arm exponent, a second aim of this paper is to establish the sharpness of the phase transition for smooth finite-range dependent Gaussian fields, and in addition verify the mean-field lower bound (MFB) for such fields. For this we adapt the celebrated argument of Duminil-Copin, Raoufi and Tassion [16] by exploiting a new ‘Russo-type inequality’ for smooth Gaussian fields (see Proposition 4.1); we expect this inequality will have further applications.

Theorem 1.14 (Sharpness of the phase transition and mean-field lower bound). *Suppose $f = q \star W$ is C^2 -smooth and satisfies (BOU). Then for every $\ell < \ell_c$ there exist $c_1, c_2 > 0$ such that, for $R \geq 1$,*

$$\mathbb{P}_\ell[\Lambda_1 \longleftrightarrow \partial\Lambda_R] \leq c_1 e^{-c_2 R}.$$

Moreover, the mean-field lower bound (MFB) holds.

Remark 1.15. For Gaussian percolation the sharpness of the phase transition was only known so far in two cases: (i) in $d = 2$ assuming Assumption 1.4 and (POS’) [38], and (ii) for certain *discrete* Gaussian fields on \mathbb{Z}^d satisfying (POS’) [13]. The mean-field lower bound (MFB) was not known for any smooth Gaussian fields. We emphasise that in Theorem 1.14 we do not assume (POS’), so this theorem holds for a class of fields lacking positive associations.

Remark 1.16. Clearly if (BOU) holds then f is finite-range dependent, but we do not know whether *every* finite-range dependent f can be represented as $q \star W$ for q with bounded support (although this seems very natural, and it is true if $d = 1$, see [19]). If we demand in addition that q be supported on half of the support of K then, rather surprisingly, this is false [19]. On the other hand, under (POS’) it is true [19, Corollary 3.2]. Moreover, it is known [45] that if f is finite-range dependent and *isotropic* (i.e. K is rotationally symmetric) it can be represented as a *countable sum* of independent $f_i = q_i \star W_i$ for q_i with *uniformly bounded* support. Since it is straightforward to extend our proof of Theorem 1.14 to handle such fields, the conclusions of Theorem 1.14 (and Corollary 1.7) also hold if f is smooth, finite-range dependent, and *isotropic*.

1.4. Other models. Other than Bernoulli percolation and level set percolation of Gaussian fields, the arguments adapt naturally to many other models in the Bernoulli percolation universality class. For instance, both Poisson-Voronoi and Poisson-Boolean percolation can be treated in a very similar way (although in the latter case the obtained bounds may depend on the decay of the radius distribution, and also some of our arguments in $d = 2$ do not apply since the model lacks self-duality). Indeed the necessary tools to apply the OSSS inequality in these settings, analogous to the arguments in Section 4, have already been developed in [15] and [17] respectively. For brevity we do not discuss details here.

While this work was being finalised we learned that similar arguments to those we use to prove $\eta_1 \leq 1/3$ if $d = 2$ were previously used in the general setting of increasing Boolean functions [8]; see Section 2.4 for a statement of the relevant result from [8] and a comparison to what we prove.

1.5. Outline of the paper. In Section 2 we study Bernoulli percolation and give the proof of Theorems 1.1 and 1.10. In Section 3 we adapt the arguments to the Gaussian setting, and give the proof of Theorem 1.5 subject to an auxiliary result (Proposition 3.9). In Section 4 we establish the Russo-type inequality for smooth Gaussian fields mentioned above, and apply it to prove Proposition 3.9 and Theorem 1.14. The appendix contains a technical result on orthogonal decompositions of Gaussian fields.

1.6. Acknowledgements. The second author was partially supported by the Australian Research Council (ARC) Discovery Early Career Researcher Award DE200101467. The authors thank Damien Gayet, Tom Hutchcroft, Ioan Manolescu and Hugo Vanneuville for helpful discussions, comments on an earlier draft, and for pointing out references [14] (Ioan) and [8, 49] (Hugo). This work was initiated while the first author was visiting Queen Mary University of London and we thank the University for its hospitality.

2. BERNOULLI PERCOLATION

In this section we focus on Bernoulli percolation, which serves as a template for the extension of the arguments to dependent percolation models.

Let us begin by introducing notation for connection events. For $k, R > 0$, define the box $B_k(R) := [-R, R] \times [-kR, kR]^{d-1} \subset \mathcal{E}$, and the ‘box-crossing event’

$$\text{Cross}_k(R) := \left\{ \{-R\} \times [-kR, kR]^{d-1} \overset{B_k(R)}{\longleftrightarrow} \{R\} \times [-kR, kR]^{d-1} \right\},$$

where $\{A \overset{E}{\longleftrightarrow} B\} := \{\text{there exists a path of open edges in } E \subset \mathcal{E} \text{ that intersects } A \text{ and } B\}$. For $R \geq 0$, define the *one-arm event*

$$A_1(R) := \{0 \longleftrightarrow \partial\Lambda_R\}.$$

Restricting for a moment to $d = 2$, we also introduce the (*polychromatic*) *two-arm event* $A_2(R)$ that was mentioned in Remark 1.3. Consider the dual lattice $(\mathbb{Z}^2)^*$; in this graph an edge is considered *open* if and only if the unique edge $e \in \mathcal{E}$ that it crosses is closed (i.e. not open). Note that each vertex $v \in \mathcal{V}$ has *four* neighbouring dual vertices, and for $A \subset \mathcal{V}$ let A^* be the union of these neighbours over $v \in A$. For $A, B \subseteq \mathcal{V}$ define

$$\{A \overset{E}{\longleftrightarrow} B\} = \{A \longleftrightarrow B\} \cap \{\text{there exists a dual path in } E \text{ that intersects } A^* \text{ and } B^*\},$$

where a *dual path in E* is a path of dual edges that cross closed edges in E , and abbreviate $\{A \longleftrightarrow B\} = \{A \overset{\mathbb{Z}^2}{\longleftrightarrow} B\}$. For $R \geq 0$, define

$$A_2(R) := \{0 \longleftrightarrow \partial\Lambda_R\} \quad \text{and} \quad \eta_2 := \liminf_{R \rightarrow \infty} \frac{-\log \mathbb{P}_{p_c}[A_2(R)]}{\log R}.$$

We make the elementary observation that

$$(2.1) \quad \eta_2 \geq 2\eta_1$$

where η_1 is defined in (1.3). To see this, note that by the FKG inequality

$$\mathbb{P}_p[A_2(R)] \leq \mathbb{P}_p[A_1(R)] \mathbb{P}_p[\{\text{there exists a dual path that intersects } 0^* \text{ and } \Lambda_R^*\}].$$

Since Bernoulli percolation on \mathbb{Z}^2 is self-dual at $p_c = 1/2$, and by translation invariance,

$$\mathbb{P}_{p_c}[\{\text{there exists a dual path that intersects } 0^* \text{ and } \Lambda_R^*\}] \leq 4\mathbb{P}_{p_c}[A_1(R-1)].$$

Hence $\mathbb{P}_{p_c}[A_2(R)] \leq 4\mathbb{P}_{p_c}[A_1(R-1)]^2$, and (2.1) follows immediately.

Let us return to the general setting of Bernoulli percolation on \mathbb{Z}^d . The case $d \geq 3$ of Theorem 1.1 is proven by combining the following result with the mean-field lower bound (1.1):

Proposition 2.1. *For $0 < p \leq q < 1$ and $R \geq 1$,*

$$\mathbb{P}_q[A_1(R)] - \mathbb{P}_p[A_1(R)] \leq \max\left\{ \frac{\sqrt{2}}{\sqrt{q(1-q)}}, \frac{\sqrt{2}}{\sqrt{p(1-p)}} \right\} (q-p) \sqrt{\mathbb{P}_q[A_1(R)] \sum_{v \in \Lambda_R} \mathbb{P}_p[0 \longleftrightarrow v]}.$$

For the case $d = 2$ of Theorem 1.1 we rely instead on the following inequalities:

Proposition 2.2. *Let $k \geq 1$. Then there exists $c > 0$ such that, for $p \in (0, 1)$ and $R \geq 1$,*

$$\frac{d}{dp} \mathbb{P}_p[\text{Cross}_k(R)] \leq \frac{cR^{d/2}}{\sqrt{p(1-p)}} \times \begin{cases} \sqrt{\mathbb{P}_p[A_2(R)]} & d = 2, \\ \sqrt{\mathbb{P}_p[A_1(R)]} & d \geq 2. \end{cases}$$

Proposition 2.3. *Let $d = 2$ and $k \geq 1$. Then there exists $c > 0$ such that, for $p \in (0, 1)$ and $R \geq 8$,*

$$(2.2) \quad \frac{d}{dp} \mathbb{P}_p[\text{Cross}_k(R)] \geq \frac{c}{p(1-p)} \frac{\mathbb{P}_p[\text{Cross}_{1/(8k)}(kR)]^4 (1 - \mathbb{P}_p[\text{Cross}_{8k}(R/8)])^2}{\mathbb{P}_p[A_2(R)]}.$$

We prove Propositions 2.1–2.3 later in the section; for now we complete the proof of our main results (Theorems 1.1 and 1.10). First we recall some standard facts:

Lemma 2.4.

(1) There exists $\delta > 0$ and $p' = p'(R) \leq p_c$ such that, for $R \geq 1$,

$$\mathbb{P}_{p'}[\text{Cross}_5(R)] = \delta.$$

(2) (RSW) Let $d = 2$ and $k > 0$. Then there exists $\delta > 0$ such that, for $R \geq 1$,

$$\mathbb{P}_{p_c}[\text{Cross}_k(R)] \in (\delta, 1 - \delta).$$

Proof. For the first statement, a classical bootstrapping argument (see, e.g., [28, Section 5.1]) shows that $\mathbb{P}_{p_c}[\text{Cross}_5(R)] > \delta$, and the result follows by continuity in p . The second statement amounts to the classical RSW estimates. \square

Proof of Theorem 1.1. In the proof $c > 0$ are constants that depend only on the dimension and may change from line to line. We begin with the case $d \geq 3$. We may assume that $\mathbb{P}_{p_c}[A_1(R)] \rightarrow 0$ as $R \rightarrow \infty$ since otherwise $\eta_1 = 0$. Define $q = q(R) > p_c$ such that

$$\mathbb{P}_q[A_1(R)] = \min\{2\mathbb{P}_{p_c}[A_1(R)], 1\},$$

which exists since $p \mapsto \mathbb{P}_p[A_1(R)]$ is continuous and strictly increasing. Note that $q(R) \rightarrow p_c$ as $R \rightarrow \infty$ since otherwise

$$\limsup_{R \rightarrow \infty} \mathbb{P}_{p_c}[A_1(R)] \geq \limsup_{R \rightarrow \infty} \theta(q(R))/2 > 0.$$

By the mean-field lower bound (1.1), for sufficiently large R

$$(2.3) \quad \mathbb{P}_{p_c}[A_1(R)] = \mathbb{P}_q[A_1(R)]/2 \geq \theta(q)/2 \geq c(q - p_c).$$

Now apply Proposition 2.1 to $p = p_c$ and $q = q(R)$; this yields

$$\mathbb{P}_{p_c}[A_1(R)] = \mathbb{P}_q[A_1(R)] - \mathbb{P}_{p_c}[A_1(R)] \leq c(q - p_c) \sqrt{\mathbb{P}_{p_c}[A_1(R)] \sum_{v \in \Lambda_R} \mathbb{P}_{p_c}[0 \longleftrightarrow v]}$$

for large R . Combining with (2.3), we deduce that

$$(2.4) \quad \mathbb{P}_{p_c}[A_1(R)] \sum_{v \in \Lambda_R} \mathbb{P}_{p_c}[0 \longleftrightarrow v] \geq c$$

for all $R \geq 1$.

We now show that $\eta_1 \leq d/3$ follows from (2.4). If $\eta_1 = 0$ there is nothing to prove, so assume $\eta_1 > 0$ and fix $\eta^* \in (0, \eta_1)$. Then by the definition of η_1

$$(2.5) \quad \mathbb{P}_{p_c}[A_1(R)] \leq R^{-\eta^*}$$

for large R ; in particular, via an integral comparison,

$$(2.6) \quad \sum_{v \in \Lambda_R} \mathbb{P}_{p_c}[A_1(\lfloor |v|_\infty / 2 \rfloor)]^2 \leq \max\{R^{d-2\eta^*}, 1\}(\log R)$$

for large R . Next observe that $\{0 \longleftrightarrow v\}$ implies the occurrence of

$$\{0 \longleftrightarrow \Lambda_{\lfloor |v|_\infty / 2 \rfloor}\} \quad \text{and} \quad \{v \longleftrightarrow v + \Lambda_{\lfloor |v|_\infty / 2 \rfloor}\}$$

which depend on disjoint subsets of edges. Hence by translation invariance and (2.6)

$$(2.7) \quad \sum_{v \in \Lambda_R} \mathbb{P}_{p_c}[0 \longleftrightarrow v] \leq \sum_{v \in \Lambda_R} \mathbb{P}_{p_c}[A_1(\lfloor |v|_\infty / 2 \rfloor)]^2 \leq \max\{R^{d-2\eta^*}, 1\}(\log R)$$

for large R , and so

$$c \leq \mathbb{P}_{p_c}[A_1(R)] \sum_{v \in \Lambda_R} \mathbb{P}_{p_c}[0 \longleftrightarrow v] \leq R^{-\eta^*} \max\{R^{d-2\eta^*}, 1\}(\log R).$$

This implies $\eta^* \leq d/3$, and since $\eta^* < \eta_1$ was arbitrary, we deduce $\eta_1 \leq d/3$.

We now turn to the case $d = 2$. By Propositions 2.2–2.3 and the RSW estimates (the second statement of Lemma 2.4), for large R

$$\frac{c}{\mathbb{P}_{p_c}[A_2(R)]} \leq \frac{d}{dp} \mathbb{P}_p[\text{Cross}_1(R)] \Big|_{p=p_c} \leq cR \sqrt{\mathbb{P}_{p_c}[A_2(R)]}$$

which yields, for large R ,

$$(2.8) \quad \mathbb{P}_{p_c}[A_2(R)] \geq cR^{-2/3}.$$

By the discussion after (2.1), this implies

$$(2.9) \quad \mathbb{P}_{p_c}[A_1(R)] \geq cR^{-1/3}$$

for large R , and hence $\eta_1 \leq 1/3$. \square

Remark 2.5. One could replace the right-hand side of (2.2) with the (perhaps simpler) expression

$$\frac{c}{p(1-p)} \frac{\mathbb{P}_p[\text{Cross}_k(R)](1 - \mathbb{P}_p[\text{Cross}_k(R)])}{\frac{1}{R^2} \sum_{i,j=0}^R \mathbb{P}_{p_c}[A_2(\min\{i, j\})]}.$$

While this suffices to prove $\eta_1 \leq 1/3$, it does not yield the stronger bounds (2.8)–(2.9).

Remark 2.6. In the case $d \geq 3$ our argument does not imply $\mathbb{P}_{p_c}[A_1(R)] \geq cR^{-d/3}$. However, as mentioned in Remark 1.3, one can obtain this by working under a ‘box-crossing’ assumption.

First, by modifying the proof of Proposition 2.3 one can prove that, for every $k \geq 1$ there exists a $c > 0$ such that, for $p \in (0, 1)$ and $R \geq 2$,

$$\frac{d}{dp} \mathbb{P}_p[\text{Cross}_k(R)] \geq \frac{c}{p(1-p)} \frac{\mathbb{P}_p[\text{Cross}_k(R)]^2 (1 - \mathbb{P}_p[\text{Cross}_{2k}(R/2)])}{\mathbb{P}_p[A_1(R)]}.$$

Next assume the following box-crossing property: For every $k \geq 1$ and $\delta_0 \in (0, 1)$ there are $\delta_1 \in (0, 1)$ and $R_0 > 0$ such that, for $R \geq R_0$ and $p \leq p_c$,

$$(BOX) \quad \mathbb{P}_p[\text{Cross}_k(R)] < 1 - \delta_0 \implies \mathbb{P}_p[\text{Cross}_{2k}(R/2)] < 1 - \delta_1.$$

Then by working on the sequence $p' = p'(R) \leq p_c$ at which $\mathbb{P}_{p'}[\text{Cross}_5(R)] = \delta$, guaranteed by the first statement of Lemma 2.4, and comparing upper and lower bounds on $\frac{d}{dp} \mathbb{P}_p[\text{Cross}_5(R)] \Big|_{p=p'}$, one deduces the result.

Note that (BOX) states roughly that if box-crossings do not occur with high probability for one aspect ratio, then they do not occur with high probability for other aspect ratios. This is known in $d = 2$ by the RSW estimates in Lemma 2.4, and is strongly believed to hold if $d < 6$ [9]. Although (BOX) seems difficult to verify, it is quite natural to work under this assumption; e.g. in [9] hyperscaling relations were proven under a version of (BOX), although interestingly they use this assumption to obtain *lower* bounds on η_1 .

Proof of Theorem 1.10. In the proof $c > 0$ are constants that depend only on the dimension and may change from line to line, and $o(1)$ denotes a quantity that decays to zero as $R \rightarrow \infty$.

We begin with the bounds $\eta_1 \leq d/(2/\beta + 1)$ and $\bar{\eta}_1 \leq (2 - \eta)/(2/\beta - 1)$ which require only a slight change to the argument used to prove $\eta_1 \leq d/3$ above. Recall (2.4) and let $q(R) \rightarrow p_c$ be defined as in (2.3). By the definition of the exponent β , one can replace (2.3) with

$$\mathbb{P}_{p_c}[A_1(R)] \geq \theta(q)/2 \geq c(q - p_c)^{\beta+o(1)},$$

which gives, in place of (2.4),

$$(2.10) \quad \mathbb{P}_{p_c}[A_1(R)]^{\frac{2}{\beta}-1+o(1)} \sum_{v \in \Lambda_R} \mathbb{P}_{p_c}[0 \longleftrightarrow v] \geq c,$$

for large R . Then using (2.7), for any $\eta^* \in (0, \eta_1)$ and large R we have

$$R^{-\eta^*(\frac{2}{\beta}-1)+o(1)} \max\{R^{d-2\eta^*}, 1\} (\log R) \geq c,$$

which implies $\eta_1 \leq d/(2/\beta + 1)$. On the other hand, by the definition of the exponent η ,

$$\sum_{v \in \Lambda_R} \mathbb{P}_{p_c}[0 \longleftrightarrow v] = R^{2-\eta+o(1)},$$

which by (2.10) implies

$$\mathbb{P}_{p_c}[A_1(R)] \geq R^{-(2-\eta)/(2/\beta-1)+o(1)}$$

and hence $\bar{\eta}_1 \leq (2-\eta)/(2/\beta-1)$.

To prove the remaining bounds we use the fact that, by a super-multiplicativity argument (see [23, Section 6.2]), there is a $c_1 > 0$ such that

$$(2.11) \quad \mathbb{P}_p[0 \longleftrightarrow v] \leq e^{-c_1|v|_\infty/\xi(p)}$$

for all $p < p_c$ and $v \in \mathbb{Z}^d$. We also recall the standard facts [23, Theorem 6.14] that $\xi(p)$ is continuous, strictly increasing, and $\xi(p) \rightarrow \infty$ as $p \uparrow p_c$.

Let $C > 0$ be a constant to be fixed later, and for R sufficiently large, let $p' = p'(R) \uparrow p_c$ be such that $R = C\xi(p') \log \xi(p')$. Since we have the a priori bound $\mathbb{P}_{p_c}[A_1(R)] \geq cR^{-(d-1)/2}$ [24, 48], we can take $C > 0$ sufficiently large so that, by (2.11) and the union bound,

$$\mathbb{P}_{p'}[A_1(R)] \leq cR^{d-1}e^{-c_1C \log \xi(p')} \leq cR^{d-1}R^{-c_1C+o(1)} \leq \mathbb{P}_{p_c}[A_1(R)]/2$$

for large R . Then applying Proposition 2.1 to $p = p'$ and $q = p_c$ gives, for large R ,

$$\mathbb{P}_{p_c}[A_1(R)]/2 \leq \mathbb{P}_{p_c}[A_1(R)] - \mathbb{P}_{p'}[A_1(R)] \leq c(p_c - p') \sqrt{\mathbb{P}_{p_c}[A_1(R)] \sum_{v \in \Lambda_R} \mathbb{P}_{p'}[0 \longleftrightarrow v]}$$

or, equivalently,

$$(2.12) \quad \mathbb{P}_{p_c}[A_1(R)] \leq c(p_c - p')^2 \sum_{v \in \Lambda_R} \mathbb{P}_{p'}[0 \longleftrightarrow v].$$

Since $\sum_{v \in \Lambda_R} \mathbb{P}_{p'}[0 \longleftrightarrow v] \leq \chi(p')$, and by the definition of the exponents ν and γ , this implies

$$\mathbb{P}_{p_c}[A_1(R)] \leq c(p_c - p')^2 \chi(p') \leq c\xi(p')^{-2/\nu+o(1)} \xi(p')^{\gamma/\nu+o(1)} = R^{-(2-\gamma)/\nu+o(1)}$$

for large R , which implies $\eta_1 \geq (2-\gamma)/\nu$.

Finally, let $\delta > 0$ be such that $\mathbb{P}_{p_c}[\text{Cross}_5(R)] \geq \delta$ for large R (possible by the first statement of Lemma 2.4), and again let $p' = p'(R) \uparrow p_c$ be such that $R = C\xi(p') \log \xi(p')$. Then

$$\mathbb{P}_{p'}[\text{Cross}_5(R)] \leq \delta/2$$

for large R , and we deduce that there exists $p'' \in (p', p_c)$ such that

$$\left. \frac{d}{dp} \mathbb{P}_p[\text{Cross}_5(R)] \right|_{p=p''} \geq \frac{\delta/2}{p_c - p'}.$$

On the other hand, by Proposition 2.2 and monotonicity in p ,

$$\left. \frac{d}{dp} \mathbb{P}_p[\text{Cross}_5(R)] \right|_{p=p''} \leq cR^{d/2} \sqrt{\mathbb{P}_{p_c}[A_1(R)]}$$

and hence

$$(p_c - p')^2 R^d \mathbb{P}_{p_c}[A_1(R)] \geq c$$

for large R . By the definition of the exponent ν , this implies

$$\xi(p')^{-2/\nu+o(1)} R^d \mathbb{P}_{p_c}[A_1(R)] = R^{d-2/\nu+o(1)} \mathbb{P}_{p_c}[A_1(R)] \geq c$$

for large R , which implies that $\bar{\eta}_1 \leq d - 2/\nu$. The bound $\bar{\eta}_1 \leq 1 - 1/\nu$ in $d = 2$ is similar, except we use two-arm events as in the proof of Theorem 1.1. \square

Remark 2.7. As mentioned in Remark 1.13, by combining the high-dimensional bounds [26, 27]

$$\xi(p) \leq c(p_c - p)^{-1/2} \quad \text{and} \quad \mathbb{P}_{p_c}[0 \longleftrightarrow v] \leq c|v|_\infty^{-d+2}$$

with (2.4) and (2.12), one arrives at a quantitative version of Corollary 1.12, namely the bounds

$$c_1 R^{-2} \leq \mathbb{P}_{p_c}[A_1(R)] \leq c_2 R^{-2} (\log R)^4.$$

2.1. Exploration algorithms. To prove Propositions 2.1–2.3 we make use of *exploration algorithms*, which we introduce in a general setting.

Definition 2.8 (Randomised algorithms). Let $X = (X_i)$ be a countable set of random variables taking values in arbitrary probability spaces. A (*randomised*) *algorithm* \mathcal{A} on X is a random adapted procedure that sequentially reveals a subset of the coordinates X_i and returns a value. We say that \mathcal{A} *determines an event* A if it returns the value $\mathbb{1}_A$ almost surely. The *revelment* $\text{Rev}(i)$ of a given coordinate X_i is the probability that \mathcal{A} reveals this coordinate.

For Bernoulli percolation we consider algorithms on $X = (X_e)_{e \in \mathcal{E}}$ for $X_e = \mathbb{1}_{e \text{ open}}$. A useful property of the events $A_1(R)$ and $\text{Cross}_k(R)$ is the existence of determining algorithms whose revealments are controlled by connection probabilities. Recall the box $B_k(R) \subset \mathcal{E}$, and define its *right half* $B_k^+(R) := [0, R] \times [-kR, kR]^{d-1} \subset \mathcal{E}$. If $d = 2$, define also its *top-right quarter* $B_k^\dagger(R) := [0, R] \times [0, kR] \subset \mathcal{E}$.

Lemma 2.9. *For every $p \in (0, 1)$ and $R \geq 1$ there is an algorithm determining $A_1(R)$ such that, under \mathbb{P}_p ,*

$$\sum_{e \in \mathcal{E}} \text{Rev}(e) \leq 2 \sum_{v \in \Lambda_R} \mathbb{P}_p[0 \longleftrightarrow v].$$

Moreover for every $k \geq 1$, $p \in (0, 1)$ and $R \geq 1$ there are algorithms determining $\text{Cross}_k(R)$ such that, under \mathbb{P}_p ,

$$\max_{e \in B_k^+(R)} \text{Rev}(e) \leq 2\mathbb{P}_p[A_1(R)],$$

and, if $d = 2$,

$$\max_{e \in B_k^\dagger(R)} \text{Rev}(e) \leq 2\mathbb{P}_p[A_2(R)].$$

We only give a sketch of proof; for more details see the proof of Lemma 3.6 which gives analogous statements in the Gaussian setting.

Proof (sketch). Recall the definition of $\{A \xleftrightarrow{E} B\}$, and for each edge $e \in \mathcal{E}$ let $\{e \xleftrightarrow{E} B\}$ be the union of $\{v \xleftrightarrow{E} B\}$ over the endpoints v of e , and $\{e \xleftrightarrow{E} B\}$ similarly.

For the first statement, let \mathcal{W} be the random subset of $B_1(R)$ defined by

$$\mathcal{W} := \left\{ e \in B_1(R) \mid \left\{ 0 \xleftrightarrow{B_1(R)} e \right\} \right\}.$$

Then consider the algorithm that sequentially reveals \mathcal{W} starting from the origin. This determines $A_1(R)$ and satisfies

$$\sum_{e \in \mathcal{E}} \text{Rev}(e) = \sum_{e \in B_1(R)} \mathbb{P}_p \left[\left\{ 0 \xleftrightarrow{B_1(R)} e \right\} \right] \leq 2 \sum_{v \in \Lambda_R} \mathbb{P}_p[\{0 \longleftrightarrow v\}].$$

For the second statement define instead

$$\mathcal{W} := \left\{ e \in B_k(R) \mid \left\{ e \xleftrightarrow{B_k(R)} \{-R\} \times [-kR, kR]^{d-1} \right\} \right\}.$$

Then consider the algorithm that sequentially reveals \mathcal{W} starting from the vertical hyperplane $\{-R\} \times [-kR, kR]^{d-1}$. This determines $\text{Cross}_k(R)$ since any crossing of $B_k(R)$ intersects the hyperplane $\{-R\} \times [-kR, kR]^{d-1}$, and the revealments for edges in $B_k^+(R)$ are bounded by

$$\max_{e \in B_k^+(R)} \mathbb{P}_p \left[\left\{ e \xleftrightarrow{B_k(R)} \{-R\} \times [-kR, kR]^{d-1} \right\} \right] \leq 2\mathbb{P}_p[A_1(R)].$$

For the third statement define instead

$$\mathcal{W} := \left\{ e \in B_k(R) \mid \left\{ e \xleftrightarrow{B_k(R)} (\{-R\} \times [-kR, kR]) \cup ([-R, R] \times \{-kR\}) \right\} \right\}$$

and consider the algorithm that sequentially reveals \mathcal{W} starting from the union of the vertical and horizontal lines $\{-R\} \times [-kR, kR]$ and $[-R, R] \times \{-kR\}$. This determines $\text{Cross}_k(R)$, since

if we reveal all interfaces that intersect these vertical and horizontal lines then we also determine $\text{Cross}_k(R)$. Moreover the revealments for edges in $B_k^\dagger(R)$ are bounded by

$$\max_{e \in B_k^\dagger(R)} \mathbb{P}_p \left[\left\{ e \xrightarrow{B_k(R)} (\{-R\} \times [-kR, kR]) \cup ([-R, R] \times \{-kR\}) \right\} \right] \leq 2\mathbb{P}_p[A_2(R)]. \quad \square$$

2.2. Proof of Propositions 2.1 and 2.2. We prove a general bound valid for arbitrary events, which extends a result from [40] (see also [46, Appendix B] and [50] for similar arguments):

Proposition 2.10. *Let $p, q \in (0, 1)$, let A be an event depending on a finite number of edges, let \mathcal{A} be an algorithm determining A , and let $\mathcal{E}' \subseteq \mathcal{E}$ be a subset of edges. Then*

$$(2.13) \quad |\mathbb{P}_{p;q}^{\mathcal{E}'}[A] - \mathbb{P}_p[A]| \leq \max \left\{ \frac{1}{\sqrt{p(1-p)}}, \frac{1}{\sqrt{q(1-q)}} \right\} |p - q| \sqrt{\max\{\mathbb{P}_p[A], \mathbb{P}_{p;q}^{\mathcal{E}'}[A]\} \mathbb{E}_p |\mathcal{W}_{\mathcal{E}'}|}$$

where $\mathbb{P}_{p;q}^{\mathcal{E}'}$ denotes the modification of \mathbb{P}_p in which the parameter on \mathcal{E}' is set to q (remaining at p on other edges), and $\mathcal{W}_{\mathcal{E}'}$ is the set of edges in \mathcal{E}' that are revealed by \mathcal{A} . In particular,

$$(2.14) \quad \left| \sum_{e \in \mathcal{E}'} \frac{\partial}{\partial p_e} \mathbb{P}_p[A] \right| \leq \frac{1}{\sqrt{p(1-p)}} \sqrt{\mathbb{P}_p[A] \mathbb{E}_p |\mathcal{W}_{\mathcal{E}'}|},$$

where $\frac{\partial}{\partial p_e}$ denotes the derivative with respect to the parameter on e .

Our proof of Proposition 2.10 is different to previous approaches in the literature (see Remark 2.14), and relies on properties of the relative entropy. For P and Q probability measures on a common measurable space, the *relative entropy* (or *Kullback-Leibler divergence*) from P to Q is defined as

$$D_{KL}(P||Q) := \int \log \left(\frac{dP}{dQ} \right) dP$$

if P is absolutely continuous with respect to Q , and $D_{KL}(P||Q) := \infty$ otherwise; $D_{KL}(P||Q)$ is non-negative by Jensen's inequality. If X and Y are random variables taking values in a common measurable space, with respective laws P and Q , we also write $D_{KL}(X||Y)$ for $D_{KL}(P||Q)$. We shall need two basic properties of the relative entropy (see [32, Theorem 2.2 and Corollary 3.2]):

- (1) (Chain rule) Let $X = (X_1, X_2)$ and $Y = (Y_1, Y_2)$ be random variables taking values in a common product measurable space. Then

$$(2.15) \quad D_{KL}(X||Y) = D_{KL}(X_1||Y_1) + \mathbb{E}_{x \sim X_1} [D_{KL}((X_2|X_1 = x)|| (Y_2|Y_1 = x))].$$

- (2) (Contraction) Let X and Y be random variables taking values in a common measurable space and let F be a measurable map from that space. Then

$$(2.16) \quad D_{KL}(X||Y) \geq D_{KL}(F(X)||F(Y)).$$

We first state a simple lemma on the relative entropy of stopped sequences of i.i.d. random variables. A *stopping time* for a real-valued sequence $X = (X_i)_{i \geq 1}$ is a positive integer $\tau = \tau(X)$ such that $\{\tau \geq n + 1\}$ is determined by $(X_i)_{i \leq n}$. We define the corresponding *stopped sequence* $X^\tau = (X_i^\tau)_{i \geq 1}$ as $X_i^\tau = X_i$ for $i \leq \tau$, and $X_i^\tau = \dagger$ for $i > \tau$, where \dagger is an arbitrary symbol.

Lemma 2.11. *Let $X = (X_i)_{i \geq 1}^n$ and $Y = (Y_i)_{i \geq 1}^n$ be finite real-valued sequences of i.i.d. random variables with respective univariate laws μ and ν , let $\tau \leq n$ be a stopping time, and let X^τ and Y^τ be the corresponding stopped sequences. Then*

$$D_{KL}(X^\tau||Y^\tau) = \mathbb{E}[\tau(X)] D_{KL}(\mu||\nu).$$

Proof. Define $X^{k \wedge \tau} = (X_i^\tau)_{i \leq k}$ and analogously for Y . By the chain rule (2.15), for $1 \leq k \leq n-1$,

$$\begin{aligned} & D_{KL}(X^{(k+1) \wedge \tau}||Y^{(k+1) \wedge \tau}) \\ &= D_{KL}(X^{k \wedge \tau}||Y^{k \wedge \tau}) + \mathbb{E}_{x \sim (X_i^\tau)_{i \leq k}} [D_{KL}(X_{k+1}^\tau|(X_i^\tau)_{i \leq k} = x||Y_{k+1}^\tau|(Y_i^\tau)_{i \leq k} = x)] \\ &= D_{KL}(X^{k \wedge \tau}||Y^{k \wedge \tau}) + \mathbb{E}_{x \sim (X_i^\tau)_{i \leq k}} [\mathbb{1}_{\tau(X) \geq k+1} D_{KL}(X_{k+1}^\tau|(X_i^\tau)_{i \leq k} = x||Y_{k+1}^\tau|(Y_i^\tau)_{i \leq k} = x)] \\ &= D_{KL}(X^{k \wedge \tau}||Y^{k \wedge \tau}) + \mathbb{P}[\tau(X) \geq k+1] D_{KL}(\mu||\nu) \end{aligned}$$

where in the last step we used that τ is a stopping time. Hence, by induction,

$$D_{KL}(X^\tau \| Y^\tau) = \sum_{1 \leq k \leq n-1} \mathbb{P}[\tau(X) \geq k+1] D_{KL}(\mu \| \nu) = \mathbb{E}[\tau(X)] D_{KL}(\mu \| \nu). \quad \square$$

We also need a variant of Pinsker's inequality:

Lemma 2.12. *Let P and Q be probability measures on a common measurable space and let A be an event. Then*

$$|P(A) - Q(A)| \leq \sqrt{2 \max\{P(A), Q(A)\} D_{KL}(P \| Q)}.$$

Proof. We use a standard reduction to the binary case. Let $\text{Ber}(x)$ and $\text{Ber}(y)$ be Bernoulli random variables with respective parameters $x := P(A)$ and $y := Q(A)$. By the contraction property (2.16) $D_{KL}(P \| Q) \geq D_{KL}(\text{Ber}(x) \| \text{Ber}(y))$, so it suffices to prove that

$$(2.17) \quad (x - y)^2 \leq 2 \max\{x, y\} D_{KL}(\text{Ber}(x) \| \text{Ber}(y)).$$

If $x \in \{0, 1\}$ or $y \in \{0, 1\}$ then (2.17) is trivial since either the right-hand side is infinite (if $x \neq y$) or both sides are zero (if $x = y$). On the other hand, if $x, y \in (0, 1)$ then

$$\begin{aligned} D_{KL}(\text{Ber}(x) \| \text{Ber}(y)) &:= x \log \frac{x}{y} + (1-x) \log \frac{1-x}{1-y} = \int_y^x \frac{x-s}{s(1-s)} ds \\ &\geq \frac{1}{\max\{x, y\}} \int_y^x (x-s) ds = \frac{1}{2 \max\{x, y\}} (x-y)^2 \end{aligned}$$

where we used that $\sup_{s \in [a, b]} s(1-s) \leq \max\{a, b\}$ for $0 \leq a \leq b \leq 1$. \square

Remark 2.13. In the proof we could replace $\max\{x, y\}$ with $\min\{\max\{x, y\}, 1/4\}$, which recovers the classical Pinsker's inequality $d_{TV}(P, Q) := \sup_A |P(A) - Q(A)| \leq \sqrt{D_{KL}(P \| Q)}/2$.

Proof of Proposition 2.10. Recall that $\mathcal{W}_{\mathcal{E}'}$ denotes the edges in \mathcal{E}' that are revealed by the algorithm, and let $W = (W_i)_{i \leq |\mathcal{W}_{\mathcal{E}'|}$ denote the configuration on $\mathcal{W}_{\mathcal{E}'}$ listed in the order of revelation. Moreover let W' denote the configuration on edges in $\mathcal{E} \setminus \mathcal{E}'$

First suppose that the algorithm \mathcal{A} depends only on the configuration (i.e. there is no auxiliary randomness). Then the event A is measurable with respect to (W, W') , and so by Lemma 2.12

$$|\mathbb{P}_{p,q}^{\mathcal{E}'}[A] - \mathbb{P}_p[A]| \leq \sqrt{2 \max\{\mathbb{P}_p[A], \mathbb{P}_{p,q}^{\mathcal{E}'}[A]\} D_{KL}((X, Z) \| (Y, Z))}$$

where (X, Z) (resp. (Y, Z)) is a random variable with the law of (W, W') under \mathbb{P}_p (resp. $\mathbb{P}_{p,q}^{\mathcal{E}'}$). Moreover, conditionally on W' , W has the law, under \mathbb{P}_p (resp. $\mathbb{P}_{p,q}^{\mathcal{E}'}$), of a sequence of i.i.d. Bernoulli random variables with parameter p (resp. q) stopped at the stopping time $|\mathcal{W}_{\mathcal{E}'|}$. Hence by the chain rule for the Kullback-Liebler divergence and Lemma 2.11,

$$D_{KL}((X, Z) \| (Y, Z)) = \mathbb{E}[D_{KL}((X | \mathcal{F}_Z) \| (Y | \mathcal{F}_Z))] = \mathbb{E}_p[|\mathcal{W}_{\mathcal{E}'|} D_{KL}(\text{Ber}(p) \| \text{Ber}(q))]$$

where \mathcal{F}_Z denotes the σ -algebra generated by Z . Combining we have

$$(2.18) \quad |\mathbb{P}_{p,q}^{\mathcal{E}'}[A] - \mathbb{P}_p[A]| \leq \sqrt{2 \max\{\mathbb{P}_p[A], \mathbb{P}_{p,q}^{\mathcal{E}'}[A]\} \mathbb{E}_p[|\mathcal{W}_{\mathcal{E}'|} D_{KL}(\text{Ber}(p) \| \text{Ber}(q))]}.$$

Finally since

$$\begin{aligned} D_{KL}(\text{Ber}(p) \| \text{Ber}(q)) &:= p \log \frac{p}{q} + (1-p) \log \frac{1-p}{1-q} = \int_q^p \frac{p-s}{s(1-s)} ds \\ &\leq \max\left\{\frac{1}{p(1-p)}, \frac{1}{q(1-q)}\right\} \int_q^p (p-s) ds = \max\left\{\frac{1}{2p(1-p)}, \frac{1}{2q(1-q)}\right\} (p-q)^2 \end{aligned}$$

the proof is complete.

The general case follows by averaging over any auxiliary randomness in the algorithm, since by Jensen's inequality $\mathbb{E}[\sqrt{\mathbb{E}[|\mathcal{W}_{\mathcal{E}'|} | \mathcal{G}]}] \leq \mathbb{E}[\sqrt{|\mathcal{W}_{\mathcal{E}'|}}]$ for any sub- σ -algebra \mathcal{G} . \square

Remark 2.14. For comparison we sketch an alternative approach which is closer to that appearing in previous works (e.g. [40]); this leads to the bound

$$(2.19) \quad \left| \sum_{e \in \mathcal{E}'} \frac{\partial}{\partial p_e} \mathbb{P}_p[A] \right| \leq \frac{1}{p(1-p)} \sqrt{\mathbb{P}_p[A] \mathbb{E}_p |\mathcal{W}_{\mathcal{E}'}|}$$

which is comparable to (2.14), although we believe it to be less general than the non-differential statement (2.13) (in particular, it does not seem straightforward to obtain (2.4) from (2.19)).

Consider Russo's formula

$$(2.20) \quad \left| \sum_{e \in \mathcal{E}'} \frac{\partial}{\partial p_e} \mathbb{P}_p[A] \right| = \frac{1}{p(1-p)} \left| \sum_{e \in \mathcal{E}'} \text{Cov}_p(\mathbb{1}_A, \mathbb{1}_{e \text{ open}}) \right|$$

and decompose the sum as

$$\sum_{e \in \mathcal{E}'} \text{Cov}_p(\mathbb{1}_A \mathbb{1}_{\text{Rev}(e)}, \mathbb{1}_{e \text{ open}}) + \sum_{e \in \mathcal{E}'} \text{Cov}_p(\mathbb{1}_A \mathbb{1}_{\text{Rev}(e)^c}, \mathbb{1}_{e \text{ open}})$$

where $\text{Rev}(e)$ denotes the event that e is revealed by the algorithm. One can check that $\mathbb{1}_A \mathbb{1}_{\text{Rev}(e)^c}$ is independent of $\mathbb{1}_{e \text{ open}}$ and so the second sum vanishes. Hence (2.20) is at most

$$\frac{1}{p(1-p)} \left| \sum_{e \in \mathcal{E}'} \text{Cov}_p(\mathbb{1}_A \mathbb{1}_{\text{Rev}(e)}, \mathbb{1}_{e \text{ open}}) \right| \leq \frac{1}{p(1-p)} \sqrt{\mathbb{P}_p[A] \mathbb{E}_p \left[\left(\sum_{e \in \mathcal{E}'} \mathbb{1}_{\text{Rev}(e)} (\mathbb{1}_{e \text{ open}} - p) \right)^2 \right]}$$

where we used the Cauchy-Schwartz inequality. For edges e and f introduce the event

$$\text{Rev}(e, f) := \text{Rev}(e) \cap \text{Rev}(f) \cap \{e \text{ is revealed before } f\}.$$

Again one checks that, for $e \neq f$, $\mathbb{1}_{\text{Rev}(e, f)} (\mathbb{1}_{e \text{ open}} - p)$ is independent of $\mathbb{1}_{f \text{ open}}$. Hence

$$\mathbb{E}_p \left[\left(\sum_{e \in \mathcal{E}'} \mathbb{1}_{\text{Rev}(e)} (\mathbb{1}_{e \text{ open}} - p) \right)^2 \right] = \sum_{e \in \mathcal{E}'} \mathbb{E}_p [\mathbb{1}_{\text{Rev}(e)} (\mathbb{1}_{e \text{ open}} - p)^2] \leq \sum_{e \in \mathcal{E}'} \mathbb{E}_p [\mathbb{1}_{\text{Rev}(e)}] = \mathbb{E}_p |\mathcal{W}_{\mathcal{E}'}|,$$

since off-diagonal terms are zero by independence, and we used that $(\mathbb{1}_{e \text{ open}} - p)^2 \leq 1$. Combining the above gives (2.19).

We can now complete the proof of Propositions 2.1 and 2.2:

Proof of Proposition 2.1. This follows directly from (2.13) (with $\mathcal{E}' = \mathcal{E}$) by considering the algorithm in Lemma 2.9 that determines $A_1(R)$ such that

$$\mathbb{E}_p |\mathcal{W}_{\mathcal{E}'}| = \sum_{e \in \mathcal{E}} \text{Rev}(e) \leq 2 \sum_{v \in \Lambda_R} \mathbb{P}_p[0 \longleftrightarrow v]. \quad \square$$

Proof of Proposition 2.2. For $d \geq 2$, recall the box $B_k(R)$ and its right half $B_k^+(R) := [0, R] \times [-kR, kR]^{d-1}$. Consider the algorithm in Lemma 2.9 that determines $\text{Cross}_k(R)$ such that

$$\sum_{e \in B_k^+(R)} \text{Rev}(e) \leq cR^d \max_{e \in B_k^+(R)} \text{Rev}(e) \leq 2cR^d \mathbb{P}_p[A_1(R)].$$

for $c = c(k) > 0$. By reflective symmetry in the vertical axis,

$$\frac{d}{dp} \mathbb{P}_p[\text{Cross}_k(R)] = \sum_{e \in B_k(R)} \frac{\partial}{\partial p_e} \mathbb{P}_p[\text{Cross}_k(R)] \leq 2 \sum_{e \in B_k^+(R)} \frac{\partial}{\partial p_e} \mathbb{P}_p[\text{Cross}_k(R)]$$

and hence, applying (2.14) (with $\mathcal{E}' = B_k^+(R)$)

$$\frac{d}{dp} \mathbb{P}_p[\text{Cross}_k(R)] \leq \frac{1}{\sqrt{p(1-p)}} \sqrt{\mathbb{E}_p |\mathcal{W}_{\mathcal{E}'}|} \leq \frac{\sqrt{2c} R^{d/2}}{\sqrt{p(1-p)}} \sqrt{\mathbb{P}_p[A_1(R)]}.$$

For $d = 2$, recall the top-right quarter $B_k^\dagger(R) := [0, R] \times [0, kR]$ of the box $B_k(R)$. Consider the algorithm in Lemma 2.9 that determines $\text{Cross}_k(R)$ such that

$$\sum_{e \in B_k^\dagger(R)} \text{Rev}(e) \leq cR^2 \max_{e \in B_k^\dagger(R)} \text{Rev}(e) \leq 2cR^2 \mathbb{P}_p[A_2(R)]$$

for $c = c(k) > 0$. Again by reflective symmetry (this time in both axes)

$$\frac{d}{dp} \mathbb{P}_p[\text{Cross}_k(R)] = \sum_{e \in B_k(R)} \frac{\partial}{\partial p_e} \mathbb{P}_p[\text{Cross}_k(R)] \leq 4 \sum_{e \in B_k^\dagger(R)} \frac{\partial}{\partial p_e} \mathbb{P}_p[\text{Cross}_k(R)],$$

and the result follows from (2.14) (with $\mathcal{E}' = B_k^\dagger(R)$) as in the previous case. \square

2.3. Proof of Proposition 2.3. We begin by introducing the OSSS inequality. Let $X = (X_i)_{i=1}^n$ be a finite sequence of independent random variables taking values in arbitrary probability spaces, and let A be an event. Then the *resampling influence* of X_i on A is

$$(2.21) \quad \text{Infl}(i) := \mathbb{P}[\mathbf{1}_{X \in A} \neq \mathbf{1}_{X^{(i)} \in A}]$$

where $X^{(i)}$ denotes X with the coordinate X_i resampled.

Theorem 2.15 (OSSS inequality [39]). *For every algorithm \mathcal{A} determining A ,*

$$\text{Var}(\mathbf{1}_A) \leq \frac{1}{2} \sum_{i=1}^n \text{Rev}(i) \text{Infl}(i)$$

where $\text{Rev}(i)$ is the revelation of X_i under \mathcal{A} .

Returning to the setting of Bernoulli percolation, combining the OSSS inequality with Russo's formula yields the following:

Proposition 2.16. *Let $p \in (0, 1)$, let A be an increasing event depending on a finite number of edges, let \mathcal{A} be an algorithm determining A , and let $\mathcal{E}' \subseteq \mathcal{E}$ be a subset of edges. Then*

$$\sum_{e \in \mathcal{E}'} \frac{\partial}{\partial p_e} \mathbb{P}_p[A] \geq \frac{4}{p(1-p)} \frac{\text{Var}_p[\mathbb{P}_p[A | \mathcal{F}_{\mathcal{E}'}]]}{\max_{e \in \mathcal{E}'} \text{Rev}(e)},$$

where $\mathcal{F}_{\mathcal{E}'}$ is the σ -algebra generated by the edges in \mathcal{E}' , and the revelations $\text{Rev}(e)$ are under \mathbb{P}_p .

Remark 2.17. If $p = 1/2$, the quantity $\text{Var}_p[\mathbb{P}_p[A | \mathcal{F}_{\mathcal{E}'}]]$ has an interpretation as the square-sum of the Fourier coefficients of $\mathbf{1}_A$ supported on non-empty subsets of \mathcal{E}' (see, e.g., [21]).

Proof. Let X_0 denote the vector of configurations on edges $e \notin \mathcal{E}'$, and $(X_e)_{e \in \mathcal{E}'}$ be the configuration on the remaining edges. Then by the OSSS inequality (Theorem 2.15) applied to $X = (X_0, (X_e)_{e \in \mathcal{E}'})$, and bounding the revelation of X_0 by 1,

$$\text{Var}_p(\mathbf{1}_A) \leq \frac{1}{2} \left(\text{Infl}(0) + \sum_{e \in \mathcal{E}'} \text{Rev}(e) \text{Infl}(e) \right)$$

where $\text{Infl}(0)$ and $\text{Infl}(e)$ are defined as in (2.21) under \mathbb{P}_p . Next observe that

$$\begin{aligned} \frac{1}{2} \text{Infl}(0) &= \frac{1}{2} \mathbb{E}_p[\mathbb{P}_p[\text{the outcome of } A \text{ changes when the edges } e \notin \mathcal{E}' \text{ are resampled} | \mathcal{F}_{\mathcal{E}'}]] \\ &= \mathbb{E}_p[\mathbb{P}_p[A | \mathcal{F}_{\mathcal{E}'}](1 - \mathbb{P}_p[A | \mathcal{F}_{\mathcal{E}'}])] = \mathbb{E}_p[\text{Var}_p[\mathbf{1}_A | \mathcal{F}_{\mathcal{E}'}]], \end{aligned}$$

and hence, by the law of total variance,

$$\text{Var}_p(\mathbf{1}_A) - \text{Infl}(0)/2 = \text{Var}_p(\mathbf{1}_A) - \mathbb{E}_p[\text{Var}_p[\mathbf{1}_A | \mathcal{F}_{\mathcal{E}'}]] = \text{Var}_p[\mathbb{P}_p[A | \mathcal{F}_{\mathcal{E}'}]].$$

This yields the following extension of the OSSS inequality

$$(2.22) \quad \text{Var}_p[\mathbb{P}_p[A | \mathcal{F}_{\mathcal{E}'}]] \leq \frac{1}{2} \sum_{e \in \mathcal{E}'} \text{Rev}(e) \text{Infl}(e) \leq \frac{\max_{e \in \mathcal{E}'} \text{Rev}(e)}{2} \sum_{e \in \mathcal{E}'} \text{Infl}(e).$$

We deduce the result by combining with Russo's formula for increasing events, namely

$$\sum_{e \in \mathcal{E}'} \frac{\partial}{\partial p_e} \mathbb{P}_p[A] = \frac{2}{p(1-p)} \sum_{e \in \mathcal{E}'} \text{Infl}(e)$$

(which coincides with (2.20) since $\text{Cov}_p(\mathbb{1}_A, \mathbb{1}_{e \text{ open}}) = 2\text{Infl}(e)$ for increasing A). \square

Proof of Proposition 2.3. Recall the top-right quarter $B_k^\dagger(R)$ and set $\mathcal{E}' = B_k^\dagger(R)$. We claim

$$(2.23) \quad \text{Var}_p[\mathbb{P}_p[\text{Cross}_k(R) \mid \mathcal{F}_{\mathcal{E}'}]] \geq \mathbb{P}_p[\text{Cross}_{1/(8k)}(kR)]^4 (1 - \mathbb{P}_p[\text{Cross}_{8k}(R/8)])^2.$$

Assuming (2.23), the statement follows by applying Proposition 2.16 to the algorithm in Lemma 2.9 that determines $\text{Cross}_k(R)$ whose revealments on $B_k^\dagger(R)$ are bounded by $2\mathbb{P}_p[A_2(R)]$.

To prove (2.23), remark first that, for any event A and sub- σ -algebra \mathcal{G} ,

$$(2.24) \quad \begin{aligned} \text{Var}[\mathbb{P}[A \mid \mathcal{G}]] &= \mathbb{E}[(\mathbb{P}[A \mid \mathcal{G}] - \mathbb{P}[A])^2] \geq \sup_{A' \in \mathcal{G}} \mathbb{E}[(\mathbb{P}[A \mid \mathcal{G}] - \mathbb{P}[A])^2 \mathbb{1}_{A'}] \\ &\geq \sup_{A' \in \mathcal{G}} \mathbb{P}[A'] (\mathbb{P}[A \mid A'] - \mathbb{P}[A])^2 \end{aligned}$$

where the second inequality is Jensen's. Hence it is enough to construct an event A' , measurable with respect to the configuration on the top-right quarter, such that $\text{Cross}_k(R)$ becomes substantially more likely if A' occurs (see Figure 1 for an illustration).

Define

$$\begin{aligned} A' := & \left\{ \{R/4\} \times [R/4, R/2] \xleftrightarrow{[R/4, R] \times [R/4, R/2]} \{R\} \times [R/4, R/2] \right\} \\ & \cap \left\{ [R/4, R/2] \times \{R/4\} \xleftrightarrow{[R/4, R/2] \times [R/4, kR]} [R/4, R/2] \times \{kR\} \right\}, \end{aligned}$$

which is measurable with respect to $\mathcal{F}_{\mathcal{E}'}$. By the FKG inequality and symmetry (and an obvious event inclusion), $\mathbb{P}_p[A'] \geq \mathbb{P}_p[\text{Cross}_{1/(8k)}(kR)]^2$. Define also the events

$$\begin{aligned} B_1 := & \left\{ \{-R\} \times [R/2, 3R/4] \xleftrightarrow{[-R, R/2] \times [R/2, 3R/4]} \{R/2\} \times [R/2, 3R/4] \right\} \\ B_2 := & \left\{ \{3R/4\} \times [-kR, kR] \xleftrightarrow{[3R/4, R] \times [-kR, kR]} \{R\} \times [-kR, kR] \right\} \end{aligned}$$

which are defined on disjoint domains and are translated copies of, respectively, $\text{Cross}_{1/6}(3R/2)$ and $\text{Cross}_{8k}(R/8)$. Finally, define

$$C := \left\{ \{-R\} \times [-kR, kR] \xleftrightarrow{B_k(R)} (\{R/2\} \times [R/2, kR]) \cup ([R/2, R] \times \{R/2\}) \cup (\{R\} \times [-kR, R/2]) \right\}$$

and observe (i) $\text{Cross}_k(R) \subseteq C$, (ii) on A' , $\text{Cross}_k(R) = C$, and (iii) $B_1 \cap B_2^c \subseteq C \setminus \text{Cross}_k(R)$. Hence

$$\begin{aligned} \mathbb{P}_p[\text{Cross}_k(R) \mid A'] - \mathbb{P}_p[\text{Cross}_k(R)] &= \mathbb{P}_p[C \mid A'] - \mathbb{P}_p[\text{Cross}_k(R)] \geq \mathbb{P}_p[C] - \mathbb{P}_p[\text{Cross}_k(R)] \\ &= \mathbb{P}_p[C \setminus \text{Cross}_k(R)] \geq \mathbb{P}_p[B_1 \cap B_2^c] = \mathbb{P}_p[B_1] (1 - \mathbb{P}_p[B_2]) \\ &\geq \mathbb{P}_p[\text{Cross}_{1/(8k)}(kR)] (1 - \mathbb{P}_p[\text{Cross}_{8k}(R/8)]), \end{aligned}$$

where the second step is by the FKG inequality, the penultimate step uses disjoint domains, and the final step is an obvious event inclusion. Applying (2.24) (with $A = \text{Cross}_k(R)$ and $\mathcal{G} = \mathcal{F}_{\mathcal{E}'}$) gives (2.23). \square

2.4. A general bound for revealments of increasing events. Combining Propositions 2.10 and 2.16 yields a general lower bound on the revealments of increasing events:

Proposition 2.18. *In the setting of Proposition 2.16 (in particular the event A is increasing),*

$$\max_{e \in \mathcal{E}'} \text{Rev}(e) \geq \frac{(4 \text{Var}_p[\mathbb{P}_p[A \mid \mathcal{F}_{\mathcal{E}'}]])^{2/3}}{(p(1-p) \mathbb{P}_p[A \mid \mathcal{E}'])^{1/3}}.$$




FIGURE 1. An illustration of the proof of (2.23). The first panel shows the event A' . The second illustrates how, on A' , the event C is equivalent to $\text{Cross}_k(R)$. The third shows how the crossing given by B_1 , combined with the dual crossing given by B_2^c , realises C but not $\text{Cross}_k(R)$.

Proof. By (2.14) and Proposition 2.16 we have

$$\begin{aligned} \frac{4}{p(1-p)} \frac{\text{Var}_p[\mathbb{P}_p[A | \mathcal{F}_{\mathcal{E}'}]]}{\max_{e \in \mathcal{E}'} \text{Rev}(e)} &\leq \sum_{e \in \mathcal{E}'} \frac{\partial}{\partial p_e} \mathbb{P}_p[A] \\ &\leq \frac{1}{\sqrt{p(1-p)}} \sqrt{\mathbb{P}_p[A] \mathbb{E}_p[|\mathcal{W}_{\mathcal{E}'}|]} \leq \frac{1}{\sqrt{p(1-p)}} \sqrt{\mathbb{P}_p[A] |\mathcal{E}'| \max_{e \in \mathcal{E}'} \text{Rev}(e)} \end{aligned}$$

and rearranging gives the result. \square

Proposition 2.18 generalises a result from [8] which considered the case $p = 1/2$ and \mathcal{E}' is the set of edges on which A depends; denoting by n the cardinality of this set of edges, this gives

$$(2.25) \quad \max_e \text{Rev}(e) \geq \frac{(8 \text{Var}_{1/2}[\mathbb{1}_A])^{2/3}}{(\mathbb{P}_{1/2}[A]n)^{1/3}}$$

which is comparable to [8, Theorem 2 (part 2)], although (2.25) has a stronger constant.

3. LEVEL SET PERCOLATION OF GAUSSIAN FIELDS

We now establish our main results in the case of Gaussian percolation; the proof will closely follow the approach for Bernoulli percolation in Section 2. For $k, R > 0$, recall the box $B_k(R) := [-R, R] \times [-kR, kR]^{d-1}$, which we now view as a subset of \mathbb{R}^d . Then define

$$\text{Cross}_k(R) := \left\{ \{-R\} \times [-kR, kR]^{d-1} \overset{B_k(R)}{\longleftrightarrow} \{R\} \times [kR, kR]^{d-1} \right\}$$

where

$$\{A \overset{E}{\longleftrightarrow} B\} := \{\text{there exists a path in } \{f \geq 0\} \cap E \text{ that intersects } A \text{ and } B\}.$$

For $0 \leq r \leq R$ define

$$A_1(r, R) := \{\Lambda_r \longleftrightarrow \partial\Lambda_R\} \quad \text{and} \quad A_2(r, R) := \{\Lambda_r \overset{\mathbb{R}^d}{\longleftrightarrow} \partial\Lambda_R\}$$

where

$$\{A \overset{E}{\longleftrightarrow} B\} = \{A \overset{E}{\longleftrightarrow} B\} \cap \{\text{there exists a path in } \{f \leq 0\} \cap E \text{ that intersects } A \text{ and } B\},$$

and $\{A \overset{\mathbb{R}^d}{\longleftrightarrow} B\} = \{A \overset{\mathbb{R}^d}{\longleftrightarrow} B\}$. By continuity of f , if $d = 2$ then $A_2(r, R)$ could equivalently be defined as

$$A_2(r, R) = \{\text{there exists a path in } \{f = 0\} \text{ that intersects } \Lambda_r \text{ and } \partial\Lambda_R\}.$$

We make the elementary observation that $\mathbb{P}_{\ell_c}[A_2(r, R)] \leq \mathbb{P}_{\ell_c}[A_1(r, R)]$, and moreover if $d = 2$ and under (POS') (so the FKG inequality is available; c.f. (2.1)) then

$$(3.1) \quad \mathbb{P}_{\ell_c}[A_2(r, R)] \leq \mathbb{P}_{\ell_c}[A_1(r, R)]^2.$$

We now state the analogues of Propositions 2.1–2.3, which concern Gaussian fields $f = q \star W$ with *finite-range dependence*. Recall the *Dini derivatives*, defined for $f : \mathbb{R} \rightarrow \mathbb{R}$ as

$$\frac{d^+}{dx} f(x) = \limsup_{\varepsilon \downarrow 0} \frac{f(x + \varepsilon) - f(x)}{\varepsilon} \quad \text{and} \quad \frac{d^-}{dx} f(x) = \liminf_{\varepsilon \downarrow 0} \frac{f(x + \varepsilon) - f(x)}{\varepsilon}.$$

Proposition 3.1. *Suppose $f = q \star W$ satisfies Assumption 1.4 and (POS)–(BOU), and let $r > 0$ be such that q is supported on Λ_r . Then for $\ell \leq \ell'$ and $R \geq r \geq 1$,*

$$\mathbb{P}_{\ell'}[A_1(1, R)] - \mathbb{P}_{\ell}[A_1(1, R)] \leq \frac{r^{d/2}(\ell' - \ell)}{\int q} \sqrt{\mathbb{P}_{\ell'}[A_1(1, R)] \sum_{v \in r\mathbb{Z}^d \cap \Lambda_{R+2r}} \mathbb{P}_{\ell}[\Lambda_1 \longleftrightarrow v + \Lambda_{6r}]}.$$

Proposition 3.2. *Suppose $f = q \star W$ satisfies Assumption 1.4 and (POS)–(BOU), and let $r > 0$ be such that q is supported on Λ_r . Then for $k \geq 1$ there exists $c = c(k) > 0$ such that, for $\ell \in \mathbb{R}$ and $R \geq 4r > 0$,*

$$\frac{d^+}{d\ell} \mathbb{P}_{\ell}[\text{Cross}_k(R)] \leq \frac{cR^{d/2}}{\int q} \begin{cases} \sqrt{\mathbb{P}_{\ell}[A_2(2r, R - 2r)]} & d = 2, \\ \sqrt{\mathbb{P}_{\ell}[A_1(2r, R - 2r)]} & d \geq 2. \end{cases}$$

Proposition 3.3. *Suppose $f = q \star W$ satisfies Assumption 1.4 and (POS)–(BOU), and let $r > 0$ be such that q is supported on Λ_r . Then for $k \geq 1$ there exists $c = c(k) > 0$ such that, for $\ell \in \mathbb{R}$ and $R \geq 8r > 0$,*

$$(3.2) \quad \frac{d^-}{d\ell} \mathbb{P}_{\ell}[\text{Cross}_k(R)] \geq \frac{c}{\|q\|_2} \frac{\mathbb{P}_{\ell}[\text{Cross}_k(R)](1 - \mathbb{P}_{\ell}[\text{Cross}_k(R)])}{\frac{r}{R} \sum_{i=2}^{R/r} \mathbb{P}_{\ell}[A_1(2r, ir)]},$$

and, if $d = 2$ and (POS') holds,

$$(3.3) \quad \frac{d^-}{d\ell} \mathbb{P}_{\ell}[\text{Cross}_k(R)] \geq \frac{c}{\|q\|_2} \frac{\mathbb{P}_{\ell}[\text{Cross}_{1/(8k)}(kR)]^4 (1 - \mathbb{P}_{\ell}[\text{Cross}_{8k}(R/8)])^2}{\mathbb{P}_{\ell}[A_2(2r, R - 2r)]}.$$

We prove Propositions 3.1–3.3 later in the section; for now we establish our main result Theorems 1.5. For this we need two auxiliary results; these are rather standard, but we give details on their proof at the end of the section. The first is the analogue of Lemma 2.4:

Lemma 3.4. *Suppose $f = q \star W$ satisfies Assumption 1.4 with parameter $\beta > d$.*

(1) *There exists $\delta > 0$ and $\ell' = \ell'(R) \leq \ell_c$ such that, for $R \geq 1$,*

$$\mathbb{P}_{\ell'}[\text{Cross}_5(R)] = \delta.$$

(2) *(RSW) Let $d = 2$ and $k > 0$ and suppose that (POS') holds. Then there exists $\delta > 0$ such that, for $R \geq 1$,*

$$\mathbb{P}_{\ell_c}[\text{Cross}_k(R)] \in (\delta, 1 - \delta).$$

The second allows us to compare a Gaussian field with an approximation that satisfies (BOU). Fix a smooth symmetric cutoff function $\varphi : \mathbb{R} \rightarrow [0, 1]$ such that $\varphi(x) = 1$ for $\|x\|_{\infty} \leq 1/2$, $\varphi(x) = 0$ for $\|x\|_{\infty} \geq 1$. For $r > 0$ define

$$(3.4) \quad f_r := q_r \star W$$

where $q_r(x) := q(x)\varphi(|x|/r)$. Note that q_r is supported on Λ_r , and also, since $q \in L^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$, as $r \rightarrow \infty$,

$$(3.5) \quad \|q_r\|_2 \rightarrow \|q\|_2 \quad \text{and} \quad \int q_r \rightarrow \int q.$$

Remark that if either Assumption 1.4 or (POS) holds for f then it holds for f_r (on the other hand, deducing this for (POS') seems difficult but we do not need it). In particular, as discussed in Section 1.1, if $d = 2$ and Assumption 1.4 holds for f then $\ell_c(f) = \ell_c(f_r) = 0$.

The following lemma, essentially taken from [38], allows us to compare f and f_r :

Lemma 3.5. *Suppose $f = q \star W$ satisfies Assumption 1.4 with parameter $\beta > d$ and (POS). Then there exist $c_1, c_2 > 0$ such that, for $r, R \geq 2$, increasing event A measurable with respect to $f|_{B(R)}$, and $\ell \in \mathbb{R}$,*

$$|\mathbb{P}_\ell[f \in A] - \mathbb{P}_\ell[f_r \in A]| \leq c_1 (R^{d/2} (\log R) r^{-(\beta-d/2)} + e^{-c_2 (\log R)^2}).$$

The same conclusion holds if A is the intersection of one increasing and one decreasing event which are both measurable with respect to $f|_{B(R)}$.

We are now ready to prove Theorem 1.5:

Proof of Theorem 1.5. In the proof $c > 0$ are constants that depend only on f (and the choice of the cutoff function φ in (3.4)) and may change from line to line. The bound $\eta_1 \leq d - 1$, and also $\eta_1 \leq 1/2$ if $d = 2$ and (POS') holds, are rather classical; in fact they are true for any $\beta > d$. For the former, combining $\mathbb{P}_{\ell_c}[\text{Cross}_5(R)] \geq \delta$ (the first statement of Lemma 3.4) with the union bound applied along the hyperplane $\{0\} \times [-kR, kR]$ gives $\mathbb{P}_{\ell_c}[A_1(1, R)] \geq cR^{-(d-1)}$. For the latter, by combining the RSW estimates (the second statement of Lemma 3.4) with Lemma 3.5 one can deduce (see [4, 38] for similar arguments)

$$\mathbb{P}_{\ell_c}[\{-R\} \times [-R, R] \stackrel{B_1(R)}{\iff} \{R\} \times [-R, R]] \geq c(1 - R^{1-(\beta-1)} (\log R)) \geq c/2$$

for sufficiently large R . By the union bound applied along $\{0\} \times [-R, R]$ this implies $\mathbb{P}_{\ell_c}[A_2(1, R)] \geq cR^{-1}$, and given (3.1) we see that $\mathbb{P}_{\ell_c}[A_1(1, R)] \geq cR^{-1/2}$.

We now prove the remaining bounds, beginning with the first statement. Fix $1 > \alpha > \frac{d/2}{\beta-d/2}$ and $\eta_1 > \eta^* > 0$ (if $\eta_1 = 0$ there is nothing to prove). Then by monotonicity in ℓ , the union bound, and the definition of η_1 ,

$$\mathbb{P}_\ell[A_1(r, R)] \leq \mathbb{P}_{\ell_c}[A_1(r, R)] \leq cr^{d-1} \mathbb{P}_{\ell_c}[A_1(1, R-r)] \leq r^{d-1} R^{-\eta^*}$$

for all $\ell \leq \ell_c$, R sufficiently large, and $r \in [1, R/2]$. Set $r = R^\alpha$. Then by an integral comparison,

$$\begin{aligned} \frac{r}{R} \sum_{i=2}^{R/r} \mathbb{P}_\ell[A_1(2r, ir)] &\leq cr^{-\eta^*+(d-1)} \times \frac{r}{R} \sum_{i=2}^{R/r} i^{-\eta^*} \\ &\leq cr^{-\eta^*+(d-1)} (R/r)^{-\min\{\eta^*, 1\}} (\log(R/r)) \\ &\leq c(\log R) (R^{\alpha(d-1-\eta^*)-(1-\alpha)\min\{\eta^*, 1\}}) \end{aligned}$$

for $\ell \leq \ell_c$ and large R . Consider the field f_r defined in (3.4). By Lemma 3.5,

$$\mathbb{P}_\ell[f_r \in A_1(r', R)] \leq \mathbb{P}_\ell[A_1(r', R)] + cR^{d/2-\alpha(\beta-d/2)} (\log R) + ce^{-c(\log R)^2}$$

for $\ell \leq \ell_c$ and $2 \leq r' \leq R$, and hence

$$\frac{r}{R} \sum_{i=2}^{R/r} \mathbb{P}_\ell[f_r \in A_1(2r, ir)] \leq c(\log R) (R^{\alpha(d-1-\eta^*)-(1-\alpha)\min\{\eta^*, 1\}} + R^{d/2-\alpha(\beta-d/2)})$$

for $\ell \leq \ell_c$ and large R . Moreover, by Lemma 3.4 there are $\delta > 0$ and $\ell' = \ell'(R) \leq \ell_c$ such that $\mathbb{P}_{\ell'}[\text{Cross}_5(R)] = \delta$. Hence, again by Lemma 3.5,

$$\mathbb{P}_{\ell'}[f_r \in \text{Cross}_5(R)] (1 - \mathbb{P}_{\ell'}[f_r \in \text{Cross}_5(R)]) \geq \delta(1 - \delta) - cR^{d/2-\alpha(\beta-d/2)} (\log R) \geq \delta(1 - \delta)/2$$

for large R , where we used that $\alpha > \frac{d/2}{\beta-d/2}$.

We now apply Propositions 3.2 and 3.3 to the field f_r at the sequence of levels $\ell'(R) \leq \ell_c$. First, by (3.2) (recalling (3.5))

$$\begin{aligned} (3.6) \quad \frac{d}{d\ell} \mathbb{P}_\ell[f_r \in \text{Cross}_5(R)] \Big|_{\ell=\ell'} &\geq \frac{c\delta(1-\delta)}{\|q_r\|_2} \left(\frac{r}{R} \sum_{i=2}^{R/r} \mathbb{P}_{\ell'}[f_r \in A_1(2r, ir)] \right)^{-1} \\ &\geq c(\log R)^{-1} (R^{\alpha(d-1-\eta^*)-(1-\alpha)\min\{\eta^*, 1\}} + R^{d/2-\alpha(\beta-d/2)})^{-1} \end{aligned}$$

for large R . Similarly, by Proposition 3.1,

$$\begin{aligned}
(3.7) \quad \frac{d}{d\ell} \mathbb{P}_{\ell} [f_r \in \text{Cross}_5(R)] \Big|_{\ell=\ell'} &\leq cR^{d/2} \left(\mathbb{P}_{\ell'} [f_r \in A_1(2r, R)] \right)^{1/2} \\
&\leq cR^{d/2} \left(R^{-\eta^* + \alpha(d-1)} + R^{d/2 - \alpha(\beta - d/2)} (\log R) \right)^{1/2} \\
&\leq c\sqrt{\log R} \left(R^{d/2 - \eta^*/2 + \alpha(d-1)/2} + R^{3d/4 - \alpha(\beta - d/2)/2} \right)
\end{aligned}$$

for large R , where we used that $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$ for $a, b > 0$. Comparing (3.6) and (3.7) and expanding the brackets we deduce that at least one of the exponents

$$\begin{aligned}
E_1 &:= (3d/4 - \alpha(\beta - d/2)/2) + (d/2 - \alpha(\beta - d/2)) \\
E_2 &:= (d/2 - \eta^*/2 + \alpha(d-1)/2) + (d/2 - \alpha(\beta - d/2)) \\
E_3 &:= (d/2 - \eta^*/2 + \alpha(d-1)/2) + (\alpha(d-1 - \eta^*) - (1-\alpha) \min\{\eta^*, 1\}) \\
E_4 &:= (3d/4 - \alpha(\beta - d/2)/2) + (\alpha(d-1 - \eta^*) - (1-\alpha) \min\{\eta^*, 1\})
\end{aligned}$$

must be non-negative. The first is equivalent to $\alpha \leq \frac{5d}{6(\beta - d/2)}$. The second implies that $\eta^* \leq \frac{d}{3} + \alpha(d-1)$, assuming that $\alpha > \frac{5d}{6(\beta - d/2)}$. The third is equivalent to either $\eta^* \leq \frac{d}{3} + \alpha(d-1)$ (if $\eta^* \leq 1$) or $\eta^* \leq \frac{d-2+\alpha(3d-1)}{1+2\alpha}$ (if $\eta^* > 1$). Finally, the fourth implies either $\eta^* \leq \frac{d}{3} + \alpha(d-1)$ (if $\eta^* \leq 1$, assuming that $\alpha > \frac{5d}{6(\beta - d/2)}$) or $\alpha \leq \frac{3d-4}{2\beta-5d+4}$ (if $\eta^* > 1$). One can check that, since $d \geq 3$, $\frac{5d}{6(\beta - d/2)} < \frac{3d-4}{2\beta-5d+4}$ and $\frac{d}{3} + \alpha(d-1) < \frac{d-2+\alpha(3d-1)}{1+2\alpha}$. Hence we conclude that if $\alpha > \frac{3d-4}{2\beta-5d+4}$ then $\eta^* \leq \frac{d-2+\alpha(3d-1)}{1+2\alpha}$. Sending $\alpha \rightarrow \frac{3d-4}{2\beta-5d+4}$ from above gives the result.

The proof of the remaining statements are similar, and closer to the arguments in Section 2. For the second statement, fix $1 > \alpha > \frac{3d/2-1}{\beta-d/2}$ and $\eta_1 > \eta^* > 0$. As in the proof of the first statement,

$$(3.8) \quad \mathbb{P}_{\ell_c} [A_1(2r, R)] \leq r^{d-1} R^{-\eta^*}$$

for large R and $r \in [1, R/4]$. Now let $r = R^\alpha$. Since we have the a priori bound $\mathbb{P}_{\ell_c} [A_1(1, R)] \geq cR^{-(d-1)}$ (from the start of the proof), by Lemma 3.5

$$\begin{aligned}
(3.9) \quad |\mathbb{P}_{\ell_c} [f_r \in A_1(r', R')] - \mathbb{P}_{\ell_c} [A_1(r', R')]| &\leq cR^{d/2 - \alpha(\beta - d/2)} (\log R) + ce^{-c(\log R)^2} \\
&\leq \mathbb{P}_{\ell_c} [A_1(r', R')]/2
\end{aligned}$$

for large R and $1 \leq r' \leq R' \leq R$, where we used that $d/2 - \alpha(\beta - d/2) \leq -(d-1)$ by the definition of α . Observe next that, for $|x| \geq 18r$, the event $\Lambda_1 \longleftrightarrow x + \Lambda_{6r}$ implies the occurrence of the events

$$\{A_1(1, |x|_\infty/3)\} \quad \text{and} \quad \{x + A_1(6r, |x|_\infty/3)\},$$

which are measurable with respect to disjoint domains separated by distance r . Since f_r is r -dependent, for large R and $18r \leq |x| \leq R + 2r$ this implies

$$\begin{aligned}
\mathbb{P}_{\ell_c} [f_r \in \Lambda_{2r} \longleftrightarrow x + \Lambda_{6r}] &\leq \mathbb{P}_{\ell_c} [f_r \in A_1(1, |x|_\infty/3)] \mathbb{P}_{\ell_c} [f_r \in A_1(6r, |x|_\infty/3)] \\
&\leq 4\mathbb{P}_{\ell_c} [A_1(1, |x|_\infty/3)] \mathbb{P}_{\ell_c} [A_1(6r, |x|_\infty/3)] \\
&\leq cr^{d-1} |x|_\infty^{-2\eta^*}
\end{aligned}$$

where we used (3.8) and then (3.9). Then by an integral comparison, for large R ,

$$\begin{aligned}
\sum_{v \in r\mathbb{Z}^d \cap \Lambda_{R+2r}} \mathbb{P}_{\ell} [f_r \in \Lambda_{2r} \longleftrightarrow v + \Lambda_{6r}] &\leq c + cr^{d-1} \sum_{v \in r\mathbb{Z}^d \cap \Lambda_{R+2r} \setminus \{0\}} |v|_\infty^{-2\eta^*} \\
&\leq c + cr^{d-1} \max\{r^{-2\eta^*} (R/r)^{d-2\eta^*} (\log(R/r)), 1\} \\
&\leq cR^{\max\{\alpha(d-1), -\alpha+d-2\eta^*\}} (\log R).
\end{aligned}$$

Next define, for large R ,

$$\ell'(R) = \inf\{\ell > \ell_c : \mathbb{P}_{\ell} [A_1(1, R)] = 2\mathbb{P}_{\ell_c} [A_1(1, R)]\},$$

which exists by continuity in ℓ (see Lemma 3.13), and since $\mathbb{P}_{\ell_c}[A_1(1, R)] > 0$ and

$$\mathbb{P}_{\ell}[A_1(1, R)] \geq \mathbb{P}\left[\sup_{x \in \Lambda_R} f(x) \leq \ell\right] \rightarrow 1$$

as $\ell \rightarrow \infty$. By the mean-field lower bound (1.1), for large R ,

$$(3.10) \quad \mathbb{P}_{\ell_c}[A_1(1, R)]/2 = \mathbb{P}_{\ell'}[A_1(1, R)]/4 \geq \theta(\ell')/4 \geq c(\ell' - \ell_c),$$

where we used that $\ell'(R) \rightarrow \ell_c$ as $R \rightarrow \infty$, since otherwise

$$\limsup_{R \rightarrow \infty} \mathbb{P}_{\ell_c}[A_1(1, R)] \geq \limsup_{R \rightarrow \infty} \theta(\ell'(R))/2 > 0$$

which contradicts (3.8). Similarly to (3.9) we also have

$$\begin{aligned} |\mathbb{P}_{\ell'}[f_r \in A_1(1, R)] - \mathbb{P}_{\ell'}[A_1(1, R)]| &\leq cR^{d/2 - \alpha(\beta - d/2)}(\log R) + ce^{-c(\log R)^2} \\ &\leq \mathbb{P}_{\ell_c}[A_1(1, R)] = \mathbb{P}_{\ell'}[A_1(1, R)]/2. \end{aligned}$$

Then applying Proposition 3.1 to the field f_r , for large R ,

$$\begin{aligned} \mathbb{P}_{\ell_c}[A_1(1, R)] &= \mathbb{P}_{\ell'}[A_1(1, R)] - \mathbb{P}_{\ell_c}[A_1(1, R)] \leq 2(\mathbb{P}_{\ell'}[f_r \in A_1(1, R)] - \mathbb{P}_{\ell_c}[f_r \in A_1(1, R)]) \\ &\leq \frac{2r^{d/2}(\ell' - \ell_c)}{\int q} \sqrt{\mathbb{P}_{\ell'}[A_1(1, R)] \sum_{v \in r\mathbb{Z}^d \cap \Lambda_{R+2r}} \mathbb{P}_{\ell_c}[\Lambda_{2r} \longleftrightarrow v + \Lambda_{6r}]} \\ &\leq \frac{2(\ell' - \ell_c)}{\int q} R^{\alpha d/2} \sqrt{R^{-\eta^*} R^{\max\{0, \alpha(d-1), -\alpha + d - 2\eta^*\}} (\log R)}. \end{aligned}$$

Comparing with (3.10) implies that $\alpha d - \eta^* + \max\{\alpha(d-1), -\alpha + d - 2\eta^*\} \geq 0$, and so $\eta^* \leq \max\{d/3 + \alpha(d-1)/3, \alpha(2d-1)\}$, and sending $\alpha \rightarrow \frac{3d/2-1}{\beta-d/2}$ from above gives the result.

Finally, consider the third statement. Fix $1 > \alpha > \frac{5}{3(\beta-1)}$ and $r = R^\alpha$. By the RSW estimates (the second statement of Lemma 3.4) and Lemma 3.5,

$$\mathbb{P}_{\ell_c}[f_r \in \text{Cross}_5(R)](1 - \mathbb{P}_{\ell_c}[f_r \in \text{Cross}_5(R)]) \leq c - cR^{1-\alpha(\beta-1)}(\log R) < c/2$$

for large R . Then by (3.3) and Proposition 3.2 we have, for large R ,

$$c\mathbb{P}_{\ell_c}[f_r \in A_2(2r, R-2r)]^{-1} \leq \frac{d}{d\ell} \mathbb{P}_{\ell}[f_r \in \text{Cross}_5(R)] \Big|_{\ell=\ell_c} \leq cR\sqrt{\mathbb{P}_{\ell_c}[f_r \in A_2(2r, R-2r)]}$$

which gives $\mathbb{P}_{\ell_c}[f_r \in A_2(2r, R-2r)] \geq cR^{-2/3}$ for large R . Applying the union bound and Lemma 3.5 (valid since $A_2(3\sqrt{2}r, R)$ is the intersection of an increasing and a decreasing event) yields

$$\mathbb{P}_{\ell_c}[A_2(1, R-2r)] \geq cr^{-1}\mathbb{P}_{\ell_c}[A_2(2r, R-2r)] \geq cR^{-\alpha}(R^{-2/3} - R^{1-\alpha(\beta-1)}(\log R)).$$

Sending $\alpha \rightarrow \frac{5}{3(\beta-1)}$ from above gives that, for every $\varepsilon > 0$,

$$(3.11) \quad \mathbb{P}_{\ell_c}[A_2(1, R)] \geq c_2 R^{-2/3-5/(3(\beta-1))-\varepsilon}.$$

for $c_2 = c_2(\varepsilon) > 0$ and large R . Hence by the FKG inequality (see (3.1))

$$(3.12) \quad \mathbb{P}_{\ell_c}[A_1(1, R)] \geq (\mathbb{P}_{\ell_c}[A_2(1, R)])^{1/2} \geq c_3 R^{-1/3-5/(6(\beta-1))-\varepsilon/2}$$

for $c_3 = c_3(\varepsilon) > 0$ and large R , which gives the result. \square

3.1. Randomised algorithms. Recall from Definition 2.8 that (*randomised*) *algorithms* are adapted procedures that sequentially reveal a subset of random variables $X = (X_i)$ and return a value. In the Bernoulli case we took $X_e = \mathbf{1}_{e \text{ open}}$ indexed by the edges of \mathbb{Z}^d . In the Gaussian setting we will instead decompose the field $f = \sum f_S$ into independent components indexed by a partition of \mathbb{R}^d into disjoint boxes S , and take $X_S = f_S$.

Fix a constant $s > 0$ and partition \mathbb{R}^d into boxes $S \in \mathcal{S}_s$ which are translations of $[0, s]^d$ by the lattice $s\mathbb{Z}^d$. Then one can decompose $f = \sum_{S \in \mathcal{S}_s} f_S$ where

$$f_S(\cdot) = (q \star W|_S)(\cdot) = \int_{y \in \mathbb{R}^d} q(\cdot - y) dW|_S(y) = \int_{y \in S} q(\cdot - y) dW(y)$$

are independent centred almost surely continuous Gaussian fields,⁵ and $W|_S = W\mathbb{1}_S$ is the restriction of the white noise W to S .⁶ We then introduce the collection \mathcal{A}_s of algorithms that adaptively reveal a subset of $(f_S)_{S \in \mathcal{S}_s}$. For brevity we say that a box $S \in \mathcal{S}_s$ is *revealed* if f_S (or equivalently $W|_S$) is revealed. As in Definition 2.8, $\text{Rev}(S)$ is the probability that S is revealed.

In the case that f satisfies (BOU), we make the important distinction between the set of boxes that are *revealed* by an algorithm, and the set $V \subset \mathbb{R}^d$ on which the field f is *determined* by an algorithm. More precisely, for $V \subset \mathbb{R}^d$ and a set of boxes $\mathcal{P} \subset \mathcal{S}_s$, we say that f is *determined on V using \mathcal{P}* if $f|_V = (\sum_{S \in \mathcal{P}} f_S)|_V$, or equivalently, if $(\bigcup_{S \in \mathcal{S}_s \setminus \mathcal{P}} \text{Supp}(q \star \mathbb{1}_S)) \cap V = \emptyset$.

We now state the analogue of Lemma 2.9. Recall the box $B_k(R) = [-R, R] \times [-kR, kR]^{d-1}$, its right half $B_k^+(R) = [0, R] \times [-kR, kR]^{d-1}$, and in the case $d = 2$, its top-right quarter $B_k^\dagger(R) = [0, R] \times [0, kR]$, all considered as subsets of \mathbb{R}^d .

Lemma 3.6. *Suppose $f = q \star W$ satisfies Assumption 1.4 and (BOU), and let $r > 0$ be such that q is supported on Λ_r . Then for every $\ell \in \mathbb{R}$ and $R \geq r$ there is an algorithm in \mathcal{A}_r determining $A(1, R)$ such that, under \mathbb{P}_ℓ ,*

$$\sum_{S \in \mathcal{S}_r} \text{Rev}(S) \leq \sum_{v \in r\mathbb{Z}^d \cap \Lambda_{R+2r}} \mathbb{P}_\ell[\Lambda_1 \longleftrightarrow v + \Lambda_{6r}].$$

Moreover for every $k \geq 1$, $\ell \in \mathbb{R}$, and $R \geq 4r > 0$, there are algorithms in \mathcal{A}_r determining $\text{Cross}_k(R)$ such that, under \mathbb{P}_ℓ , these algorithms satisfy respectively

$$\max_{S \in \mathcal{S}_r} \text{Rev}(S) \leq \frac{4r}{R} \sum_{i=2}^{R/r} \mathbb{P}_\ell[A_1(2r, ir)], \quad \max_{S \in \mathcal{S}_r: d(S, B_k^+(R)) < r} \text{Rev}(S) \leq \mathbb{P}_\ell[A_1(2r, R - 2r)],$$

and, if $d = 2$,

$$\max_{S \in \mathcal{S}_r: d(S, B_k^\dagger(R)) < r} \text{Rev}(S) \leq \mathbb{P}_\ell[A_2(2r, R - 2r)].$$

Proof. We begin by introducing some notation. Distinct boxes $S, S' \in \mathcal{S}_r$ are *adjacent* if their closures have non-empty intersection. For a set of boxes $\mathcal{P} \subset \mathcal{S}_r$ define its *outer boundary*

$$\partial^+ \mathcal{P} := \{S \in \mathcal{S}_r \setminus \mathcal{P} : S \text{ is adjacent to a box } S' \in \mathcal{P}\},$$

so in particular $\partial^+ \{S\}$ are the boxes adjacent to S . Define also the *interior* $\text{int}(\mathcal{P}) := \{S \in \mathcal{P} : \partial^+ \{S\} \subseteq \mathcal{P}\}$. Note that, since q is supported on Λ_r , f is determined on $\text{int}(\mathcal{P})$ using \mathcal{P} . A *primal (resp. dual) path* will designate a path in $\{f \geq 0\}$ (resp. $\{f \leq 0\}$) and a *level line* will designate a path in $\{f = 0\}$; these paths are *contained in* a set of boxes $\mathcal{P} \subset \mathcal{S}_r$ if they are contained in $\bigcup_{S \in \mathcal{P}} S$. The *left* and *right* sides of $B_k(R)$ are respectively $\{-R\} \times [-kR, kR]^{d-1}$ and $\{R\} \times [-kR, kR]^{d-1}$, and if $d = 2$ the *top* and *bottom* sides are defined similarly.

For the first statement consider the following algorithm:

- Reveal every box that intersects Λ_1 as well as all adjacent boxes.
- Iterate the following steps:
 - Let $\mathcal{W} \subset \mathcal{S}_r$ be the boxes that have been revealed.
 - Identify the set $\mathcal{U} \subseteq \partial^+(\text{int}(\mathcal{W}))$ such that, for each $S \in \mathcal{U}$, there is a primal path contained in $\text{int}(\mathcal{W}) \cap \Lambda_R$ between Λ_1 and the boundary of S (measurable since f is determined on $\text{int}(\mathcal{W})$). In other words, \mathcal{U} contains all boxes on which f is not yet determined but which are connected to Λ_1 by a primal path in Λ_R that *has* been determined.
 - If \mathcal{U} is empty end the loop. Otherwise reveal the boxes in $\partial^+ \mathcal{U} \setminus \mathcal{W}$.

⁵For fixed $s > 0$ we can suppose they are simultaneously continuous almost surely by countability.

⁶More precisely $(W|_S)_{S \in \mathcal{S}_s}$ are defined by setting, for $g \in L^2(\mathbb{R}^d)$, $\int_{y \in \mathbb{R}^d} g(y) dW|_S(y)$ to be jointly centred Gaussian random variables with covariance

$$\mathbb{E} \left[\int_{y \in \mathbb{R}^d} g_1(y) dW|_{S_1}(y) \int_{y \in \mathbb{R}^d} g_2(y) dW|_{S_2}(y) \right] = \begin{cases} \int_{y \in S_1} g_1(y) g_2(y) dy & \text{if } S_1 = S_2, \\ 0 & \text{else.} \end{cases}$$

- If $\text{int}(\mathcal{W}) \cap \Lambda_R$ contains a primal path between Λ_1 and $\partial\Lambda_R$ output 1, otherwise output 0.

This algorithm determines $A(1, R)$ since $\text{int}(\mathcal{W})$ eventually contains all the components of $\{f \geq 0\} \cap \Lambda_R$ that intersect Λ_1 . To estimate the sum of revealments $\text{Rev}(S)$, a box S is revealed if and only if either (i) it is adjacent to a box that intersects Λ_1 , or (ii) there is a primal path in Λ_R between Λ_1 and a box adjacent to S . If $S = v + [0, r]^2$ and $\Lambda_1 \cap (v + \Lambda_{6r}) = \emptyset$ then the latter implies the occurrence of $\Lambda_1 \longleftrightarrow v + \Lambda_{6r}$. Summing over S gives

$$\sum_{S \in \mathcal{S}_r} \text{Rev}(S) \leq \sum_{v \in r\mathbb{Z}^d \cap \Lambda_{R+2r}} \mathbb{P}_p[\Lambda_1 \longleftrightarrow v + \Lambda_{6r}].$$

For the second statement, the first algorithm is:

- Draw a random integer i uniformly in $[-R/r, 0]$, define $L = \{ir\} \times [-kR, kR]^{d-1}$, and reveal every box that intersects $L \cap B_k(R)$, as well as all adjacent boxes.
- Iterate the following steps:
 - Let $\mathcal{W} \subset \mathcal{S}_r$ be the boxes that have been revealed.
 - Identify the set $\mathcal{U} \subseteq \partial^+(\text{int}(\mathcal{W}))$ such that, for each $S \in \mathcal{U}$, there is a primal path contained in $\text{int}(\mathcal{W}) \cap B_k(R)$ between $L \cap B_k(R)$ and the boundary of S .
 - If \mathcal{U} is empty end the loop. Otherwise reveal the boxes in $\partial^+\mathcal{U} \setminus \mathcal{W}$.
- If $\text{int}(\mathcal{W}) \cap B_k(R)$ contains a primal path between the left and right sides of $B_k(R)$ output 1, otherwise output 0.

This algorithm determines $\text{Cross}_k(R)$ since $\text{int}(\mathcal{W})$ eventually contains all the components of $\{f \geq 0\} \cap B_k(R)$ that intersect $L \cap B_k(R)$, and any primal path in $B_k(R)$ between its left and right sides must intersect $L \cap B_k(R)$. To estimate the revealments $\text{Rev}(S)$ of this algorithm, a box S is revealed if and only if either (i) it is adjacent to a box that intersects $L \cap B_k(R)$, or (ii) there is a primal path in $B_k(R)$ between L and a box adjacent to S . If d' denotes the distance from the centre of S to L , this implies the occurrence of (a translation of) the event $A_1(2r, d')$. Averaging over $i \in [-R/r, 0]$ gives

$$\text{Rev}(S) \leq \frac{r}{R} \left(4 + 2 \sum_{i=3}^{R/r} \mathbb{P}_\ell[A_1(2r, ir)] \right) \leq \frac{4r}{R} \sum_{i=2}^{R/r} P_\ell[A_1(2r, ir)].$$

For the second algorithm we modify the above by setting L as $\{-R\} \times [-kR, kR]^{d-1}$, and repeating all other steps. A box $S \in \mathcal{S}_r$ such that $d(S, B_k^+(R)) < r$ is only revealed if there is a primal path in $B_k(R)$ between L and a box adjacent to S , which as before implies the occurrence of (a translation of) the event $A_1(2r, d)$, where d is the distance from the centre of S to L . Since d is at least $R - 2r$, $\text{Rev}(S) \leq \mathbb{P}_\ell[A_1(2r, R - 2r)]$ as required.

The final algorithm (specific to $d = 2$) is:

- Define $L_1 = [-R, R] \times \{-kR\}$ and $L_2 = \{-R\} \times [-kR, kR]$, and reveal every box that intersects $(L_1 \cup L_2) \cap B_k(R)$, as well as all adjacent boxes.
- Iterate the following steps:
 - Let $\mathcal{W} \subset \mathcal{S}_r$ be the boxes that have been revealed.
 - Identify the set $\mathcal{U} \subseteq \partial^+(\text{int}(\mathcal{W}))$ such that, for each $S \in \mathcal{U}$, there is a level line contained in $\text{int}(\mathcal{W}) \cap B_k(R)$ between $(L_1 \cup L_2) \cap B_k(R)$ and the boundary of S .
 - If \mathcal{U} is empty end the loop. Otherwise reveal the boxes in $\partial^+\mathcal{U} \setminus \mathcal{W}$.
- If $\text{int}(\mathcal{W}) \cap B_k(R)$ contains a primal (resp. dual) path between the left and right (resp. top and bottom) sides of $B_k(R)$ terminate with output 1 (resp. 0).
- Since $\text{int}(\mathcal{W})$ contains all components of $\{f = 0\} \cap B_k(R)$ that intersect $L_1 \cup L_2$ and the algorithm has not yet terminated, exactly one of $\{f \geq 0\} \cap B_k(R)$ or $\{f \leq 0\} \cap B_k(R)$ has a component that intersects all four sides of $B_k(R)$. Partition $B_k(R)$ into regions (P_i) using the components of $\{f = 0\} \cap B_k(R)$ that intersect $L_1 \cup L_2$. Let \mathbb{A} to be the region P_i which contains the top-left corner of $B_k(R)$, and set $\mathcal{C} = 1$ (resp. $\mathcal{C} = 0$) if f is positive (resp. negative) on P_i . Then iterate the following:
 - If \mathbb{A} contains a path in $B_k(R)$ between its left and right sides terminate with output \mathcal{C} .




FIGURE 2. The final loop of the algorithm in the proof of the third statement of Lemma 3.6; this loop occurs when there is no left-right or top-bottom paths in $\{f = 0\} \cap B_k(R)$. In this example the loop expands the area \mathbb{A} three times in order to determine the sign \mathcal{C} of the crossing.

- Change the value of \mathcal{C} (from 0 to 1 or 1 to 0), and add to \mathbb{A} the region P_i that is adjacent to it.

The final loop is illustrated in Figure 2; it terminates almost surely since there are a finite number of connected components of $\{f = 0\} \cap B_k(R)$ (recall that f is C^1 -smooth). Note that the algorithm does not necessarily reveal all components of $\{f = 0\}$ inside $B_k(R)$ – any components which are closed loops or only touch the top and right sides of $B_k(R)$ are not revealed – but these do not affect whether $\text{Cross}_k(R)$ occurs.

To estimate the revealments of this algorithm, a box $S \in \mathcal{S}_r$ such that $d(S, B_k^\dagger(R)) < r$ is only revealed if there is a level line in $B_k(R)$ between $L_1 \cap L_2$ and a box adjacent to S , which implies the occurrence of (a translation of) the event $A_2(2r, d')$, where d' is the distance from the centre of S to $L_1 \cap L_2$. Since d' is at least $R - 2r$, $\text{Rev}(S) \leq \mathbb{P}_\ell[A_2(2r, R - 2r)]$ as required. \square

3.2. Proof of Propositions 3.1–3.3. Before proving Propositions 3.1 and 3.2 we give the analogue of Proposition 2.10, which applies to continuous stationary Gaussian fields $f = q \star W$ (note that we do not need to assume Assumption 1.4):

Proposition 3.7. *Suppose $f = q \star W$ is continuous. Then for every $\ell \in \mathbb{R}$, event A , $s > 0$, algorithm $\mathcal{A} \in \mathcal{A}_s$ that determines A , set of boxes $\mathcal{S}' \subseteq \mathcal{S}_s$, and $\varepsilon \geq 0$,*

$$|\mathbb{P}_\ell[f + \varepsilon \sum_{S \in \mathcal{S}'} q \star \mathbf{1}_S \in A] - \mathbb{P}_\ell[f \in A]| \leq \varepsilon s^{d/2} \sqrt{\max\{\mathbb{P}_\ell[A], \mathbb{P}_\ell[f + \varepsilon \sum_{S \in \mathcal{S}'} q \star \mathbf{1}_S \in A]\} \mathbb{E}_\ell |\mathcal{W}_{\mathcal{S}'}|},$$

where $\mathcal{W}_{\mathcal{S}'}$ is the set of boxes in \mathcal{S}' that are revealed by \mathcal{A} . In particular, if (POS) holds,

$$(3.13) \quad |\mathbb{P}_{\ell+\varepsilon}[A] - \mathbb{P}_\ell[f \in A]| \leq \frac{\varepsilon s^{d/2}}{\int q} \sqrt{\max\{\mathbb{P}_\ell[A], \mathbb{P}_{\ell+\varepsilon}[A]\} \mathbb{E}_\ell |\mathcal{W}|},$$

where \mathcal{W} is the set of all boxes in \mathcal{S}_s that are revealed by \mathcal{A} .

Proof. Consider $S \in \mathcal{S}_s$. We use the decomposition (see Proposition A.1 in the appendix)

$$f_S(\cdot) \stackrel{d}{=} \frac{Z_S(q \star \mathbf{1}_S)(\cdot)}{s^{d/2}} + g_S(\cdot),$$

where Z_S is a standard normal random variable and g_S is a continuous Gaussian field independent of Z_S , which implies also that

$$f_S(\cdot) + \varepsilon(q \star \mathbf{1}_S)(\cdot) \stackrel{d}{=} \frac{(Z_S + \varepsilon s^{d/2})(q \star \mathbf{1}_S)(\cdot)}{s^{d/2}} + g_S(\cdot).$$

The same argument that led to (2.18) (this time with W equal to $(Z_S)_{S \in \mathcal{W}_{S'}}$ in the order of revealment, and W' containing Z_S on $S \notin \mathcal{S}'$ as well as g_S for all S) yields in this case

$$\begin{aligned} & |\mathbb{P}_\ell[f + \varepsilon \sum_{S \in \mathcal{S}'} q \star \mathbf{1}_S \in A] - \mathbb{P}_\ell[f \in A]| \\ & \leq \sqrt{2 \max \left\{ \mathbb{P}_\ell[A], \mathbb{P}_\ell \left[f + \varepsilon \sum_{S \in \mathcal{S}'} q \star \mathbf{1}_S \in A \right] \right\} \mathbb{E}_\ell | \mathcal{W}_{\mathcal{S}'} | D_{KL}(Z \| Z + \varepsilon s^{d/2})} \end{aligned}$$

where Z is a standard normal random variable. Since $D_{KL}(Z \| Z + \varepsilon s^{d/2}) = \varepsilon^2 s^d / 2$ we have the first statement. For second statement, notice that $\sum_{S \in \mathcal{S}_s} (q \star \mathbf{1}_S) = (q \star \mathbf{1})(x) = \int q$. Then set $\mathcal{S}' = \mathcal{S}_s$ and replace $\varepsilon \mapsto \varepsilon / \int q$ in the first statement. \square

Proof of Proposition 3.1. This follows directly from (3.13) by considering the algorithm in Lemma 3.6 that determines $A_1(1, R)$ such that

$$\mathbb{E}_\ell |\mathcal{W}| = \sum_{S \in \mathcal{S}_r} \text{Rev}(S) \leq \sum_{v \in r\mathbb{Z}^d \cap \Lambda_{R+2r}} \mathbb{P}_\ell[\Lambda_1 \longleftrightarrow v + \Lambda_{6r}]. \quad \square$$

Proof of Proposition 3.2. We begin with the general case $d \geq 2$. We first partition the set of boxes $\{S \in \mathcal{S}_r : d(S, B_k(R)) < r\}$ that cover $B_k(R)$ into the disjoint sets

$$\mathcal{S}'_1 = \{S \in \mathcal{S}_r : d(S, B_k^+(R)) < r\} \quad \text{and} \quad \mathcal{S}'_2 = \{S \in \mathcal{S}_r : d(S, B_k(R)) < r\} \setminus \mathcal{S}'_1$$

Note that \mathcal{S}'_1 and \mathcal{S}'_2 correspond roughly to boxes which cover, respectively, the right-half $B_k^+(R)$ and its complement $B_k(R) \setminus B_k^+(R)$, except that we enforce disjointness (see Remark 3.8 for an explanation) so we do not have exact reflective symmetry. However the reflection of \mathcal{S}'_2 in the hyperplane $\{0\} \times \mathbb{R}^{d-1}$ is contained in \mathcal{S}'_1 .

By disjointness and since q is supported on Λ_r , for every $x \in B_k(R)$ we have

$$\sum_{i=1,2} \sum_{S \in \mathcal{S}'_i} (q \star \mathbf{1}_S)(x) = \sum_{S \in \mathcal{S}'_1 \cup \mathcal{S}'_2} (q \star \mathbf{1}_S)(x) = (q \star \mathbf{1})(x) = \int q.$$

Then by the multivariate chain rule for Dini derivatives

$$\begin{aligned} \frac{\partial^+}{\partial \ell} \mathbb{P}_\ell[\text{Cross}_k(R)] &= \frac{1}{\int q} \frac{\partial^+}{\partial \varepsilon} \mathbb{P}_\ell \left[f + \varepsilon \sum_{i=1,2} \sum_{S \in \mathcal{S}'_i} q \star \mathbf{1}_S \in \text{Cross}_k(R) \right] \Big|_{\varepsilon=0} \\ &\leq \frac{1}{\int q} \sum_{i=1,2} \frac{\partial^+}{\partial \varepsilon} \mathbb{P}_\ell \left[f + \varepsilon \sum_{S \in \mathcal{S}'_i} q \star \mathbf{1}_S \in \text{Cross}_k(R) \right] \Big|_{\varepsilon=0}. \end{aligned}$$

Now consider the algorithm in Lemma 3.6 that determines $\text{Cross}_k(R)$ such that, under \mathbb{P}_ℓ ,

$$\max_{S \in \mathcal{S}'_1} \text{Rev}(S) \leq \mathbb{P}_\ell[A_1(2r, R - 2r)].$$

By reflective symmetry, there is also an algorithm determining $\text{Cross}_k(R)$ such that, under \mathbb{P}_ℓ ,

$$\max_{S \in \mathcal{S}'_2} \text{Rev}(S) \leq \mathbb{P}_\ell[A_1(2r, R - 2r)].$$

Since also $\max_{i=1,2} |\mathcal{S}'_i| \leq c_1(R/r)^d$ for some $c_1 > 0$ depending only on k and d , applying Proposition 3.7 gives

$$\frac{\partial^+}{\partial \ell} \mathbb{P}_\ell[\text{Cross}_k(R)] \leq \frac{r^{d/2}}{2 \int q} \sqrt{c_1(R/r)^d \mathbb{P}_\ell[A_1(2r, R - 2r)]} = \frac{c_2 R^{d/2}}{\int q} \sqrt{\mathbb{P}_\ell[A_1(2r, R - 2r)]}$$

for some $c_2 = c_2(k, d) > 0$, as required.

For $d = 2$ we consider the top-right quadrant $B_k^\dagger(R)$ and the algorithm in Lemma 3.6 that determines $\text{Cross}_k(R)$ such that

$$\max_{\{S \in \mathcal{S}_r : d(S, B_k^\dagger(R)) < r\}} \text{Rev}(S) \leq \mathbb{P}_\ell[A_2(2r, R - 2r)].$$

Then a similar argument to in the previous case (except partitioning $\{S \in \mathcal{S}_r : d(S, B_k(R)) < r\}$ into $\cup_{i=1,\dots,4} \mathcal{S}'_i$ into four disjoint sets that approximate the four quadrants of $B_k^+(R)$ and using reflective symmetry in both axes) yields the result. \square

Remark 3.8. Since we do not assume $q \geq 0$, it is not necessarily true that

$$\frac{\partial^+}{\partial \varepsilon} \mathbb{P}_\ell[f + \varepsilon q \star \mathbf{1}_S \in \text{Cross}_k(R)] \geq 0$$

for every $S \in \mathcal{S}_r$. Hence in the proof of Proposition 3.2 it was crucial that we partitioned $\{S \in \mathcal{S}_r : d(S, B_k(R)) < r\}$ disjointly into $\cup_i \mathcal{S}'_i$, since otherwise we could not deduce that

$$\frac{\partial^+}{\partial \ell} \mathbb{P}_\ell[\text{Cross}_k(R)] \leq \frac{1}{\int q} \frac{\partial^+}{\partial \varepsilon} \mathbb{P}_\ell[f + \varepsilon \sum_{i=1,2} \sum_{S \in \mathcal{S}'_i} q \star \mathbf{1}_S \in \text{Cross}_k(R)] \Big|_{\varepsilon=0}.$$

To prove Proposition 3.3 we need the analogue of Proposition 2.16. We say that an event A is *compactly supported* if A is measurable with respect to $f|_D$ for a compact $D \subset \mathbb{R}^d$, and is a *continuity event* if $\ell \mapsto \mathbb{P}_\ell[f + g \in A]$ is continuous for every smooth function $g : \mathbb{R}^d \rightarrow \mathbb{R}$; for example, $\text{Cross}_k(R)$ and $A_i(r, R)$, $i = 1, 2$, are compactly supported continuity events by Lemma 3.13 below.

Proposition 3.9. *Suppose $f = q \star W$ satisfies Assumption 1.4 and (BOU), and let $r > 0$ be such that q is supported on Λ_r . Then there exists a $c > 0$ depending only on d such that, for every $\ell \in \mathbb{R}$, increasing compactly supported continuity event A , $s > 0$, algorithm $\mathcal{A} \in \mathcal{A}_s$ determining A , and set of boxes $\mathcal{S}' \subseteq \mathcal{S}_s$,*

$$(3.14) \quad \frac{d^-}{d\ell} \mathbb{P}_\ell[A] \geq \frac{c \min\{1, (s/r)^d\}}{\|q\|_2} \frac{\text{Var}_\ell[\mathbb{P}_\ell[A|\mathcal{F}_{\mathcal{S}'}]]}{\max_{S \in \mathcal{S}'} \text{Rev}(S)},$$

where $\mathcal{F}_{\mathcal{S}'}$ denotes the σ -algebra generated by $(f_S)_{S \in \mathcal{S}'}$, and the revealments $\text{Rev}(S)$ are under \mathbb{P}_ℓ .

Remark 3.10. The proof of (3.14) shows that it can be strengthened by replacing $\frac{d^-}{d\ell} \mathbb{P}_\ell[A] = \frac{d^-}{d\varepsilon} \mathbb{P}_\ell[f + \varepsilon \in A]$ with $\frac{d^-}{d\varepsilon} \mathbb{P}_\ell[f + \varepsilon g_{\mathcal{S}'} \in A]$ where $g_{\mathcal{S}'}(\cdot) := \mathbf{1}_{d(\cdot, \mathcal{S}') \leq 2r}$, but we do not need this.

Remark 3.11. In [38] a similar result (in the case $\mathcal{S}' = \mathcal{S}_s$) was proven for an approximation of the field f in which the white noise is replaced with its *discretisation* at scale $\varepsilon \ll 1$. However, since one needed to take $\varepsilon \ll 1$ in the approximation (e.g. in $d = 2$ one needs $\varepsilon \ll 1/R$ if the event is supported on $B(R)$), this approach results in a prefactor $\varepsilon^{d/2}$ that decays rapidly in the scale of the event. Although this prefactor is also present in (3.14) as $s \rightarrow 0$, the difference is that one can work with fixed s .

We prove Proposition 3.9 in Section 4 below. Let us complete the proof of Proposition 3.3:

Proof of Proposition 3.3. For the first statement we apply Proposition 3.9 (in the case $s = r$, $\mathcal{S}' = \mathcal{S}_s$, and $A = \text{Cross}_k(R)$, which is a continuity event by Lemma 3.13) to the algorithm in Lemma 3.6 that determines $\text{Cross}_k(R)$ whose revealments are bounded by $\frac{4r}{R} \sum_{i=2}^{R/r} \mathbb{P}_\ell[A_1(2r, ir)]$.

For the second statement we follow the proof of Proposition 2.3 in the Bernoulli case, except using Proposition 3.9 (in the case $s = r$, $\mathcal{S}' = \{S \in \mathcal{S}_s : d(S, B_k^+(R)) < r\}$, and $A = \text{Cross}_k(R)$) and the algorithm in Lemma 3.6 that determines $\text{Cross}_k(R)$ whose revealments on \mathcal{S}' are bounded by $\mathbb{P}_\ell[A_2(2r, R-2r)]$. To control the conditional variances in (3.14) we use the same argument as in the proof of Proposition 2.3; in particular the FKG inequality is available and, since $R \geq 8r$, the events B_1 and B_2 are independent as in the Bernoulli case. \square

Remark 3.12. Similarly to in Section 2.4, combining Propositions 3.7 and 3.9 yields a general lower bound on the revealments of increasing events. We omit the proof, but the result is the following. Suppose $f = q \star W$ satisfies Assumption 1.4 and (POS)–(BOU). Let r be such that q is supported on Λ_r , let $\ell \in \mathbb{R}$, let $R \geq r$, let A be an increasing continuity event supported

on $B(R)$, let $s > 0$, and let $\mathcal{A} \in \mathcal{A}_s$ be an algorithm determining A . Then there exists a $c > 0$ depending only on d such that

$$\max_{S \in \mathcal{S}_s} \text{Rev}(S) \geq \frac{c(\|q\|_2 \min\{1, (s/r)^d\} \text{Var}_\ell[\mathbb{1}_A])^{2/3}}{\mathbb{P}_\ell[A]^{1/3} R^{d/3}}$$

where the revealments $\text{Rev}(S)$ are under \mathbb{P}_ℓ .

One can also prove a lower bound on $\max_{S \in \mathcal{S}'} \text{Rev}(S)$ for general $\mathcal{S}' \subset \mathcal{S}_s$, analogous to Proposition 2.18, however in that case we would need $q \geq 0$ (for the same reason as explained in Remark 3.8 above) and also the refinement to Proposition 3.9 mentioned in Remark 3.10.

3.3. Proof of auxiliary results. To finish the section we prove Lemmas 3.4 and 3.5:

Proof of Lemma 3.5. We first observe that $g := f - f_r = (q - q_r) \star W$ is a C^1 -smooth stationary Gaussian field satisfying

$$\mathbb{E}[g(0)^2] = \int_{x \in \mathbb{R}^d} (q - q_r)^2(x) dx = \int_{|x| > r/2} q(x)^2 (1 - \varphi(|x/r|))^2 dx \leq \int_{|x| > r/2} q(x)^2 \leq c_1 r^{d-2\beta},$$

for some $c_1 > 0$ and we used that $|q(x)| \leq c|x|^{-\beta}$ by Assumption 1.4. Similarly, for every direction $v \in \mathbb{S}^1$,

$$\mathbb{E}[(\partial_v g(0))^2] = \int_{|x| > r/2} (\partial_v(q(x)(1 - \varphi(|x/r|)))^2 dx \leq c_2 r^{d-2\beta},$$

for some $c_2 > 0$ that depends on the (uniformly bounded) derivatives of φ , and we used that $|\nabla q(x)| \leq c|x|^{-\beta}$ by Assumption 1.4. Then by a Borell-TIS argument (see [38, Proposition 3.11] for the case $d = 2$, and the proof is identical in all dimensions) there exist $c_3, c_4 > 0$ such that, for all $R, r \geq 2$,

$$(3.15) \quad \mathbb{P}[\|f - f_r\|_{\infty, B(R)} > c_3(\log R)r^{-(\beta-d/2)}] \leq c_3 e^{-c_4(\log R)^2}$$

We also note the following consequence of (POS) which can be proved with a Cameron-Martin argument (see [38, Proposition 3.6] for the case $d = 2$, and the proof is identical in all dimensions): there exists a $c_5 > 0$ such that, for $R \geq 1$, increasing event A' measurable with respect to $f|_{B(R)}$, $\ell \in \mathbb{R}$ and $t > 0$,

$$(3.16) \quad \mathbb{P}_\ell[\{f + t \in A'\} \setminus \{f \in A'\}] = \mathbb{P}_\ell[f + t \in A'] - \mathbb{P}_\ell[f \in A'] \leq c_5 t R^{d/2}.$$

We now complete the proof, for which we may assume that $\ell = 0$. Consider $A = A_1 \cap A_2$ where A_1 is increasing, A_2 is decreasing, and both A_1 and A_2 are measurable with respect to $f|_{B(R)}$. Abbreviate $t = c_3(\log R)r^{-(\beta-d/2)}$ and define $E = \{\|f - f_r\|_{\infty, B(R)} > t\}$. Then

$$\begin{aligned} \mathbb{P}[f_r \in A_1 \cap A_2] &\leq \mathbb{P}[f_r \in A_1 \cap A_2 \cap E^c] + \mathbb{P}[E] \\ &\leq \mathbb{P}[\{f + t \in A_1\} \cap \{f - t \in A_2\}] + \mathbb{P}[E] \\ &\leq \mathbb{P}[f \in A_1 \cap A_2] + \mathbb{P}[\{f + t \in A_1\} \setminus \{f \in A_1\}] + \mathbb{P}[\{f - t \in A_2\} \setminus \{f \in A_2\}] + \mathbb{P}[E] \\ &\leq \mathbb{P}[f \in A_1 \cap A_2] + 2c_5 t R^{d/2} + c_3 e^{-c_4(\log R)^2} \end{aligned}$$

where in the second inequality we used that A_1 (resp. A_2) is increasing (resp. decreasing) and measurable with respect to $f|_{B(R)}$, and the final inequality was by (3.15) and (3.16). Similarly

$$\begin{aligned} \mathbb{P}[f_r \in A_1 \cap A_2] &\geq \mathbb{P}[\{f - t \in A_1\} \cap \{f + t \in A_2\} \cap E^c] \\ &\geq \mathbb{P}[f \in A_1 \cap A_2] - \mathbb{P}[\{f \in A_1\} \setminus \{f - t \in A_1\}] - \mathbb{P}[\{f \in A_2\} \setminus \{f + t \in A_2\}] - \mathbb{P}[E] \\ &\geq \mathbb{P}[f \in A_1 \cap A_2] - 2c_5 t R^{d/2} - c_3 e^{-c_4(\log R)^2} \end{aligned}$$

which gives the result. \square

Proof of Lemma 3.4. For the first statement, it is enough to prove that

$$(3.17) \quad \liminf_{R \rightarrow \infty} \mathbb{P}_{\ell_c}[\text{Cross}_5(R)] > 0$$

since then the result follows by the continuity of $\ell \mapsto \mathbb{P}_\ell[\text{Cross}_5(R)]$ (by Lemma 3.13 below for instance). By a classical bootstrapping argument [28, Section 5.1] and Lemma 3.5, there are $c_1, \varepsilon > 0$ such that

$$(3.18) \quad \mathbb{P}_\ell[\text{Cross}_5(3R)] \leq c_1 (\mathbb{P}_\ell[\text{Cross}_5(R)]^2 + R^{-\varepsilon})$$

for $\ell \in \mathbb{R}$ and R sufficiently large. A consequence of (3.18) and the continuity of $\ell \mapsto \mathbb{P}_\ell[\text{Cross}_5(R)]$ is that

$$\liminf_{R \rightarrow \infty} \mathbb{P}_\ell[\text{Cross}_5(R)] < 1/c_1 \quad \implies \quad \liminf_{R \rightarrow \infty} \mathbb{P}_{\ell'}[\text{Cross}_5(R)] = 0 \quad \text{for some } \ell' > \ell.$$

Covering the annulus $\Lambda_{5R} \setminus \Lambda_{3R}$ with $2d$ symmetric copies of $B_5(R)$, one can find a finite collection of copies A_i of $\text{Cross}_5(R)$ such that $\{\Lambda_1 \longleftrightarrow \infty\} \subseteq A_1(3R, 5R) \subseteq \cup_i A_i$. Hence we also have

$$\liminf_{R \rightarrow \infty} \mathbb{P}_{\ell'}[\text{Cross}_5(R)] = 0 \quad \implies \quad \mathbb{P}_{\ell'}[\Lambda_1 \longleftrightarrow \infty] = 0 \quad \implies \quad \ell' \leq \ell_c,$$

and so we deduce (3.17).

For the second statement we refer to [38] where it is shown that the RSW estimates hold under Assumption 1.4 and (POS') (indeed the recent work [30] shows that the correlation decay in Assumption 1.4 is not even needed). \square

We also state a continuity result that we used in the section:

Lemma 3.13. *Let f be a C^2 -smooth Gaussian field on \mathbb{R}^d such that $(f(x), \nabla f(x), \nabla^2 f(x))$ is non-degenerate for every $x \in \mathbb{R}^d$. Then for every $k \geq 1$ and $R \geq r > 0$,*

$$\mathbb{P}_\ell[\text{Cross}_k(R)] \quad \text{and} \quad \mathbb{P}_\ell[A_i(r, R)], \quad i = 1, 2$$

are continuous functions of $\ell \in \mathbb{R}$.

Proof. Since f is C^2 -smooth and $(f(x), \nabla f(x), \nabla^2 f(x))$ is non-degenerate, by Bulinskaya's lemma [1, Lemma 11.2.10] the critical points of f , as well as its restriction to a smooth hypersurface, are almost surely locally finite and have distinct critical levels. Since the events $\{f + \ell \in \text{Cross}_k(R)\}$ and $\{f + \ell \in A_i(r, R)\}$ depend only on the (stratified) diffeomorphism class of the level set $\{f + \ell = 0\}$ restricted to, respectively, $B_k(R)$ and $\Lambda_R \setminus \Lambda_r$, by the (stratified) Morse lemma [25, Theorem 7] almost surely there is a $\delta > 0$ such that $\mathbb{1}_{\{f + \ell + s \in \text{Cross}_k(R)\}}$ and $\mathbb{1}_{\{f + \ell + s \in A_i(r, R)\}}$ are constant on $s \in (-\delta, \delta)$, which is equivalent to the claimed continuity. \square

4. THE OSSS INEQUALITY FOR SMOOTH GAUSSIAN FIELDS AND APPLICATIONS

In this section we establish a new Russo-type inequality for smooth Gaussian fields which we use to prove Proposition 3.9, with Theorem 1.14 following as an application. We consider a field $f = q \star W$ which is C^2 -smooth and satisfies (BOU), and let $r > 0$ be such that q is supported on Λ_r . In particular this implies that $(f(0), \nabla f(0), \nabla^2 f(0))$ is non-degenerate. We emphasise that in this section neither (POS) nor (POS') play any role.

As in Section 3.1, fix $s > 0$ and consider the orthogonal decomposition $f = \sum_{S \in \mathcal{S}_s} f_S$ where

$$f_S(\cdot) = (q \star W|_S)(\cdot) = \int_{y \in \mathbb{R}^d} q(\cdot - y) dW_S(y) = \int_{y \in S} q(\cdot - y) dW(y).$$

The proof of Proposition 3.9 is based on an application of the OSSS inequality (Theorem 2.15) to the independent fields $(f_S)_{S \in \mathcal{S}_s}$. In this context the *resampling influences* (c.f. (2.21)) are defined, for each $S \in \mathcal{S}_s$, as

$$\text{Infl}_A(S) := \mathbb{P}_\ell[\mathbb{1}_{\{f \in A\}} \neq \mathbb{1}_{\{f^{(S)} \in A\}}]$$

where $f^{(S)}$ denotes the field $f = \sum_{S \in \mathcal{S}_s} f_S$ with f_S resampled. Just as for other recent applications of the OSSS inequality in percolation theory [16, 15, 17], the crucial mechanism is that $\frac{d^-}{d\ell} \mathbb{P}_\ell[A]$ is bounded below by the sum of the resampling influences. Recall the definition of compactly supported continuity events from the statement of Proposition 3.9.

Proposition 4.1 (Russo-type inequality). *There exists a constant $c > 0$ depending only on d such that, for every $\ell \in \mathbb{R}$, $s > 0$, and increasing compactly supported continuity event A ,*

$$\frac{d^-}{d\ell} \mathbb{P}_\ell[A] \geq \frac{c \min\{1, (s/r)^d\}}{\|q\|_2} \sum_{S \in \mathcal{S}_s} \text{Infl}_A(S)$$

where the resampling influences $\text{Infl}_A(S)$ are under \mathbb{P}_ℓ .

Before proving Proposition 4.1, let us complete the proof of Proposition 3.9.

Proof of Proposition 3.9. The OSSS inequality (Theorem 2.15), combined with the reasoning leading to (2.22), gives

$$\text{Var}_\ell[\mathbb{P}_\ell[A|\mathcal{F}_{S'}]] \leq \frac{1}{2} \sum_{S \in \mathcal{S}'} \text{Rev}(S) \text{Infl}_A(S)$$

and hence (true for arbitrary event A)

$$\sum_{S \in \mathcal{S}_s} \text{Infl}_A(S) \geq \sum_{S \in \mathcal{S}'} \text{Infl}_A(S) \geq \frac{2 \text{Var}_\ell[\mathbb{P}_\ell[A|\mathcal{F}_{S'}]]}{\max_{S \in \mathcal{S}'} \text{Rev}(S)}.$$

Combining with Proposition 4.1 yields the result. \square

The main idea in the proof of Proposition 4.1, which distinguishes it from the discretisation approach in [38], is to use an orthonormal decomposition of each f_S to interpret $\frac{d^-}{d\ell} \mathbb{P}_\ell[A]$ and the resampling influences $\text{Infl}_A(S)$ as measuring, respectively, the ‘boundary’ and ‘volume’ of certain sets in Gaussian space. Then we can apply Gaussian isoperimetry to deduce the result. For a set $E \subset \mathbb{R}^n$ we denote

$$E^{+\varepsilon} := \{x \in \mathbb{R}^n : \text{there exists } y \in E \text{ s.t. } |x - y|_2 \leq \varepsilon\}$$

to be the ε -thickening of E .

Proposition 4.2 (Gaussian isoperimetry). *There exists a constant $c > 0$ such that, for every measurable $E \subset \mathbb{R}^n$ and $\varepsilon \geq 0$,*

$$\mathbb{P}[X \in E^{+\varepsilon} \setminus E] \geq \sqrt{\frac{2}{\pi}} \mathbb{P}[X \in E](1 - \mathbb{P}[X \in E])\varepsilon - c\varepsilon^2$$

where X is an n -dimensional standard Gaussian vector.

Proof. Let $\varphi(x)$ and $\Phi(x)$ denote the standard normal pdf and cdf respectively. The classical Gaussian isoperimetric inequality states that

$$\liminf_{\varepsilon \downarrow 0} \varepsilon^{-1} \mathbb{P}[X \in E^{+\varepsilon} \setminus E] \geq \varphi(\Phi^{-1}(\mathbb{P}[X \in E])).$$

A simple consequence (see, e.g., [33, Eq. (3)]) is that, for any $\varepsilon \geq 0$,

$$(4.1) \quad \mathbb{P}[X \in E^{+\varepsilon}] \geq \Phi(\Phi^{-1}(\mathbb{P}[X \in E]) + \varepsilon).$$

By Taylor expanding Φ on the right-hand side of (4.1) we have

$$\mathbb{P}[X \in E^{+\varepsilon} \setminus E] \geq \varepsilon \varphi(\Phi^{-1}(\mathbb{P}[X \in E])) - \frac{1}{2} \sup_{x \in \mathbb{R}} |\varphi'(x)| \varepsilon^2,$$

and the result follows since, for all $x \in \mathbb{R}$, $\varphi(x) \geq \sqrt{\frac{2}{\pi}} \Phi(x)(1 - \Phi(x))$ (as can be seen from the fact that the Mill’s ratio $(1 - \Phi(x))/\varphi(x)$ is decreasing on $x \geq 0$), and since $|\varphi'(x)|$ is uniformly bounded on $x \in \mathbb{R}$. \square

We use the following orthogonal decomposition of f_S (see Proposition A.1 in the appendix). Let $Z = (Z_i)_{i \geq 1}$ be a sequence of i.i.d. standard normal random variables and let $(\varphi_i)_{i \geq 1}$ be an orthonormal basis of $L^2(S)$. Then

$$(4.2) \quad f_S^n := \sum_{i \geq 1}^n Z_i (q \star \varphi_i) \Rightarrow f_S,$$

in law with respect to the C^0 -topology.

Proof of Proposition 4.1. By linear rescaling, we may suppose without loss of generality that $\ell = 0$, $\|q\|_2 = 1$, and that q is supported on Λ_1 (i.e. $r = 1$). For each $S \in \mathcal{S}_s$, let $g_S : \mathbb{R}^d \rightarrow [0, 1]$ be a smooth function such that $g_S(x) = 1$ on $\{x : d(x, S) \leq 1\}$ and $g_S(x) = 0$ on $\{x : d(x, S) \geq 2\}$. Then $\sum_{S \in \mathcal{S}_s} g_S(x) \leq c_1 \max\{1, s^{-d}\}$ for some constant $c_1 > 0$ depending only on d . Therefore, since A is increasing, and by the multivariate chain rule for Dini derivatives,

$$(4.3) \quad \frac{d^-}{d\varepsilon} \mathbb{P}[f + \varepsilon \in A] \Big|_{\varepsilon=0} \geq \frac{1}{c_1 \max\{1, s^{-d}\}} \sum_{S \in \mathcal{S}_s} \frac{d^-}{d\varepsilon} \mathbb{P}[f + \varepsilon g_S \in A] \Big|_{\varepsilon=0}.$$

For each $S \in \mathcal{S}_s$, let f'_S denote an independent copy of f_S , define $h_S = f - f_S$, and let \mathcal{F}_{h_S} be the σ -algebra generated by h_S . We next claim that, almost surely over \mathcal{F}_{h_S} ,

$$(4.4) \quad \frac{d^-}{d\varepsilon} \mathbb{P}[f_S + h_S + \varepsilon g_S \in A | \mathcal{F}_{h_S}] \Big|_{\varepsilon=0} \geq c_2 \mathbb{P}[\mathbb{1}_{\{f_S + h_S \in A\}} \neq \mathbb{1}_{\{f'_S + h_S \in A\}} | \mathcal{F}_{h_S}]$$

for some universal $c_2 > 0$. Together with (4.3), this will complete the proof of Proposition 4.1 since

$$\begin{aligned} \frac{d^-}{d\varepsilon} \mathbb{P}[f_S + \varepsilon g_S \in A] \Big|_{\varepsilon=0} &\geq \mathbb{E} \left[\frac{d^-}{d\varepsilon} \mathbb{P}[f_S + h_S + \varepsilon g_S \in A | \mathcal{F}_{h_S}] \Big|_{\varepsilon=0} \right] \\ &\geq c_2 \mathbb{E} [\mathbb{P}[\mathbb{1}_{\{f_S + h_S \in A\}} \neq \mathbb{1}_{\{f'_S + h_S \in A\}} | \mathcal{F}_{h_S}]] \\ &=: c_2 \text{Infl}_A(S). \end{aligned}$$

where the first inequality is Fatou's lemma, and the second inequality is by (4.4).

It remains to prove (4.4). Henceforth we fix $S \in \mathcal{S}_s$, condition on h_S , and drop \mathcal{F}_{h_S} from the notation. Let $(\varphi_i)_{i \geq 1}$ be an orthonormal basis of $L^2(S)$ and recall the decomposition (4.2). Fixing $n \in \mathbb{N}$ and viewing $\{f'_S + h_S \in A\}$ as a Borel set E in the n -dimensional Gaussian space generated by the standard Gaussian vector $Z^n = (Z_i)_{1 \leq i \leq n}$, by Proposition 4.2

$$(4.5) \quad \mathbb{P}[Z^n \in E^{+\varepsilon} \setminus E] \geq c_3 \varepsilon \mathbb{P}[Z^n \in E] (1 - \mathbb{P}[Z^n \in E]) - c_4 \varepsilon^2$$

for some $c_3, c_4 > 0$ and every $\varepsilon \geq 0$. Consider $y = (y_i) \in \mathbb{R}^n$ such that $\|y\|_2 = \varepsilon$. By Young's convolution inequality, and since φ_i are an orthonormal basis,

$$\left\| \sum_{i \leq n} y_i (q \star \varphi_i) \right\|_\infty \leq \|q\|_2 \left\| \sum_{i \leq n} y_i \varphi_i \right\|_2 = \|y\|_2 = \varepsilon.$$

Since $q \star \varphi_i$ is supported on $\{x : d(x, S) \leq 1\}$, and recalling that $g_S(\cdot) := \mathbb{1}_{d(\cdot, S) \leq 1}$, this gives

$$\sup_{y: \|y\|_2 \leq \varepsilon} \sum_{i \leq n} (Z_i + y_i) (q \star \varphi_i) - f'_S = \sup_{y: \|y\|_2 \leq \varepsilon} \sum_{i \leq n} y_i (q \star \varphi_i) \leq \varepsilon g_S.$$

Therefore, since A is increasing,

$$\begin{aligned} \mathbb{P}[f'_S + h_S + \varepsilon g_S \in A] - \mathbb{P}[f'_S + h_S \in A] &\geq \mathbb{P}[\cup_{y: \|y\|_2 \leq \varepsilon} \{Z^n + y \in E\}] - \mathbb{P}[Z^n \in E] \\ &= \mathbb{P}[Z^n \in E^{+\varepsilon} \setminus E]. \end{aligned}$$

Combining with (4.5),

$$(4.6) \quad \mathbb{P}[f'_S + h_S + \varepsilon g_S \in A] - \mathbb{P}[f'_S + h_S \in A] \geq c_3 \varepsilon \mathbb{P}[f'_S + h_S \in A] (1 - \mathbb{P}[f'_S + h_S \in A]) - c_4 \varepsilon^2.$$

It remains to prove that almost surely (with respect to h_S), as $n \rightarrow \infty$,

$$(4.7) \quad \mathbb{P}[f'_S + h_S \in A] \rightarrow \mathbb{P}[f_S + h_S \in A] \quad \text{and} \quad \mathbb{P}[f'_S + h_S + \varepsilon g_S \in A] \rightarrow \mathbb{P}[f_S + h_S + \varepsilon g_S \in A],$$

since then sending $n \rightarrow \infty$ in (4.6) yields

$$\mathbb{P}[f_S + h_S + \varepsilon g_S \in A] - \mathbb{P}[f_S + h_S \in A] \geq c_3 \varepsilon \mathbb{P}[f_S + h_S \in A] (1 - \mathbb{P}[f_S + h_S \in A]) - c_4 \varepsilon^2,$$

which gives (4.4) after sending $\varepsilon \rightarrow 0$.

So let us justify (4.7). Recall that A is an increasing continuity event; this means that for almost every $f = f_S + h_S$ there exists $\delta > 0$ such that

$$\mathbb{1}_{\{f_S + h_S + s \in A\}} \quad \text{and} \quad \mathbb{1}_{\{f_S + h_S + \varepsilon g_S + s \in A\}}$$

are constant for $s \in (-\delta, \delta)$. Then since $f_S^n \rightarrow f_S$ in law with respect to the C^0 -topology, we have (4.7) (by the Portmanteau lemma for instance). \square

Remark 4.3. Note that in the proof of Proposition 4.1 we did not require that the Borel set E in the n -dimensional Gaussian space generated by Z^n be increasing, since Gaussian isoperimetry is valid for arbitrary sets. This allows us to avoid any requirement that $q \star \varphi_i$ be a positive function, in contrast to the discretisation approach in [38].

4.1. Application to the sharpness of the phase transition for finite-range Gaussian fields. We conclude the section by proving Theorem 1.14, following the approach in [16]. For this we only need the special case $s = r$ and $\mathcal{S}' = \mathcal{S}_s$ of Proposition 3.9.

Proof of Theorem 1.14. By linear rescaling and adjusting constants, we may assume without loss of generality that q is supported on Λ_1 , and prove the theorem for Λ_2 replacing Λ_1 .

For $R \geq 0$ define $g_R(\ell) := \mathbb{P}_\ell[A_1(2, R)]$ (recall that this means $g_R(\ell) := 1$ if $R \in [0, 2]$), and its limit $g_R := \lim_{R \rightarrow \infty} g_R(\ell) = \mathbb{P}_\ell[\Lambda_2 \longleftrightarrow \infty]$. We will first establish the differential inequality

$$(4.8) \quad \frac{d^-}{d\ell} g_R(\ell) \geq \frac{c_1 g_R(\ell)(1 - g_R(\ell))}{\frac{1}{R} \sum_{i=0}^{R-1} g_i(\ell)}$$

for some $c_1 > 0$, every R sufficiently large, and every $\ell \in \mathbb{R}$. Recall the notation from the beginning of the proof of Lemma 3.6 and for $R \geq 2$ consider the following algorithm in \mathcal{A}_1 (essentially taken from [16]):

- Draw a random integer i uniformly in $[2, R]$, and reveal every box that intersects $\partial\Lambda_i$, as well as all adjacent boxes.
- Iterate the following steps:
 - Let $\mathcal{W} \subset \mathcal{S}_1$ be the boxes that have been revealed.
 - Identify the set $\mathcal{U} \subseteq \partial^+(\text{int}(\mathcal{W}))$ such that, for each $S \in \mathcal{U}$, there is a primal path contained in $\text{int}(\mathcal{W}) \cap \Lambda_R$ between $\partial\Lambda_i$ and the boundary of S .
 - If \mathcal{U} is empty end the loop. Otherwise reveal the boxes in $\partial^+\mathcal{U} \setminus \mathcal{W}$.
- If $\text{int}(\mathcal{W})$ contains a primal path between Λ_2 and Λ_R output 1, otherwise output 0.

This algorithm determines $A_1(2, R)$ since $\text{int}(\mathcal{W})$ eventually contains all the components of $\{f \geq 0\} \cap \Lambda_R$ that intersect $\partial\Lambda_i$, and any primal path between Λ_2 and Λ_R must intersect $\partial\Lambda_i$. To estimate the revealments $\text{Rev}(S)$ under \mathbb{P}_ℓ , note that a box S is revealed if and only if either (i) it intersects, or is adjacent to a box that intersects, $\partial\Lambda_i$, or (ii) there is a primal path in Λ_R between $\partial\Lambda_i$ and a box adjacent to S . If d' denotes the distance from the centre of S to Λ_i , this implies the occurrence of (a translation of) the event $A_1(2, d')$. Averaging on $i \in [2, R]$, we have

$$\text{Rev}(S) \leq \frac{1}{R-1} \left(4 + 2 \sum_{i=3}^{R-1} \mathbb{P}_\ell[A_1(2, i)] \right) \leq \frac{4}{R-1} \sum_{i=0}^{R-1} g_i(\ell) \leq \frac{5}{R} \sum_{i=0}^{R-1} g_i(\ell)$$

for sufficiently large R . Applying Proposition 3.9 (with $s = 1$ and $\mathcal{S}' = \mathcal{S}_1$, recalling that $A_1(2, R)$ is a continuity event by Lemma 3.13) gives that

$$\frac{d^-}{d\ell} g_R(\ell) \geq \frac{c_2 g_R(\ell)(1 - g_R(\ell))}{\max_{S \in \mathcal{S}_1} \text{Rev}(S)} \geq \frac{c_2 g_R(\ell)(1 - g_R(\ell))}{\frac{5}{R} \sum_{i=0}^{R-1} g_i(\ell)}$$

for some $c_2 > 0$ and sufficiently large R , which gives (4.8).

We now argue that (4.8) implies the result. First assume that there exists a $\ell_0 > \ell_c$ such that $g(\ell_0) < 1$ (this is clear if f satisfies (POS'), since then $\mathbb{P}[\inf_{x \in \Lambda_3} f(x) \geq \ell] > 0$ for every $\ell \in \mathbb{R}$, but not in general). Then by monotonicity $1 - g_\ell(R) > (1 - g(\ell_0))/2$ for all $\ell < \ell_0$ and large R . Hence setting $c_3 = c_1(1 - g(\ell_0))/2 > 0$ and defining $f_R(\ell) = g_R(\ell)/c_3$ we have

$$\frac{d^-}{d\ell} f_R(\ell) \geq \frac{f_R(\ell)}{\frac{1}{R} \sum_{i=0}^{R-1} f_i(\ell)}.$$

for all $\ell < \ell_0$ and large R , and applying [16, Lemma 3.1]⁷ yields the result. On the other hand, if $g(\ell_0) = 1$ for every $\ell_0 > \ell_c$ then the second statement of the theorem is immediate. To prove the first statement, instead choose a $\ell_0 < \ell_c$ and repeat the above argument. This implies the statement for $\ell < \ell_0$, and taking $\ell_0 \uparrow \ell_c$ gives the claim. \square

APPENDIX A. ORTHOGONAL DECOMPOSITION OF f_S

For completeness we present a classical orthogonal decomposition of the Gaussian field

$$f_S(\cdot) = (q \star W|_S)(\cdot) = \int_{y \in S} q(\cdot - y) dW(y)$$

where $S \subset \mathbb{R}^d$ is a compact domain, $q \in L^2(\mathbb{R}^d)$, and W is the white noise on \mathbb{R}^d . In this section we shall assume only that f_S is continuous, all other conditions on q being irrelevant.

Proposition A.1 (Orthogonal decomposition of f_S). *Let $(Z_i)_{i \geq 1}$ be a sequence of i.i.d. standard normal random variables and let $(\varphi_i)_{i \geq 1}$ be an orthonormal basis of $L^2(S)$. Then, as $n \rightarrow \infty$,*

$$f_S^n := \sum_{i \geq 1}^n Z_i (q \star \varphi_i) \Rightarrow f_S$$

in law with respect to the C^0 -topology on compact sets. In particular,

$$f_S(\cdot) \stackrel{d}{=} \frac{Z_1 (q \star \mathbf{1}_S)(\cdot)}{\sqrt{\text{Vol}(S)}} + g(\cdot)$$

where g is an continuous Gaussian field independent of Z_1 .

Proof. Remark that, for each $x \in \mathbb{R}^d$, $f_S^n(x) \Rightarrow f_S(x)$ in law since they are centred Gaussian random variables and

$$\mathbb{E} \left[\left(\sum_{i \geq 1}^n Z_i (q \star \varphi_i)(x) \right)^2 \right] = \sum_{i \geq 1}^n \left(\int_S q(x-s) \varphi_i(s) dx \right)^2 \rightarrow \int_S q(x-s)^2 dx = \mathbb{E}[f_S(x)^2]$$

by Parseval's identity. Note also that the functions $q \star \varphi_i$ are continuous (as a convolution of L^2 functions), and so each f_S^n is continuous. Hence the first statement of the proposition follows by an application of Lemma A.2 below. For the second statement, set φ_1 to be constant on S . \square

Lemma A.2. *Let $(f_i)_{i \geq 1}$ be a sequence of independent continuous centred Gaussian fields on \mathbb{R}^d and define $g_n := \sum_{i \geq 1}^n f_i$. Suppose there exists a continuous Gaussian field g on \mathbb{R}^d such that, for each $x \in \mathbb{R}^d$, $g_n(x) \Rightarrow g(x)$ in law. Then $g_n \Rightarrow g$ in law with respect to the C^0 -topology on compact sets.*

Proof. We follow the proof of [1, Theorem 3.1.2]. Since $g_n(x)$ is a sum of independent random variables converging in law, by Levy's equivalence theorem we may define $g(x)$ as the almost sure limit of $g_n(x)$. Fix a compact set $\Omega \subset \mathbb{R}^d$, and consider $(g_n)_{n \geq 1}$ as elements of the Banach space $C(\Omega)$ of continuous functions on Ω equipped with the C_0 -topology. By the Itô-Nisio theorem [1, Theorem 3.1.3], it suffices to show that

$$\int_{\Omega} g_n d\mu \rightarrow \int_{\Omega} g d\mu$$

in mean (and so in probability) for every finite signed Borel measure μ on Ω . Define the continuous functions $u_n(x) := \mathbb{E}[g_n(x)^2]$ and $u(x) := \mathbb{E}[g(x)^2]$. Then

$$\mathbb{E} \left[\left| \int_{\Omega} g d\mu - \int_{\Omega} g_n d\mu \right| \right] \leq \int_{\Omega} \left(\mathbb{E}[(g(x) - g_n(x))^2] \right)^{1/2} |\mu|(dx) \leq \int_{\Omega} \left(u(x) - u_n(x) \right)^{1/2} |\mu|(dx).$$

Since $u_n \rightarrow u$ monotonically, by Dini's theorem the convergence is uniform on Ω , so we have that $\mathbb{E} \left[\left| \int_{\Omega} g d\mu - \int_{\Omega} g_n d\mu \right| \right] \rightarrow 0$ as required. \square

⁷Although this lemma is stated for differentiable functions, it is easy to check that the proof goes through without differentiability since it only uses $f(b) - f(a) \geq \int_a^b \frac{d^-}{dx} f(x) dx$.

REFERENCES

- [1] R.J. Adler and J.E. Taylor. *Random fields and geometry*. Springer, 2007.
- [2] M. Aizenman and D.J. Barsky. Sharpness of the phase transition in percolation models. *Comm. Math. Phys.*, 108(3):489–526, 1987.
- [3] M. Aizenman and C.M. Newman. Tree graph inequalities and critical behavior in percolation models. *J. Stat. Phys.*, 36:107–143, 1984.
- [4] V. Beffara and D. Gayet. Percolation of random nodal lines. *Publ. Math. IHES*, 126(1):131–176, 2017.
- [5] D. Beliaev, M. McAuley, and S. Muirhead. Fluctuations in the number of excursion sets of planar gaussian fields. *arXiv preprint arXiv:1908.10708*, 2019.
- [6] D. Beliaev, M. McAuley, and S. Muirhead. Smoothness and monotonicity of the excursion set density of planar gaussian fields. *Electron. J. Probab.*, 25(93):1–37, 2020.
- [7] D. Beliaev, S. Muirhead, and A. Rivera. A covariance formula for topological events of smooth Gaussian fields. *Ann. Probab. (to appear)*, 2020.
- [8] I. Benjamini, O. Schramm, and D.B. Wilson. Balanced Boolean functions that can be evaluated so that every input bit is unlikely to be read. In *STOC'05: Proceedings of the 37th Annual ACM Symposium on Theory of Computing*, pages 244–250, 2005.
- [9] C. Borgs, J. T. Chayes, H. Kesten, and J. Spencer. Uniform boundedness of critical crossing probabilities implies hyperscaling. *Random Struct. Algo.*, 15(3–4):368–413, 1999.
- [10] J. T. Chayes and L. Chayes. Finite-size scaling and correlation lengths for disordered systems. *Phys. Rev. Lett.*, 57(24):2999–3002, 1986.
- [11] J. T. Chayes and L. Chayes. Inequality for the infinite-cluster density in Bernoulli percolation. *Phys. Rev. Lett.*, 56(16):1619–1622, 1986.
- [12] V. Dewan and D. Gayet. Random pseudometrics and applications. *arXiv preprint arXiv:2004.05057*, 2020.
- [13] H. Duminil-Copin, S. Goswami, P.-F. Rodriguez, and F. Severo. Equality of critical parameters for percolation of Gaussian free field level-sets. *arXiv preprint arXiv:2002.07735*, 2020.
- [14] H. Duminil-Copin, I. Manolescu, and V. Tassion. Planar random-cluster model: fractal properties of the critical phase. *arXiv preprint arXiv:2007.14707*, 2020.
- [15] H. Duminil-Copin, A. Raoufi, and V. Tassion. Exponential decay of connection probabilities for subcritical Voronoi percolation in \mathbb{R}^d . *Probab. Theory Related Fields*, 173(1–2):479–490, 2019.
- [16] H. Duminil-Copin, A. Raoufi, and V. Tassion. Sharp phase transition for the random-cluster and Potts models via decision trees. *Ann. Math.*, 189(1):75–99, 2019.
- [17] H. Duminil-Copin, A. Raoufi, and V. Tassion. Subcritical phase of d -dimensional Poisson-Boolean percolation and its vacant set. *Ann. H. Lebesgue*, 3:677–700, 2020.
- [18] H. Duminil-Copin and V. Tassion. A new proof of the sharpness of the phase transition for Bernoulli percolation and the Ising model. *Comm. Math. Phys.*, 343:725–745, 2016.
- [19] W. Ehm, T. Gneiting, and D. Richards. Convolution roots of radial positive definite function with compact support. *Trans. Am. Math. Soc.*, 356(11):4655–4685, 2004.
- [20] R. Fitzner and R. van der Hofstad. Mean-field behavior for nearest-neighbor percolation in $d > 10$. *Electron. J. Probab.*, 22:65 pp., 2017.
- [21] C. Garban, G. Pete, and O. Schramm. The Fourier spectrum of critical percolation. *Acta Math.*, 205(1):19–104, 2010.
- [22] C. Garban and H. Vanneuville. Bargmann-Fock percolation is noise sensitive. *arXiv preprint arXiv:1906.02666*, 2019.
- [23] G.R. Grimmett. *Percolation*. Springer, 1999.
- [24] J.M. Hammersley. Percolation processes: Lower bounds for the critical probability. *Ann. Math. Statist.*, 28:790–795, 1957.
- [25] D.G. Handron. Generalized billiard paths and Morse theory for manifolds with corners. *Topology Appl.*, 126(1-2):83–118, 2002.
- [26] T. Hara. Mean-field critical behaviour for correlation length for percolation in high dimensions. *Probab. Theory Related Fields*, 86:337–385, 1990.
- [27] T. Hara. Decay of correlations in nearest-neighbor self-avoiding walk, percolation, lattice trees and animals. *Ann. Probab.*, 36(2):530–593, 2008.
- [28] H. Kesten. *Percolation theory for mathematicians*. Progress in Probability and Statistics Vol. 2. Springer, 1982.
- [29] H. Kesten. Scaling relations for 2D-percolation. *Comm. Math. Phys.*, 109:109–156, 1987.
- [30] L. Köhler-Schindler and V. Tassion. Crossing probabilities for planar percolation. *arXiv preprint arXiv:2011.04618*, 2020.
- [31] G. Kozma and A. Nachmias. Arm exponents in high dimensional percolation. *J. Amer. Math. Soc.*, 24(2):375–409, 2011.
- [32] S. Kullback. *Information theory and statistics*. Dover, 1978.
- [33] M. Ledoux. A short proof of the Gaussian isoperimetric inequality. In E. Eberlein, M. Hahn, and M. Talagrand, editors, *High Dimensional Probability. Progress in Probability, vol 43.*, pages 229–232. Birkhäuser, Basel, 1998.

- [34] M. Menshikov. Coincidence of critical points in percolation problems. *Sov. Math. Dokl.*, 33:856–859, 1986.
- [35] S.A. Molchanov and A.K. Stepanov. Percolation in random fields. I. *Theor. Math. Phys.*, 55(2):478–484, 1983.
- [36] S.A. Molchanov and A.K. Stepanov. Percolation in random fields. II. *Theor. Math. Phys.*, 55(3):592–599, 1983.
- [37] S. Muirhead, A. Rivera, and H. Vanneuille (with an appendix by L. Köhler-Schindler). The phase transition for planar Gaussian percolation models without FKG. *arXiv preprint arXiv:2010.11770*, 2020.
- [38] S. Muirhead and H. Vanneuille. The sharp phase transition for level set percolation of smooth planar Gaussian fields. *Ann. I. Henri Poincaré Probab. Stat. (to appear)*, 2020.
- [39] R. O’Donnell, M. Saks, O. Schramm, and R.A. Servedio. Every decision tree has an influential variable. In *46th Annual IEEE Symposium on Foundations of Computer Science (FOCS’05)*, pages 31–39, 2005.
- [40] R. O’Donnell and R.A. Servedio. Learning monotone decision trees in polynomial time. *SIAM J. Comput.*, 37(3):827–844, 2007.
- [41] L.D. Pitt. Positively correlated normal variables are associated. *Ann. Probab.*, 10(2):496–499, 1982.
- [42] A. Rivera. Talagrand’s inequality in planar Gaussian field percolation. *arXiv preprint arXiv:1905.13317*, 2019.
- [43] A. Rivera and H. Vanneuille. Quasi-independence for nodal lines. *Ann. Inst. H. Poincaré Probab. Statist.*, 55(3):1679–1711, 2019.
- [44] A. Rivera and H. Vanneuille. The critical threshold for Bargmann-Fock percolation. *Ann. Henri Lebesgue*, 3, 2020.
- [45] W. Rudin. An extension theorem for positive-definite functions. *Duke Math. J.*, 37(1):49–53, 1970.
- [46] O. Schramm and S. Smirnov (with an appendix by C. Garban). On the scaling limits of planar percolation. *Ann. Probab.*, 39(5):1768–1814, 2011.
- [47] S. Smirnov and W. Werner. Critical exponents for two-dimensional percolation. *Math. Res. Lett.*, 8(5):729–744, 2001.
- [48] H. Tasaki. Hyperscaling inequalities for percolation. *Comm. Math. Phys.*, 113(1):49–65, 1987.
- [49] R. van den Berg and H. Don. A lower bound for point-to-point connection probabilities in critical percolation. *Electron. Comm. Probab.*, (47), 2020.
- [50] R. van den Berg and P. Nolin. On the four-arm exponent for $2d$ percolation at criticality. *arXiv preprint arXiv:2008.01606*, 2020.
- [51] A. Weinrib. Long-range correlated percolation. *Phys. Rev. B*, 29(1):387, 1984.