



**HAL**  
open science

## Estrogenic activity of surface waters using zebrafish- and human-based in vitro assays: The Danube as a case-study

Hélène Serra, François Brion, Clémence Chardon, Hélène Budzinski, Tobias Schulze, Werner Brack, Selim Aït-Aïssa

### ► To cite this version:

Hélène Serra, François Brion, Clémence Chardon, Hélène Budzinski, Tobias Schulze, et al.. Estrogenic activity of surface waters using zebrafish- and human-based in vitro assays: The Danube as a case-study. *Environmental Toxicology and Pharmacology*, 2020, 78, pp.art. 103401. 10.1016/j.etap.2020.103401 . hal-03159755

**HAL Id: hal-03159755**

**<https://hal.science/hal-03159755v1>**

Submitted on 9 Aug 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Estrogenic activity of surface waters using zebrafish- and human-based *in vitro* assays:**  
2 **the Danube as a case-study**

3 Hélène Serra<sup>1,2</sup>, François Brion<sup>1</sup>, Clémence Chardon<sup>1</sup>, Hélène Budzinski<sup>2</sup>, Tobias Schulze<sup>3</sup>,  
4 Werner Brack<sup>3,4</sup>, Selim Aït-Aïssa<sup>1\*</sup>

5 <sup>1</sup>Unité Ecotoxicologie *in vitro* et *in vivo*, UMR-I 02 SEBIO, Institut National de  
6 l'Environnement Industriel et des risques (INERIS), Verneuil-en-Halatte, France

7 <sup>2</sup>UMR-CNRS EPOC/LPTC, Université de Bordeaux, Talence, France

8 <sup>3</sup>UFZ– Helmholtz Centre for Environmental Research, Leipzig, Germany

9 <sup>4</sup>RWTH Aachen University, Aachen, Germany

10 \* corresponding author: [selim.ait-aissa@ineris.fr](mailto:selim.ait-aissa@ineris.fr)

11

12

13 **HIGHLIGHTS**

14

- 15 • *In vitro* evaluation of the estrogenicity of 25 Danube water samples
- 16 • Most samples have a low estrogenic activity
- 17 • Both human and zebrafish bioassays allow the identification of two hot spot sites
- 18 • Estrogenic effect is confirmed *in vivo* using the zebrafish EASZY assay
- 19 • Established effect-based trigger values for human bioassays allow prioritisation

20

21

22 **ABSTRACT**

23 Most *in vitro* reporter gene assays used to assess estrogenic contamination are based on  
24 human estrogen receptor  $\alpha$  (hER $\alpha$ ) activation. However, fish bioassays can have distinct  
25 response to estrogenic chemicals and mixtures, questioning the relevance of human-based  
26 bioassays for assessing risk to this species. In this study, zebrafish liver cells stably expressing  
27 zebrafish ER $\beta$ 2 (ZELH $\beta$ 2) and human breast cancer cells expressing hER $\alpha$  (MELN) were  
28 used to quantify the estrogenic activity of 25 surface water samples of the Danube River, for  
29 which chemicals have been previously quantified. Most samples had a low estrogenic activity  
30 below 0.1 ng/L 17 $\beta$ -estradiol-equivalents that was more often detected by MELN cells, while  
31 ZELH $\beta$ 2 response tend to be lower than predicted based on the chemicals identified.  
32 Nevertheless, both bioassays quantified well a higher estrogenic activity at two sites, which  
33 was confirmed *in vivo* using a transgenic zebrafish assay. The results are discussed  
34 considering the effect-based trigger values proposed for water quality monitoring.

35

## 36 1. INTRODUCTION

37 The evidence that estrogenic chemicals occurring in the aquatic environment may adversely  
38 affect the endocrine system of fish has led to a growing concern regarding their long-term  
39 reproductive effects (Sumpter and Jobling 2013). In 2012, the European Commission  
40 proposed to set environmental quality standards (EQS) for estradiol (E2), estrone (E1) and  
41 ethynyl estradiol (EE2) at 0.4 ng/L, 3.6 ng/L and 0.035 ng/L, respectively (European  
42 Commission 2015). However, most current analytical methods are not sensitive enough to  
43 quantify E2 and EE2 at these very low concentrations, below the ng/L (Kase et al. 2018). For  
44 instance, the results of the European Union (EU)-wide monitoring exercise for chemicals on  
45 the Watch List showed that only half of EU member states achieved to implement a method  
46 with a limit of quantification at the EQS level for EE2 (Loos et al. 2018). Consequently, the  
47 lack of sufficiently sensitive and cost-effective analytical methods for monitoring estrogenic  
48 chemicals considerably hinders their prioritization under the European Water Framework  
49 Directive (WFD).

50 *In vitro* bioassays based on the activation of the estrogen receptor (ER) are sensitive effect-  
51 based methods to assess estrogenic chemicals (Wernersson et al. 2015). Integrated with  
52 analytical tools, they have been successfully applied to quantify the estrogenic activity of  
53 surface and wastewater, and they showed to have a good sensitivity to detect E1, E2 or EE2 at  
54 environmental levels, highlighting their potential to support chemical monitoring (Leusch et  
55 al. 2010, Könemann et al. 2018, Kunz et al. 2015). Furthermore, they enable to detect active  
56 chemicals in complex mixtures without *a priori* knowledge, as they take into account the  
57 combined effects of both known and unknown substances targeting the ER, and, thus, they  
58 provide a holistic evaluation of estrogenic mixtures (Altenburger et al. 2019).

59 Most *in vitro* bioassays used in biomonitoring are mammalian or yeast-based system  
60 expressing the human ER $\alpha$  (hER $\alpha$ ), which controls the expression of a reporter gene used to  
61 quantify the biological activity. Compared to the human genome that encodes only for two ER  
62 subtypes, hER $\alpha$  and hER $\beta$ , most teleost fish express at least 3 ER subtypes, ER $\alpha$ , ER $\beta$ 1 and  
63 ER $\beta$ 2 (Tohyama et al. 2016, Menuet et al. 2002) that can have distinct sensitivities to  
64 estrogenic chemicals compared to human isoforms (Cosnefroy et al. 2009, Matthews et al.  
65 2000, Pinto et al. 2014). For instance, zebrafish zfER $\beta$  subtypes are one order on magnitude  
66 more sensitive to E2 than zfER $\alpha$ , while the reverse is observed for human ERs (Pinto et al.  
67 2014). Among zfER subtypes, zfER $\beta$ 2 is the most sensitive and responsive one to various  
68 estrogenic substances (Cosnefroy et al., 2012). In addition to structural differences at the  
69 receptor level, the response to estrogens is influenced by the cell context, such as the presence  
70 of transcriptional co-factors or metabolic pathways, which confer a tissue-specific response  
71 (Ohtake et al. 2003). In the aquatic environment, many studies reported that fish are among  
72 the most sensitive organisms to xenoestrogens (Tyler et al. 1998, Matthiessen et al 2018) and  
73 the question of the relevance of human-based bioassays to predict an effect in fish can thus be  
74 raised (Hotchkiss et al 2008).

75 Recent field studies showed that *in vitro* estradiol-equivalents (EEQs) measured in surface  
76 and waste waters correlated to *in vivo* responses measured in fish models (Ihara et al. 2015,  
77 Brion et al. 2019). Nevertheless, some studies suggested that a better correlation is found  
78 with fish-based bioassays. Using wastewater samples, Ihara et al. (2015) showed that the  
79 induction of vtg1/chgH in male medaka was better correlated with *in vitro* medaka ER $\alpha$   
80 activation than with human ER $\alpha$ . Interactions between chemicals leading to co-occurrence of  
81 estrogenic and anti-estrogenic activities were suggested to drive the observed differences  
82 (Ihara et al. 2014). In a previous field study, we also reported inter-specific qualitative and

83 quantitative differences between human and fish-based cell lines for surface water extracts,  
84 with some samples being selectively active in zebrafish liver cells expressing zfER $\beta$ 2  
85 (ZELH $\beta$ 2 cells) and in an *in vivo* zebrafish assay but not in human MELN cells (Sonavane et  
86 al. 2016). Furthermore, ZELH $\beta$ 2 and MELN cells have distinct sensitivity to anti-estrogenic  
87 chemicals, which can significantly influence the response to estrogenic chemicals in a cell-  
88 specific manner (Serra et al. 2019). These results suggest the presence of fish-specific factors  
89 influencing the *in vitro* estrogenic response to environmental extracts that warrants further  
90 research.

91 This study was carried out to gain further knowledge on possible differences between fish-  
92 and human-based bioassays in the assessment of the estrogenic activity of river water sample  
93 extracts, including the evaluation of the contribution of known estrogenic chemicals to the  
94 measured activity. For that purpose, we used water extracts from the Danube River, which  
95 have been previously characterized chemically and are representative of a large river with  
96 high dilution capacity and highly diverse chemical contamination pattern (Liška et al. 2015).  
97 The water was sampled using on-site large-volume solid phase extraction (LVSPE) during the  
98 Join Danube Survey 3 (Neale et al. 2015) and in Novi Sad city (Serbia) that discharges  
99 untreated wastewater directly into the Danube River (König et al. 2017). The biological  
100 responses of both *in vitro* bioassays were compared with the estrogenic chemicals identified  
101 by target chemical analyses. In addition, we assessed the anti-estrogenic activity of Novi Sad  
102 samples and evaluated the *in vivo* estrogenic response in transgenic zebrafish embryos using  
103 the EASZY assay (Brion et al. 2012). The response of the bioassays is discussed considering  
104 the effect-based trigger values (EBT) proposed to implement the bioassays in a regulatory  
105 context.

106

## 107 2. MATERIAL AND METHODS

### 108 2.1. Chemicals and reagents

109 17 $\beta$ -estradiol (E2, CAS 50-28-2, purity of >98%), ethynyl estradiol (EE2, CAS n° 57-63-6,  
110 purity of >98%) and hydroxy-tamoxifen (OH-TAM, CAS 68392-35-8, purity of >98%),  
111 dimethylsulfoxide (DMSO), Leibovitz 15 culture medium (L-15), fetal calf serum (FCS), 4-  
112 (2-hydroxy-ethyl)-1-piperazineethanesulfonic acid (HEPES), epidermal growth factor (EGF),  
113 G418, 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl-tetrazoliumbromide (MTT) and D-luciferin  
114 were purchased from Sigma Aldrich (St-Quentin Fallavier, France). Dulbecco's Modified  
115 Eagle Medium High Glucose (DMEM HG) powder, F-12 nutrient mixture (Ham's F12)  
116 powder, penicillin and streptomycin were purchased from Gibco (France). Insulin,  
117 hygromycin B and sodium bicarbonate were purchased from Dominique Dutscher (France).

### 118 2.2. Study sites, sampling and extraction procedure

119 Two sets of Danube River water samples were collected and prepared during the Joint  
120 Danube Survey 3 (JDS, 22 samples) and Novi Sad (NS, 3 samples) campaigns as detailed  
121 previously (Neale et al. 2015, König et al. 2017). In brief, sampling of the JDS was carried  
122 out between August and September 2013 in Danube River and some tributaries, from Austria  
123 to Romania (Liška et al. 2015). The city of Novi Sad city (300 000 inhabitants, Serbia),  
124 located on the Danube River, was selected to investigate the impact of untreated municipal  
125 effluent release on the Danube River (König et al. 2017) and to identify drivers of endocrine  
126 disruption including estrogenicity and androgenicity (Hashmi et al. 2018) and progestogenic  
127 and glucocorticoid receptor-mediated effects (Hashmi et al. 2020). The Novi Sad sites  
128 investigated were selected as follows: NS1 site 2 km upstream the effluent release, NS2 site  
129 200 m downstream the effluent release, and NS3 site 7 km downstream the effluent release. In



130 both case-studies, the samples were collected using an on-site solid phase extraction (SPE)  
131 device designed to sample and extract large volume (LV) of water on site (LVSPE, Schulze et  
132 al. 2017). About 500 L of water were extracted and concentrated at JDS sites, and 850 L to  
133 1000 L at NS sites supporting a large set of investigations (Schulze et al. 2017) while for the  
134 present study aliquots representing 6 to 12 L water equivalents were used. The organic  
135 extracts were reduced to dryness before shipping and were then resuspended in DMSO and  
136 stored at -20 °C before bioanalysis.

### 137 **2.3. *In vitro* bioassays**

138 The zebrafish *in vitro* assay derived from the zebrafish liver (ZFL) cell line. ZFL cells were  
139 stably transfected first by an ERE-driven luciferase gene, yielding the ZELH cell line, and  
140 then by zfER $\beta$ 2 subtype yielding the ZELH $\beta$ 2 cell lines (Cosnefroy et al. 2012). The  
141 establishment of this cell model and its response to different classes of well-known  
142 xenoestrogens have been previously described (Cosnefroy et al. 2012, Serra et al. 2019, Le  
143 Fol et al. 2017, Sonavane et al. 2016). In addition, we used the human-derived MELN cell  
144 line (Balaguer et al. 1999) kindly provided by Dr. Patrick Balaguer (INSERM Montpellier,  
145 France). The MELN cells are derived from the breast cancer MCF-7 cells that endogenously  
146 express the hER $\alpha$ , but no functional hER $\beta$  (P. Balaguer, *personal communication*). MCF-7  
147 cells were stably transfected with an ERE-driven firefly luciferase reporter gene to yield  
148 MELN cells.

149 Conditions for routine cell culture and exposure to chemicals have been detailed previously  
150 (Sonavane et al. 2016). Briefly, ZELH $\beta$ 2 cells were seeded in 96-well white opaque culture  
151 plates (Greiner CellStar™, Dutscher, France) at 25,000 cells per well in phenol red free LDF-  
152 DCC medium (containing L-15 50%, DMEM HG 35%, Ham's F12 15%, HEPES 15 mM,

153 0.15 g/L sodium bicarbonate, 0.01 mg/mL insulin, 50 ng/mL EGF, 50 U/mL penicillin and  
154 streptomycin antibiotics, 5% v/v stripped FCS). MELN were seeded at 80,000 cells per well  
155 in steroid- and phenol red-free DMEM medium. Cells were left to adhere for 24 h, and then  
156 exposed for either 72 h at 28 °C for zebrafish cells or 16h at 37°C for MELN cells. Cells were  
157 exposed in triplicates to serial dilutions of sample extracts to obtain a final DMSO  
158 concentration in the well plate of 0.5% (v/v). To assess anti-estrogenic activity at selected  
159 sites, ZELHβ2 and MELN cells were co-exposed to the sample in presence of 0.1 nM E2, i.e.  
160 a concentration saturating 80% to 100% of ER. Each plate included both solvent (DMSO) and  
161 positive (E2 for estrogenic activity, OH-TAM for anti-estrogenic activity) controls. After  
162 exposure, culture medium was removed and replaced by 50 µl per well of medium containing  
163 0.3 mM D-luciferin. The luminescence signal was measured in living cells using a microtiter  
164 plate luminometer (Synergy H4, BioTek). The effect of samples on cell viability was assessed  
165 by using the 3-(4,5-dimethyl-thiazol-2-yl)-2,5diphenyl tetrazolium bromide (MTT) assay  
166 (Mosmann 1983). After cell exposure, culture medium was removed and replaced by 100 µL  
167 of medium containing 0.5 mg/mL MTT. Cells were incubated for 1 h to 3 h. In metabolically  
168 active cells, MTT is reduced onto a blue formazan precipitate, which is dissolved by adding  
169 100 µL of DMSO after removal of MTT containing medium. Plates were read at 570 nm  
170 against a 640 nm reference wavelength on a microplate reader (KC-4, BioTek Instruments,  
171 France) and results are expressed as absorbance relative to control cells.

#### 172 ***2.4. Zebrafish embryo-based bioassay (EASZY assay)***

173 The estrogenic activity of Novi Sad samples was assessed *in vivo* using the EASZY assay  
174 based on transgenic *cyp19a1b*-GFP zebrafish line (Brion et al. 2012). The assay procedure for  
175 sample testing has been described in Sonavane et al. (2016) and Brion et al. (2019). In brief,

176 15 newly fertilized transgenic eggs (< 4h post fertilization) were selected for each  
177 experimental group and exposed for 96 h in 15 mL of acclimated water in glass crystallizers.  
178 Serial dilutions were tested with a final volume of solvent (DMSO) of 0.1% v/v, a  
179 concentration without effects on embryo development or GFP expression. In each  
180 experimental series, positive (EE2 0.05 nM) and DMSO controls were included as separate  
181 experimental groups. Exposed embryos were incubated at 28 °C, under semi-static conditions  
182 with daily complete renewal of medium. At the end of the exposure period, each zebrafish  
183 embryo was photographed using a Zeiss AxioImager.Z1 microscope equipped with an  
184 AxioCam Mrm camera (Zeiss GmbH, Gottingen, Germany) to measure GFP expression in the  
185 brain. Image analysis was performed using the ImageJ software, and fluorescence data was  
186 treated exactly as previously described (Brion et al. 2012).

## 187 **2.5. Data analysis**

188 Sample concentration data are expressed in relative enrichment factor (REF) that considers  
189 both sample enrichment in LVSPE and further dilution in the test system. For all *in vitro* data,  
190 the luciferase activity was normalized between 0 and 1, as described in equation (1), with  
191  $signal_{sample}$  the signal of the tested sample,  $signal_{control}$  the signal of the solvent control and  
192  $signal_{positive}$  the signal of the positive control (E2):

$$193 \quad Response = \frac{signal_{sample} - signal_{control}}{signal_{positive} - signal_{control}} \quad (1)$$

194 Concentration-response curves of bioassay data were fitted to all replicates pooled together  
195 with a minimum of two independent experiments (each with three technical replicates) using  
196 Hill equation in the RegTox 7.5 Microsoft Excel™ macro (freely available at  
197 [http://www.normalesup.org/~vindimian/fr\\_download.html](http://www.normalesup.org/~vindimian/fr_download.html)). Positive control 20 (PC20, a

198 concentration corresponding to 20% of the response of the positive control) were estimated  
199 for samples reaching at least 20% effect by fixing the minimum and the maximum of the  
200 sample response to 0 and 1, respectively, and fixing the slope to the one of E2. The biological  
201 estradiol-equivalent (Bio-EEQ) was then calculated as the ratio of EC20 of positive control to  
202 the PC20 of the sample, expressed in ng/L EEQ, as presented in equation (2):

$$203 \quad BioEEQ = \frac{EC20_{positive\ control}}{PC20_{sample}} \quad (2)$$

204 The MELN raw data have been previously analyzed using PC10 value and linear regression  
205 model (Neale et al. 2015). The data were re-evaluated in the present study using PC20 and  
206 Hill model and a very good agreement between both Bio-EEQ calculation methods was  
207 observed (Figure SI 1).

208 To assess the contribution of detected chemicals to the observed biological response, chemical  
209 equivalents (Chem-EEQ) were calculated for the estrogenic chemicals as the sum of the  
210 concentration of each active chemical weighted by their relative estrogenic potency (REP),  
211 based on the equation (3):

$$212 \quad ChemEEQ = \sum_i^n REP_i \times C_i \quad (3)$$

213 The REP is defined as the ratio of PC20 of E2 to the PC20 of the chemical, or as the ratio of  
214 EC50s, when no PC20 is available. The contribution of the quantified chemicals to the  
215 biological response was assessed by dividing the Chem-EEQ by the Bio-EEQ, expressed in  
216 %.

217 In addition, the concentration addition (CA) model was used to predict the additive effects of  
218 identified estrogenic chemicals along the entire dose-response curve, to facilitate graphical

219 comparison of observed and predicted estrogenic activity. For that purpose, single chemical  
220 data were fitted with logit non-linear regression model in GraphPad Prism v.5 and the logit  
221 parameters derived were entered into an Excel template to predict the additive response using  
222 the concentrations and mixture ratios of the active chemicals identified in each sample (Serra  
223 et al. 2019, Altenburger et al. 2018). For simplification purpose, only the active chemicals  
224 contributing to more than 5% of the predicted response at REF of 1 were included in CA  
225 prediction. Logit equation and the Excel template were kindly provided by Martin Scholze  
226 (Brunel University, UK).

227

### 228 **3. RESULTS**

#### 229 *3.1. The estrogenic activity of Danube water extracts is overall low*

230 The results of the estrogenic activity of JDS and Novi Sad samples measured in MELN and  
231 ZELH $\beta$ 2 bioassays are presented in Table 1, and the individual concentration-response curves  
232 are provided in supplementary information (figures SI 2 and SI 3 for MELN and ZELH $\beta$ 2  
233 cells, respectively).

234 Overall, the estrogenic activity measured in the JDS samples was relatively low, in both cell  
235 models. In MELN cells, 15 out of 22 JDS samples had an estrogenic activity above the LOQ  
236 (0.002 ng/L EEQ). The response measured was below 0.3 ng/L EEQ in all samples, except for  
237 JDS41 that had an estrogenic activity of 0.74 ng/L EEQ. A similar pattern was observed with  
238 the ZELH $\beta$ 2 bioassay, with, however, only 8 samples being positively quantified. In all  
239 samples, the estrogenic activity measured in ZELH $\beta$ 2 was below 0.08 ng/L EEQ, except for  
240 JDS41 that reached 2.3 ng/L EEQ. The comparison of both MELN and ZELH $\beta$ 2 showed that  
241 10 samples were equally detected in both cell lines (being either active or inactive), while 2

242 samples were selectively detected in ZELH $\beta$ 2 cells (JDS63, JDS67) and 10 samples only in  
243 MELN cells (e.g. JDS22, JDS27, JDS30).

244 The concentration-response curves of Novi Sad samples are presented in Figure 1, and the  
245 Bio-EEQs are given in Table 1. The estrogenic activity pattern of Novi Sad samples was  
246 similar between MELN and ZELH $\beta$ 2 bioassays, with a peak of activity detected just after the  
247 effluent release at NS2 site. However, the NS2 sample had a higher estrogenic activity in  
248 MELN cells (1.52 ng/L EEQ) compared to the one measured in ZELH $\beta$ 2 cells (0.19 ng/L  
249 EEQ).

250 When quantified by the two assays, Bio-EEQs provided by the two models were overall  
251 correlated (Figure SI 4) although MELN assay tended to quantify higher levels (Table 1).

### 252 *3.2. The identified estrogenic chemicals only partially explained the fish-specific* 253 *response*

254 We further assessed the contributions of estrogenic chemicals identified in the samples to the  
255 activity measured in fish and human-based bioassays. More than 100 chemicals were  
256 quantified in JDS and Novi Sad samples (König et al. 2017, Neale et al. 2015), and  
257 information about the activity of the chemicals on MELN and ZELH $\beta$ 2 cells was available  
258 only for 36 substances, among which 7 were estrogenic on ZELH $\beta$ 2 cells and 9 on MELN  
259 cells. The Relative Potency (REP) of these xenoestrogens for each cell line are indicated in  
260 Table 2, and the concentrations of the active chemicals across all sampling sites are presented  
261 in Table SI 1. These chemicals include natural steroid estrogens, bisphenols, phytoestrogens  
262 and pesticides.

263 The Chem-EEQs of the 25 samples and their contribution to the biological response (Bio-  
264 EEQs) are presented in Table 1. Overall, the same main contributors to the estrogenic activity  
265 were detected for both cell lines: estrone and genistein, confirming our previous report (Neale  
266 et al. 2015). It is worth to note that EE2 was included in the analyses but was never detected  
267 (LOQ 0.4 ng/L). Regarding JDS samples, estrone and genistein together explained 19% to  
268 124% of the biological response observed in MELN cells for most samples, except at JDS27  
269 and JDS60 (<1%, no estrone detected), and at JDS64 (710%, very weak estrogenic response  
270 measured). Furthermore, 5 samples were expected to have an estrogenic activity ranging from  
271 0.18 to 0.58 ng/L EEQ in MELN cells based on the chemical quantified, but were not detected  
272 by the bioassay (JDS35, JDS57, JDS59, JDS63 and JDS65), as confirmed by comparing CA  
273 prediction curve with the observed response (Figure SI 2). Conversely, JDS27 and JDS60  
274 were active in MELN cells albeit no significant contributors could be identified. In ZELH $\beta$ 2  
275 cells, estrone and genistein explained 11% to 136% of the observed biological response of  
276 JDS samples, except at 3 sites for which the biological activity was lower than expected based  
277 on the estrogenic chemicals quantified (JDS29: 503%, JDS39: 306% and JDS67: 214%). In  
278 contrast, 11 samples were predicted to have an estrogenic activity in ZELH $\beta$ 2 cells, ranging  
279 from 0.022 to 0.065 ng/L EEQ, as predicted by CA model, but were not detected by the  
280 bioassay (Table 1 and Figure SI 3).

281 Because estrone is a major driver of estrogenicity in JDS samples, it may thus contribute to  
282 estrogenic activity of the extracts even if present at concentration below or at its LOQ (i.e. 0.1  
283 ng/L). Thus, for those few sites with very low contribution of Chem-EEQ in MELN cells,  
284 namely JDS 27 and JDS60, including estrone LOQ in the mass balance calculation increases  
285 the contribution of Chem-EEQs up to 13% and 16%, respectively. In all cases, maximized

286 Chem-EEQs remain below Bio-EEQ and suggest that other non-detected active compounds  
287 are present in the samples at these sites.

288 As for most JDS samples, genistein was detected in all Novi Sad samples but explained only a  
289 small fraction (less than 10%) of the observed response, in both MELN and ZELH $\beta$ 2 cells  
290 (Figure 2). In NS2 sample, the steroidal estrogens (E1, E2, E3) were the main drivers  
291 identified. They were adequately detected in MELN cells (contribution of 88% to the  
292 biological response); however, the activity of NS2 measured in ZELH $\beta$ 2 cells was about 7  
293 times lower than predicted based on Chem-EEQ (Table 1, Figure 1). To investigate possible  
294 negative interactions on the estrogenic response, we assessed the anti-estrogenic activity of  
295 Novi Sad samples by co-exposing the cells to the samples with E2 at a concentration inducing  
296 80% of E2 maximal response. As presented in Figure 1, a decrease in E2-induced luciferase  
297 activity was noted but only at the highest non-cytotoxic concentration tested. Thus, no  
298 significant anti-estrogenic activity could be evidenced, especially regarding NS2 sample  
299 extract in ZELH $\beta$ 2 cells.

### 300 ***3.3. In vivo estrogenic activity of Novi Sad samples***

301 In order to further investigate the toxicological relevance of detected estrogenic activities at  
302 Novi Sad sites, we assessed *in vivo* effect of the samples in zebrafish embryos that express  
303 GFP under control of the ER-regulated *cyp19a1b* promoter, applying the EASZY assay  
304 (Brion et al., 2012, Brion et al., 2019). As presented in Figure 3A, a significant response was  
305 observed at NS2 site, while no effect was detected at the upstream and downstream sites. The  
306 NS2 sample had an estrogenic activity of 3.3 ng/L EEQ in transgenic zebrafish embryos, in  
307 very good agreement with the chemicals identified, E1, E2 and E3 being the main  
308 contributors *in vivo* (Figure 3B).



## 310 4. DISCUSSION

### 311 4.1. Comparative evaluation of estrogenic activity in fish and human-based bioassays

312 Using an integrated analytical-bioanalytical method, we assessed the estrogenic activity of the  
313 Danube River combining zebrafish ZELH $\beta$ 2 and human MELN cells. Our results show that,  
314 with the exception of two sites, the estrogenic activity of the Danube River was overall low.  
315 The majority of samples had a Bio-EEQ below 0.1 ng/L EEQ both in MELN cells (18/25  
316 samples) and in ZELH $\beta$ 2 cells (23/25 samples). The low estrogenicity measured is consistent  
317 with the high dilution capacity of the large Danube River, resulting in trace levels of  
318 contaminants. Estrone was the main estrogenic chemical detected in the Danube samples, in  
319 line with previous studies evaluating surface water estrogenicity in other contexts (Conley et  
320 al. 2017, Alvarez et al. 2013). Overall, we show that, although both cell lines are sensitive,  
321 levels of estrogenic activity below 0.1 ng/L EEQ were more often detected by MELN cells  
322 than ZELH $\beta$ 2 cells. Indeed, the estrogenic activity predicted in ZELH $\beta$ 2 cells tend to be  
323 higher than the biological response observed, whenever the sample was detected.

324 The latter outcome contrasts with the results of a previous study that compared the estrogenic  
325 activity of POCIS extracts deployed on 20 French river sites (Sonavane et al. 2016). In the  
326 latter, ZELH $\beta$ 2 cells revealed an estrogenic activity at 8 sites that was not detected by MELN  
327 cells. The authors hypothesized the presence of fish-selective ER active chemicals, although  
328 the samples were not characterized chemically. In the current study, we cannot argue in  
329 favour of bioassay-specific estrogenic substances, as genistein and estrone, identified as main  
330 estrogenic drivers, are active in both MELN and ZELH $\beta$ 2 cells. The lower response of  
331 zebrafish cells might be explained by a higher sensitivity of ZELH $\beta$ 2 cells over MELN cells

332 to non-ER chemicals present in the mixture and modulating the zER $\beta$ 2 response (Serra et al.  
333 2019). In the highly diluted context of Danube, estrogenic compounds are at very low  
334 concentrations and in presence of a universe of other co-occurring compounds that are also  
335 caught and concentrated by LVSPE. Distinct sampling approaches and pollution patterns may  
336 account for the differences observed between the outcomes of both studies.

337 Despite different ability to pick up low estrogenicity levels, both bioassays quantified a higher  
338 estrogenic activity at JDS41 and at NS2. JDS41 site is located at the confluence of the Molika  
339 Morava (Serbia) and the Danube River. This site was identified as a highly contaminated site  
340 in the Danube survey (Liška et al. 2015), and had, indeed, the highest load in chemicals of all  
341 the JDS samples (Neale et al. 2015). The detected estrogenic chemicals together explained  
342 less than 30% of estrogenic activity at this site. This finding illustrates the complementarity of  
343 chemical and bioanalytical tools to assess environmental xenoestrogens, as the *in vitro*  
344 bioassays were able to detect chemicals not targeted by chemical analysis. In the Novi Sad  
345 case study, the detection of a higher estrogenic activity at NS2 site confirms that the  
346 wastewater release is a point source of environmental xenoestrogens (Könemann et al. 2018,  
347 Sonavane et al. 2018). For instance, the steroidal estrogens E1, E2 and E3 that drove the  
348 estrogenic activity of NS2 sample were well diluted few kilometres downstream of the  
349 release, as shown by both analytical and bioanalytical tools at NS3 site. Furthermore, our  
350 results are coherent with the estrogenic evaluation of the same Novi Sad samples performed  
351 using two hER $\alpha$ -based reporter gene bioassays (König et al. 2017). The estrogenic activity of  
352 NS2 of 1.52 ng/L EEQ measured in MELN cells was higher than the one reported in hER $\alpha$ -  
353 GenBLAzer (0.26 ng/L EEQ) and in BG1-Luc4E (0.67 ng/L EEQ) bioassays, although well  
354 explained by the chemicals quantified (mass balance of 115%). The tendency of MELN cells  
355 to provide higher Bio-EEQ than other reporter gene assays has been evidenced in previous

356 studies, and is believed to account for a higher sensitivity of MELN cells to estrone  
357 (Könemann et al. 2018).

358 Compared to MELN cells, the Bio-EEQ of NS2 sample in ZELH $\beta$ 2 cells was about 7 times  
359 lower than the Chem-EEQ. The lower biological activity in zebrafish cells suggests the  
360 occurrence of negative interactions within the environmental mixture modifying the zfER $\beta$ 2  
361 response to xenoestrogens. We previously showed that bisphenol A and genistein had additive  
362 effects in both ZELH $\beta$ 2 and in MELN cells, however certain environmental contaminants,  
363 such as propiconazole and triphenyl phosphate, were able to selectively decrease the  
364 estrogenic response of ZELH $\beta$ 2 cells but not MELN cells, resulting in deviation from  
365 expected additive response (Serra et al. 2019). These results demonstrated a different  
366 sensitivity to anti-estrogenic chemicals between zebrafish and human ER cell bioassays. Ihara  
367 et al. (2014) documented a similar outcome using wastewater effluents, by showing that the  
368 anti-estrogenic activity detected by medaka ER $\alpha$  bioassay was much more prevalent in  
369 secondary wastewater effluents, while the primary wastewater effluent had much higher anti-  
370 estrogenic activity on human ER $\alpha$  activation. In this study, we did not evidence a strong anti-  
371 estrogenic activity of NS2 extract, and the anti-estrogenic chemicals identified in Serra et al.  
372 (2019) on ZELH $\beta$ 2 cells were not occurring at concentrations expected to have an effect in  
373 the sample, whenever detected. However, the co-occurrence of both ER agonists and  
374 antagonists within the same sample could impede a clear assessment of anti-estrogenic  
375 compounds that may be masked by ER active compounds.

376 Further experiments using sample fractionation and biotesting of the fractions, for instance,  
377 would be warranted to unravel the mixture effect and to evidence possible masking effects, as  
378 previously demonstrated for (anti)androgens in sediments (Weiss et al. 2009),

379 (anti)mineralocorticoids in surface water (Creusot et al. 2014) and progestogen and  
380 glucocorticoid-receptor mediated effects in the same samples (Hashmi et al. 2020).

381 Finally, the ZELH $\beta$ 2 cells originate from zebrafish liver cells that remain metabolically active  
382 (Creusot et al. 2015, Le Fol et al. 2015). Thus, bioassay-specific factors, such as distinct  
383 metabolic capacity and sensitivity of zfER $\beta$ 2, may have also contributed to lower the  
384 estrogenic response of steroidal estrogens in ZELH $\beta$ 2 cells.

#### 385 **4.2. Identification of hot spots and comparison with effect-based trigger values**

386 *In vitro* bioassays are sensitive and specific tools to assess the estrogenic activity of  
387 environmental samples (Könemann et al. 2018). The implementation of bioassays in a  
388 regulatory context, such as the WFD, relies upon the derivation of a biological threshold of  
389 response to enable water managers to discriminate between water bodies that are at low or  
390 high risk (Escher et al. 2018). Several effect-based trigger values (EBT) have been proposed  
391 for *in vitro* estrogenicity bioassays, with different protection goals and construction methods,  
392 as presented in Table 3. Most of the EBT are constructed using primarily fish toxicity data.  
393 For instance, Kase et al. (2018) and Escher et al. (2018) read across from the proposed EQS  
394 derived from chronic toxicity of E1, E2 or EE2 on fish, and Jarošová et al. (2014) used  
395 published long-term PNEC for E1, E2, E3 and EE2 in fish. Brion et al. (2019) compared  
396 empirically the *in vitro* activity of environmental samples with the *in vivo* ER-regulated  
397 *cyp19a1b* expression in transgenic zebrafish embryos. Alternatively, the proposed EBT for  
398 ER-CALUX as part of SIMONI strategy is based on all acute and chronic data on fish,  
399 insects, algae and crustaceans (van der Oost et al. 2017). It is worth noticing that only two  
400 approaches actually take into account xenoestrogen mixture effects, either assuming additive  
401 effects and average environmental occurrence ratio (Escher et al. 2018) or assuming all

402 possible mixture effects based on empirical response of fish embryos to environmental  
403 samples (Brion et al. 2019). With the exception of the EBT proposed by Kunz et al. (2017)  
404 which is a direct translation of the EQS of E2 into EEQ (0.4 ng/L EEQ), all other approaches  
405 consider the bioassay-specific sensitivity (REP) to estrogenic chemicals in deriving the EBT.

406 The EBTs proposed for MELN cells are very consistent one to another, ranging from 0.3 ng/L  
407 to 0.56 ng/L EEQ (Table 3). In the current study, only JDS41 (0.74 ng/L EEQ) and NS2 (1.52  
408 ng/L EEQ) samples exceeded the EBT, including the highest one of 0.56 ng/L EEQ. Thus, the  
409 application of these thresholds for MELN cells would have enabled to identify JDS41 and  
410 NS2 as polluted sites. The risk quotient, defined as the sum of ratios of measured  
411 concentrations of E1, E2 and EE2 to their respective EQS, reaches 0.95 for NS2 sample, as  
412 both estrone and estradiol were at concentrations very close to their EQS. In comparison,  
413 estrone was the only steroidal estrogen quantified in JDS41 sample and had a concentration of  
414 0.196 ng/L, below the EQS level proposed of 3.6 ng/L (risk quotient of 0.05). Thus, JDS41  
415 site would have not been prioritised based only on the steroidal estrogens identified, as  
416 proposed in the WFD.

417 It is noticeable that *in vivo* estrogenic activity at the three NS sites was correctly predicted by  
418 the *in vitro* bioassays in the present study. Since the EBT value of 0.56 ng/L was built based  
419 on a comparison of MELN and EASZY data (Brion et al. 2019), the present dataset confirms  
420 the suitability of this EBT value in another environmental context. Furthermore, the use of the  
421 *in vivo* EASZY assay enabled us to confirm the estrogenic activity observed *in vitro* for NS2  
422 sample at the organism level. It showed that the estrogenic pollutants could target the ER and  
423 induce the expression of the brain aromatase in the developing zebrafish, hence adding further  
424 toxicological relevance to the environmental diagnosis.

425 The prioritization of steroidal estrogens under the WFD relies upon the development of  
426 enough sensitive and cost-effective monitoring methods. In the current study, we show that  
427 both the MELN and ZELH $\beta$ 2 bioassays performed well in detecting a higher estrogenic  
428 activity at two hot spot sites. Furthermore, this study showed that MELN cells detected both  
429 sites exceeding the proposed EBT, allowing a correct prioritisation of sites with a risk. In  
430 contrast, JDS41 site would not have been prioritized based only on the steroidal estrogens  
431 detected, as estrone was quantified at a level 20-time lower than the proposed EQS. Although  
432 both human and fish-based bioassays identified well the two hot spot sites, the relevance of  
433 the deviation from additivity observed in ZELH $\beta$ 2 cells, notably at trace levels, remains to be  
434 addressed.

435

#### 436 **ACKNOWLEDGMENTS**

437 This study was supported by the EU Seventh Framework Programme as a part of  
438 SOLUTIONS project (FP7-ENV-2013-two-stage) under grant agreement number 603437, and  
439 by the French Ministry of Ecology (P190-Ecotoxicologie, P181-DRC60). We wish to thank  
440 Emmanuelle Maillot-Maréchal for her excellent technical support with the cell lines, Martin  
441 Scholze for providing tools and methods to predict additive effects of xenoestrogens, Margit  
442 Petre, Melis Muz and Riccardo Massei for sample preparation and Peter Tarabek and Jörg  
443 Ahlheim for technical help with sample collection. We are grateful to Jaroslav Slobodnik and  
444 his team at Environmental Institute who organised all logistics with the JDS3 sampling.

445

#### 446 **REFERENCES**

447 Altenburger R., Brack W., Burgess R.M., Busch W., Escher B.I., Focks A., Hewitt L.M., Jacobsen B.N.,  
448 Lopez de Alda M., Ait-Aissa S., Backhaus T., Ginebreda A., Hilscherova K., Hollender J., Hollert H.,  
449 Neale P.A., Schulze T., Schymanski E.L., Teodorovic I., Tindall A.J., de Aragao Umbuzeiro G., Vrana  
450 B., Zonja B. and Krauss M. (2019). "Future water quality monitoring: improving the balance

451 between exposure and toxicity assessments of real-world pollutant mixtures." *Environmental*  
452 *Sciences Europe* 31.

453 Altenburger R., Scholze M., Busch W., Escher B.I., Jakobs G., Krauss M., Krueger J., Neale P.A., Ait-  
454 Aissa S., Almeida A.C., Seiler T.-B., Brion F., Hilscherova K., Hollert H., Novak J., Schlichting R., Serra  
455 H., Shao Y., Tindall A., Tolefsen K.-E., Umbuzeiro G., Williams T.D. and Kortenkamp A. (2018).  
456 "Mixture effects in samples of multiple contaminants - An inter-laboratory study with manifold  
457 bioassays." *Environment International* 114: 95-106.

458 Alvarez D.A., Shappell N.W., Billey L.O., Bermudez D.S., Wilson V.S., Kolpin D.W., Perkins S.D., Evans  
459 N., Foreman W.T., Gray J.L., Shipitalo M.J. and Meyer M.T. (2013). "Bioassay of estrogenicity and  
460 chemical analyses of estrogens in streams across the United States associated with livestock  
461 operations." *Water Research* 47(10): 3347-3363.

462 Balaguer P., François F., Comunale F., Fenet H., Boussioux A.-M., Pons M., Nicolas J.-C. and Casellas C.  
463 (1999). "Reporter cell lines to study the estrogenic effects of xenoestrogens." *Sci. Tot. Environ.*  
464 233: 47-56.

465 Brion F., De Gussem V., Buchinger S., Hollert H., Carere M., Porcher J.-M., Piccini B., Feray C., Dulio V.,  
466 Konemann S., Simon E., Werner I., Kase R. and Ait-Aissa S. (2019). "Monitoring estrogenic  
467 activities of waste and surface waters using a novel in vivo zebrafish embryonic (EASZY) assay:  
468 Comparison with in vitro cell-based assays and determination of effect-based trigger values."  
469 *Environment International* 130.

470 Brion F., Le Page Y., Piccini B., Cardoso O., Tong S.-K., Chung B.-c. and Kah O. (2012). "Screening  
471 Estrogenic Activities of Chemicals or Mixtures In Vivo Using Transgenic (cyp19a1b-GFP) Zebrafish  
472 Embryos." *Plos One* 7(5).

473 Conley J.M., Evans N., Cardon M.C., Rosenblum L., Iwanowicz L.R., Hartig P.C., Schenck K.M., Bradley  
474 P.M. and Wilson V.S. (2017). "Occurrence and In Vitro Bioactivity of Estrogen, Androgen, and  
475 Glucocorticoid Compounds in a Nationwide Screen of United States Stream Waters."  
476 *Environmental Science & Technology* 51(9): 4781-4791.

477 Cosnefroy A., Brion F., Guillet B., Laville N., Porcher J.M., Balaguer P. and Ait-Aissa S. (2009). "A stable  
478 fish reporter cell line to study estrogen receptor transactivation by environmental  
479 (xeno)estrogens." *Toxicology in Vitro* 23(8): 1450-1454.

480 Cosnefroy A., Brion F., Maillot-Marechal E., Porcher J.M., Pakdel F., Balaguer P. and Ait-Aissa S.  
481 (2012). "Selective Activation of Zebrafish Estrogen Receptor Subtypes by Chemicals by Using  
482 Stable Reporter Gene Assay Developed in a Zebrafish Liver Cell Line." *Toxicological Sciences*  
483 125(2): 439-449.

484 Creusot N., Ait-Aissa S., Tapie N., Pardon P., Brion F., Sanchez W., Thybaud E., Porcher J.M. and  
485 Budzinski H. (2014). "Identification of Synthetic Steroids in River Water Downstream from  
486 Pharmaceutical Manufacture Discharges Based on a Bioanalytical Approach and Passive  
487 Sampling." *Environmental Science & Technology* 48(7): 3649-3657.

488 Creusot N., Brion F., Piccini B., Budzinski H., Porcher J.M. and Ait-Aissa S. (2015). "BFCOD activity in  
489 fish cell lines and zebrafish embryos and its modulation by chemical ligands of human aryl  
490 hydrocarbon and nuclear receptors." *Environmental Science and Pollution Research* 22(21):  
491 16393-16404.

492 Escher B.I., Ait-Aissa S., Behnisch P.A., Brack W., Brion F., Brouwer A., Buchinger S., Crawford S.E., Du  
493 Pasquier D., Hamers T., Hettwer K., Hilscherova K., Hollert H., Kase R., Kienle C., Tindall A.J., Tuerk  
494 J., van der Oost R., Vermeirssen E. and Neale P.A. (2018). "Effect-based trigger values for in vitro

495 and in vivo bioassays performed on surface water extracts supporting the environmental quality  
496 standards (EQS) of the European Water Framework Directive." *Science of the Total Environment*  
497 628-629: 748-765.

498 European commission EU (2015). "Commission implementing decision (EU) 2015/495 of 20 March  
499 2015 establishing a watch list of substances for Union-wide monitoring in the field of water  
500 policy."

501 Hashmi M.A.K., Escher B.I., Krauss M., Teodorovic I. and Brack W. (2018). "Effect-directed analysis  
502 (EDA) of Danube River water sample receiving untreated municipal wastewater from Novi Sad,  
503 Serbia." *Science of the Total Environment* 624: 1072-1081.

504 Hashmi M.A.K., Krauss M., Escher B.I., Teodorovic I. and Brack W. (2020). "Effect-Directed Analysis of  
505 Progestogens and Glucocorticoids at Trace Concentrations in River Water." *Environmental*  
506 *Toxicology and Chemistry* 39(1): 189-199.

507 Hotchkiss A.K., Rider C.V., Blystone C.R., Wilson V.S., Hartig P.C., Ankley G.T., Foster P.M., Gray C.L.,  
508 Gray L.E. (2008). Fifteen years after "Wingspread" - environmental endocrine disrupters and  
509 human and wildlife health: where we are today and where we need to go. *Toxicol. Sci.* 105, 235-  
510 259.

511 Ihara M., Ihara M.O., Kumar V., Narumiya M., Hanamoto S., Nakada N., Yamashita N., Miyagawa S.,  
512 Iguchi T. and Tanaka H. (2014). "Co-occurrence of Estrogenic and Antiestrogenic Activities in  
513 Wastewater: Quantitative Evaluation of Balance by in Vitro ER  $\alpha$  Reporter Gene Assay and  
514 Chemical Analysis." *Environmental Science & Technology* 48(11): 6366-6373.

515 Ihara M., Kitamura T., Kumar V., Park C.-B., Ihara M.O., Lee S.-J., Yamashita N., Miyagawa S., Iguchi T.,  
516 Okamoto S., Suzuki Y. and Tanaka H. (2015). "Evaluation of Estrogenic Activity of Wastewater:  
517 Comparison Among In Vitro ER $\alpha$  Reporter Gene Assay, In Vivo Vitellogenin Induction, and  
518 Chemical Analysis." *Environmental Science & Technology* 49(10): 6319-6326.

519 Jarosova B., Blaha L., Giesy J.P. and Hilscherova K. (2014). "What level of estrogenic activity  
520 determined by in vitro assays in municipal waste waters can be considered as safe?" *Environment*  
521 *International* 64: 98-109.

522 Kase R., Javurkova B., Simon E., Swart K., Buchinger S., Koenemann S., Escher B.I., Carere M., Dulio V.,  
523 Ait-Aissa S., Hollert H., Valsecchi S., Polesello S., Behnisch P., di Paolo C., Olbrich D., Sychrova E.,  
524 Gundlach M., Schlichting R., Leborgne L., Clara M., Scheffknecht C., Marneffe Y., Chalon C., Tusil  
525 P., Soldan P., von Danwitz B., Schwaiger J., Moran Palao A., Bersani F., Perceval O., Kienle C.,  
526 Vermeirssen E., Hilscherova K., Reifferscheid G. and Werner I. (2018). "Screening and risk  
527 management solutions for steroidal estrogens in surface and wastewater." *Trac-Trends in*  
528 *Analytical Chemistry* 102: 343-358.

529 Koenemann S., Kase R., Simon E., Swart K., Buchinger S., Schluesener M., Hollert H., Escher B.I.,  
530 Werner I., Ait-Aissa S., Vermeirssen E., Dulio V., Valsecchi S., Polesello S., Behnisch P., Javurkova  
531 B., Perceval O., Di Paolo C., Olbrich D., Sychrova E., Schlichting R., Leborgne L., Clara M.,  
532 Scheffknecht C., Marneffe Y., Chalon C., Tusil P., Soldan P., von Danwitz B., Schwaiger J., San  
533 Martin Becares M.I., Bersani F., Hilscherova K., Reifferscheid G., Ternes T. and Carere M. (2018).  
534 "Effect-based and chemical analytical methods to monitor estrogens under the European Water  
535 Framework Directive." *Trac-Trends in Analytical Chemistry* 102: 225-235.

536 Koenig M., Escher B.I., Neale P.A., Krauss M., Hilscherova K., Novak J., Teodorovic I., Schulze T.,  
537 Seidensticker S., Hashmi M.A.K., Ahlheim J. and Brack W. (2017). "Impact of untreated  
538 wastewater on a major European river evaluated with a combination of in vitro bioassays and  
539 chemical analysis." *Environmental Pollution* 220: 1220-1230.



540 Kunz P.Y., Kienle C., Carere M., Homazava N. and Kase R. (2015). "In vitro bioassays to screen for  
541 endocrine active pharmaceuticals in surface and waste waters." *Journal of Pharmaceutical and*  
542 *Biomedical Analysis* 106: 107-115.

543 Kunz P.Y., Simon E., Creusot N., Jayasinghe B.S., Kienle C., Maletz S., Schifferli A., Schonlau C., Ait-  
544 Aissa S., Denslow N.D., Hollert H., Werner I. and Vermeirssen E.L.M. (2017). "Effect-based tools for  
545 monitoring estrogenic mixtures: Evaluation of five in vitro bioassays." *Water Research* 110: 378-  
546 388.

547 Le Fol V., Ait-Aissa S., Cabaton N., Dolo L., Grimaldi M., Balaguer P., Perdu E., Debrauwer L., Brion F.  
548 and Zalko D. (2015). "Cell-Specific Biotransformation of Benzophenone-2 and Bisphenol-S in  
549 Zebrafish and Human in Vitro Models Used for Toxicity and Estrogenicity Screening."  
550 *Environmental Science & Technology* 49(6): 3860-3868.

551 Le Fol V., Ait-Aissa S., Sonavane M., Porcher J.M., Balaguer P., Cravedi J.P., Zalko D. and Brion F.  
552 (2017). "In vitro and in vivo estrogenic activity of BPA, BPF and BPS in zebrafish-specific assays."  
553 *Ecotoxicology and Environmental Safety* 142: 150-156.

554 Leusch F.D.L., De Jager C., Levi Y., Lim R., Puijker L., Sacher F., Tremblay L.A., Wilson V.S. and  
555 Chapman H.F. (2010). "Comparison of Five in Vitro Bioassays to Measure Estrogenic Activity in  
556 Environmental Waters." *Environmental Science & Technology* 44(10): 3853-3860.

557 Liska I., Wagner F., Sengl M., Deutsch K. and Slonodnik J. (2015). Joint Danube Survey 3. A  
558 comprehensive analysis of Danube Water Quality. International Commission for the Protection of  
559 the Danube River (ICPDR).

560 Loos R., Marinov D., Sanseverino I., Napierska D. and Lettieri T. (2018). Review of the 1st Watch List  
561 under the Water Framework Directive and recommendations for the 2nd Watch List.

562 Matthews J., Celius T., Halgren R. and Zacharewski T. (2000). "Differential estrogen receptor binding  
563 of estrogenic substances: a species comparison." *J. Steroid Biochem. Mol. Biol.* 74: 223-234.

564 Matthiessen P., Wheeler J.R., Weltje L. (2018). A review of the evidence for endocrine disrupting  
565 effects of current-use chemicals on wildlife populations. *Critical reviews in Toxicology*, 48, 195-  
566 216

567 Menuet A., Pellegrini E., Anglade I., Blaise O., Laudet V., Kah O. and Pakdel F. (2002). "Molecular  
568 characterization of three estrogen receptor forms in zebrafish: Binding characteristics,  
569 transactivation properties, and tissue distributions." *Biology of Reproduction* 66(6): 1881-1892.

570 Molina-Molina J.-M., Amaya E., Grimaldi M., Saenz J.-M., Real M., Fernandez M.F., Balaguer P. and  
571 Olea N. (2013). "In vitro study on the agonistic and antagonistic activities of bisphenol-S and other  
572 bisphenol-A congeners and derivatives via nuclear receptors." *Toxicology and Applied*  
573 *Pharmacology* 272(1): 127-136.

574 Molina-Molina J.M., Escande A., Pillon A., Gomez E., Pakdel F., Cavailles V., Olea N., Ait-Aissa S. and  
575 Balaguer P. (2008). "Profiling of benzophenone derivatives using fish and human estrogen  
576 receptor-specific in vitro bioassays." *Toxicology and Applied Pharmacology* 232(3): 384-395.

577 Mossman T. (1983). "Rapid colorimetric assay for cellular growth and survival: application to  
578 proliferation and cytotoxicity assays." *J. Immunol.* 65: 55-63.

579 Neale P.A., Ait-Aissa S., Brack W., Creusot N., Denison M.S., Deutschmann B., Hilscherova K., Hollert  
580 H., Krauss M., Novak J., Schulze T., Seiler T.B., Serra H., Shao Y. and Escher B.I. (2015). "Linking in  
581 Vitro Effects and Detected Organic Micropollutants in Surface Water Using Mixture-Toxicity  
582 Modeling." *Environmental Science & Technology* 49(24): 14614-14624.

583 Neale P.A., Altenburger R., Ait-Aissa S., Brion F., Busch W., Umbuzeiro G.D., Denison M.S., Du  
584 Pasquier D., Hilscherova K., Hollert H., Morales D.A., Novak J., Schlichting R., Seiler T.B., Serra H.,  
585 Shao Y., Tindall A.J., Tollefsen K.E., Williams T.D. and Escher B.I. (2017). "Development of a  
586 bioanalytical test battery for water quality monitoring: Fingerprinting identified micropollutants  
587 and their Contribution to effects in surface water." *Water Research* 123: 734-750.

588 Ohtake F., Takeyama K., Matsumoto T., Kitagawa H., Yamamoto Y., Nohara K., Tohyama C., Krust A.,  
589 Mimura J., Chambon P., Yanagisawa J., Fujii-Kuriyama Y. and Kato S. (2003). "Modulation of  
590 oestrogen receptor signalling by association with the activated dioxin receptor." *Nature*  
591 423(6939): 545-550.

592 Pinto C., Grimaldi M., Boulahtouf A., Pakdel F., Brion F., Ait-Aissa S., Cavailles V., Bourguet W.,  
593 Gustafsson J.A., Bondesson M. and Balaguer P. (2014). "Selectivity of natural, synthetic and  
594 environmental estrogens for zebrafish estrogen receptors." *Toxicology and Applied Pharmacology*  
595 280(1): 60-69.

596 Schulze T., Ahel M., Ahlheim J., Ait-Aissa S., Brion F., Di Paolo C., Froment J., Hidas A.O., Hollender J.,  
597 Hollert H., Hu M., Klolss A., Koprivica S., Krauss M., Muz M., Oswald P., Petre M., Schollee J.E.,  
598 Seiler T.B., Shao Y., Slobodnik J., Sonavane M., Suter M.J.F., Tollefsen K.E., Tousova Z., Walz K.H.  
599 and Brack W. (2017). "Assessment of a novel device for onsite integrative large-volume solid  
600 phase extraction of water samples to enable a comprehensive chemical and effect-based  
601 analysis." *Science of the Total Environment* 581: 350-358.

602 Serra H., Scholze M., Altenburger R., Busch W., Budzinski H., Brion F. and Ait-Aissa S. (2019).  
603 "Combined effects of environmental xeno-estrogens within multi-component mixtures:  
604 Comparison of in vitro human- and zebrafish-based estrogenicity bioassays." *Chemosphere* 227:  
605 334-344.

606 Sonavane M., Creusot N., Maillot-Marechal E., Pery A., Brion F. and Ait-Aissa S. (2016). "Zebrafish-  
607 based reporter gene assays reveal different estrogenic activities in river waters compared to a  
608 conventional human-derived assay." *Science of the Total Environment* 550: 934-939.

609 Sonavane M., Schollee J.E., Hidas A.O., Creusot N., Brion F., Suter M.J.F., Hollender J. and Ait-Aissa S.  
610 (2018). "An integrative approach combining passive sampling, bioassays, and effect-directed  
611 analysis to assess the impact of wastewater effluent." *Environmental Toxicology and Chemistry*  
612 37(8): 2079-2088.

613 Sumpter J.P. and Jobling S. (2013). "The occurrence, causes, and consequences of estrogens in the  
614 aquatic environment." *Environmental Toxicology and Chemistry* 32(2): 249-251.

615 Tohyama S., Miyagawa S., Lange A., Ogino Y., Mizutani T., Ihara M., Tanaka H., Tatarazako N.,  
616 Kobayashi T., Tyler C.R. and Iguchi T. (2016). "Evolution of estrogen receptors in ray-finned fish  
617 and their comparative responses to estrogenic substances." *Journal of Steroid Biochemistry and*  
618 *Molecular Biology* 158: 189-197.

619 Tong S.-K., Mouriec K., Kuo M.-W., Pellegrini E., Gueguen M.-M., Brion F., Kah O. and Chung B.-c.  
620 (2009). "A cyp19a1b-GFP (Aromatase B) Transgenic Zebrafish Line That Expresses GFP in Radial  
621 Glial Cells." *Genesis* 47(2): 67-73.

622 Tyler C.R., Jobling S., Sumpter J.P. (1998). Endocrine disruption in wildlife: A critical review of the  
623 evidence. *Critical reviews in Toxicology*, 28, 319-361.

624 van der Oost R., Sileno G., Suarez-Munoz M., Nguyen M.T., Besselink H. and Brouwer A. (2017).  
625 "SIMONI (Smart Integrated Monitoring) as a novel bioanalytical strategy for water quality

626 assessment: Part I-Model design and effect-based trigger values." *Environmental Toxicology and*  
627 *Chemistry* 36(9): 2385-2399.

628 Weiss J.M., Hamers T., Thomas K.V., van der Linden S., Leonards P.E.G. and Lamoree M.H. (2009).  
629 "Masking effect of anti-androgens on androgenic activity in European river sediment unveiled by  
630 effect-directed analysis." *Analytical and Bioanalytical Chemistry* 394(5): 1385-1397.

631 Wernersson A.-S., Carere M., Maggi C., Tusil P., Soldan P., James A., Sanchez W., Dulio V., Broeg K.,  
632 Reifferscheid G., Buchinger S., Maas H., Van Der Grinten E., O'Toole S., Ausili A., Manfra L.,  
633 Marziali L., Polesello S., Lacchetti I., Mancini L., Lilja K., Linderoth M., Lundeberg T., Fjallborg B.,  
634 Porsbring T., Larsson D.G.J., Bengtsson-Palme J., Forlin L., Kienle C., Kunz P., Vermeirssen E.,  
635 Werner I., Robinson C.D., Lyons B., Katsiadaki I., Whalley C., den Haan K., Messiaen M., Clayton H.,  
636 Lettieri T., Carvalho R.N., Gawlik B.M., Hollert H., Di Paolo C., Brack W., Kammann U. and Kase R.  
637 (2015). The European technical report on aquatic effect-based monitoring tools under the water  
638 framework directive. *Environmental Sciences Europe*. 27: 1-11.

639

640 **Figure captions**

641

642 **Figure 1:** Estrogenic and anti-estrogenic activities of Novi Sad samples on (A) MELN and  
643 (B) ZELH $\beta$ 2 cells. The filled symbols represent the estrogenic activity, and the empty  
644 symbols represent the anti-estrogenic activity. The green line represents concentration  
645 addition (CA) prediction based on the main active xenoestrogens identified in the sample  
646 (contribution >5% of the response at REF=1).

647

648 **Figure 2:** Contribution of identified xeno-estrogens to the estrogenic activity measured in  
649 Novi Sad samples for (A) MELN and (B) ZELH $\beta$ 2 cells. 100% corresponds to Bio-EEQ  
650 measured experimentally.

651

652 **Figure 3:** In vivo estrogenic activity of Novi Sad sites in transgenic zebrafish embryos using  
653 the EASZY assay. (A) *In vivo* induction of GFP in 4-days old zebrafish embryos (results of  
654 one experiment with n=15 embryos per condition, representative of two independent  
655 experiments). The concentrations are expressed in Relative Enrichment Factor (REF). The  
656 positive control is 0.05 nM EE2. The mortality was below 20% in all groups. (\*) denotes a  
657 statistically significant induction in GFP intensity compared with the DMSO control group  
658 (Mann-Whitney test,  $\alpha$ : 5%). (B) Contribution of identified xeno-estrogens to *in vivo*  
659 estrogenic activity of Novi Sad samples. 100% corresponds to Bio-EEQ measured  
660 experimentally. n.a.: no activity.

661 **Table 1: Estrogenic activity of the Joint Danube Survey (JDS) and Novi Sad (NS)**  
662 **samples on MELN and ZELH $\beta$ 2 cells.** The Chem-EEQ for MELN cell line originate from  
663 Neale *et al.* (2017). The Bio-EEQ and Chem-EEQ are expressed in ng/L EEQ. The % effect  
664 indicates the contribution of identified chemicals (Chem-EEQ) to the biological response  
665 (Bio-EEQ). CI95: 95% confidence intervals, n.a.: not active, LOQ: limit of quantification.

	MELN			ZELH $\beta$ 2			
	LOQ	Bio-EEQ [CI95]	Chem-EEQ	% effect	Bio-EEQ [CI95]	Chem-EEQ	% effect
JDS	<b>LOQ</b>	<b>0.002</b>	-	-	<b>0.002</b>	-	-
	8	0.015 [0.015 ; 0.020]	0.019	124%	0.031 [0.031 ; 0.031]	0.024	77%
	22	0.048 [0.047 ; 0.061]	0.021	43%	n.a.	0.026	-
	27	0.10 [0.096 ; 0.13]	<LOQ	1%	n.a.	< LOQ	-
	29	0.11 [0.11 ; 0.17]	0.050	44%	0.012 [0.010 ; 0.014]	0.062	503%
	30	0.22 [0.19 ; 0.32]	0.040	19%	n.a.	0.051	-
	32	0.064 [0.061 ; 0.086]	0.040	62%	n.a.	0.054	-
	33	0.033 [0.032 ; 0.042]	0.040	122%	n.a.	0.051	-
	35	n.a.	0.034	n.a.	n.a.	0.042	-
	36	0.054 [0.052 ; 0.066]	0.021	39%	n.a.	0.026	-
	37	n.a.	0.024	n.a.	n.a.	0.030	-
	39	0.15 [0.14 ; 0.21]	0.051	34%	0.021 [0.017 ; 0.026]	0.063	306%
	41	0.74 [0.65 ; 0.84]	0.20	27%	2.3 [2.05 ; 2.9]	0.244	11%
	44	0.070 [0.069 ; 0.089]	0.026	37%	0.039 [0.039 ; 0.063]	0.032	82%
	53	n.a.	0.053	n.a.	n.a.	0.065	-
	55	n.a.	<LOQ	n.a.	n.a.	<LOQ	-
	57	n.a.	0.018	n.a.	n.a.	0.022	-
	59	0.053	0.050	94%	n.a.	0.062	-
	60	0.078 [0.066 ; 0.092]	<LOQ	0%	n.a.	<LOQ	-
	63	n.a.	0.058	n.a.	0.053 [0.039 ; 0.064]	0.072	136%
	64	0.0026 [0.26 ; 0.33]	0.019	710%	n.a.	0.023	-
	65	0.12 [0.11 ; 0.13]	0.066	55%	0.071 [0.062 ; 0.082]	0.082	116%
	67	n.a.	0.045	n.a.	0.026 [0.023 ; 0.034]	0.056	214%
Novi Sad	<b>LOQ</b>	<b>0.005</b>	-	-	<b>0.008</b>	-	-
	NS1	0.079 [0.073 ; 0.092]	<LOQ	2%	0.050 [0.044 ; 0.053]	<LOQ	2%
	NS2	1.52 [1.44 ; 1.83]	1.76	115%	0.19 [0.17 ; 0.22]	1.39	743%
	NS3	0.12 [0.11 ; 0.15]	0.021	18%	0.051 [0.039 ; 0.060]	0.025	48%

666

667







668 **Table 2: Relative potency (REP) of the chemicals quantified for each bioassay.** The REP  
 669 values were calculated as the ratio of EC20 of E2 to that of test compound. They were derived  
 670 from initial data published by <sup>(a)</sup>Neale et al. (2017), <sup>(b)</sup>Molina-Molina et al. (2008) and  
 671 Molina-Molina et al. (2013), <sup>(c)</sup>LeFol et al. (2017), <sup>(d)</sup>Brion et al. (2012), <sup>(e)</sup>Brion et al. (2019).

<b>Chemical</b>	<b>MELN</b>	<b>ZELH<math>\beta</math>2</b>	<b>EASZY Assay</b>
Estrone	0.110 <sup>a</sup>	0.136 <sup>a</sup>	0.78 <sup>e</sup>
Estriol	0.077 <sup>a</sup>	0.039 <sup>a</sup>	0.06 <sup>d</sup>
Bisphenol F	1.43 $\times 10^{-5}$ <sup>b</sup>	5.14 $\times 10^{-6}$ <sup>c</sup>	4.00 $\times 10^{-4}$ <sup>c</sup>
Bisphenol S	1.18 $\times 10^{-6}$ <sup>a</sup>	6.69 $\times 10^{-6}$ <sup>c</sup>	7.80 $\times 10^{-6}$ <sup>c</sup>
Bisphenol A	3.50 $\times 10^{-5}$ <sup>a</sup>	3.57 $\times 10^{-6}$ <sup>c</sup>	1.44 $\times 10^{-4}$ <sup>c</sup>
Daidzein	6.50 $\times 10^{-5}$ <sup>a</sup>	9.22 $\times 10^{-6}$ <sup>a</sup>	Inactive <sup>a</sup>
Diazinon	2.80 $\times 10^{-7}$ <sup>a</sup>	Inactive <sup>a</sup>	Inactive <sup>a</sup>
Genistein	2.00 $\times 10^{-4}$ <sup>a</sup>	3.47 $\times 10^{-4}$ <sup>a</sup>	1.91 $\times 10^{-4}$ <sup>a</sup>
Terbutylazine	1.60 $\times 10^{-7}$ <sup>a</sup>	Inactive <sup>a</sup>	Inactive <sup>a</sup>

672


**Table 3: Comparison of effect-based trigger values proposed for estrogenic activity.** The characteristics of each EBT are presented with a focus on the point of departure, inclusion of bioassay-specific and mixture considerations. n.a.: not applicable, n.i.: not included, REP: relative potency, CI: confidence interval, EQS: environmental quality standard. <sup>(a)</sup>: according to option G in Escher et al. (2018).

<b>Point of Departure</b>		<b>Special considerations</b>			<b>EB</b>
<b>Hazard</b>	<b>Chemicals</b>	<b>Occurrence</b>	<b>REP</b>	<b>Mixture effects</b>	<b>All</b>
Chronic toxicity data on fish and amphibian (based on EQS)	E2	n.i.	n.i.	n.i.	0.4
Acute and chronic toxicity data on fish, crustacean, algae, insects	7 chemicals with REP > 0.001 in ER-CALUX	n.i.	Yes, only ER-CALUX	n.i.	0.5 (95)
Chronic toxicity data on fish (PNEC)	E1, E2, E3, EE2	Proportion in waste water and surface water	Yes, 8 in vitro bioassays compared	n.i.	0.1 (me)
Chronic toxicity data on fish and amphibian (based on EQS)	E1, E2, EE2	Proportion in waste water and surface water	Yes, 9 in vitro bioassays compared	Yes (additive effects)	0.1 (me)
in vitro-in vivo comparison of 33 water samples using cyp19a1b-GFP transgenic embryos	All active chemicals in the samples	Based on empirical data	Yes, 5 in vitro bioassays compared	Yes (additive and non-additive effects)	0.1 (me)






# A MELN



## Chem-TEQ (%):

- Bisphenol F
- Bisphenol S
- Bisphenol A
- Daidzein
- Diazinon
- Estradiol
- Estriol
- Estrone
- Genistein
- Terbutylazine

# B ZELH $\beta$ 2



## Chem-TEQ (%):

- Bisphenol S
- Bisphenol F
- Bisphenol A
- Daidzein
- Estradiol
- Estriol
- Estrone
- Genistein

