



Exogenous acquisition of *Pseudomonas aeruginosa* in intensive care units: a prospective multi-centre study (DYNAPYO study)

Maidier Coppry, Camille Leroyer, Marion Saly, Anne Gaëlle Venier, Celine Slekovec, Xavier Bertrand, Sylvie Parer, Serge Alfandari, Emmanuelle Cambau, Bruno Mégarbane, et al.

► To cite this version:

Maidier Coppry, Camille Leroyer, Marion Saly, Anne Gaëlle Venier, Celine Slekovec, et al.. Exogenous acquisition of *Pseudomonas aeruginosa* in intensive care units: a prospective multi-centre study (DYNAPYO study). *Journal of Hospital Infection*, 2020, 104 (1), pp.40-45. <10.1016/j.jhin.2019.08.008>. <hal-03161089v2>

HAL Id: hal-03161089

<https://hal.science/hal-03161089v2>

Submitted on 21 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

Exogenous acquisition of *Pseudomonas aeruginosa* in intensive care units: a prospective multicentre study, DYNAPYO study.

Authors: Maïder Coppry^{1,2}, Camille Leroyer³, Marion Saly³, Anne-Gaëlle Venier^{2,4}, Céline Slekovec⁵, Xavier Bertrand⁵, Sylvie Parer⁶, Serge Alfandari⁷, Emmanuelle Cambau⁸, Bruno Megarbane⁹, Christine Lawrence¹⁰, Bernard Clair¹¹, Alain Lepape¹², Pierre Cassier¹³, Dominique Trivier¹⁴, Alexandre Boyer¹⁵, Hélène Boulestreau³, Julien Asselineau¹⁸, Véronique Dubois^{16,17}, Rodolphe Thiébaud^{2,18}, Anne-Marie Rogues^{1,2}.

Laboratory or institution of origin

1. Univ. Bordeaux, CHU Bordeaux, Hygiène hospitalière, F-33000 Bordeaux, France
2. Univ. Bordeaux, Inserm, Bordeaux Population Health Research Center, team Pharmacoepidemiology, UMR 1219, F-33000, France
3. CHU Bordeaux, Hygiène hospitalière, F-33000 Bordeaux, France
4. CHU Bordeaux, CPIAS Nouvelle-Aquitaine, F-33000 Bordeaux, France
5. CHU Besançon, Hygiène hospitalière, F-25000 Besançon, France
6. CHU Montpellier, Hygiène hospitalière, F-34000 Montpellier, France
7. CH Tourcoing, Réanimation et maladies infectieuses, F-59200 Tourcoing, France
8. CHU Lariboisière, AP-HP, Hygiène hospitalière, F-75000 Paris, France
9. CHU Lariboisière, AP-HP, Réanimation médicale, F-75000 Paris, France
10. CHU Poincaré, AP-HP, Hygiène hospitalière, F-92380 Garches, France
11. CHU Poincaré, AP-HP, Réanimation médicale, F-92380 Garches, France
12. CHU Lyon Sud, Réanimation médicale, F- 69000 Lyon, France

- 26 13. CHU Lyon Hôpital E. Herriot, Laboratoire d'hygiène, F- 69437 Lyon, France
- 27 14. CH Lens, Hygiène hospitalière, F- 62300 Lens, France
- 28 15. Univ.Bordeaux, CHU Bordeaux, Réanimation médicale, F-33000 Bordeaux,
- 29 France
- 30 16. Univ. Bordeaux, CHU Bordeaux, Laboratoire de bactériologie, F-33000
- 31 Bordeaux, France
- 32 17. Univ. Bordeaux, UMR 5234 CNRS, Université de Bordeaux, F-33000
- 33 Bordeaux, France
- 34 18. CHU Bordeaux, Service d'information médicale, Unité de Soutien
- 35 Méthodologique à la Recherche Clinique et Epidémiologique, F-33000
- 36 Bordeaux, France

37

38 **Corresponding author:** Maider Coppry, Hygiène hospitalière, CHU Pellegrin, place

39 Raba-Léon, 33076 Bordeaux cedex. Tel.: +33 556 795 553, Fax: +33 556 794 997.

40 E-mail maider.coppry@chu-bordeaux.fr

41

42 Alternate corresponding author: **Anne-Marie Rogues, Service d'Hygiène**

43 **hospitalière, CHU Pellegrin, place Raba-Léon, 33076 Bordeaux cedex.** Tel.:

44 +33 556 795 553, Fax: +33 556 794 997. E-mail anne-marie.rogues@chu-

45 bordeaux.fr

Summary

Pseudomonas aeruginosa remains one of the most common nosocomial pathogens in intensive care units (ICUs). Although exogenous acquisition has been widely documented in outbreaks, its importance is unclear in non-epidemic situations. We aimed to elucidate the role of exogenous origin of *P. aeruginosa* in ICU patients.

We performed a chronological analysis of the acquisition of *P. aeruginosa* using samples collected in 2009 in DYNAPYO cohort study during which patients and tap water were weekly screened. Molecular relatedness of *P. aeruginosa* isolates was investigated by pulsed-field gel electrophoresis. Exogenous acquisition was defined as identification of a *P. aeruginosa* pulsotype previously isolated from another patient or tap water in the ICU.

DYNAPYO cohort included 1,808 patients (10,402 samples) and 233 water taps (4,946 samples). Typing of 1,515 isolates from 373 patients and 375 isolates from 81 tap water samples identified 296 pulsotypes. Analysis showed an exogenous acquisition in 170 (45.6%) of 373 patients. The pulsotype identified was previously isolated from another patient and from a tap water sample for 86 and 29 patients respectively. The results differed according to the ICU.

The exogenous acquisition of *P. aeruginosa* could be prevented in a half proportion of patients. The overall findings of this survey supports the need for studies on routes of transmission and risk assessment approach to better define how to control exogenous acquisition in ICUs.

Introduction

Pseudomonas aeruginosa remains one of the most common hospital-acquired pathogens, and is endemic in many intensive care units (ICUs).[1,2] Infections caused by *P. aeruginosa*, especially ventilator-associated pneumonia or bloodstream infections, are often severe and associated with considerable mortality in ICU patients; morbidity and mortality rates are higher still in cases of multidrug resistance. [3–6] In ICUs, these infections are usually considered to be endogenous, arising from pre-existing colonization of patients. In addition, the roles of exogenous reservoirs and of patient-to-patient transmission have been convincingly documented during outbreaks. However, the importance of exogenously acquired *P. aeruginosa* in non-epidemic situations remains uncertain.[7,8] Several epidemiological studies have indicated that colonization pressure seems to be a more relevant risk factor than exposure to antibiotics for the acquisition of *P. aeruginosa*. [9–12]

A previous monocentric study showed a clear genetic temporal and spatial relationship between *P. aeruginosa* strains isolated from tap water samples and ICU inpatients.[13] Moreover, the first part of the multicentre DYNAPYO (Dynamics of *Pseudomonas aeruginosa* acquisition in ICU) study showed that exposure to contaminated taps water was a risk factor for *P. aeruginosa* colonisation in ICU patients.[14,15] It is essential to better understand the true contribution of the exogenous acquisition of *P. aeruginosa* in ICU to inform infection prevention and control strategies. We performed a chronological analysis within DYNAPYO participating ICUs to assess the respective contributions of *P. aeruginosa* exogenous acquisition by patient-to-patient transmission and from contaminated taps.

METHODS

Study design and study population

DYNAPYO was a prospective five-month observational survey performed in 2009 in ten ICUs (four medical, two surgical and four mixed medical and surgical ICUs) from eight French health care facilities: University hospital of Besançon and Lyon which included two ICUs; University hospital of Bordeaux, Garches, Montpellier and Paris and general hospitals of Lens and Tourcoing which included one ICU. These ICUs had 9 to 20 beds with an average length of stay of 8 to 16 days (Table I). The ICUs did not implement changes during the course of the study, such as infection control or antimicrobial stewardship initiatives. The ICUs had to follow French recommendations in terms of water use for care and measures to prevent patient-to-patient transmission. Outlet taps were equipped with antibacterial filters in one ICU (ICU 1).

The DYNAPYO cohort included all adult patients admitted for more than 24 h during the study inclusion period. Demographic and epidemiological data (admission dates and discharge, room number) were collected prospectively. The type of taps (electronic or conventional) was recorded, and patients and tap water were monitored for contamination with *P. aeruginosa* weekly over the study period. Specific trained healthcare workers were identified in each centre for water sampling and for data collection on a secured online case report form. This study was approved by the local ethics committee and the data underlying this study restricted by the French data protection commission (Commission Nationale Informatique et Liberté - France).

Surveillance culture and microbiological analysis

During the data collection period, patients were screened on admission (within the first 48 h of ICU stay), and then once a week and on discharge or death.

Screening samples were oropharyngeal, rectal swabs and tracheobronchial aspirate (or sputum). Others clinical specimens were performed as clinically indicated (as appropriate).

Cold water samples were taken weekly from the 233 taps in the ten ICUs (patients' rooms and other sites) for testing for *P. aeruginosa* (without colony count). Taps were opened and the first 250 mL of flush of water were immediately collected in a sterile flask with sodium thiosulfate. The aerator was swabbed and the swab broken into the water sample. Tap water samples were processed by membrane filtration. A volume of 100 mL was filtered through a 0.45 µm pore size membrane filter (Millipore Microfil, Molsheim, France). Swabs and filters were cultured on cefrimide-agar plates (Bio-Rad, Marnes-la-Coquette, France) at 37°C and examined for growth of colonies after 24 and 48 hours. Any colony that grew on cefrimide-agar plate was identified using the API20 NE identification system (bioMérieux, Marcy l'Etoile, France). All *P. aeruginosa* isolates were sent to the coordinating centre (Bordeaux) on semi-solid agar.

Genotyping

Molecular relatedness of *P. aeruginosa* isolates was investigated by pulsed-field gel electrophoresis (PFGE). Clonality of strains was investigated by PFGE with DraI digestion as previously described.[16] The banding patterns were analysed by scanning photographic negatives. GelCompar software was used for analysis of PFGE patterns (Applied Maths, Kortrijk, Belgium). Pulsotypes were defined according to international recommendations.[17] All patients' first isolates from an anatomic site were analysed. Tap water isolates selected for comparison were:

142 isolates identified in water on the previous week and the week after of a newly
143 identified patient.

144 **Definitions**

145 Exogenous acquisition was defined as colonisation or infection by a strain of
146 *P. aeruginosa* with a pulsotype previously isolated from another patient (i.e. patient-
147 to-patient transmission) or from tap water sample in the ICU. Patient-to-patient
148 transmission was considered possible when a similar pulsotype was isolated in more
149 than two patients hospitalised during overlapping period without similar pulsotype
150 isolated from tap water. An exogenous origin from tap water was considered possible
151 when a similar pulsotype was isolated in a patient and in at least one ICU tap water
152 prior to *P. aeruginosa* identification in the patient.

RESULTS

Study population

Of the 1,808 patients included in DYNAPYO cohort, 206 were excluded because screening at admission was not carried out. A total of 10,402 screening samples were performed and 427 patients were positive for *P. aeruginosa*; of these 41 were found on entering the study. The average incidence of *P. aeruginosa* in the ten ICUs was 12.7 per 1 000 days hospitalisation (Table II).

Water samples

A total of 4,946 water samples were obtained. Among the 233 taps screened, 81(35%) were positive for *P. aeruginosa* at least once during the study, including 51 at the beginning of the study. The median duration of contamination was 5 weeks (range 1 to 13 weeks). The median duration of contamination differed between electronic and conventional taps (12.6 vs. 8 weeks respectively; $p = 0.003$).

Genotyping and chronological epidemiological analysis

Typing of 1,880 non-replicate isolates (1,515 from 373 patients and 375 from 81 water samples) identified 296 pulsotypes. A total of 270 different pulsotypes were found in patients: 201 (74%) were sporadic, 52 were shared by patients and 17 were shared by water and patient. Variations according to the ICU are shown in Table II.

The chronological epidemiological analysis showed an exogenous acquisition in 170 (45.6%) patients out of the 373 for which at least one isolate was available for typing with variation according to the ICU (from 16.3% in ICU 7 and 85.7 % in ICU 5; Table II). There was a patient-to-patient transmission for 86 of the 170 patients (50.6%) and an exogenous origin from tap water for 29 others patients (17.1%). Moreover, for 55 patients from the two ICUs with higher rate of positive tap water (ICU 5, ICU10) it

was not possible to conclude because pulsotypes were shared by many patients and tap water samples.

DISCUSSION

To our knowledge DYNAPYO is the largest cohort study intending to assess the relative contribution of exogenous acquisition of *P. aeruginosa* in ICUs.[14] We showed an exogenous origin of *P. aeruginosa* in nearly one in two patients. Patient-to-patient transmission was more frequent than acquisition from the tap water. At least half of colonisation or infection by *P. aeruginosa* could be preventable. Furthermore, our study showed discrepancies in the rates of exogenous origin of *P. aeruginosa* according to the ICU that could explain the differences in results in previous monocentric studies.

Patient-to-patient transmission occurs by carriage on the hands of healthcare workers or through contaminated medical equipment.[18–20] There is a considerable body of published literature for patient-to-patient transmission of multidrug resistant *P. aeruginosa* from outbreaks reports.[12,21,22] The strict maintenance of infection control measures is essential to limit the spread of this bacteria. Infection control strategies to decrease the incidence of infection due to *P. aeruginosa* in ICUs mostly includes bundle approaches involving general measures, disinfection and replacing reservoirs.[24,25]

Previous studies observed that clinical strains of *P. aeruginosa* were genetically related to the strains found in the patient's environment, such as in tap water, P-traps, sinks, handwashing stations, faucet aerators or washbasins drain; but a causal link was controversial.[25] In a systematic review, seven monocentric studies were assessed as providing plausible evidence of a link between tap water as a

202 reservoir for *P. aeruginosa* and colonization/infection in patient in an endemic setting.
203 In these studies rates of exogenous acquisition of *P. aeruginosa* varied from 29% to
204 81% of patients as in our different ICUs.[13,19,26–30]

205 In our study, there was a wide heterogeneity in tap water contamination by *P.*
206 *aeruginosa* among the ICUs. Like others authors, we showed that *P. aeruginosa* may
207 persist in tap water over prolonged periods and that electronic taps are potential
208 reservoirs of *P. aeruginosa* in ICUs.[31–33] Tap water could become positive for
209 *P. aeruginosa* through contamination of the water supply or retrograde contamination
210 (e.g. from splashing on to the faucet when water is drawn, especially if the water flow
211 directly impacts on the drain outlet, or if fluids are inappropriately discarded in
212 handwash basins.[34,35]

213 Discrepancies observed among ICUs may be explained by various factors:
214 compliance to infection control measures, contamination load of the environment,
215 biological features of the pathogen (intrinsic fitness factors).[36] Based on our finding
216 we suggest monitoring tap water in ICUs with high rates of colonisation or infection
217 with *P. aeruginosa*; although there are no recommendations for a systematic
218 screening in search of *P. aeruginosa* in ICUs. Furthermore these ICUs should
219 consider eliminating work processes involving sinks in proximity patients and favour
220 compliance to alcoholic hand disinfection in order to limit the spread of *P. aeruginosa*
221 among patients. Some guidelines recommend sampling outlets in ICUs on a six-
222 month basis and taking remedial action for outlets that are positive for *P.*
223 *aeruginosa*.[37]

224 The strengths of this study are the prospective multicentre design with a study
225 population in accordance with most of the previous studies analysing *P. aeruginosa*
226 colonisation/infection in ICUs; the use of methods enabling temporal relationship to

be identified between water taps and identification of colonisation/infection in patients; repeated sampling during the five months of the study and the huge number of isolates typed by PFGE. Some limitations should also be noted. First, we selected isolates for genotyping and then performed genotyping from one colony of each positive culture; however isolates were not available for some patients. It may not accurately represent the whole epidemiology. A robust methodology should be to type up to 4-10 different colonies from each culture. In a study of more than 1,600 isolates of *P. aeruginosa*, the authors found that over 60% of the tap water samples were contaminated by *P. aeruginosa* and overall 83% of the patient strains were classified as exogenous. They typed of at least four colonies that were representative of the different morphological types of *P. aeruginosa* present on each culture plate.[29] Second, the limitation of water samples may have underestimated the number of exogenous sources as we did not performed extensive microbiological samples of the environment other than water taps and patients. The range of reservoirs in healthcare environments from which *P. aeruginosa* has been isolated is wide, including respiratory therapy equipment, ice makers, endoscopes, and cleaning equipment. [38,39]

Conclusion

This multicentre study conducted in ICUs suggests that exogenous origin of *P. aeruginosa* could be prevented in a substantial proportion of patients. Given the possible consequences of *P. aeruginosa* infection in ICU, it is clear that strategies to prevent *P. aeruginosa* acquisition should become a key priority. We support the need for studies on routes of transmission and risk assessment approach to better define how to control exogenous acquisition in ICUs.

Acknowledgements

252 Authors thank the ICUs, microbiology laboratories and infection control teams of the
253 participating centres.

254 **Conflict of interest:** None to declare.

255 **Funding sources:** This work was supported by the French Ministry of Health (PHRC
256 national 2008). CHUBX 2008/15, 03/12/2008.

257

Reference

- [1] Álvarez-Lerma F, Olaechea-Astigarraga P, Palomar-Martínez M, Catalan M, Nuvials X, Gimeno R, et al. Invasive device-associated infections caused by *Pseudomonas aeruginosa* in critically ill patients: evolution over 10 years. *J Hosp Infect* 2018;100:e204–8. doi:10.1016/j.jhin.2018.04.027.
- [2] Cohen R, Babushkin F, Cohen S, Afraimov M, Shapiro M, Uda M, et al. A prospective survey of *Pseudomonas aeruginosa* colonization and infection in the intensive care unit. *Antimicrob Resist Infect Control* 2017;6:7. doi:10.1186/s13756-016-0167-7.
- [3] Vincent J-L, Rello J, Marshall J, Silva E, Anzueto A, Martin CD, et al. International study of the prevalence and outcomes of infection in intensive care units. *JAMA* 2009;302:2323–9. doi:10.1001/jama.2009.1754.
- [4] Fernández-Barat L, Ferrer M, De Rosa F, Gabarrús A, Esperatti M, Terraneo S, et al. Intensive care unit-acquired pneumonia due to *Pseudomonas aeruginosa* with and without multidrug resistance. *Journal of Infection* 2017;74:142–52. doi:10.1016/j.jinf.2016.11.008.
- [5] Micek ST, Wunderink RG, Kollef MH, Chen C, Rello J, Chastre J, et al. An international multicenter retrospective study of *Pseudomonas aeruginosa* nosocomial pneumonia: impact of multidrug resistance. *Crit Care* 2015;19:219. doi:10.1186/s13054-015-0926-5.
- [6] Rosenthal VD, Maki DG, Mehta Y, Leblebicioglu H, Memish ZA, Al-Mousa HH, et al. International Nosocomial Infection Control Consortium (INICC) report, data summary of 43 countries for 2007-2012. Device-associated module. *Am J Infect Control* 2014;42:942–56. doi:10.1016/j.ajic.2014.05.029.

- [7] Cuttelod M, Senn L, Terletskiy V, Nahimana I, Petignat C, Eggimann P, et al. Molecular epidemiology of *Pseudomonas aeruginosa* in intensive care units over a 10-year period (1998-2007). *Clin Microbiol Infect* 2011;17:57–62. doi:10.1111/j.1469-0691.2010.03164.x.
- [8] Loveday HP, Wilson JA, Kerr K, Pitchers R, Walker JT, Browne J. Association between healthcare water systems and *Pseudomonas aeruginosa* infections: a rapid systematic review. *J Hosp Infect* 2014;86:7–15. doi:10.1016/j.jhin.2013.09.010.
- [9] Bonten MJM, Weinstein RA. Transmission pathways of *Pseudomonas aeruginosa* in intensive care units: don't go near the water. *Crit Care Med* 2002;30:2384–5. doi:10.1097/01.CCM.0000025906.15627.D2.
- [10] Boyer A, Doussau A, Thiébault R, Venier AG, Tran V, Boulestreau H, et al. *Pseudomonas aeruginosa* acquisition on an intensive care unit: relationship between antibiotic selective pressure and patients' environment. *Crit Care* 2011;15:R55. doi:10.1186/cc10026.
- [11] Cobos-Trigueros N, Solé M, Castro P, Torres JL, Hernández C, Rinaudo M, et al. Acquisition of *Pseudomonas aeruginosa* and its resistance phenotypes in critically ill medical patients: role of colonization pressure and antibiotic exposure. *Crit Care* 2015;19:218. doi:10.1186/s13054-015-0916-7.
- [12] DalBen MF, Basso M, Garcia CP, Costa SF, Toscano CM, Jarvis WR, et al. Colonization pressure as a risk factor for colonization by multiresistant *Acinetobacter* spp and carbapenem-resistant *Pseudomonas aeruginosa* in an intensive care unit. *Clinics (Sao Paulo)* 2013;68:1128–33. doi:10.6061/clinics/2013(08)11.
- [13] Rogues A-M, Boulestreau H, Lashéras A, Boyer A, Gruson D, Merle C, et al. Contribution of tap water to patient colonisation with *Pseudomonas aeruginosa* in a

306 medical intensive care unit. J Hosp Infect 2007;67:72–8.
 307 doi:10.1016/j.jhin.2007.06.019.

308 [14] Venier AG, Gruson D, Lavigne T, Jarno P, L’hériveau F, Coignard B, et al.
 309 Identifying new risk factors for *Pseudomonas aeruginosa* pneumonia in intensive
 310 care units: experience of the French national surveillance, REA-RAISIN. J Hosp
 311 Infect 2011;79:44–8. doi:10.1016/j.jhin.2011.05.007.

312 [15] Hoang S, Georget A, Asselineau J, Venier A-G, Leroyer C, Rogues AM, et al.
 313 Risk factors for colonization and infection by *Pseudomonas aeruginosa* in patients
 314 hospitalized in intensive care units in France. PLoS ONE 2018;13:e0193300.
 315 doi:10.1371/journal.pone.0193300.

316 [16] Slekovec C, Plantin J, Cholley P, Thouverez M, Talon D, Bertrand X, et al.
 317 Tracking down antibiotic-resistant *Pseudomonas aeruginosa* isolates in a wastewater
 318 network. PLoS ONE 2012;7:e49300. doi:10.1371/journal.pone.0049300.

319 [17] Tenover FC, Arbeit RD, Goering RV, Mickelsen PA, Murray BE, Persing DH,
 320 et al. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel
 321 electrophoresis: criteria for bacterial strain typing. J Clin Microbiol 1995;33:2233–9.

322 [18] Reuter S, Sigge A, Wiedeck H, Trautmann M. Analysis of transmission
 323 pathways of *Pseudomonas aeruginosa* between patients and tap water outlets. Crit
 324 Care Med 2002;30:2222–8. doi:10.1097/01.CCM.0000025907.12833.10.

325 [19] Moolenaar RL, Crutcher JM, San Joaquin VH, Sewell LV, Hutwagner LC,
 326 Carson LA, et al. A prolonged outbreak of *Pseudomonas aeruginosa* in a neonatal
 327 intensive care unit: did staff fingernails play a role in disease transmission? Infect
 328 Control Hosp Epidemiol 2000;21:80–5. doi:10.1086/501739.

329 [20] DiazGranados CA, Jones MY, Kongphet-Tran T, White N, Shapiro M, Wang
 330 YF, et al. Outbreak of *Pseudomonas aeruginosa* infection associated with

331 contamination of a flexible bronchoscope. *Infect Control Hosp Epidemiol*
332 2009;30:550–5. doi:10.1086/597235.

333 [21] Suarez C, Peña C, Arch O, Dominguez MA, Tubau F, Juan C, et al. A large
334 sustained endemic outbreak of multiresistant *Pseudomonas aeruginosa*: a new
335 epidemiological scenario for nosocomial acquisition. *BMC Infect Dis* 2011;11:272.
336 doi:10.1186/1471-2334-11-272.

337 [22] Cholley P, Gbaguidi-Haore H, Bertrand X, Thouverez M, Plésiat P, Hocquet D,
338 et al. Molecular epidemiology of multidrug-resistant *Pseudomonas aeruginosa* in a
339 French university hospital. *J Hosp Infect* 2010;76:316–9.
340 doi:10.1016/j.jhin.2010.06.007.

341 [23] Gbaguidi-Haore H, Varin A, Cholley P, Thouverez M, Hocquet D, Bertrand X.
342 A Bundle of Measures to Control an Outbreak of *Pseudomonas aeruginosa*
343 Associated With P-Trap Contamination. *Infect Control Hosp Epidemiol* 2018;39:164–
344 9. doi:10.1017/ice.2017.304.

345 [24] Petignat C, Francioli P, Nahimana I, Wenger A, Bille J, Schaller M-D, et al.
346 Exogenous sources of *pseudomonas aeruginosa* in intensive care unit patients:
347 implementation of infection control measures and follow-up with molecular typing.
348 *Infect Control Hosp Epidemiol* 2006;27:953–7. doi:10.1086/506409.

349 [25] Tran-Dinh A, Neulier C, Amara M, Nebot N, Troché G, Breton N, et al. Impact
350 of intensive care unit relocation and role of tap water on an outbreak of
351 *Pseudomonas aeruginosa* expressing OprD-mediated resistance to imipenem. *J*
352 *Hosp Infect* 2018;100:e105–14. doi:10.1016/j.jhin.2018.05.016.

353 [26] Berthelot P, Grattard F, Mahul P, Pain P, Jospé R, Venet C, et al. Prospective
354 study of nosocomial colonization and infection due to *Pseudomonas aeruginosa* in

355 mechanically ventilated patients. *Intensive Care Med* 2001;27:503–12.
 356 doi:10.1007/s001340100870.

357 [27] Trautmann M, Michalsky T, Wiedeck H, Radosavljevic V, Ruhnke M. Tap
 358 Water Colonization With *Pseudomonas aeruginosa* in a Surgical Intensive Care Unit
 359 (ICU) and Relation to *Pseudomonas* Infections of ICU Patients. *Infection Control &*
 360 *Hospital Epidemiology* 2001;22:49–52. doi:10.1086/501828.

361 [28] Trautmann M, Bauer C, Schumann C, Hahn P, Höher M, Haller M, et al.
 362 Common RAPD pattern of *Pseudomonas aeruginosa* from patients and tap water in
 363 a medical intensive care unit. *International Journal of Hygiene and Environmental*
 364 *Health* 2006;209:325–31. doi:10.1016/j.ijheh.2006.04.001.

365 [29] Vallés J, Mariscal D, Cortés P, Coll P, Villagrà A, Díaz E, et al. Patterns of
 366 colonization by *Pseudomonas aeruginosa* in intubated patients: a 3-year prospective
 367 study of 1,607 isolates using pulsed-field gel electrophoresis with implications for
 368 prevention of ventilator-associated pneumonia. *Intensive Care Med* 2004;30:1768–
 369 75. doi:10.1007/s00134-004-2382-6.

370 [30] Cholley P, Thouverez M, Floret N, Bertrand X, Talon D. The role of water
 371 fittings in intensive care rooms as reservoirs for the colonization of patients with
 372 *Pseudomonas aeruginosa*. *Intensive Care Med* 2008;34:1428. doi:10.1007/s00134-
 373 008-1110-z.

374 [31] Walker J, Moore G. *Pseudomonas aeruginosa* in hospital water systems:
 375 biofilms, guidelines, and practicalities. *J Hosp Infect* 2015;89:324–7.
 376 doi:10.1016/j.jhin.2014.11.019.

377 [32] Costa D, Bousseau A, Thevenot S, Dufour X, Laland C, Burucoa C, et al.
 378 Nosocomial outbreak of *Pseudomonas aeruginosa* associated with a drinking water
 379 fountain. *J Hosp Infect* 2015;91:271–4. doi:10.1016/j.jhin.2015.07.010.

- 380 [33] Garvey MI, Bradley CW, Tracey J, Oppenheim B. Continued transmission of
381 *Pseudomonas aeruginosa* from a wash hand basin tap in a critical care unit. *J Hosp*
382 *Infect* 2016;94:8–12. doi:10.1016/j.jhin.2016.05.004.
- 383 [34] Berthelot P, Chord F, Mallaval F, Grattard F, Brajon D, Pozzetto B. Magnetic
384 valves as a source of faucet contamination with *Pseudomonas aeruginosa*? *Intensive*
385 *Care Med* 2006;32:1271. doi:10.1007/s00134-006-0206-6.
- 386 [35] Kizny Gordon AE, Mathers AJ, Cheong EYL, Gottlieb T, Kotay S, Walker AS,
387 et al. The Hospital Water Environment as a Reservoir for Carbapenem-Resistant
388 Organisms Causing Hospital-Acquired Infections-A Systematic Review of the
389 Literature. *Clin Infect Dis* 2017;64:1435–44. doi:10.1093/cid/cix132.
- 390 [36] Blanc DS, Francioli P, Zanetti G. Molecular Epidemiology of *Pseudomonas*
391 *aeruginosa* in the Intensive Care Units - A Review. *Open Microbiol J* 2007;1:8–11.
392 doi:10.2174/1874285800701010008.
- 393 [37] Department of Health. Water Systems, Health Technical Memorandum 04-01:
394 Addendum. *Pseudomonas aeruginosa* – advice for augmented care units. 2013. n.d.
395 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/14010](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/140105/Health_Technical_Memorandum_04-01_Addendum.pdf)
396 [5/Health_Technical_Memorandum_04-01_Addendum.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/140105/Health_Technical_Memorandum_04-01_Addendum.pdf) (accessed March 4, 2019).
- 397 [38] Ferranti G, Marchesi I, Favale M, Borella P, Bargellini A. Aetiology, source and
398 prevention of waterborne healthcare-associated infections: a review. *J Med Microbiol*
399 2014;63:1247–59. doi:10.1099/jmm.0.075713-0.
- 400 [39] Kerr KG, Snelling AM. *Pseudomonas aeruginosa*: a formidable and ever-
401 present adversary. *J Hosp Infect* 2009;73:338–44. doi:10.1016/j.jhin.2009.04.020.

Table I. Description of the intensive care units

	Type	Rooms (n)	Length of stay in days (mean)
ICU 1*	Mixed	20	14
ICU 2	Mixed	16	12
ICU 3	Medical	12	10
ICU 4	Surgical	14	14
ICU 5	Medical	18	8
ICU 6	Medical	12	10
ICU 7	Mixed	18	11
ICU 8	Medical	15	11
ICU 9	Surgical	15	11
ICU 10	Mixed	15	16

* ICU with filtered taps.

Table II. Data of surveillance, typing of isolates and chronological analysis

	ICU 1*	ICU 2	ICU 3	ICU 4	ICU 5	ICU 6	ICU 7	ICU 8	ICU 9	ICU 10	Total
Surveillance											
Patients included	203	135	152	138	295	190	200	123	204	168	1808
Patients colonised or infected	51	29	59	45	48	37	51	18	43	46	427
Patients colonised or infected at the beginning of the study	6	4	5	7	2	1	1	0	5	10	41
Incidence of patients colonised or infected (<i>patient per 1000 days of hospitalisation</i>)	14.8	14.6	13.3	9.6	15.9	11.5	9.4	9.5	11.2	15.5	12.7
Taps water screened	45	15	28	31	25	15	29	10	15	20	233
Taps water positives	3	2	10	5	22	11	9	1	0	18	81
Taps water positives at the beginning of the study	5	3	9	5	19	0	2	0	0	8	51
Typing											
Patients isolates	69	90	223	155	166	72	227	82	100	331	1515
Water taps isolates	11	13	69	34	112	18	20	0	0	98	375
Patients with at least one isolate typed	34	26	54	44	42	27	49	18	38	41	373
Pulsotypes within patients	31	21	37	34	16	27	45	10	28	21	270
Sporadic pulsotypes	22	17	32	22	11	22	37	4	21	13	201
Shared by patients	8	2	4	9	0	4	8	6	7	4	52
Shared by water and patient	1	2	1	3	5	1	0	0	0	4	17
Pulsotypes in water taps	3	4	7	5	6	4	3	0	0	11	43
Sporadic pulsotypes	2	2	6	2	1	3	3	0	0	7	26
Chronological analysis											
Exogenous origin	17	10	23	23	36	6	8	8	10	29	170
Patient-to-patient transmission	14	6	4	20	1	5	8	8	10	10	86
Exogenous origin from water tap	3	3	19	3	/	1	0	0	0	/	29
Origin not concluded	/	1	/	/	35	/	/	/	/	19	55

* ICU with filtered taps.